

102(e)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/57190 A2(51) International Patent Classification⁷: C12N

(21) International Application Number: PCT/US01/04098

(22) International Filing Date: 5 February 2001 (05.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

09/496,914	3 February 2000 (03.02.2000)	US
09/560,875	27 April 2000 (27.04.2000)	US
09/598,075	20 June 2000 (20.06.2000)	US
09/620,325	19 July 2000 (19.07.2000)	US
09/654,936	1 September 2000 (01.09.2000)	US
09/663,561	15 September 2000 (15.09.2000)	US
09/693,325	20 October 2000 (20.10.2000)	US
09/728,422	30 November 2000 (30.11.2000)	US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US	09/496,914 (CIP)
Filed on	3 February 2000 (03.02.2000)
US	09/560,875 (CIP)
Filed on	27 April 2000 (27.04.2000)
US	09/598,075 (CIP)
Filed on	20 June 2000 (20.06.2000)
US	09/620,325 (CIP)
Filed on	19 July 2000 (19.07.2000)
US	09/654,936 (CIP)
Filed on	1 September 2000 (01.09.2000)
US	09/663,561 (CIP)
Filed on	15 September 2000 (15.09.2000)
US	09/693,325 (CIP)
Filed on	20 October 2000 (20.10.2000)
US	09/728,422 (CIP)
Filed on	30 November 2000 (30.11.2000)

(71) Applicant (for all designated States except US): HYSEQ, INC. [US/US]; 670 Almanor Avenue, Sunnyvale, CA 94086 (US).

WO 01/57190 A2

(72) Inventors; and

(75) Inventors/Applicants (for US only): TANG, Y., Tom [US/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). LIU, Chenghua [CN/US]; 1125 Ranchero Way, Apt. #14, San Jose, CA 95117 (US). DRMANAC, Radoje, T. [YU/US]; 850 Greenwich Place, Palo Alto, CA 94303 (US). ASUNDI, Vinod [US/US]; 709 Foster City

Boulevard, Foster City, CA 94404 (US). ZHOU, Ping [CN/US]; 7595 Newcastle Drive, Cupertino, CA 95014 (US). XU, Chongjun [US/US]; 4918 Manitoba Drive, San Jose, CA 95130 (US). CAO, Yicheng [CN/US]; 260 N. Mathilda Avenue, #E6, Sunnyvale, CA 94086 (US). MA, Yunqing [CN/US]; 280 W. California Avenue, #206, Sunnyvale, CA 94086 (US). ZHAO, Qing, A. [CN/US]; 1556 Kooser Road, San Jose, CA 95118 (US). WANG, Dunrui [CN/US]; 932 La Palma Place, Milpitas, CA 95035 (US). WANG, Jian-Rui [CN/US]; 744 Stendahl Lane, Cupertino, CA 95014 (US). ZHANG, Jie [CN/US]; 4930 Poplar Terrace, Campbell, CA 95008 (US). REN, Feiyan [US/US]; 7703 Oak Meadow Court, Cupertino, CA 95014 (US). CHEN, Rui-hong [US/US]; 1031 Flying Fish Street, Foster City, CA 94404 (US). WANG, Zhi, Wei [CN/US]; 836 Alturas Avenue, #B6, Sunnyvale, CA 94085 (US). XUE, Aidong, J. [CN/US]; 1621 S. Mary Avenue, Sunnyvale, CA 94087 (US). YANG, Yonghong [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). WEJHRMAN, Tom [US/US]; CCSR Mol Pharm 3210, 269 W. Campus Drive, Stanford, CA 94305 (US). GOODRICH, Ryle [US/US]; 4896 Sandy Lane, San Jose, CA 95124 (US).

(74) Agent: ELRIFI, Ivor, R.; Mintz, Levin, Cohn, Ferris, Glovsky, and Popeo, P.C., One Financial Center, Boston, MA 02111 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CL, CN, CZ, DE, DK, DM, DZ, EE, EG, ES, FI, GB, GD, GE, GR, GM, GU, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL NUCLEIC ACIDS AND POLYPEPTIDES

(57) Abstract: The present invention provides novel nucleic acids, novel polypeptide sequences encoded by these nucleic acids and uses thereof.

NOVEL NUCLEIC ACIDS AND POLYPEPTIDES

1. TECHNICAL FIELD

The present invention provides novel polynucleotides and proteins encoded by such
5 polynucleotides, along with uses for these polynucleotides and proteins, for example in
therapeutic, diagnostic and research methods.

2. BACKGROUND

Technology aimed at the discovery of protein factors (including e.g., cytokines, such as
10 lymphokines, interferons, CSFs, chemokines, and interleukins) has matured rapidly over the past
decade. The now routine hybridization cloning and expression cloning techniques clone novel
polynucleotides "directly" in the sense that they rely on information directly related to the
discovered protein (i.e., partial DNA/amino acid sequence of the protein in the case of
hybridization cloning; activity of the protein in the case of expression cloning). More recent
15 "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences
based on the presence of a now well-recognized secretory leader sequence motif, as well as
various PCR-based or low stringency hybridization-based cloning techniques, have advanced the
state of the art by making available large numbers of DNA/amino acid sequences for proteins
that are known to have biological activity, for example, by virtue of their secreted nature in the
20 case of leader sequence cloning, by virtue of their cell or tissue source in the case of PCR-based
techniques, or by virtue of structural similarity to other genes of known biological activity.

Identified polynucleotide and polypeptide sequences have numerous applications in, for
example, diagnostics, forensics, gene mapping; identification of mutations responsible for
genetic disorders or other traits, to assess biodiversity, and to produce many other types of data
25 and products dependent on DNA and amino acid sequences.

3. SUMMARY OF THE INVENTION

The compositions of the present invention include novel isolated polypeptides, novel
isolated polynucleotides encoding such polypeptides, including recombinant DNA molecules,
30 cloned genes or degenerate variants thereof, especially naturally occurring variants such as allelic
variants, antisense polynucleotide molecules, and antibodies that specifically recognize one or more
epitopes present on such polypeptides, as well as hybridomas producing such antibodies.

The compositions of the present invention additionally include vectors, including expression
vectors, containing the polynucleotides of the invention, cells genetically engineered to contain such
35 polynucleotides and cells genetically engineered to express such polynucleotides.

The present invention relates to a collection or library of at least one novel nucleic acid sequence assembled from expressed sequence tags (ESTs) isolated mainly by sequencing by hybridization (SBH), and in some cases, sequences obtained from one or more public databases. The invention relates also to the proteins encoded by such polynucleotides, along with therapeutic, diagnostic and research utilities for these polynucleotides and proteins. These nucleic acid sequences are designated as SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954. The polypeptides sequences are designated SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960. The nucleic acids and polypeptides are provided in the Sequence Listing. In the nucleic acids provided in the Sequence Listing, A is adenosine; C is cytosine; G is guanine; T is thymine; and N is any of the four bases. In the amino acids provided in the Sequence Listing, * corresponds to the stop codon.

The nucleic acid sequences of the present invention also include, nucleic acid sequences that hybridize to the complement of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 under stringent hybridization conditions; nucleic acid sequences which are allelic variants or species homologues of any of the nucleic acid sequences recited above, or nucleic acid sequences that encode a peptide comprising a specific domain or truncation of the peptides encoded by SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954. A polynucleotide comprising a nucleotide sequence having at least 90% identity to an identifying sequence of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 or a degenerate variant or fragment thereof. The identifying sequence can be 100 base pairs in length.

The nucleic acid sequences of the present invention also include the sequence information from the nucleic acid sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954. The sequence information can be a segment of any one of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 that uniquely identifies or represents the sequence information of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954.

A collection as used in this application can be a collection of only one polynucleotide. The collection of sequence information or identifying information of each sequence can be provided on a nucleic acid array. In one embodiment, segments of sequence information is provided on a nucleic acid array to detect the polynucleotide that contains the segment. The array can be designed to detect full-match or mismatch to the polynucleotide that contains the segment. The collection can also be provided in a computer-readable format.

This invention also includes the reverse or direct complement of any of the nucleic acid sequences recited above; cloning or expression vectors containing the nucleic acid sequences; and host cells or organisms transformed with these expression vectors. Nucleic acid sequences (or their reverse or direct complements) according to the invention have numerous applications in a variety

of techniques known to those skilled in the art of molecular biology, such as use as hybridization probes, use as primers for PCR, use in an array, use in computer-readable media, use in sequencing full-length genes, use for chromosome and gene mapping, use in the recombinant production of protein, and use in the generation of anti-sense DNA or RNA, their chemical analogs and the like.

5 In a preferred embodiment, the nucleic acid sequences of SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954 or novel segments or parts of the nucleic acids of the invention are used as primers in expression assays that are well known in the art. In a particularly preferred embodiment, the nucleic acid sequences of SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954 or novel segments or parts of the nucleic acids provided herein are used in diagnostics for identifying
10 expressed genes or, as well known in the art and exemplified by Vollrath et al., Science 258:52-59 (1992), as expressed sequence tags for physical mapping of the human genome.

The isolated polynucleotides of the invention include, but are not limited to, a polynucleotide comprising any one of the nucleotide sequences set forth in SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954; a polynucleotide comprising any of the full length protein
15 coding sequences of SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954; and a polynucleotide comprising any of the nucleotide sequences of the mature protein coding sequences of SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent hybridization conditions to (a) the complement of any one of the nucleotide sequences set forth in SEQ ID NO:1-
20 984, 1969-2952, 3937-3942 or 3949-3954; (b) a nucleotide sequence encoding any one of the amino acid sequences set forth in the Sequence Listing; (c) a polynucleotide which is an allelic variant of any polynucleotides recited above, (d) a polynucleotide which encodes a species homolog (e.g. orthologs) of any of the proteins recited above; or (e) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of any of the polypeptides comprising an
25 amino acid sequence set forth in the Sequence Listing.

The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising any of the amino acid sequences set forth in SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960; or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides with biological activity that are encoded by (a) any of the
30 polynucleotides having a nucleotide sequence set forth in SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954; or (b) polynucleotides that hybridize to the complement of the polynucleotides of (a) under stringent hybridization conditions. Biologically or immunologically active variants of any of the polypeptide sequences in the Sequence Listing, and "substantial equivalents" thereof (e.g., with at least about 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% amino acid sequence
35 identity) that preferably retain biological activity are also contemplated. The polypeptides of the

invention may be wholly or partially chemically synthesized but are preferably produced by recombinant means using the genetically engineered cells (e.g. host cells) of the invention.

The invention also provides compositions comprising a polypeptide of the invention. Polypeptide compositions of the invention may further comprise an acceptable carrier, such as a hydrophilic, e.g., pharmaceutically acceptable, carrier.

The invention also provides host cells transformed or transfected with a polynucleotide of the invention.

The invention also relates to methods for producing a polypeptide of the invention comprising growing a culture of the host cells of the invention in a suitable culture medium under conditions permitting expression of the desired polypeptide, and purifying the polypeptide from the culture or from the host cells. Preferred embodiments include those in which the protein produced by such process is a mature form of the protein.

Polynucleotides according to the invention have numerous applications in a variety of techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use as oligomers, or primers, for PCR, use for chromosome and gene mapping, use in the recombinant production of protein, and use in generation of anti-sense DNA or RNA, their chemical analogs and the like. For example, when the expression of an mRNA is largely restricted to a particular cell or tissue type, polynucleotides of the invention can be used as hybridization probes to detect the presence of the particular cell or tissue mRNA in a sample using, e.g., *in situ* hybridization.

In other exemplary embodiments, the polynucleotides are used in diagnostics as expressed sequence tags for identifying expressed genes or, as well known in the art and exemplified by Vollrath et al., Science 258:52-59 (1992), as expressed sequence tags for physical mapping of the human genome.

The polypeptides according to the invention can be used in a variety of conventional procedures and methods that are currently applied to other proteins. For example, a polypeptide of the invention can be used to generate an antibody that specifically binds the polypeptide. Such antibodies, particularly monoclonal antibodies, are useful for detecting or quantitating the polypeptide in tissue. The polypeptides of the invention can also be used as molecular weight markers, and as a food supplement.

Methods are also provided for preventing, treating, or ameliorating a medical condition which comprises the step of administering to a mammalian subject a therapeutically effective amount of a composition comprising a polypeptide of the present invention and a pharmaceutically acceptable carrier.

In particular, the polypeptides and polynucleotides of the invention can be utilized, for example, in methods for the prevention and/or treatment of disorders involving aberrant protein expression or biological activity.

The present invention further relates to methods for detecting the presence of the
5 polynucleotides or polypeptides of the invention in a sample. Such methods can, for example, be utilized as part of prognostic and diagnostic evaluation of disorders as recited herein and for the identification of subjects exhibiting a predisposition to such conditions. The invention provides a method for detecting the polynucleotides of the invention in a sample, comprising contacting the sample with a compound that binds to and forms a complex with the polynucleotide of
10 interest for a period sufficient to form the complex and under conditions sufficient to form a complex and detecting the complex such that if a complex is detected, the polynucleotide of interest is detected. The invention also provides a method for detecting the polypeptides of the invention in a sample comprising contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex
15 and detecting the formation of the complex such that if a complex is formed, the polypeptide is detected.

The invention also provides kits comprising polynucleotide probes and/or monoclonal antibodies, and optionally quantitative standards, for carrying out methods of the invention. Furthermore, the invention provides methods for evaluating the efficacy of drugs, and
20 monitoring the progress of patients, involved in clinical trials for the treatment of disorders as recited above.

The invention also provides methods for the identification of compounds that modulate (i.e., increase or decrease) the expression or activity of the polynucleotides and/or polypeptides of the invention. Such methods can be utilized, for example, for the identification of compounds
25 that can ameliorate symptoms of disorders as recited herein. Such methods can include, but are not limited to, assays for identifying compounds and other substances that interact with (*e.g.*, bind to) the polypeptides of the invention. The invention provides a method for identifying a compound that binds to the polypeptides of the invention comprising contacting the compound with a polypeptide of the invention in a cell for a time sufficient to form a polypeptide/compound
30 complex, wherein the complex drives expression of a reporter gene sequence in the cell; and detecting the complex by detecting the reporter gene sequence expression such that if expression of the reporter gene is detected the compound the binds to a polypeptide of the invention is identified.

The methods of the invention also provides methods for treatment which involve the
35 administration of the polynucleotides or polypeptides of the invention to individuals exhibiting

symptoms or tendencies. In addition, the invention encompasses methods for treating diseases or disorders as recited herein comprising administering compounds and other substances that modulate the overall activity of the target gene products. Compounds and other substances can effect such modulation either on the level of target gene/protein expression or target protein activity.

The polypeptides of the present invention and the polynucleotides encoding them are also useful for the same functions known to one of skill in the art as the polypeptides and polynucleotides to which they have homology (set forth in Tables 2 and 9); for which they have a signature region (as set forth in Tables 3 and 10); or for which they have homology to a gene family (as set forth in Tables 4 and 11). If no homology is set forth for a sequence, then the polypeptides and polynucleotides of the present invention are useful for a variety of applications, as described herein, including use in arrays for detection.

4. DETAILED DESCRIPTION OF THE INVENTION

4.1 DEFINITIONS

It must be noted that as used herein and in the appended claims, the singular forms "a", "an" and "the" include plural references unless the context clearly dictates otherwise.

The term "active" refers to those forms of the polypeptide which retain the biologic and/or immunologic activities of any naturally occurring polypeptide. According to the invention, the terms "biologically active" or "biological activity" refer to a protein or peptide having structural, regulatory or biochemical functions of a naturally occurring molecule. Likewise "immunologically active" or "immunological activity" refers to the capability of the natural, recombinant or synthetic polypeptide to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "activated cells" as used in this application are those cells which are engaged in extracellular or intracellular membrane trafficking, including the export of secretory or enzymatic molecules as part of a normal or disease process.

The terms "complementary" or "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence 5'-AGT-3' binds to the complementary sequence 3'-TCA-5'. Complementarity between two single-stranded molecules may be "partial" such that only some of the nucleic acids bind or it may be "complete" such that total complementarity exists between the single stranded molecules. The degree of complementarity between the nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands.

The term "embryonic stem cells (ES)" refers to a cell that can give rise to many differentiated cell types in an embryo or an adult, including the germ cells. The term "germ line stem cells (GSCs)" refers to stem cells derived from primordial stem cells that provide a steady and continuous source of germ cells for the production of gametes. The term "primordial germ cells (PGCs)" refers to a small population of cells set aside from other cell lineages particularly from the yolk sac, mesenteries, or gonadal ridges during embryogenesis that have the potential to differentiate into germ cells and other cells. PGCs are the source from which GSCs and ES cells are derived. The PGCs, the GSCs and the ES cells are capable of self-renewal. Thus these cells not only populate the germ line and give rise to a plurality of terminally differentiated cells that comprise the adult specialized organs, but are able to regenerate themselves.

The term "expression modulating fragment," EMF, means a series of nucleotides which modulates the expression of an operably linked ORF or another EMF.

As used herein, a sequence is said to "modulate the expression of an operably linked sequence" when the expression of the sequence is altered by the presence of the EMF. EMFs include, but are not limited to, promoters, and promoter modulating sequences (inducible elements). One class of EMFs are nucleic acid fragments which induce the expression of an operably linked ORF in response to a specific regulatory factor or physiological event.

The terms "nucleotide sequence" or "nucleic acid" or "polynucleotide" or "oligonucleotide" are used interchangeably and refer to a heteropolymer of nucleotides or the sequence of these nucleotides. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA) or to any DNA-like or RNA-like material. In the sequences herein A is adenine, C is cytosine, T is thymine, G is guanine and N is A, C, G or T (U). It is contemplated that where the polynucleotide is RNA, the T (thymine) in the sequences provided herein is substituted with U (uracil). Generally, nucleic acid segments provided by this invention may be assembled from fragments of the genome and short oligonucleotide linkers, or from a series of oligonucleotides, or from individual nucleotides, to provide a synthetic nucleic acid which is capable of being expressed in a recombinant transcriptional unit comprising regulatory elements derived from a microbial or viral operon, or a eukaryotic gene.

The terms "oligonucleotide fragment" or a "polynucleotide fragment", "portion," or "segment" or "probe" or "primer" are used interchangeably and refer to a sequence of nucleotide residues which are at least about 5 nucleotides, more preferably at least about 7 nucleotides, more preferably at least about 9 nucleotides, more preferably at least about 11 nucleotides and most preferably at least about 17 nucleotides. The fragment is preferably less than about 500 nucleotides, preferably less than about 200 nucleotides, more preferably less than about 100

nucleotides, more preferably less than about 50 nucleotides and most preferably less than 30 nucleotides. Preferably the probe is from about 6 nucleotides to about 200 nucleotides, preferably from about 15 to about 50 nucleotides, more preferably from about 17 to 30 nucleotides and most preferably from about 20 to 25 nucleotides. Preferably the fragments can be used in polymerase chain reaction (PCR), various hybridization procedures or microarray procedures to identify or amplify identical or related parts of mRNA or DNA molecules. A fragment or segment may uniquely identify each polynucleotide sequence of the present invention. Preferably the fragment comprises a sequence substantially similar to any one of SEQ ID NOs:1-20.

Probes may, for example, be used to determine whether specific mRNA molecules are present in a cell or tissue or to isolate similar nucleic acid sequences from chromosomal DNA as described by Walsh et al. (Walsh, P.S. et al., 1992, PCR Methods Appl 1:241-250). They may be labeled by nick translation, Klenow fill-in reaction, PCR, or other methods well known in the art. Probes of the present invention, their preparation and/or labeling are elaborated in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, NY; or Ausubel, F.M. et al., 1989, Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, both of which are incorporated herein by reference in their entirety.

The nucleic acid sequences of the present invention also include the sequence information from the nucleic acid sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954. The sequence information can be a segment of any one of SEQ ID NO:1-1-984, 1969-2952, 3937-3942 or 3949-3954 that uniquely identifies or represents the sequence information of that sequence of SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954. One such segment can be a twenty-mer nucleic acid sequence because the probability that a twenty-mer is fully matched in the human genome is 1 in 300. In the human genome, there are three billion base pairs in one set of chromosomes. Because 4^{20} possible twenty-mers exist, there are 300 times more twenty-mers than there are base pairs in a set of human chromosomes. Using the same analysis, the probability for a seventeen-mer to be fully matched in the human genome is approximately 1 in 5. When these segments are used in arrays for expression studies, fifteen-mer segments can be used. The probability that the fifteen-mer is fully matched in the expressed sequences is also approximately one in five because expressed sequences comprise less than approximately 5% of the entire genome sequence.

Similarly, when using sequence information for detecting a single mismatch, a segment can be a twenty-five mer. The probability that the twenty-five mer would appear in a human genome with a single mismatch is calculated by multiplying the probability for a full match ($1+4^{25}$) times the

increased probability for mismatch at each nucleotide position (3 x 25). The probability that an eighteen mer with a single mismatch can be detected in an array for expression studies is approximately one in five. The probability that a twenty-mer with a single mismatch can be detected in a human genome is approximately one in five.

5 The term "open reading frame," ORF, means a series of nucleotide triplets coding for amino acids without any termination codons and is a sequence translatable into protein.

 The terms "operably linked" or "operably associated" refer to functionally related nucleic acid sequences. For example, a promoter is operably associated or operably linked with a coding sequence if the promoter controls the transcription of the coding sequence. While operably
10 linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements e.g. repressor genes are not contiguously linked to the coding sequence but still control transcription/translation of the coding sequence.

 The term "pluripotent" refers to the capability of a cell to differentiate into a number of differentiated cell types that are present in an adult organism. A pluripotent cell is restricted in its
15 differentiation capability in comparison to a totipotent cell.

 The terms "polypeptide" or "peptide" or "amino acid sequence" refer to an oligopeptide, peptide, polypeptide or protein sequence or fragment thereof and to naturally occurring or synthetic molecules. A polypeptide "fragment," "portion," or "segment" is a stretch of amino acid residues of at least about 5 amino acids, preferably at least about 7 amino acids, more
20 preferably at least about 9 amino acids and most preferably at least about 17 or more amino acids. The peptide preferably is not greater than about 500 amino acids, more preferably less than 200 amino acids more preferably less than 150 amino acids and most preferably less than 100 amino acids. Preferably the peptide is from about 5 to about 200 amino acids. To be active, any polypeptide must have sufficient length to display biological and/or immunological activity.

25 The term "naturally occurring polypeptide" refers to polypeptides produced by cells that have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including, but not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

 The term "translated protein coding portion" means a sequence which encodes for the full
30 length protein which may include any leader sequence or any processing sequence.

 The term "mature protein coding sequence" means a sequence which encodes a peptide or protein without a signal or leader sequence. The "mature protein portion" means that portion of the protein which does not include a signal or leader sequence. The peptide may have been produced by processing in the cell which removes any leader/signal sequence. The mature
35 protein portion may or may not include the initial methionine residue. The methionine residue

may be removed from the protein during processing in the cell. The peptide may be produced synthetically or the protein may have been produced using a polynucleotide only encoding for the mature protein coding sequence.

The term "derivative" refers to polypeptides chemically modified by such techniques as ubiquitination, labeling (e.g., with radionuclides or various enzymes), covalent polymer attachment such as pegylation (derivatization with polyethylene glycol) and insertion or substitution by chemical synthesis of amino acids such as ornithine, which do not normally occur in human proteins.

The term "variant" (or "analog") refers to any polypeptide differing from naturally occurring polypeptides by amino acid insertions, deletions, and substitutions, created using, e.g., recombinant DNA techniques. Guidance in determining which amino acid residues may be replaced, added or deleted without abolishing activities of interest, may be found by comparing the sequence of the particular polypeptide with that of homologous peptides and minimizing the number of amino acid sequence changes made in regions of high homology (conserved regions) or by replacing amino acids with consensus sequence.

Alternatively, recombinant variants encoding these same or similar polypeptides may be synthesized or selected by making use of the "redundancy" in the genetic code. Various codon substitutions, such as the silent changes which produce various restriction sites, may be introduced to optimize cloning into a plasmid or viral vector or expression in a particular prokaryotic or eukaryotic system. Mutations in the polynucleotide sequence may be reflected in the polypeptide or domains of other peptides added to the polypeptide to modify the properties of any part of the polypeptide, to change characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate.

Preferably, amino acid "substitutions" are the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, i.e., conservative amino acid replacements. "Conservative" amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. "Insertions" or "deletions" are preferably in the range of about 1 to 20 amino acids, more preferably 1 to 10 amino acids. The variation allowed may be experimentally determined by systematically making

insertions, deletions, or substitutions of amino acids in a polypeptide molecule using recombinant DNA techniques and assaying the resulting recombinant variants for activity.

Alternatively, where alteration of function is desired, insertions, deletions or non-conservative alterations can be engineered to produce altered polypeptides. Such alterations can, for example, alter one or more of the biological functions or biochemical characteristics of the polypeptides of the invention. For example, such alterations may change polypeptide characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate. Further, such alterations can be selected so as to generate polypeptides that are better suited for expression, scale up and the like in the host cells chosen for expression. For example, cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

The terms "purified" or "substantially purified" as used herein denotes that the indicated nucleic acid or polypeptide is present in the substantial absence of other biological macromolecules, *e.g.*, polynucleotides, proteins, and the like. In one embodiment, the polynucleotide or polypeptide is purified such that it constitutes at least 95% by weight, more preferably at least 99% by weight, of the indicated biological macromolecules present (but water, buffers, and other small molecules, especially molecules having a molecular weight of less than 1000 daltons, can be present).

The term "isolated" as used herein refers to a nucleic acid or polypeptide separated from at least one other component (*e.g.*, nucleic acid or polypeptide) present with the nucleic acid or polypeptide in its natural source. In one embodiment, the nucleic acid or polypeptide is found in the presence of (if anything) only a solvent, buffer, ion, or other component normally present in solution of the same. The terms "isolated" and "purified" do not encompass nucleic acids or polypeptides present in their natural source.

The term "recombinant," when used herein to refer to a polypeptide or protein, means that a polypeptide or protein is derived from recombinant (*e.g.*, microbial, insect, or mammalian) expression systems. "Microbial" refers to recombinant polypeptides or proteins made in bacterial or fungal (*e.g.*, yeast) expression systems. As a product, "recombinant microbial" defines a polypeptide or protein essentially free of native endogenous substances and unaccompanied by associated native glycosylation. Polypeptides or proteins expressed in most bacterial cultures, *e.g.*, *E. coli*, will be free of glycosylation modifications; polypeptides or proteins expressed in yeast will have a glycosylation pattern in general different from those expressed in mammalian cells.

The term "recombinant expression vehicle or vector" refers to a plasmid or phage or virus or vector, for expressing a polypeptide from a DNA (RNA) sequence. An expression vehicle can

comprise a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an amino terminal methionine residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

The term "recombinant expression system" means host cells which have stably integrated a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit extrachromosomally. Recombinant expression systems as defined herein will express heterologous polypeptides or proteins upon induction of the regulatory elements linked to the DNA segment or synthetic gene to be expressed. This term also means host cells which have stably integrated a recombinant genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers. Recombinant expression systems as defined herein will express polypeptides or proteins endogenous to the cell upon induction of the regulatory elements linked to the endogenous DNA segment or gene to be expressed. The cells can be prokaryotic or eukaryotic.

The term "secreted" includes a protein that is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence when it is expressed in a suitable host cell. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins that are transported across the membrane of the endoplasmic reticulum. "Secreted" proteins are also intended to include proteins containing non-typical signal sequences (e.g. Interleukin-1 Beta, see Krasney, P.A. and Young, P.R. (1992) Cytokine 4(2):134 -143) and factors released from damaged cells (e.g. Interleukin-1 Receptor Antagonist, see Arend, W.P. et. al. (1998) Annu. Rev. Immunol. 16:27-55)

Where desired, an expression vector may be designed to contain a "signal or leader sequence" which will direct the polypeptide through the membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous protein sources by recombinant DNA techniques.

The term "stringent" is used to refer to conditions that are commonly understood in the art as stringent. Stringent conditions can include highly stringent conditions (i.e., hybridization

to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1X SSC/0.1% SDS at 68°C), and moderately stringent conditions (i.e., washing in 0.2X SSC/0.1% SDS at 42°C). Other exemplary hybridization conditions are described herein in the examples.

- 5 In instances of hybridization of deoxyoligonucleotides, additional exemplary stringent hybridization conditions include washing in 6X SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligonucleotides), 48°C (for 17-base oligos), 55°C (for 20-base oligonucleotides), and 60°C (for 23-base oligonucleotides).

- As used herein, "substantially equivalent" can refer both to nucleotide and amino acid
10 sequences, for example a mutant sequence, that varies from a reference sequence by one or more substitutions, deletions, or additions, the net effect of which does not result in an adverse functional dissimilarity between the reference and subject sequences. Typically, such a substantially equivalent sequence varies from one of those listed herein by no more than about 35% (i.e., the number of individual residue substitutions, additions, and/or deletions in a
15 substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of residues in the substantially equivalent sequence is about 0.35 or less). Such a sequence is said to have 65% sequence identity to the listed sequence. In one embodiment, a substantially equivalent, e.g., mutant, sequence of the invention varies from a listed sequence by no more than 30% (70% sequence identity); in a variation of this embodiment,
20 by no more than 25% (75% sequence identity); and in a further variation of this embodiment, by no more than 20% (80% sequence identity) and in a further variation of this embodiment, by no more than 10% (90% sequence identity) and in a further variation of this embodiment, by no more than 5% (95% sequence identity). Substantially equivalent, e.g., mutant, amino acid sequences according to the invention preferably have at least 80% sequence identity with a listed
25 amino acid sequence, more preferably at least 85% sequence identity, more preferably at least 90% sequence identity, more preferably at least 95% sequence identity, more preferably at least 98% sequence identity and most preferably at least 98% identity. Substantially equivalent nucleotide sequences of the invention can have lower percent sequence identities, taking into account, for example, the redundancy or degeneracy of the genetic code. Preferably, nucleotide
30 sequence has at least about 65% identity, more preferably at least about 75% identity, more preferably at least about 80% identity, more preferably at least about 85% identity, more preferably at least about 90% identity, and most preferably at least about 95% identity, more preferably at least 98% and most preferably at least about 99% identity. For the purposes of the present invention, sequences having substantially equivalent biological activity and substantially
35 equivalent expression characteristics are considered substantially equivalent. For the purposes of

determining equivalence, truncation of the mature sequence (e.g., via a mutation which creates a spurious stop codon) should be disregarded. Sequence identity may be determined, e.g., using the Jotun Hein method (Hein, J. (1990) *Methods Enzymol.* 183:626-645). Identity between sequences can also be determined by other methods known in the art, e.g. by varying

5 hybridization conditions.

The term "totipotent" refers to the capability of a cell to differentiate into all of the cell types of an adult organism.

The term "transformation" means introducing DNA into a suitable host cell so that the DNA is replicable, either as an extrachromosomal element, or by chromosomal integration. The
10 term "transfection" refers to the taking up of an expression vector by a suitable host cell, whether or not any coding sequences are in fact expressed. The term "infection" refers to the introduction of nucleic acids into a suitable host cell by use of a virus or viral vector.

As used herein, an "uptake modulating fragment," UMF, means a series of nucleotides which mediate the uptake of a linked DNA fragment into a cell. UMFs can be readily identified
15 using known UMFs as a target sequence or target motif with the computer-based systems described below. The presence and activity of a UMF can be confirmed by attaching the suspected UMF to a marker sequence. The resulting nucleic acid molecule is then incubated with an appropriate host under appropriate conditions and the uptake of the marker sequence is determined. As described above, a UMF will increase the frequency of uptake of a linked
20 marker sequence.

Each of the above terms is meant to encompass all that is described for each, unless the context dictates otherwise.

4.2 NUCLEIC ACIDS OF THE INVENTION

25 Nucleotide sequences of the invention are set forth in the Sequence Listing.

The isolated polynucleotides of the invention include a polynucleotide comprising the nucleotide sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954; a polynucleotide encoding any one of the peptide sequences of SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960; and a polynucleotide comprising the nucleotide sequence encoding the
30 mature protein coding sequence of the polypeptides of any one of SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent conditions to (a) the complement of any of the nucleotides sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954; (b) nucleotide sequences encoding any one of the amino acid sequences set forth
35 in the Sequence Listing as SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960; (c) a

polynucleotide which is an allelic variant of any polynucleotide recited above; (d) a polynucleotide which encodes a species homolog of any of the proteins recited above; or (e) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptides of SEQ ID NO:985-1968, 2953-3936, 3943-3948 or 3955-3960. Domains of interest may depend on the nature of the encoded polypeptide; e.g., domains in receptor-like polypeptides include ligand-binding, extracellular, transmembrane, or cytoplasmic domains, or combinations thereof; domains in immunoglobulin-like proteins include the variable immunoglobulin-like domains; domains in enzyme-like polypeptides include catalytic and substrate binding domains; and domains in ligand polypeptides include receptor-binding domains.

The polynucleotides of the invention include naturally occurring or wholly or partially synthetic DNA, e.g., cDNA and genomic DNA, and RNA, e.g., mRNA. The polynucleotides may include all of the coding region of the cDNA or may represent a portion of the coding region of the cDNA.

The present invention also provides genes corresponding to the cDNA sequences disclosed herein. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. Further 5' and 3' sequence can be obtained using methods known in the art. For example, full length cDNA or genomic DNA that corresponds to any of the polynucleotides of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 can be obtained by screening appropriate cDNA or genomic DNA libraries under suitable hybridization conditions using any of the polynucleotides of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 or a portion thereof as a probe. Alternatively, the polynucleotides of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 may be used as the basis for suitable primer(s) that allow identification and/or amplification of genes in appropriate genomic DNA or cDNA libraries.

The nucleic acid sequences of the invention can be assembled from ESTs and sequences (including cDNA and genomic sequences) obtained from one or more public databases, such as dbEST, gbpri, and UniGene. The EST sequences can provide identifying sequence information, representative fragment or segment information, or novel segment information for the full-length gene.

The polynucleotides of the invention also provide polynucleotides including nucleotide sequences that are substantially equivalent to the polynucleotides recited above. Polynucleotides according to the invention can have, e.g., at least about 65%, at least about 70%, at least about

75%, at least about 80%, 81%, 82%, 83%, 84%, more typically at least about 85%, 86%, 87%, 88%, 89%, and more typically at least about 90%, 91%, 92%, 93%, 94%, and even more typically at least about 95%, 96%, 97%, 98%, 99%, sequence identity to a polynucleotide recited above.

5 Included within the scope of the nucleic acid sequences of the invention are nucleic acid sequence fragments that hybridize under stringent conditions to any of the nucleotide sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, or complements thereof, which fragment is greater than about 5 nucleotides, preferably 7 nucleotides, more preferably greater than 9 nucleotides and most preferably greater than 17 nucleotides. Fragments of, e.g. 15, 17, or
10 20 nucleotides or more that are selective for (i.e. specifically hybridize to any one of the polynucleotides of the invention) are contemplated. Probes capable of specifically hybridizing to a polynucleotide can differentiate polynucleotide sequences of the invention from other polynucleotide sequences in the same family of genes or can differentiate human genes from genes of other species, and are preferably based on unique nucleotide sequences.

15 The sequences falling within the scope of the present invention are not limited to these specific sequences, but also include allelic and species variations thereof. Allelic and species variations can be routinely determined by comparing the sequence provided SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, a representative fragment thereof, or a nucleotide sequence at least 90% identical, preferably 95% identical, to SEQ ID NO: 1-984, 1969-2952, 3937-3942 or
20 3949-3954 with a sequence from another isolate of the same species. Furthermore, to accommodate codon variability, the invention includes nucleic acid molecules coding for the same amino acid sequences as do the specific ORFs disclosed herein. In other words, in the coding region of an ORF, substitution of one codon for another codon that encodes the same amino acid is expressly contemplated.

25 The nearest neighbor or homology result for the nucleic acids of the present invention, including SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, can be obtained by searching a database using an algorithm or a program. Preferably, a BLAST which stands for Basic Local Alignment Search Tool is used to search for local sequence alignments (Altschul, S.F. J Mol. Evol. 36 290-300 (1993) and Altschul S.F. et al. J. Mol. Biol. 21:403-410 (1990)). Alternatively a
30 FASTA version 3 search against Genpept, using Fastxy algorithm.

Species homologs (or orthologs) of the disclosed polynucleotides and proteins are also provided by the present invention. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species.

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotide which also encode proteins which are identical, homologous or related to that encoded by the polynucleotides.

- 5 The nucleic acid sequences of the invention are further directed to sequences which encode variants of the described nucleic acids. These amino acid sequence variants may be prepared by methods known in the art by introducing appropriate nucleotide changes into a native or variant polynucleotide. There are two variables in the construction of amino acid sequence variants: the location of the mutation and the nature of the mutation. Nucleic acids
10 encoding the amino acid sequence variants are preferably constructed by mutating the polynucleotide to encode an amino acid sequence that does not occur in nature. These nucleic acid alterations can be made at sites that differ in the nucleic acids from different species (variable positions) or in highly conserved regions (constant regions). Sites at such locations will typically be modified in series, *e.g.*, by substituting first with conservative choices (*e.g.*,
15 hydrophobic amino acid to a different hydrophobic amino acid) and then with more distant choices (*e.g.*, hydrophobic amino acid to a charged amino acid), and then deletions or insertions may be made at the target site. Amino acid sequence deletions generally range from about 1 to 30 residues, preferably about 1 to 10 residues, and are typically contiguous. Amino acid insertions include amino- and/or carboxyl-terminal fusions ranging in length from one to one
20 hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions may range generally from about 1 to 10 amino residues, preferably from 1 to 5 residues. Examples of terminal insertions include the heterologous signal sequences necessary for secretion or for intracellular targeting in different host cells and sequences such as FLAG or poly-histidine sequences useful for purifying the expressed protein.
- 25 In a preferred method, polynucleotides encoding the novel amino acid sequences are changed via site-directed mutagenesis. This method uses oligonucleotide sequences to alter a polynucleotide to encode the desired amino acid variant, as well as sufficient adjacent nucleotides on both sides of the changed amino acid to form a stable duplex on either side of the site of being changed. In general, the techniques of site-directed mutagenesis are well known to
30 those of skill in the art and this technique is exemplified by publications such as, Edelman et al., *DNA* 2:183 (1983). A versatile and efficient method for producing site-specific changes in a polynucleotide sequence was published by Zoller and Smith, *Nucleic Acids Res.* 10:6487-6500 (1982). PCR may also be used to create amino acid sequence variants of the novel nucleic acids. When small amounts of template DNA are used as starting material, primer(s) that differs
35 slightly in sequence from the corresponding region in the template DNA can generate the desired

amino acid variant. PCR amplification results in a population of product DNA fragments that differ from the polynucleotide template encoding the polypeptide at the position specified by the primer. The product DNA fragments replace the corresponding region in the plasmid and this gives a polynucleotide encoding the desired amino acid variant.

5 A further technique for generating amino acid variants is the cassette mutagenesis technique described in Wells et al., *Gene* 34:315 (1985); and other mutagenesis techniques well known in the art, such as, for example, the techniques in Sambrook et al., *supra*, and *Current Protocols in Molecular Biology*, Ausubel et al. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent
10 amino acid sequence may be used in the practice of the invention for the cloning and expression of these novel nucleic acids. Such DNA sequences include those which are capable of hybridizing to the appropriate novel nucleic acid sequence under stringent conditions.

Polynucleotides encoding preferred polypeptide truncations of the invention can be used to generate polynucleotides encoding chimeric or fusion proteins comprising one or more
15 domains of the invention and heterologous protein sequences.

The polynucleotides of the invention additionally include the complement of any of the polynucleotides recited above. The polynucleotide can be DNA (genomic, cDNA, amplified, or synthetic) or RNA. Methods and algorithms for obtaining such polynucleotides are well known to those of skill in the art and can include, for example, methods for determining hybridization
20 conditions that can routinely isolate polynucleotides of the desired sequence identities.

In accordance with the invention, polynucleotide sequences comprising the mature protein coding sequences corresponding to any one of SEQ ID NO: 1-984, 1969-2957, 3937-3942 or 3949-3954, or functional equivalents thereof, may be used to generate recombinant DNA molecules that direct the expression of that nucleic acid, or a functional equivalent thereof,
25 in appropriate host cells. Also included are the cDNA inserts of any of the clones identified herein.

A polynucleotide according to the invention can be joined to any of a variety of other nucleotide sequences by well-established recombinant DNA techniques (see Sambrook J et al. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, NY). Useful
30 nucleotide sequences for joining to polynucleotides include an assortment of vectors, e.g., plasmids, cosmids, lambda phage derivatives, phagemids, and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide. In general, the vector contains an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a
35 selectable marker for the host cell. Vectors according to the invention include expression

vectors, replication vectors, probe generation vectors, and sequencing vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a multicellular organism.

The present invention further provides recombinant constructs comprising a nucleic acid
5 having any of the nucleotide sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 or a fragment thereof or any other polynucleotides of the invention. In one embodiment, the recombinant constructs of the present invention comprise a vector, such as a plasmid or viral vector, into which a nucleic acid having any of the nucleotide sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 or a fragment thereof is inserted, in a forward or reverse
10 orientation. In the case of a vector comprising one of the ORFs of the present invention, the vector may further comprise regulatory sequences, including for example, a promoter, operably linked to the ORF. Large numbers of suitable vectors and promoters are known to those of skill in the art and are commercially available for generating the recombinant constructs of the present invention. The following vectors are provided by way of example. Bacterial: pBs, phagescript, PsiX174, pBluescript SK, pBs KS, pNH8a, pNH16a, pNH18a, pNH46a (Stratagene); pTrc99A,
15 pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLneo, pSV2cat, pOG44, PX1, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia).

The isolated polynucleotide of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al.,
20 *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector
25 or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt,
30 lambda PR, and trc. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art. Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly-expressed gene to direct
35

transcription of a downstream structural sequence. Such promoters can be derived from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), a-factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an amino terminal identification peptide imparting desired characteristics, *e.g.*, stabilization or simplified purification of expressed recombinant product. Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM 1 (Promega Biotech, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced or derepressed by appropriate means (*e.g.*, temperature shift or chemical induction) and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Polynucleotides of the invention can also be used to induce immune responses. For example, as described in Fan et al., *Nat. Biotech.* 17:870-872 (1999), incorporated herein by reference, nucleic acid sequences encoding a polypeptide may be used to generate antibodies against the encoded polypeptide following topical administration of naked plasmid DNA or following injection, and preferably intramuscular injection of the DNA. The nucleic acid sequences are preferably inserted in a recombinant expression vector and may be in the form of naked DNA.

4.3 ANTISENSE

Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of a protein of any of SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960 or antisense nucleic acids complementary to a nucleic acid sequence of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence of the invention. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence of the invention. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding a nucleic acid disclosed herein (*e.g.*, SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of a mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of a mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of a mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a protein according to the invention to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the

strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett* 215: 327-330).

5

4.4 RIBOZYMES AND PNA MOIETIES

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme.

Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as a mRNA, to which they have a complementary region.

10 Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave a mRNA transcripts to thereby inhibit translation of a mRNA. A ribozyme having specificity for a nucleic acid of the invention can be designed based upon the nucleotide sequence of a DNA disclosed herein (*i.e.*, SEQ ID NO: 1-1984, 1969-2952, 3937-3942 or 3949-3954). For example, a derivative of a Tetrahymena L-19

15 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a SECX-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Pat. No. 4,987,071; and Cech *et al.* U.S. Pat. No. 5,116,742. Alternatively, SECX mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel *et al.*, (1993) *Science* 261:1411-1418.

20 Alternatively, gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region (*e.g.*, promoter and/or enhancers) to form triple helical structures that prevent transcription of the gene in target cells. See generally, Helene. (1991) *Anticancer Drug Des.* 6: 569-84; Helene. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14: 807-15.

25 In various embodiments, the nucleic acids of the invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorg Med Chem* 4: 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid

30 mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996) above;

35 Perry-O'Keefe *et al.* (1996) *PNAS* 93: 14670-675.

PNAs of the invention can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of the invention can also be used, *e.g.*, in the analysis of single base pair mutations in a
5 gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup B. (1996) above); or as probes or primers for DNA sequence and hybridization (Hyrup *et al.* (1996), above; Perry-O'Keefe (1996), above).

In another embodiment, PNAs of the invention can be modified, *e.g.*, to enhance their
10 stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras can be generated that may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA
15 portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup (1996) above). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996) above and Finn *et al.* (1996) *Nucl Acids Res* 24: 3357-63. For example, a DNA chain can be synthesized on a solid support using standard
20 phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA (Mag *et al.* (1989) *Nucl Acid Res* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.* (1996) above). Alternatively, chimeric molecules can be synthesized
25 with a 5' DNA segment and a 3' PNA segment. See, Petersen *et al.* (1975) *Bioorg Med Chem Lett* 5: 1119-1124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556;
30 Lemaitre *et al.*, 1987, *Proc. Natl. Acad. Sci.* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (See, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon, 1988, *Pharm. Res.* 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a

peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

4.5 HOSTS

5 The present invention further provides host cells genetically engineered to contain the polynucleotides of the invention. For example, such host cells may contain nucleic acids of the invention introduced into the host cell using known transformation, transfection or infection methods. The present invention still further provides host cells genetically engineered to express the polynucleotides of the invention, wherein such polynucleotides are in operative association
10 with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell.

Knowledge of nucleic acid sequences allows for modification of cells to permit, or increase, expression of endogenous polypeptide. Cells can be modified (e.g., by homologous recombination) to provide increased polypeptide expression by replacing, in whole or in part, the
15 naturally occurring promoter with all or part of a heterologous promoter so that the cells express the polypeptide at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to the encoding sequences. See, for example, PCT International Publication No. WO94/12650, PCT International Publication No. WO92/20808, and PCT International Publication No. WO91/09955. It is also contemplated that, in addition to heterologous promoter
20 DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the desired protein coding sequences in the cells.

25 The host cell can be a higher eukaryotic host cell, such as a mammalian cell, a lower eukaryotic host cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the recombinant construct into the host cell can be effected by calcium phosphate transfection, DEAE, dextran mediated transfection, or electroporation (Davis, L. et al., *Basic Methods in Molecular Biology* (1986)). The host cells containing one of the
30 polynucleotides of the invention, can be used in conventional manners to produce the gene product encoded by the isolated fragment (in the case of an ORF) or can be used to produce a heterologous protein under the control of the EMF.

Any host/vector system can be used to express one or more of the ORFs of the present invention. These include, but are not limited to, eukaryotic hosts such as HeLa cells, Cv-1 cell,
35 COS cells, 293 cells, and Sf9 cells, as well as prokaryotic host such as *E. coli* and *B. subtilis*.

The most preferred cells are those which do not normally express the particular polypeptide or protein or which expresses the polypeptide or protein at low natural level. Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using
5 RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, New York (1989), the disclosure of which is hereby incorporated by reference.

Various mammalian cell culture systems can also be employed to express recombinant
10 protein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, *Cell* 23:175 (1981). Other cell lines capable of expressing a compatible vector are, for example, the C127, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived
15 from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example,
20 SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements. Recombinant polypeptides and proteins produced in bacterial culture are usually isolated by initial extraction from cell pellets, followed by one or more salting-out, aqueous ion exchange or size exclusion chromatography steps. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein.
25 Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast
30 or insects or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it
35 may be necessary to modify the protein produced therein, for example by phosphorylation or

glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by targeting. These sequence include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the host cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (gpt) gene.

The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No. 5 PCT/US90/06436 (WO91/06667) by Skoultchi et al., each of which is incorporated by reference herein in its entirety.

4.6 POLYPEPTIDES OF THE INVENTION

The isolated polypeptides of the invention include, but are not limited to, a polypeptide 10 comprising: the amino acid sequences set forth as any one of SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960 or an amino acid sequence encoded by any one of the nucleotide sequences SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides preferably with biological or immunological activity that are encoded by: (a) a polynucleotide having any one of 15 the nucleotide sequences set forth in SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 or (b) polynucleotides encoding any one of the amino acid sequences set forth as SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960 or (c) polynucleotides that hybridize to the complement of the polynucleotides of either (a) or (b) under stringent hybridization conditions. The invention also provides biologically active or immunologically active variants of any of the 20 amino acid sequences set forth as SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960 or the corresponding full length or mature protein; and "substantial equivalents" thereof (e.g., at least about 65%, at least about 70%, at least about 75%, at least about 80%, 81%, 82%, 83%, 84%, more typically at least about 85%, 86%, 87%, 88%, 89%, and more typically at least about 90%, 91%, 92%, 93%, 94%, and even more typically at least about 95%, 96%, 97%, 98%, 99%, 25 sequence identity that retain biological activity. Polypeptides encoded by allelic variants may have a similar, increased, or decreased activity compared to polypeptides comprising SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may 30 be in linear form or they may be cyclized using known methods, for example, as described in H. U. Saragovi, et al., Bio/Technology 10, 773-778 (1992) and in R. S. McDowell, et al., J. Amer. Chem. Soc. 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites.

The present invention also provides both full-length and mature forms (for example, without a signal sequence or precursor sequence) of the disclosed proteins. The protein coding sequence is identified in the sequence listing by translation of the disclosed nucleotide sequences. The mature form of such protein may be obtained by expression of a full-length polynucleotide in a suitable mammalian cell or other host cell. The sequence of the mature form of the protein is also determinable from the amino acid sequence of the full-length form. Where proteins of the present invention are membrane bound, soluble forms of the proteins are also provided. In such forms, part or all of the regions causing the proteins to be membrane bound are deleted so that the proteins are fully secreted from the cell in which they are expressed.

Protein compositions of the present invention may further comprise an acceptable carrier, such as a hydrophilic, *e.g.*, pharmaceutically acceptable, carrier.

The present invention further provides isolated polypeptides encoded by the nucleic acid fragments of the present invention or by degenerate variants of the nucleic acid fragments of the present invention. By "degenerate variant" is intended nucleotide fragments which differ from a nucleic acid fragment of the present invention (*e.g.*, an ORF) by nucleotide sequence but, due to the degeneracy of the genetic code, encode an identical polypeptide sequence. Preferred nucleic acid fragments of the present invention are the ORFs that encode proteins.

A variety of methodologies known in the art can be utilized to obtain any one of the isolated polypeptides or proteins of the present invention. At the simplest level, the amino acid sequence can be synthesized using commercially available peptide synthesizers. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins that possess biological properties in common therewith, including protein activity. This technique is particularly useful in producing small peptides and fragments of larger polypeptides. Fragments are useful, for example, in generating antibodies against the native polypeptide. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The polypeptides and proteins of the present invention can alternatively be purified from cells which have been altered to express the desired polypeptide or protein. As used herein, a cell is said to be altered to express a desired polypeptide or protein when the cell, through genetic manipulation, is made to produce a polypeptide or protein which it normally does not produce or which the cell normally produces at a lower level. One skilled in the art can readily adapt procedures for introducing and expressing either recombinant or synthetic sequences into eukaryotic or prokaryotic cells in order to generate a cell which produces one of the polypeptides or proteins of the present invention.

The invention also relates to methods for producing a polypeptide comprising growing a culture of host cells of the invention in a suitable culture medium, and purifying the protein from the cells or the culture in which the cells are grown. For example, the methods of the invention include a process for producing a polypeptide in which a host cell containing a suitable
5 expression vector that includes a polynucleotide of the invention is cultured under conditions that allow expression of the encoded polypeptide. The polypeptide can be recovered from the culture, conveniently from the culture medium, or from a lysate prepared from the host cells and further purified. Preferred embodiments include those in which the protein produced by such process is a full length or mature form of the protein.

10 In an alternative method, the polypeptide or protein is purified from bacterial cells which naturally produce the polypeptide or protein. One skilled in the art can readily follow known methods for isolating polypeptides and proteins in order to obtain one of the isolated polypeptides or proteins of the present invention. These include, but are not limited to, immunochromatography, HPLC, size-exclusion chromatography, ion-exchange chromatography,
15 and immuno-affinity chromatography. See, e.g., Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag (1994); Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*; Ausubel et al., *Current Protocols in Molecular Biology*. Polypeptide fragments that retain biological/immunological activity include fragments comprising greater than about 100 amino acids, or greater than about 200 amino acids, and fragments that encode specific protein
20 domains.

The purified polypeptides can be used in *in vitro* binding assays which are well known in the art to identify molecules which bind to the polypeptides. These molecules include but are not limited to, for e.g., small molecules, molecules from combinatorial libraries, antibodies or other proteins. The molecules identified in the binding assay are then tested for antagonist or agonist
25 activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

In addition, the peptides of the invention or molecules capable of binding to the peptides may be complexed with toxins, e.g., ricin or cholera, or with other compounds that are toxic to
30 cells. The toxin-binding molecule complex is then targeted to a tumor or other cell by the specificity of the binding molecule for SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3960.

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized
35 by somatic or germ cells containing a nucleotide sequence encoding the protein.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications, in the peptide or DNA sequence, can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, *e.g.*, U.S. Pat. No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein. Regions of the protein that are important for the protein function can be determined by various methods known in the art including the alanine-scanning method which involved systematic substitution of single or strings of amino acids with alanine, followed by testing the resulting alanine-containing variant for biological activity. This type of analysis determines the importance of the substituted amino acid(s) in biological activity. Regions of the protein that are important for protein function may be determined by the eMATRIX program.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and are useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are encompassed by the present invention.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *e.g.*, Invitrogen, San Diego, Calif., U.S.A. (the MaxBat™ kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl™ or Cibacrom blue 3GA Sepharose™; one or more steps involving

hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX), or as a His tag. Kits for expression and purification of such fusion proteins are commercially available from New England BioLab (Beverly, Mass.), Pharmacia (Piscataway, N.J.) and Invitrogen, respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("FLAG®") is commercially available from Kodak (New Haven, Conn.).

Finally, one or more reverse-phase high performance liquid chromatography (RP- HPLC) steps employing hydrophobic RP-HPLC media, *e.g.*, silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The polypeptides of the invention include analogs (variants). This embraces fragments, as well as peptides in which one or more amino acids has been deleted, inserted, or substituted. Also, analogs of the polypeptides of the invention embrace fusions of the polypeptides or modifications of the polypeptides of the invention, wherein the polypeptide or analog is fused to another moiety or moieties, *e.g.*, targeting moiety or another therapeutic agent. Such analogs may exhibit improved properties such as activity and/or stability. Examples of moieties which may be fused to the polypeptide or an analog include, for example, targeting moieties which provide for the delivery of polypeptide to pancreatic cells, *e.g.*, antibodies to pancreatic cells, antibodies to immune cells such as T-cells, monocytes, dendritic cells, granulocytes, etc., as well as receptor and ligands expressed on pancreatic or immune cells. Other moieties which may be fused to the polypeptide include therapeutic agents which are used for treatment, for example, immunosuppressive drugs such as cyclosporin, SK506, azathioprine, CD3 antibodies and steroids. Also, polypeptides may be fused to immune modulators, and other cytokines such as alpha or beta interferon.

4.6.1 DETERMINING POLYPEPTIDE AND POLYNUCLEOTIDE IDENTITY AND SIMILARITY

Preferred identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in computer programs including, but are not limited to, the GCG program package, including GAP (Devereux, J., et al., *Nucleic Acids Research* 12(1):387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, BLASTX, FASTA (Altschul, S.F. et al., *J. Molec. Biol.* 215:403-410 (1990), PSI-BLAST (Altschul S.F. et al., *Nucleic Acids Res.* vol. 25, pp. 3389-3402, herein incorporated by reference), eMatrix software (Wu et al., *J. Comp. Biol.*, Vol. 6, pp. 219-235 (1999), herein incorporated by reference), eMotif software (Nevill-Manning et al, *ISMB-97*, Vol. 4, pp. 202-209, herein incorporated by reference), pFam software (Sonnhammer et al., *Nucleic Acids Res.*, Vol. 26(1), pp. 320-322 (1998), herein incorporated by reference) and the Kyte-Doolittle hydrophobicity prediction algorithm (*J. Mol Biol*, 157, pp. 105-31 (1982), incorporated herein by reference). The BLAST programs are publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul, S., et al. NCB NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215:403-410 (1990).

4.7 CHIMERIC AND FUSION PROTEINS

The invention also provides chimeric or fusion proteins. As used herein, a "chimeric protein" or "fusion protein" comprises a polypeptide of the invention operatively linked to another polypeptide. Within a fusion protein the polypeptide according to the invention can correspond to all or a portion of a protein according to the invention. In one embodiment, a fusion protein comprises at least one biologically active portion of a protein according to the invention. In another embodiment, a fusion protein comprises at least two biologically active portions of a protein according to the invention. Within the fusion protein, the term "operatively linked" is intended to indicate that the polypeptide according to the invention and the other polypeptide are fused in-frame to each other. The polypeptide can be fused to the N-terminus or C-terminus.

For example, in one embodiment a fusion protein comprises a polypeptide according to the invention operably linked to the extracellular domain of a second protein. In another embodiment, the fusion protein is a GST-fusion protein in which the polypeptide sequences of the invention are fused to the C-terminus of the GST (i.e., glutathione S-transferase) sequences.

In another embodiment, the fusion protein is an immunoglobulin fusion protein in which the polypeptide sequences according to the invention comprise one or more domains fused to sequences derived from a member of the immunoglobulin protein family. The immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and

administered to a subject to inhibit an interaction between a ligand and a protein of the invention on the surface of a cell, to thereby suppress signal transduction *in vivo*. The immunoglobulin fusion proteins can be used to affect the bioavailability of a cognate ligand. Inhibition of the ligand/protein interaction may be useful therapeutically for both the treatment of proliferative and differentiative disorders, *e.g.*, cancer as well as modulating (*e.g.*, promoting or inhibiting) cell survival. Moreover, the immunoglobulin fusion proteins of the invention can be used as immunogens to produce antibodies in a subject, to purify ligands, and in screening assays to identify molecules that inhibit the interaction of a polypeptide of the invention with a ligand.

A chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, Ausubel et al. (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A nucleic acid encoding a polypeptide of the invention can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the protein of the invention.

4.8 GENE THERAPY

Mutations in the polynucleotides of the invention gene may result in loss of normal function of the encoded protein. The invention thus provides gene therapy to restore normal activity of the polypeptides of the invention; or to treat disease states involving polypeptides of the invention. Delivery of a functional gene encoding polypeptides of the invention to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and more particularly viral vectors (*e.g.*, adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (*e.g.*, liposomes or chemical treatments). See, for example, Anderson, Nature, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, Science, 244: 1275-1281 (1989); Verma, Scientific American: 68-84 (1990); and Miller, Nature, 357: 455-460 (1992). Introduction of any one of

the nucleotides of the present invention or a gene encoding the polypeptides of the present invention can also be accomplished with extrachromosomal substrates (transient expression) or artificial chromosomes (stable expression). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or
5 activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes. Alternatively, it is contemplated that in other human disease states, preventing the expression of or inhibiting the activity of polypeptides of the invention will be useful in treating the disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of polypeptides of the invention.

10 Other methods inhibiting expression of a protein include the introduction of antisense molecules to the nucleic acids of the present invention, their complements, or their translated RNA sequences, by methods known in the art. Further, the polypeptides of the present invention can be inhibited by using targeted deletion methods, or the insertion of a negative regulatory element such as a silencer, which is tissue specific.

15 The present invention still further provides cells genetically engineered *in vivo* to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell. These methods can be used to increase or decrease the expression of the polynucleotides of the present invention.

20 Knowledge of DNA sequences provided by the invention allows for modification of cells to permit, increase, or decrease, expression of endogenous polypeptide. Cells can be modified (e.g., by homologous recombination) to provide increased polypeptide expression by replacing, in whole or in part, the naturally occurring promoter with all or part of a heterologous promoter so that the cells express the protein at higher levels. The heterologous promoter is inserted in such a manner that it is
25 operatively linked to the desired protein encoding sequences. See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955. It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or
30 intron DNA may be inserted along with the heterologous promoter DNA. If linked to the desired protein coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the desired protein coding sequences in the cells.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of
35 inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may

be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by targeting. These sequences include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (*gpt*) gene.

The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No. PCT/US90/06436 (WO91/06667) by Skoultchi et al., each of which is incorporated by reference herein in its entirety.

4.9 TRANSGENIC ANIMALS

In preferred methods to determine biological functions of the polypeptides of the invention *in vivo*, one or more genes provided by the invention are either over expressed or inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals, preferably non-human mammals, can be prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference. Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model systems to identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals, are produced using methods as described in U.S. Patent No 5,489,743 and PCT Publication No. WO94/28122, incorporated herein by reference.

Transgenic animals can be prepared wherein all or part of a promoter of the polynucleotides of the invention is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using homologous recombination methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased protein expression. The homologous promoter can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

The polynucleotides of the present invention also make possible the development, through, e.g., homologous recombination or knock out strategies, of animals that fail to express polypeptides of the invention or that express a variant polypeptide. Such animals are useful as models for studying the *in vivo* activities of polypeptide as well as for studying modulators of the polypeptides of the invention.

In preferred methods to determine biological functions of the polypeptides of the invention *in vivo*, one or more genes provided by the invention are either over expressed or inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals, preferably non-human mammals, can be prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference. Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model systems to

identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals, are produced using methods as described in U.S. Patent No 5,489,743 and PCT Publication No. WO94/28122, incorporated herein by reference.

Transgenic animals can be prepared wherein all or part of the polynucleotides of the invention promoter is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using homologous recombination methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased protein expression. The homologous promoter can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

4.10 USES AND BIOLOGICAL ACTIVITY

The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified herein. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA). The mechanism underlying the particular condition or pathology will dictate whether the polypeptides of the invention, the polynucleotides of the invention or modulators (activators or inhibitors) thereof would be beneficial to the subject in need of treatment. Thus, "therapeutic compositions of the invention" include compositions comprising isolated polynucleotides (including recombinant DNA molecules, cloned genes and degenerate variants thereof) or polypeptides of the invention (including full length protein, mature protein and truncations or domains thereof), or compounds and other substances that modulate the overall activity of the target gene products, either at the level of target gene/protein expression or target protein activity. Such modulators include polypeptides, analogs, (variants), including fragments and fusion proteins, antibodies and other binding proteins; chemical compounds that directly or indirectly activate or inhibit the polypeptides of the invention (identified, e.g., via drug screening assays as described herein); antisense polynucleotides and polynucleotides suitable for triple helix formation; and in particular antibodies or other binding partners that specifically recognize one or more epitopes of the polypeptides of the invention.

The polypeptides of the present invention may likewise be involved in cellular activation or in one of the other physiological pathways described herein.

4.10.1 RESEARCH USES AND UTILITIES

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The polypeptides provided by the present invention can similarly be used in assays to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding polypeptide is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

4.10.2 NUTRITIONAL USES

Polynucleotides and polypeptides of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the polypeptide or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the polypeptide or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

4.10.3 CYTOKINE AND CELL PROLIFERATION/DIFFERENTIATION ACTIVITY

A polypeptide of the present invention may exhibit activity relating to cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of therapeutic compositions of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+(preB M+), 2E8, RB, DA1, 123, T1165, HT2, CTLL2, TF-1, M-1e, CMK, HUVEC, and Caco. Therapeutic compositions of the invention can be used in the following:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992; Bowman et al., J. Immunol. 152:1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A. M. and Shevach, E. M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse

and human interleukin- γ , Schreiber, R. D. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L. S. and Lipsky, P. E. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6--Nordan, R. In Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11--Bennett, F., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9--Ciarletta, A., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic Studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6095-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

4.10.4 STEM CELL GROWTH FACTOR ACTIVITY

A polypeptide of the present invention may exhibit stem cell growth factor activity and be involved in the proliferation, differentiation and survival of pluripotent and totipotent stem cells including primordial germ cells, embryonic stem cells, hematopoietic stem cells and/or germ line stem cells. Administration of the polypeptide of the invention to stem cells *in vivo* or *ex vivo* is expected to maintain and expand cell populations in a totipotential or pluripotential state which would be useful for re-engineering damaged or diseased tissues, transplantation, manufacture of bio-pharmaceuticals and the development of bio-sensors. The ability to produce large quantities of human cells has important working applications for the production of human proteins which currently must be obtained from non-human sources or donors, implantation of

cells to treat diseases such as Parkinson's, Alzheimer's and other neurodegenerative diseases; tissues for grafting such as bone marrow, skin, cartilage, tendons, bone, muscle (including cardiac muscle), blood vessels, cornea, neural cells, gastrointestinal cells and others; and organs for transplantation such as kidney, liver, pancreas (including islet cells), heart and lung.

5 It is contemplated that multiple different exogenous growth factors and/or cytokines may be administered in combination with the polypeptide of the invention to achieve the desired effect, including any of the growth factors listed herein, other stem cell maintenance factors, and specifically including stem cell factor (SCF), leukemia inhibitory factor (LIF), Flt-3 ligand (Flt-3L), any of the interleukins, recombinant soluble IL-6 receptor fused to IL-6, macrophage
10 inflammatory protein 1-alpha (MIP-1-alpha), G-CSF, GM-CSF, thrombopoietin (TPO), platelet factor 4 (PF-4), platelet-derived growth factor (PDGF), neural growth factors and basic fibroblast growth factor (bFGF).

Since totipotent stem cells can give rise to virtually any mature cell type, expansion of these cells in culture will facilitate the production of large quantities of mature cells. Techniques
15 for culturing stem cells are known in the art and administration of polypeptides of the invention, optionally with other growth factors and/or cytokines, is expected to enhance the survival and proliferation of the stem cell populations. This can be accomplished by direct administration of the polypeptide of the invention to the culture medium. Alternatively, stroma cells transfected with a polynucleotide that encodes for the polypeptide of the invention can be used as a feeder
20 layer for the stem cell populations in culture or in vivo. Stromal support cells for feeder layers may include embryonic bone marrow fibroblasts, bone marrow stromal cells, fetal liver cells, or cultured embryonic fibroblasts (see U.S. Patent No. 5,600,926).

Stem cells themselves can be transfected with a polynucleotide of the invention to induce
autocrine expression of the polypeptide of the invention. This will allow for generation of
25 undifferentiated totipotent/pluripotent stem cell lines that are useful as is or that can then be differentiated into the desired mature cell types. These stable cell lines can also serve as a source of undifferentiated totipotent/pluripotent mRNA to create cDNA libraries and templates for polymerase chain reaction experiments. These studies would allow for the isolation and identification of differentially expressed genes in stem cell populations that regulate stem cell
30 proliferation and/or maintenance.

Expansion and maintenance of totipotent stem cell populations will be useful in the treatment of many pathological conditions. For example, polypeptides of the present invention may be used to manipulate stem cells in culture to give rise to neuroepithelial cells that can be used to augment or replace cells damaged by illness, autoimmune disease, accidental damage or
35 genetic disorders. The polypeptide of the invention may be useful for inducing the proliferation

of neural cells and for the regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders which involve degeneration, death or trauma to neural cells or nerve tissue. In addition, the expanded stem cell populations can also be genetically altered for gene therapy purposes and to decrease host rejection of replacement tissues after grafting or implantation.

Expression of the polypeptide of the invention and its effect on stem cells can also be manipulated to achieve controlled differentiation of the stem cells into more differentiated cell types. A broadly applicable method of obtaining pure populations of a specific differentiated cell type from undifferentiated stem cell populations involves the use of a cell-type specific promoter driving a selectable marker. The selectable marker allows only cells of the desired type to survive. For example, stem cells can be induced to differentiate into cardiomyocytes (Wobus et al., *Differentiation*, 48: 173-182, (1991); Klug et al., *J. Clin. Invest.*, 98(1): 216-224, (1998)) or skeletal muscle cells (Browder, L. W. In: *Principles of Tissue Engineering eds.* Lanza et al., Academic Press (1997)). Alternatively, directed differentiation of stem cells can be accomplished by culturing the stem cells in the presence of a differentiation factor such as retinoic acid and an antagonist of the polypeptide of the invention which would inhibit the effects of endogenous stem cell factor activity and allow differentiation to proceed.

In vitro cultures of stem cells can be used to determine if the polypeptide of the invention exhibits stem cell growth factor activity. Stem cells are isolated from any one of various cell sources (including hematopoietic stem cells and embryonic stem cells) and cultured on a feeder layer, as described by Thompson et al. *Proc. Natl. Acad. Sci, U.S.A.*, 92: 7844-7848 (1995), in the presence of the polypeptide of the invention alone or in combination with other growth factors or cytokines. The ability of the polypeptide of the invention to induce stem cells proliferation is determined by colony formation on semi-solid support e.g. as described by Bernstein et al., *Blood*, 77: 2316-2321 (1991).

4.10.5 HEMATOPOIESIS REGULATING ACTIVITY

A polypeptide of the present invention may be involved in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell disorders. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e.,

traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or
5 complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment
10 post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

Therapeutic compositions of the invention can be used in the following:

Suitable assays for proliferation and differentiation of various hematopoietic lines are
15 cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al., Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood 81:2903-2915, 1993.

20 Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M. G. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, N.Y. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells
25 with high proliferative potential, McNiece, I. K. and Briddell, R. A. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, N.Y. 1994; Neben et al., Experimental Hematology 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R. E. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, N.Y. 1994; Long term bone marrow cultures in the presence of
30 stromal cells, Spooncer, E., Dexter, M. and Allen, T. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, N.Y. 1994; Long term culture initiating cell assay, Sutherland, H. J. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, N.Y. 1994.

35 4.10.6 TISSUE GROWTH ACTIVITY

A polypeptide of the present invention also may be involved in bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as in wound healing and tissue repair and replacement, and in healing of burns, incisions and ulcers.

5 A polypeptide of the present invention which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Compositions of a polypeptide, antibody, binding partner, or other modulator of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an osteogenic agent contributes to the repair
10 of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A polypeptide of this invention may also be involved in attracting bone-forming cells, stimulating growth of bone-forming cells, or inducing differentiation of progenitors of bone-forming cells. Treatment of osteoporosis, osteoarthritis, bone degenerative disorders, or
15 periodontal disease, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes may also be possible using the composition of the invention.

Another category of tissue regeneration activity that may involve the polypeptide of the
20 present invention is tendon/ligament formation. Induction of tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as
25 use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may
30 provide environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include
35 an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The compositions of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a composition may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a composition of the invention.

Compositions of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

Compositions of the present invention may also be involved in the generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring may allow normal tissue to regenerate. A polypeptide of the present invention may also exhibit angiogenic activity.

A composition of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A composition of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

Therapeutic compositions of the invention can be used in the following:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, H. I. and Rovee, D. T., eds.), Year Book

Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

4.10.7 IMMUNE STIMULATING OR SUPPRESSING ACTIVITY

5 A polypeptide of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A polynucleotide of the invention can encode a polypeptide exhibiting such activities. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and
10 proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses,
15 herpes viruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, proteins of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus,
20 rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease, and autoimmune inflammatory eye disease. Such a protein (or antagonists thereof, including antibodies) of the present invention may also be useful in the treatment of allergic reactions and conditions (e.g., anaphylaxis, serum sickness, drug reactions, food allergies, insect
25 venom allergies, mastocytosis, allergic rhinitis, hypersensitivity pneumonitis, urticaria, angioedema, eczema, atopic dermatitis, allergic contact dermatitis, erythema multiforme, Stevens-Johnson syndrome, allergic conjunctivitis, atopic keratoconjunctivitis, venereal keratoconjunctivitis, giant papillary conjunctivitis and contact allergies), such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune
30 suppression is desired (including, for example, organ transplantation), may also be treatable using a protein (or antagonists thereof) of the present invention. The therapeutic effects of the polypeptides or antagonists thereof on allergic reactions can be evaluated by in vivo animals models such as the cumulative contact enhancement test (Lastbom et al., Toxicology 125: 59-66, 1998), skin prick test (Hoffmann et al., Allergy 54: 446-54, 1999), guinea pig skin sensitization

test (Vohr et al., Arch. Toxicol. 73: 501-9), and murine local lymph node assay (Kimber et al., J. Toxicol. Environ. Health 53: 563-79).

Using the proteins of the invention it may also be possible to modulate immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as, for example, B7)), e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a therapeutic composition of the invention may prevent cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, a lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular therapeutic compositions in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins in vivo as described in Lenschow et al., Science 257:789-792 (1992) and Turka et al., Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of therapeutic compositions of the invention on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may
5 reduce or eliminate disease symptoms. Administration of reagents which block stimulation of T cells can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating
10 autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp.
15 840-856).

Upregulation of an antigen function (e.g., a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response may be useful in cases of viral
20 infection, including systemic viral diseases such as influenza, the common cold, and encephalitis.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the *in vitro* activated T cells into the
25 patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells *in vivo*.

A polypeptide of the present invention may provide the necessary stimulation signal to T
30 cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient mounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an
35 MHC class I alpha chain protein and β_2 microglobulin protein or an MHC class II alpha chain

protein and an MHC class II beta chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bowman et al., J. Virology 61:1992-1998; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J. J. and Brunswick, M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery

et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

4.10.8 ACTIVIN/INHIBIN ACTIVITY

A polypeptide of the present invention may also exhibit activin- or inhibin-related activities. A polynucleotide of the invention may encode a polypeptide exhibiting such characteristics. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a polypeptide of the present invention, alone or in heterodimers with a member of the inhibin family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the polypeptide of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, U.S. Pat. No. 4,798,885. A polypeptide of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as, but not limited to, cows, sheep and pigs.

The activity of a polypeptide of the invention may, among other means, be measured by the following methods.

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., *Endocrinology* 91:562-572, 1972; Ling et al., *Nature* 321:779-782, 1986; Vale et al., *Nature* 321:776-779, 1986; Mason et al., *Nature* 318:659-663, 1985; Forage et al., *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986.

4.10.9 CHEMOTACTIC/CHEMOKINETIC ACTIVITY

A polypeptide of the present invention may be involved in chemotactic or chemokinetic activity for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Chemotactic and chemokinetic receptor activation can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic compositions (e.g. proteins, antibodies, binding partners, or modulators of the invention) provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

Therapeutic compositions of the invention can be used in the following:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margules, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. *J. Clin. Invest.* 95:1370-1376, 1995; Lind et al. *APMIS* 103:140-146, 1995; Muller et al. *Eur. J. Immunol.* 25:1744-1748; Gruber et al. *J. of Immunol.* 152:5860-5867, 1994; Johnston et al. *J. of Immunol.* 153:1762-1768, 1994.

4.10.10 HEMOSTATIC AND THROMBOLYTIC ACTIVITY

A polypeptide of the invention may also be involved in hemostasis or thrombolysis or thrombosis. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Compositions may be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A composition of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

Therapeutic compositions of the invention can be used in the following:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

4.10.11 CANCER DIAGNOSIS AND THERAPY

Polypeptides of the invention may be involved in cancer cell generation, proliferation or metastasis. Detection of the presence or amount of polynucleotides or polypeptides of the invention may be useful for the diagnosis and/or prognosis of one or more types of cancer. For example, the presence or increased expression of a polynucleotide/polypeptide of the invention may indicate a hereditary risk of cancer, a precancerous condition, or an ongoing malignancy. Conversely, a defect in the gene or absence of the polypeptide may be associated with a cancer condition. Identification of single nucleotide polymorphisms associated with cancer or a predisposition to cancer may also be useful for diagnosis or prognosis.

Cancer treatments promote tumor regression by inhibiting tumor cell proliferation, inhibiting angiogenesis (growth of new blood vessels that is necessary to support tumor growth) and/or prohibiting metastasis by reducing tumor cell motility or invasiveness. Therapeutic compositions of the invention may be effective in adult and pediatric oncology including in solid phase tumors/malignancies, locally advanced tumors, human soft tissue sarcomas, metastatic cancer, including lymphatic metastases, blood cell malignancies including multiple myeloma, acute and chronic leukemias, and lymphomas, head and neck cancers including mouth cancer, larynx cancer and thyroid cancer, lung cancers including small cell carcinoma and non-small cell cancers, breast cancers including small cell carcinoma and ductal carcinoma, gastrointestinal cancers including esophageal cancer, stomach cancer, colon cancer, colorectal cancer and polyps associated with colorectal neoplasia, pancreatic cancers, liver cancer, urologic cancers including

bladder cancer and prostate cancer, malignancies of the female genital tract including ovarian carcinoma, uterine (including endometrial) cancers, and solid tumor in the ovarian follicle, kidney cancers including renal cell carcinoma, brain cancers including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in the central
5 nervous system, bone cancers including osteomas, skin cancers including malignant melanoma, tumor progression of human skin keratinocytes, squamous cell carcinoma, basal cell carcinoma, hemangiopericytoma and Kaposi's sarcoma.

Polypeptides, polynucleotides, or modulators of polypeptides of the invention (including inhibitors and stimulators of the biological activity of the polypeptide of the invention) may be
10 administered to treat cancer. Therapeutic compositions can be administered in therapeutically effective dosages alone or in combination with adjuvant cancer therapy such as surgery, chemotherapy, radiotherapy, thermotherapy, and laser therapy, and may provide a beneficial effect, e.g. reducing tumor size, slowing rate of tumor growth, inhibiting metastasis, or otherwise improving overall clinical condition, without necessarily eradicating the cancer.

15 The composition can also be administered in therapeutically effective amounts as a portion of an anti-cancer cocktail. An anti-cancer cocktail is a mixture of the polypeptide or modulator of the invention with one or more anti-cancer drugs in addition to a pharmaceutically acceptable carrier for delivery. The use of anti-cancer cocktails as a cancer treatment is routine. Anti-cancer drugs that are well known in the art and can be used as a treatment in combination
20 with the polypeptide or modulator of the invention include: Actinomycin D, Aminoglutethimide, Asparaginase, Bleomycin, Busulfan, Carboplatin, Carmustine, Chlorambucil, Cisplatin (cis-DDP), Cyclophosphamide, Cytarabine HCl (Cytosine arabinoside), Dacarbazine, Daunorubicin HCl, Doxorubicin HCl, Estramustine phosphate sodium, Etoposide (V16-213), Floxuridine, 5-Fluorouracil (5-Fu), Flutamide, Hydroxyurea (hydroxycarbamide), Ifosfamide,
25 Interferon Alpha-2a, Interferon Alpha-2b, Leuprolide acetate (LHRH-releasing factor analog), Lomustine, Mechlorethamine HCl (nitrogen mustard), Melphalan, Mercaptopurine, Mesna, Methotrexate (MTX), Mitomycin, Mitoxantrone HCl, Octreotide, Plicamycin, Procarbazine HCl, Streptozocin, Tamoxifen citrate, Thioguanine, Thiotepa, Vinblastine sulfate, Vincristine sulfate, Amsacrine, Azacitidine, Hexamethylmelamine, Interleukin-2, Mitoguazone, Pentostatin,
30 Semustine, Teniposide, and Vindesine sulfate.

In addition, therapeutic compositions of the invention may be used for prophylactic treatment of cancer. There are hereditary conditions and/or environmental situations (e.g. exposure to carcinogens) known in the art that predispose an individual to developing cancers. Under these circumstances, it may be beneficial to treat these individuals with therapeutically
35 effective doses of the polypeptide of the invention to reduce the risk of developing cancers.

In vitro models can be used to determine the effective doses of the polypeptide of the invention as a potential cancer treatment. These *in vitro* models include proliferation assays of cultured tumor cells, growth of cultured tumor cells in soft agar (see Freshney, (1987) Culture of Animal Cells: A Manual of Basic Technique, Wiley-Liss, New York, NY Ch 18 and Ch 21),
5 tumor systems in nude mice as described in Giovanella et al., J. Natl. Can. Inst., 52: 921-30 (1974), mobility and invasive potential of tumor cells in Boyden Chamber assays as described in Pilkington et al., Anticancer Res., 17: 4107-9 (1997), and angiogenesis assays such as induction of vascularization of the chick chorioallantoic membrane or induction of vascular endothelial cell migration as described in Ribatta et al., Intl. J. Dev. Biol., 40: 1189-97 (1999) and Li et al.,
10 Clin. Exp. Metastasis, 17:423-9 (1999), respectively. Suitable tumor cells lines are available, e.g. from American Type Tissue Culture Collection catalogs.

4.10.12 RECEPTOR/LIGAND ACTIVITY

A polypeptide of the present invention may also demonstrate activity as receptor,
15 receptor ligand or inhibitor or agonist of receptor/ligand interactions. A polynucleotide of the invention can encode a polypeptide exhibiting such characteristics. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins,
20 integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses. Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand
25 interactions.

The activity of a polypeptide of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley- Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1- 7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

By way of example, the polypeptides of the invention may be used as a receptor for a ligand(s) thereby transmitting the biological activity of that ligand(s). Ligands may be identified through binding assays, affinity chromatography, dihybrid screening assays, BIAcore assays, gel overlay assays, or other methods known in the art.

5 Studies characterizing drugs or proteins as agonist or antagonist or partial agonists or a partial antagonist require the use of other proteins as competing ligands. The polypeptides of the present invention or ligand(s) thereof may be labeled by being coupled to radioisotopes, colorimetric molecules or a toxin molecules by conventional methods. ("Guide to Protein Purification" Murray P. Deutscher (ed) Methods in Enzymology Vol. 182 (1990) Academic Press, Inc. San Diego). Examples of radioisotopes include, but are not limited to, tritium and
10 carbon-14. Examples of colorimetric molecules include, but are not limited to, fluorescent molecules such as fluorescamine, or rhodamine or other colorimetric molecules. Examples of toxins include, but are not limited, to ricin.

15 4.10.13 DRUG SCREENING

This invention is particularly useful for screening chemical compounds by using the novel polypeptides or binding fragments thereof in any of a variety of drug screening techniques. The polypeptides or fragments employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. One method of drug screening
20 utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or a fragment thereof. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between polypeptides of the invention or fragments and the agent being tested or examine the
25 diminution in complex formation between the novel polypeptides and an appropriate cell line, which are well known in the art.

Sources for test compounds that may be screened for ability to bind to or modulate (i.e., increase or decrease) the activity of polypeptides of the invention include (1) inorganic and organic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries
30 comprised of either random or mimetic peptides, oligonucleotides or organic molecules.

Chemical libraries may be readily synthesized or purchased from a number of commercial sources, and may include structural analogs of known compounds or compounds that are identified as "hits" or "leads" via natural product screening.

The sources of natural product libraries are microorganisms (including bacteria and
35 fungi), animals, plants or other vegetation, or marine organisms, and libraries of mixtures for

screening may be created by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of the organisms themselves. Natural product libraries include polyketides, non-ribosomal peptides, and (non-naturally occurring) variants thereof. For a review, see *Science* 282:63-68 (1998).

- 5 Combinatorial libraries are composed of large numbers of peptides, oligonucleotides or organic compounds and can be readily prepared by traditional automated synthesis methods, PCR, cloning or proprietary synthetic methods. Of particular interest are peptide and oligonucleotide combinatorial libraries. Still other libraries of interest include peptide, protein, peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries.
- 10 For a review of combinatorial chemistry and libraries created therefrom, see Myers, *Curr. Opin. Biotechnol.* 8:701-707 (1997). For reviews and examples of peptidomimetic libraries, see Al-Obeidi et al., *Mol. Biotechnol.*, 9(3):205-23 (1998); Hruby et al., *Curr Opin Chem Biol*, 1(1):114-19 (1997); Dorner et al., *Bioorg Med Chem*, 4(5):709-15 (1996) (alkylated dipeptides).

- Identification of modulators through use of the various libraries described herein permits
- 15 modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to bind a polypeptide of the invention. The molecules identified in the binding assay are then tested for antagonist or agonist activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

- 20 The binding molecules thus identified may be complexed with toxins, e.g., ricin or cholera, or with other compounds that are toxic to cells such as radioisotopes. The toxin-binding molecule complex is then targeted to a tumor or other cell by the specificity of the binding molecule for a polypeptide of the invention. Alternatively, the binding molecules may be complexed with imaging agents for targeting and imaging purposes.

25

4.10.14 ASSAY FOR RECEPTOR ACTIVITY

- The invention also provides methods to detect specific binding of a polypeptide e.g. a ligand or a receptor. The art provides numerous assays particularly useful for identifying previously unknown binding partners for receptor polypeptides of the invention. For example,
- 30 expression cloning using mammalian or bacterial cells, or dihybrid screening assays can be used to identify polynucleotides encoding binding partners. As another example, affinity chromatography with the appropriate immobilized polypeptide of the invention can be used to isolate polypeptides that recognize and bind polypeptides of the invention. There are a number of different libraries used for the identification of compounds, and in particular small molecules,
- 35 that modulate (*i.e.*, increase or decrease) biological activity of a polypeptide of the invention.

Ligands for receptor polypeptides of the invention can also be identified by adding exogenous ligands, or cocktails of ligands to two cells populations that are genetically identical except for the expression of the receptor of the invention: one cell population expresses the receptor of the invention whereas the other does not. The response of the two cell populations to the addition of ligands(s) are then compared. Alternatively, an expression library can be co-expressed with the polypeptide of the invention in cells and assayed for an autocrine response to identify potential ligand(s). As still another example, BIAcore assays, gel overlay assays, or other methods known in the art can be used to identify binding partner polypeptides, including, (1) organic and inorganic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules.

The role of downstream intracellular signaling molecules in the signaling cascade of the polypeptide of the invention can be determined. For example, a chimeric protein in which the cytoplasmic domain of the polypeptide of the invention is fused to the extracellular portion of a protein, whose ligand has been identified, is produced in a host cell. The cell is then incubated with the ligand specific for the extracellular portion of the chimeric protein, thereby activating the chimeric receptor. Known downstream proteins involved in intracellular signaling can then be assayed for expected modifications i.e. phosphorylation. Other methods known to those in the art can also be used to identify signaling molecules involved in receptor activity.

4.10.15 ANTI-INFLAMMATORY ACTIVITY

Compositions of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Compositions with such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation intimation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Compositions of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material. Compositions of this invention may be utilized to prevent or treat conditions such as, but not limited to, sepsis, acute pancreatitis, endotoxin shock, cytokine induced shock, rheumatoid

arthritis, chronic inflammatory arthritis, pancreatic cell damage from diabetes mellitus type 1, graft versus host disease, inflammatory bowel disease, inflammation associated with pulmonary disease, other autoimmune disease or inflammatory disease, an antiproliferative agent such as for acute or chronic myelogenous leukemia or in the prevention of premature labor secondary to intrauterine infections.

4.10.16 LEUKEMIAS

Leukemias and related disorders may be treated or prevented by administration of a therapeutic that promotes or inhibits function of the polynucleotides and/or polypeptides of the invention. Such leukemias and related disorders include but are not limited to acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia, chronic leukemia, chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia (for a review of such disorders, see Fishman et al., 1985, Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia).

4.10.17 NERVOUS SYSTEM DISORDERS

Nervous system disorders, involving cell types which can be tested for efficacy of intervention with compounds that modulate the activity of the polynucleotides and/or polypeptides of the invention, and which can be treated upon thus observing an indication of therapeutic utility, include but are not limited to nervous system injuries, and diseases or disorders which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the invention include but are not limited to the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems:

- (i) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries;
- (ii) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia;
- (iii) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis;

- (iv) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis;
- 5 (v) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration;
- 10 (vi) neurological lesions associated with systemic diseases including but not limited to diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis;
- (vii) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and
- 15 (viii) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including but not limited to multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

Therapeutics which are useful according to the invention for treatment of a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, therapeutics which elicit any of the following effects may be useful according to the invention:

- (i) increased survival time of neurons in culture;
- (ii) increased sprouting of neurons in culture or *in vivo*;
- 25 (iii) increased production of a neuron-associated molecule in culture or *in vivo*, *e.g.*, choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or
- (iv) decreased symptoms of neuron dysfunction *in vivo*.

Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may be measured by the method set forth in Arakawa et al. (1990, J. Neurosci. 10:3507-3515); increased sprouting of neurons may be detected by methods set forth in Pestronk et al. (1980, Exp. Neurol. 70:65-82) or Brown et al. (1981, Ann. Rev. Neurosci. 4:17-42); increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, *etc.*, depending on the molecule to be measured; and motor neuron dysfunction may be measured by

assessing the physical manifestation of motor neuron disorder, *e.g.*, weakness, motor neuron conduction velocity, or functional disability.

In specific embodiments, motor neuron disorders that may be treated according to the invention include but are not limited to disorders such as infarction, infection, exposure to toxin, 5 trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including but not limited to progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), 10 poliomyelitis and the post polio syndrome, and Hereditary Motorsensory Neuropathy (Charcot-Marie-Tooth Disease).

4.10.18 OTHER ACTIVITIES

A polypeptide of the invention may also exhibit one or more of the following additional 15 activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); 20 effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, co-factors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression 25 (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such 30 as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

4.10.19 IDENTIFICATION OF POLYMORPHISMS

The demonstration of polymorphisms makes possible the identification of such polymorphisms in human subjects and the pharmacogenetic use of this information for diagnosis and treatment. Such polymorphisms may be associated with, e.g., differential predisposition or susceptibility to various disease states (such as disorders involving inflammation or immune response) or a differential response to drug administration, and this genetic information can be used to tailor preventive or therapeutic treatment appropriately. For example, the existence of a polymorphism associated with a predisposition to inflammation or autoimmune disease makes possible the diagnosis of this condition in humans by identifying the presence of the polymorphism.

Polymorphisms can be identified in a variety of ways known in the art which all generally involve obtaining a sample from a patient, analyzing DNA from the sample, optionally involving isolation or amplification of the DNA, and identifying the presence of the polymorphism in the DNA. For example, PCR may be used to amplify an appropriate fragment of genomic DNA which may then be sequenced. Alternatively, the DNA may be subjected to allele-specific oligonucleotide hybridization (in which appropriate oligonucleotides are hybridized to the DNA under conditions permitting detection of a single base mismatch) or to a single nucleotide extension assay (in which an oligonucleotide that hybridizes immediately adjacent to the position of the polymorphism is extended with one or more labeled nucleotides). In addition, traditional restriction fragment length polymorphism analysis (using restriction enzymes that provide differential digestion of the genomic DNA depending on the presence or absence of the polymorphism) may be performed. Arrays with nucleotide sequences of the present invention can be used to detect polymorphisms. The array can comprise modified nucleotide sequences of the present invention in order to detect the nucleotide sequences of the present invention. In the alternative, any one of the nucleotide sequences of the present invention can be placed on the array to detect changes from those sequences.

Alternatively a polymorphism resulting in a change in the amino acid sequence could also be detected by detecting a corresponding change in amino acid sequence of the protein, e.g., by an antibody specific to the variant sequence.

4.10.20 ARTHRITIS AND INFLAMMATION

The immunosuppressive effects of the compositions of the invention against rheumatoid arthritis is determined in an experimental animal model system. The experimental model system is adjuvant induced arthritis in rats, and the protocol is described by J. Holoshitz, et al., 1983, Science, 219:56, or by B. Waksman et al., 1963, Int. Arch. Allergy Appl. Immunol., 23:129.

Induction of the disease can be caused by a single injection, generally intradermally, of a

suspension of killed *Mycobacterium tuberculosis* in complete Freund's adjuvant (CFA). The route of injection can vary, but rats may be injected at the base of the tail with an adjuvant mixture. The polypeptide is administered in phosphate buffered solution (PBS) at a dose of about 1-5 mg/kg. The control consists of administering PBS only.

5 The procedure for testing the effects of the test compound would consist of intradermally injecting killed *Mycobacterium tuberculosis* in CFA followed by immediately administering the test compound and subsequent treatment every other day until day 24. At 14, 15, 18, 20, 22, and 24 days after injection of *Mycobacterium* CFA, an overall arthritis score may be obtained as described by J. Holoskitz above. An analysis of the data would reveal that the test compound
10 would have a dramatic affect on the swelling of the joints as measured by a decrease of the arthritis score.

4.11 THERAPEUTIC METHODS

15 The compositions (including polypeptide fragments, analogs, variants and antibodies or other binding partners or modulators including antisense polynucleotides) of the invention have numerous applications in a variety of therapeutic methods. Examples of therapeutic applications include, but are not limited to, those exemplified herein.

4.11.1 EXAMPLE

20 One embodiment of the invention is the administration of an effective amount of the polypeptides or other composition of the invention to individuals affected by a disease or disorder that can be modulated by regulating the peptides of the invention. While the mode of administration is not particularly important, parenteral administration is preferred. An exemplary mode of administration is to deliver an intravenous bolus. The dosage of the
25 polypeptides or other composition of the invention will normally be determined by the prescribing physician. It is to be expected that the dosage will vary according to the age, weight, condition and response of the individual patient. Typically, the amount of polypeptide administered per dose will be in the range of about 0.01 µg/kg to 100 mg/kg of body weight, with the preferred dose being about 0.1 µg/kg to 10 mg/kg of patient body weight. For parenteral
30 administration, polypeptides of the invention will be formulated in an injectable form combined with a pharmaceutically acceptable parenteral vehicle. Such vehicles are well known in the art and examples include water, saline, Ringer's solution, dextrose solution, and solutions consisting of small amounts of the human serum albumin. The vehicle may contain minor amounts of additives that maintain the isotonicity and stability of the polypeptide or other active ingredient.
35 The preparation of such solutions is within the skill of the art.

4.12 PHARMACEUTICAL FORMULATIONS AND ROUTES OF ADMINISTRATION

A protein or other composition of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources and including antibodies and other binding partners of the polypeptides of the invention) may be administered to a patient in need, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a variety of disorders. Such a composition may optionally contain (in addition to protein or other active ingredient and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the disease or disorder in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet-derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), insulin-like growth factor (IGF), as well as cytokines described herein.

The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or other active ingredient or complement its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein or other active ingredient of the invention, or to minimize side effects. Conversely, protein or other active ingredient of the present invention may be included in formulations of the particular clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent (such as IL-1Ra, IL-1 Hy1, IL-1 Hy2, anti-TNF, corticosteroids, immunosuppressive agents). A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

As an alternative to being included in a pharmaceutical composition of the invention including a first protein, a second protein or a therapeutic agent may be concurrently administered with the first protein (e.g., at the same time, or at differing times provided that therapeutic concentrations of the combination of agents is achieved at the treatment site).

- 5 Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms, *e.g.*, treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or
- 10 amelioration of such conditions. When applied to an individual active ingredient, administered alone, a therapeutically effective dose refers to that ingredient alone. When applied to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.
- 15 In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein or other active ingredient of the present invention is administered to a mammal having a condition to be treated. Protein or other active ingredient of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other
- 20 hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein or other active ingredient of the present invention may be administered ~~either simultaneously~~ with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein or other
- 25 active ingredient of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

4.12.1 ROUTES OF ADMINISTRATION

- Suitable routes of administration may, for example, include oral, rectal, transmucosal, or
- 30 intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections. Administration of protein or other active ingredient of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral

ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a arthritic joints or in fibrotic tissue, often in a depot or sustained release formulation. In order to prevent the scarring process frequently occurring as complication of glaucoma surgery, the compounds may be administered topically, for example, as eye drops. Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with a specific antibody, targeting, for example, arthritic or fibrotic tissue. The liposomes will be targeted to and taken up selectively by the afflicted tissue.

The polypeptides of the invention are administered by any route that delivers an effective dosage to the desired site of action. The determination of a suitable route of administration and an effective dosage for a particular indication is within the level of skill in the art. Preferably for wound treatment, one administers the therapeutic compound directly to the site. Suitable dosage ranges for the polypeptides of the invention can be extrapolated from these dosages or from similar studies in appropriate animal models. Dosages can then be adjusted as necessary by the clinician to provide maximal therapeutic benefit.

4.12.2 COMPOSITIONS/FORMULATIONS

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. These pharmaceutical compositions may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of protein or other active ingredient of the present invention is administered orally, protein or other active ingredient of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein or other active ingredient of the present invention, and preferably from about 25 to 90% protein or other active ingredient of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the

pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein or other active ingredient of the present invention, and preferably from about 1 to 50% protein or other active ingredient of the present invention.

When a therapeutically effective amount of protein or other active ingredient of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein or other active ingredient of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein or other active ingredient solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein or other active ingredient of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art. For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained from a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic,

talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

5 Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in
10 suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration. For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

 For administration by inhalation, the compounds for use according to the present
15 invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in
20 an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch. The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for ~~the~~ injection may be presented in unit dosage form, *e.g.*, in ampules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or
25 emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

 Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or
30 vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated

solutions. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides. In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a co-solvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The co-solvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics. Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, *e.g.* polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose. Alternatively, other delivery systems for hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various types of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein or other active ingredient stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium

carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols. Many of the active ingredients of the invention may be provided as salts with pharmaceutically compatible counter ions. Such pharmaceutically acceptable base addition salts are those salts which retain the biological effectiveness and properties of the free acids and which are obtained by reaction with inorganic or organic bases such as sodium hydroxide, magnesium hydroxide, ammonia, trialkylamine, dialkylamine, monoalkylamine, dibasic amino acids, sodium acetate, potassium benzoate, triethanol amine and the like.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) or other active ingredient(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithins, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent Nos. 4,235,871; 4,501,728; 4,837,028; and 4,737,323, all of which are incorporated herein by reference.

The amount of protein or other active ingredient of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein or other active ingredient of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein or other active ingredient of the present

invention and observe the patient's response. Larger doses of protein or other active ingredient of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should
5 contain about 0.01 μg to about 100 mg (preferably about 0.1 μg to about 10 mg, more preferably about 0.1 μg to about 1 mg) of protein or other active ingredient of the present invention per kg body weight. For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic
10 composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein or other active ingredient of the invention which may also optionally be included in the composition as
15 described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing or other active ingredient-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally
20 capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions
25 may be biodegradable and chemically defined calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass,
30 aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalcium phosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability. Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and
35 glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns.

In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions from disassociating from the matrix.

- A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol).
- The amount of sequestering agent useful herein is 0.5-20 wt %, preferably 1-10 wt % based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells. In further compositions, proteins or other active ingredients of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- α and TGF- β), and insulin-like growth factor (IGF).
- The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins or other active ingredients of the present invention. The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, *e.g.*, amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (*e.g.*, bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

- Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either in vivo or ex vivo into cells for expression in a

mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or
5 activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

4.12.3 EFFECTIVE DOSAGE

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its
10 intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from
15 appropriate *in vitro* assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that can be used to more accurately determine useful doses in humans. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of the protein's biological activity).
20 Such information can be used to more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD_{50} (the dose lethal to 50% of the
25 population) and the ED_{50} (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD_{50} and ED_{50} . Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range
30 of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. See, *e.g.*, Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1. Dosage amount and interval may be adjusted
35 individually to provide plasma levels of the active moiety which are sufficient to maintain the

desired effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

5 Dosage intervals can also be determined using MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

10 An exemplary dosage regimen for polypeptides or other compositions of the invention will be in the range of about 0.01 µg/kg to 100 mg/kg of body weight daily, with the preferred dose being about 0.1 µg/kg to 25 mg/kg of patient body weight daily, varying in adults and children. Dosing may be once daily, or equivalent doses may be delivered at longer or shorter intervals.

15 The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's age and weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

4.12.4 PACKAGING

20 The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an
25 appropriate container, and labeled for treatment of an indicated condition.

4.13 ANTIBODIES

Also included in the invention are antibodies to proteins, or fragments of proteins of the invention. The term "antibody" as used herein refers to immunoglobulin molecules and
30 immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F_{ab}, F_{ab}' and F_{(ab)2} fragments, and an F_{ab} expression library. In general, an antibody molecule obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another
35 by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well,

such as IgG₁, IgG₂, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.

An isolated related protein of the invention may be intended to serve as an antigen, or a portion or fragment thereof, and additionally can be used as an immunogen to generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein, such as an amino acid sequence shown in SEQ ID NO:985, and encompasses an epitope thereof such that an antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of -related protein that is located on the surface of the protein, *e.g.*, a hydrophilic region. A hydrophobicity analysis of the human related protein sequence will indicate which regions of a related protein are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody production. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier transformation. See, *e.g.*, Hopp and Woods, 1981, *Proc. Nat. Acad. Sci. USA* 78: 3824-3828; Kyte and Doolittle 1982, *J. Mol. Biol.* 157: 105-142, each of which is incorporated herein by reference in its entirety. Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

A protein of the invention, or a derivative, fragment, analog, homolog or ortholog thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind these protein components.

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives, fragments, analogs homologs or orthologs thereof (see, for example, *Antibodies: A Laboratory*

Manual, Harlow E, and Lane D, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Some of these antibodies are discussed below.

5.13.1 Polyclonal Antibodies

5 For the production of polyclonal antibodies, various suitable host animals (e.g., rabbit, goat, mouse or other mammal) may be immunized by one or more injections with the native protein, a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, the naturally occurring immunogenic protein, a chemically synthesized polypeptide representing the immunogenic protein, or a
10 recombinantly expressed immunogenic protein. Furthermore, the protein may be conjugated to a second protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not
15 limited to, Freund's (complete and incomplete), mineral gels (e.g., aluminum hydroxide), surface active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), adjuvants usable in humans such as Bacille Calmette-Guerin and *Corynebacterium parvum*, or similar immunostimulatory agents. Additional examples of adjuvants which can be employed include MPL-TDM adjuvant (monophosphoryl Lipid A,
20 synthetic trehalose dicorynomycolate).

The polyclonal antibody molecules directed against the immunogenic protein can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as affinity chromatography using protein A or protein G, which provide primarily the IgG fraction of immune serum. Subsequently, or alternatively, the specific antigen which is the
25 target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17, 2000), pp. 25-28).

30 5.13.2 Monoclonal Antibodies

The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal
35 antibody are identical in all the molecules of the population. MAbs thus contain an antigen

binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes can be immunized in vitro. The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63).

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the

Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). Preferably, antibodies having a high degree of specificity and a high binding affinity for the target antigen are isolated.

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium.

Alternatively, the hybridoma cells can be grown in vivo as ascites in a mammal.

The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, Nature 368: 812-13 (1994)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

5.13.2 Humanized Antibodies

The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. These antibodies are suitable for administration to humans without engendering an immune response by the human against the administered immunoglobulin. Humanized forms of antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that are principally comprised of the sequence of a human

immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin. Humanization can be performed following the method of Winter and co-workers (Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeven et al., Science, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the
5 corresponding sequences of a human antibody. (See also U.S. Patent No. 5,225,539.) In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies can also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable
10 domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin (Jones et al., 1986; Riechmann et al., 1988; and Presta, Curr. Op. Struct. Biol.,
15 2:593-596 (1992)).

5.13.3 Human Antibodies

Fully human antibodies relate to antibody molecules in which essentially the entire sequences of both the light chain and the heavy chain, including the CDRs, arise from human
20 genes. Such antibodies are termed "human antibodies", or "fully human antibodies" herein. Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique (see Kozbor, et al., 1983 Lancet Today 4: 72) and the EBV hybridoma
technique to produce human monoclonal antibodies (see Cole, et al., 1985 In: MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96). Human monoclonal
25 antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cote, et al., 1983. Proc Natl Acad Sci USA 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus in vitro (see Cole, et al., 1985 In: MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96).

In addition, human antibodies can also be produced using additional techniques,
30 including phage display libraries (Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon
challenge, human antibody production is observed, which closely resembles that seen in humans
35 in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach

is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in Marks et al. (Bio/Technology 10, 779-783 (1992)); Lonberg et al. (Nature 368 856-859 (1994)); Morrison (Nature 368, 812-13 (1994)); Fishwild et al. (Nature Biotechnology 14, 845-51 (1996)); Neuberger (Nature Biotechnology 14, 826 (1996)); and
5 Lonberg and Huszar (Intern. Rev. Immunol. 13 65-93 (1995)).

Human antibodies may additionally be produced using transgenic nonhuman animals which are modified so as to produce fully human antibodies rather than the animal's endogenous antibodies in response to challenge by an antigen. (See PCT publication WO94/02602). The endogenous genes encoding the heavy and light immunoglobulin chains in the nonhuman host
10 have been incapacitated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. The
15 preferred embodiment of such a nonhuman animal is a mouse, and is termed the XenomouseTM as disclosed in PCT publications WO 96/33735 and WO 96/34096. This animal produces B cells which secrete fully human immunoglobulins. The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells derived from the animal, such as
20 hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly, or can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

An example of a method of producing a nonhuman host, exemplified as a mouse, lacking
25 expression of an endogenous immunoglobulin heavy chain is disclosed in U.S. Patent No. 5,939,598. It can be obtained by a method including deleting the J segment genes from at least one endogenous heavy chain locus in an embryonic stem cell to prevent rearrangement of the locus and to prevent formation of a transcript of a rearranged immunoglobulin heavy chain locus, the deletion being effected by a targeting vector containing a gene encoding a selectable marker; and producing from the embryonic stem cell a transgenic mouse whose somatic and germ cells
30 contain the gene encoding the selectable marker.

A method for producing an antibody of interest, such as a human antibody, is disclosed in U.S. Patent No. 5,916,771. It includes introducing an expression vector that contains a nucleotide sequence encoding a heavy chain into one mammalian host cell in culture, introducing
35 an expression vector containing a nucleotide sequence encoding a light chain into another

mammalian host cell, and fusing the two cells to form a hybrid cell. The hybrid cell expresses an antibody containing the heavy chain and the light chain.

In a further improvement on this procedure, a method for identifying a clinically relevant epitope on an immunogen, and a correlative method for selecting an antibody that binds immunospecifically to the relevant epitope with high affinity, are disclosed in PCT publication WO 99/53049.

5.13.4 F_{ab} Fragments and Single Chain Antibodies

According to the invention, techniques can be adapted for the production of single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of F_{ab} expression libraries (see e.g., Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective identification of monoclonal F_{ab} fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the idiotypes to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F_{(ab)₂} fragment produced by pepsin digestion of an antibody molecule; (ii) an F_{ab} fragment generated by reducing the disulfide bridges of an F_{(ab)₂} fragment; (iii) an F_{ab} fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F_v fragments.

5.13.5 Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for an antigenic protein of the invention. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, *Nature*, 305:537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, 1991 *EMBO J.*, 10:3655-3659.

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion

preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to dinitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally, Fab' fragments can be directly recovered from E. coli and chemically coupled to form bispecific antibodies. Shalaby et al., J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from E. coli and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., J. Immunol. 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., Proc. Natl. Acad. Sci. USA 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., J. Immunol. 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., J. Immunol. 147:60 (1991). Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in the protein antigen of the invention. Alternatively, an anti-antigenic arm of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific antibodies can also be used to direct cytotoxic agents to cells which express a particular antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the protein antigen described herein and further binds tissue factor (TF).

5.13.6 Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/200373; EP 03089). It is contemplated that the antibodies can be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins

can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

5 5.13.7 Effector Function Engineering

It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved
10 internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., J. Exp Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al. Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that
15 has dual Fc regions and can thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., Anti-Cancer Drug Design, 3: 219-230 (1989).

5.13.8 Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a
20 cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radiocoujugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include
25 diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolacca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of
30 radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL),
35 active esters (such as disuccinimidyl suberate), aldehydes (such as glutarelddehyde), bis-azido

compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science, 238: 1098 (1987).

- 5 Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is
10 administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn conjugated to a cytotoxic agent.

4.14 COMPUTER READABLE SEQUENCES

- 15 In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media" refers to any medium which can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM
20 and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer readable mediums can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide sequence of the present invention. As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the
25 presently known methods for recording information on computer readable medium to generate manufactures comprising the nucleotide sequence information of the present invention.

A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen
30 to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase,
35 Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring

formats (e.g. text file or database) in order to obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing any of the nucleotide sequences SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 or a representative fragment thereof; or a nucleotide sequence at least 95% identical to any of the nucleotide sequences of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 in computer readable form, a skilled artisan can routinely access the sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium. The examples which follow demonstrate how software which implements the BLAST (Altschul et al., J. Mol. Biol. 215:403-410 (1990)) and BLAZE (Brutlag et al., Comp. Chem. 17:203-207 (1993)) search algorithms on a Sybase system is used to identify open reading frames (ORFs) within a nucleic acid sequence. Such ORFs may be protein encoding fragments and may be useful in producing commercially important proteins such as enzymes used in fermentation reactions and in the production of commercially useful metabolites.

As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are suitable for use in the present invention. As stated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory which can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention.

As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of a known sequence which match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, Smith-Waterman, MacPattern (EMBL), BLASTN and BLASTA (NPOLYPEPTIDEIA). A skilled artisan can readily recognize that any one of the available algorithms or implementing

software packages for conducting homology searches can be adapted for use in the present computer-based systems. As used herein, a "target sequence" can be any nucleic acid or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 300 amino acids, more preferably from about 30 to 100 nucleotide residues. However, it is well recognized that searches for commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are chosen based on a three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzyme active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures and inducible expression elements (protein binding sequences).

4.15 TRIPLE HELIX FORMATION

In addition, the fragments of the present invention, as broadly described, can be used to control gene expression through triple helix formation or antisense DNA or RNA, both of which methods are based on the binding of a polynucleotide sequence to DNA or RNA. Polynucleotides suitable for use in these methods are preferably 20 to 40 bases in length and are designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 152:41:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Olmno, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide.

4.16 DIAGNOSTIC ASSAYS AND KITS

The present invention further provides methods to identify the presence or expression of one of the ORFs of the present invention, or homolog thereof, in a test sample, using a nucleic

acid probe or antibodies of the present invention, optionally conjugated or otherwise associated with a suitable label.

In general, methods for detecting a polynucleotide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polynucleotide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polynucleotide of the invention is detected in the sample. Such methods can also comprise contacting a sample under stringent hybridization conditions with nucleic acid primers that anneal to a polynucleotide of the invention under such conditions, and amplifying annealed polynucleotides, so that if a polynucleotide is amplified, a polynucleotide of the invention is detected in the sample.

In general, methods for detecting a polypeptide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polypeptide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polypeptide of the invention is detected in the sample.

In detail, such methods comprise incubating a test sample with one or more of the antibodies or one or more of the nucleic acid probes of the present invention and assaying for binding of the nucleic acid probes or antibodies to components within the test sample.

Conditions for incubating a nucleic acid probe or antibody with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid probe or antibody used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or immunological assay formats can readily be adapted to employ the nucleic acid probes or antibodies of the present invention. Examples of such assays can be found in Chard, T., An Introduction to Radioimmunoassay and Related Techniques, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G.R. et al., Techniques in Immunocytochemistry, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., Practice and Theory of immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1985). The test samples of the present invention include cells, protein or membrane extracts of cells, or biological fluids such as sputum, blood, serum, plasma, or urine. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention. Specifically, the invention provides a compartment kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the probes or antibodies of the present invention; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound probe or antibody.

In detail, a compartment kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers or strips of plastic or paper. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the antibodies used in the assay, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound antibody or probe. Types of detection reagents include labeled nucleic acid probes, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. One skilled in the art will readily recognize that the disclosed probes and antibodies of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

MEDICAL IMAGING

The novel polypeptides and binding partners of the invention are useful in medical imaging of sites expressing the molecules of the invention (e.g., where the polypeptide of the invention is involved in the immune response, for imaging sites of inflammation or infection). See, e.g., Kunkel et al., U.S. Pat. NO. 5,413,778. Such methods involve chemical attachment of a labeling or imaging agent, administration of the labeled polypeptide to a subject in a pharmaceutically acceptable carrier, and imaging the labeled polypeptide *in vivo* at the target site.

30

4.18 SCREENING ASSAYS

Using the isolated proteins and polynucleotides of the invention, the present invention further provides methods of obtaining and identifying agents which bind to a polypeptide encoded by an ORF corresponding to any of the nucleotide sequences set forth in SEQ ID NO:

1-984, 1969-2952, 3937-3942 or 3949-3954, or bind to a specific domain of the polypeptide encoded by the nucleic acid. In detail, said method comprises the steps of:

- (a) contacting an agent with an isolated protein encoded by an ORF of the present invention, or nucleic acid of the invention; and
- 5 (b) determining whether the agent binds to said protein or said nucleic acid.

In general, therefore, such methods for identifying compounds that bind to a polynucleotide of the invention can comprise contacting a compound with a polynucleotide of the invention for a time sufficient to form a polynucleotide/compound complex, and detecting the complex, so that if a polynucleotide/compound complex is detected, a compound that binds
10 to a polynucleotide of the invention is identified.

Likewise, in general, therefore, such methods for identifying compounds that bind to a polypeptide of the invention can comprise contacting a compound with a polypeptide of the invention for a time sufficient to form a polypeptide/compound complex, and detecting the complex, so that if a polypeptide/compound complex is detected, a compound that binds to a
15 polynucleotide of the invention is identified.

Methods for identifying compounds that bind to a polypeptide of the invention can also comprise contacting a compound with a polypeptide of the invention in a cell for a time sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a receptor gene sequence in the cell, and detecting the complex by detecting reporter gene
20 sequence expression, so that if a polypeptide/compound complex is detected, a compound that binds a polypeptide of the invention is identified.

Compounds identified via such methods can include compounds which modulate the activity of a polypeptide of the invention (that is, increase or decrease its activity, relative to activity observed in the absence of the compound). Alternatively, compounds identified via such
25 methods can include compounds which modulate the expression of a polynucleotide of the invention (that is, increase or decrease expression relative to expression levels observed in the absence of the compound). Compounds, such as compounds identified via the methods of the invention, can be tested using standard assays well known to those of skill in the art for their ability to modulate activity/expression.

30 The agents screened in the above assay can be, but are not limited to, peptides, carbohydrates, vitamin derivatives, or other pharmaceutical agents. The agents can be selected and screened at random or rationally selected or designed using protein modeling techniques.

For random screening, agents such as peptides, carbohydrates, pharmaceutical agents and the like are selected at random and are assayed for their ability to bind to the protein encoded by
35 the ORF of the present invention. Alternatively, agents may be rationally selected or designed.

As used herein, an agent is said to be "rationally selected or designed" when the agent is chosen based on the configuration of the particular protein. For example, one skilled in the art can readily adapt currently available procedures to generate peptides, pharmaceutical agents and the like, capable of binding to a specific peptide sequence, in order to generate rationally designed antipeptide peptides, for example see Hurby et al., Application of Synthetic Peptides: Antisense Peptides," In Synthetic Peptides, A User's Guide, W.H. Freeman, NY (1992), pp. 289-307, and Kaspiczak et al., Biochemistry 28:9230-8 (1989), or pharmaceutical agents, or the like.

In addition to the foregoing, one class of agents of the present invention, as broadly described, can be used to control gene expression through binding to one of the ORFs or EMFs of the present invention. As described above, such agents can be randomly screened or rationally designed/selected. Targeting the ORF or EMF allows a skilled artisan to design sequence specific or element specific agents, modulating the expression of either a single ORF or multiple ORFs which rely on the same EMF for expression control. One class of DNA binding agents are agents which contain base residues which hybridize or form a triple helix formation by binding to DNA or RNA. Such agents can be based on the classic phosphodiester, ribonucleic acid backbone, or can be a variety of sulfhydryl or polymeric derivatives which have base attachment capacity.

Agents suitable for use in these methods preferably contain 20 to 40 bases and are designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide and other DNA binding agents.

Agents which bind to a protein encoded by one of the ORFs of the present invention can be used as a diagnostic agent. Agents which bind to a protein encoded by one of the ORFs of the present invention can be formulated using known techniques to generate a pharmaceutical composition.

4.19 USE OF NUCLEIC ACIDS AS PROBES

Another aspect of the subject invention is to provide for polypeptide-specific nucleic acid hybridization probes capable of hybridizing with naturally occurring nucleotide sequences. The

hybridization probes of the subject invention may be derived from any of the nucleotide sequences SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954. Because the corresponding gene is only expressed in a limited number of tissues, a hybridization probe derived from any of the nucleotide sequences SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954 can be used as an indicator of the presence of RNA of cell type of such a tissue in a sample.

Any suitable hybridization technique can be employed, such as, for example, *in situ* hybridization. PCR as described in US Patents Nos. 4,683,195 and 4,965,188 provides additional uses for oligonucleotides based upon the nucleotide sequences. Such probes used in PCR may be of recombinant origin, may be chemically synthesized, or a mixture of both. The probe will comprise a discrete nucleotide sequence for the detection of identical sequences or a degenerate pool of possible sequences for identification of closely related genomic sequences.

Other means for producing specific hybridization probes for nucleic acids include the cloning of nucleic acid sequences into vectors for the production of mRNA probes. Such vectors are known in the art and are commercially available and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate radioactively labeled nucleotides. The nucleotide sequences may be used to construct hybridization probes for mapping their respective genomic sequences. The nucleotide sequence provided herein may be mapped to a chromosome or specific regions of a chromosome using well known genetic and/or chromosomal mapping techniques. These techniques include *in situ* hybridization, linkage analysis against known chromosomal markers, hybridization screening with libraries or flow-sorted chromosomal preparations specific to known chromosomes, and the like. The technique of fluorescent *in situ* hybridization of chromosome spreads has been described, among other places, in Verma et al (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York NY.

Fluorescent *in situ* hybridization of chromosomal preparations and other physical chromosome mapping techniques may be correlated with additional genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of a nucleic acid on a physical chromosomal map and a specific disease (or predisposition to a specific disease) may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier or affected individuals.

4.20 PREPARATION OF SUPPORT BOUND OLIGONUCLEOTIDES

Oligonucleotides, i.e., small nucleic acid segments, may be readily prepared by, for example, directly synthesizing the oligonucleotide by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer.

- 5 Support bound oligonucleotides may be prepared by any of the methods known to those of skill in the art using any suitable support such as glass, polystyrene or Teflon. One strategy is to precisely spot oligonucleotides synthesized by standard synthesizers. Immobilization can be achieved using passive adsorption (Inouye & Hondo, (1990) J. Clin. Microbiol. 28(6) 1469-72); using UV light (Nagata *et al.*, 1985; Dahlen *et al.*, 1987; Morrissey & Collins, (1989) Mol. Cell
10 Probes 3(2) 189-207) or by covalent binding of base modified DNA (Keller *et al.*, 1988; 1989); all references being specifically incorporated herein.

- Another strategy that may be employed is the use of the strong biotin-streptavidin interaction as a linker. For example, Broude *et al.* (1994) Proc. Natl. Acad. Sci. USA 91(8) 3072-6, describe the use of biotinylated probes, although these are duplex probes, that are immobilized on
15 streptavidin-coated magnetic beads. Streptavidin-coated beads may be purchased from Dynal, Oslo. Of course, this same linking chemistry is applicable to coating any surface with streptavidin. Biotinylated probes may be purchased from various sources, such as, e.g., Operon Technologies (Alameda, CA).

- Nunc Laboratories (Naperville, IL) is also selling suitable material that could be used. Nunc
20 Laboratories have developed a method by which DNA can be covalently bound to the microwell surface termed CovaLink NH. CovaLink NH is a polystyrene surface grafted with secondary amino groups (>NH) that serve as bridge-heads for further covalent coupling. CovaLink Modules may be purchased from Nunc Laboratories. DNA molecules may be bound to CovaLink exclusively at the 5'-end by a phosphoramidate bond, allowing immobilization of more than 1 pmol of DNA
25 (Rasmussen *et al.*, (1991) Anal. Biochem. 198(1) 138-42).

- The use of CovaLink NH strips for covalent binding of DNA molecules at the 5'-end has been described (Rasmussen *et al.*, (1991). In this technology, a phosphoramidate bond is employed (Chu *et al.*, (1983) Nucleic Acids Res. 11(8) 6513-29). This is beneficial as immobilization using only a single covalent bond is preferred. The phosphoramidate bond joins the DNA to the
30 CovaLink NH secondary amino groups that are positioned at the end of spacer arms covalently grafted onto the polystyrene surface through a 2 nm long spacer arm. To link an oligonucleotide to CovaLink NH via an phosphoramidate bond, the oligonucleotide terminus must have a 5'-end phosphate group. It is, perhaps, even possible for biotin to be covalently bound to CovaLink and then streptavidin used to bind the probes.

More specifically, the linkage method includes dissolving DNA in water (7.5 ng/ul) and denaturing for 10 min. at 95°C and cooling on ice for 10 min. Ice-cold 0.1 M 1-methylimidazole, pH 7.0 (1-MeIm₇), is then added to a final concentration of 10 mM 1-MeIm₇. A ss DNA solution is then dispensed into Covalink NH strips (75 ul/well) standing on ice.

5 Carbodiimide 0.2 M 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC), dissolved in 10 mM 1-MeIm₇, is made fresh and 25 ul added per well. The strips are incubated for 5 hours at 50°C. After incubation the strips are washed using, e.g., Nunc-Immuno Wash; first the wells are washed 3 times, then they are soaked with washing solution for 5 min., and finally they are washed 3 times (where in the washing solution is 0.4 N NaOH, 0.25% SDS heated to 50°C).

10 It is contemplated that a further suitable method for use with the present invention is that described in PCT Patent Application WO 90/03382 (Southern & Maskos), incorporated herein by reference. This method of preparing an oligonucleotide bound to a support involves attaching a nucleoside 3'-reagent through the phosphate group by a covalent phosphodiester link to aliphatic hydroxyl groups carried by the support. The oligonucleotide is then synthesized on the supported
15 nucleoside and protecting groups removed from the synthetic oligonucleotide chain under standard conditions that do not cleave the oligonucleotide from the support. Suitable reagents include nucleoside phosphoramidite and nucleoside hydrogen phosphate.

An on-chip strategy for the preparation of DNA probe for the preparation of DNA probe arrays may be employed. For example, addressable laser-activated photodeprotection may be
20 employed in the chemical synthesis of oligonucleotides directly on a glass surface, as described by Fodor *et al.* (1991) Science 251(4995) 767-73, incorporated herein by reference. Probes may also be immobilized on nylon supports as described by Van Ness *et al.* (1991) Nucleic Acids Res. 19(12) 3345-50; or linked to Teflon using the method of Duncan & Cavalier (1988) Anal. Biochem. 169(1) 104-8; all references being specifically incorporated herein.

25 To link an oligonucleotide to a nylon support, as described by Van Ness *et al.* (1991), requires activation of the nylon surface via alkylation and selective activation of the 5'-amine of oligonucleotides with cyanuric chloride.

One particular way to prepare support bound oligonucleotides is to utilize the light-generated synthesis described by Pease *et al.*, (1994) PNAS USA 91(11) 5022-6, incorporated
30 herein by reference). These authors used current photolithographic techniques to generate arrays of immobilized oligonucleotide probes (DNA chips). These methods, in which light is used to direct the synthesis of oligonucleotide probes in high-density, miniaturized arrays, utilize photolabile 5'-protected *N*-acyl-deoxynucleoside phosphoramidites, surface linker chemistry and versatile combinatorial synthesis strategies. A matrix of 256 spatially defined oligonucleotide probes may be
35 generated in this manner.

4.21 PREPARATION OF NUCLEIC ACID FRAGMENTS

The nucleic acids may be obtained from any appropriate source, such as cDNAs, genomic DNA, chromosomal DNA, microdissected chromosome bands, cosmid or YAC inserts, and RNA, including mRNA without any amplification steps. For example, Sambrook *et al.* (1989) describes three protocols for the isolation of high molecular weight DNA from mammalian cells (p. 9.14-9.23).

DNA fragments may be prepared as clones in M13, plasmid or lambda vectors and/or prepared directly from genomic DNA or cDNA by PCR or other amplification methods. Samples may be prepared or dispensed in multiwell plates. About 100-1000 ng of DNA samples may be prepared in 2-500 ml of final volume.

The nucleic acids would then be fragmented by any of the methods known to those of skill in the art including, for example, using restriction enzymes as described at 9.24-9.28 of Sambrook *et al.* (1989), shearing by ultrasound and NaOH treatment.

Low pressure shearing is also appropriate, as described by Schriefer *et al.* (1990) Nucleic Acids Res. 18(24) 7455-6, incorporated herein by reference). In this method, DNA samples are passed through a small French pressure cell at a variety of low to intermediate pressures. A lever device allows controlled application of low to intermediate pressures to the cell. The results of these studies indicate that low-pressure shearing is a useful alternative to sonic and enzymatic DNA fragmentation methods.

One particularly suitable way for fragmenting DNA is contemplated to be that using the two base recognition endonuclease, *Cvi*JI, described by Fitzgerald *et al.* (1992) Nucleic Acids Res. 20(14) 3753-62. These authors described an approach for the rapid fragmentation and fractionation of DNA into particular sizes that they contemplated to be suitable for shotgun cloning and sequencing.

The restriction endonuclease *Cvi*JI normally cleaves the recognition sequence PuGCPy between the G and C to leave blunt ends. Atypical reaction conditions, which alter the specificity of this enzyme (*Cvi*JI**), yield a quasi-random distribution of DNA fragments from the small molecule pUC19 (2688 base pairs). Fitzgerald *et al.* (1992) quantitatively evaluated the randomness of this fragmentation strategy, using a *Cvi*JI** digest of pUC19 that was size fractionated by a rapid gel filtration method and directly ligated, without end repair, to a lac Z minus M13 cloning vector. Sequence analysis of 76 clones showed that *Cvi*JI** restricts pyGCPy and PuGCPu, in addition to PuGCPy sites, and that new sequence data is accumulated at a rate consistent with random fragmentation.

As reported in the literature, advantages of this approach compared to sonication and agarose gel fractionation include: smaller amounts of DNA are required (0.2-0.5 ug instead of 2-5

ug); and fewer steps are involved (no preligation, end repair, chemical extraction, or agarose gel electrophoresis and elution are needed)

Irrespective of the manner in which the nucleic acid fragments are obtained or prepared, it is important to denature the DNA to give single stranded pieces available for hybridization. This is achieved by incubating the DNA solution for 2-5 minutes at 80-90°C. The solution is then cooled quickly to 2°C to prevent renaturation of the DNA fragments before they are contacted with the chip. Phosphate groups must also be removed from genomic DNA by methods known in the art.

4.22 PREPARATION OF DNA ARRAYS

Arrays may be prepared by spotting DNA samples on a support such as a nylon membrane.

Spotting may be performed by using arrays of metal pins (the positions of which correspond to an array of wells in a microtiter plate) to repeated by transfer of about 20 nl of a DNA solution to a nylon membrane. By offset printing, a density of dots higher than the density of the wells is achieved. One to 25 dots may be accommodated in 1 mm², depending on the type of label used. By avoiding spotting in some preselected number of rows and columns, separate subsets (subarrays) may be formed. Samples in one subarray may be the same genomic segment of DNA (or the same gene) from different individuals, or may be different, overlapped genomic clones. Each of the subarrays may represent replica spotting of the same samples. In one example, a selected gene segment may be amplified from 64 patients. For each patient, the amplified gene segment may be in one 96-well plate (all 96 wells containing the same sample). A plate for each of the 64 patients is prepared. By using a 96-pin device, all samples may be spotted on one 8 x 12 cm membrane. Subarrays may contain 64 samples, one from each patient. Where the 96 subarrays are identical, the dot span may be 1 mm² and there may be a 1 mm space between subarrays.

Another approach is to use membranes or plates (available from NUNC, Naperville, Illinois) which may be partitioned by physical spacers e.g. a plastic grid molded over the membrane, the grid being similar to the sort of membrane applied to the bottom of multiwell plates, or hydrophobic strips. A fixed physical spacer is not preferred for imaging by exposure to flat phosphor-storage screens or x-ray films.

The present invention is illustrated in the following examples. Upon consideration of the present disclosure, one of skill in the art will appreciate that many other embodiments and variations may be made in the scope of the present invention. Accordingly, it is intended that the broader aspects of the present invention not be limited to the disclosure of the following examples. The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and compositions and methods which are functionally equivalent are within the scope of the invention. Indeed, numerous modifications and

variations in the practice of the invention are expected to occur to those skilled in the art upon consideration of the present preferred embodiments. Consequently, the only limitations which should be placed upon the scope of the invention are those which appear in the appended claims.

All references cited within the body of the instant specification are hereby incorporated by
5 reference in their entirety.

5.0 EXAMPLES

5.1 EXAMPLE 1

Novel Nucleic Acid Sequences Obtained From Various Libraries

A plurality of novel nucleic acids were obtained from cDNA libraries prepared from various
10 human tissues and in some cases isolated from a genomic library derived from human chromosome using standard PCR, SBH sequence signature analysis and Sanger sequencing techniques. The inserts of the library were amplified with PCR using primers specific for the vector sequences which flank the inserts. Clones from cDNA libraries were spotted on nylon membrane filters and screened with oligonucleotide probes (e.g., 7-mers) to obtain signature sequences. The clones were clustered
15 into groups of similar or identical sequences. Representative clones were selected for sequencing.

In some cases, the 5' sequence of the amplified inserts was then deduced using a typical Sanger sequencing protocol. PCR products were purified and subjected to fluorescent dye terminator cycle sequencing. Single pass gel sequencing was done using a 377 Applied Biosystems (ABI) sequencer to obtain the novel nucleic acid sequences. In some cases RACE (Random
20 Amplification of cDNA Ends) was performed to further extend the sequence in the 5' direction.

5.2 EXAMPLE 2

Assemblage of Novel Nucleic Acids

The contigs or nucleic acids of the present invention, designated as SEQ ID NO: 1969-2951, and 3949-3954 were assembled using an EST sequence as a seed. Then a recursive algorithm was
25 used to extend the seed EST into an extended assemblage, by pulling additional sequences from different databases (i.e., Hyseq's database containing EST sequences, dbEST version 114, gb pri 114, and UniGene version 101) that belong to this assemblage. The algorithm terminated when there was no additional sequences from the above databases that would extend the assemblage. Inclusion of component sequences into the assemblage was based on a BLASTN hit to the
30 extending assemblage with BLAST score greater than 300 and percent identity greater than 95%.

Tables 6 and 8 sets forth the novel predicted polypeptides (including proteins) encoded by the novel polynucleotides (SEQ ID NO:2953-3936, and 3949-3954) of the present invention, and their corresponding nucleotide locations to each of SEQ ID NO: 2953-3936 and 3955-3960. Tables

6 and 8 also indicates the method by which the polypeptide was predicted. Method A refers to a polypeptide obtained by using a software program called FASTY (available from <http://fasta.bioch.virginia.edu>) which selects a polypeptide based on a comparison of the translated novel polynucleotide to known polynucleotides (W.R. Pearson, Methods in Enzymology, 183:63-98 (1990), herein incorporated by reference). Method B refers to a polypeptide obtained by using a software program called GenScan for human/vertebrate sequences (available from Stanford University, Office of Technology Licensing) that predicts the polypeptide based on a probabilistic model of gene structure/compositional properties (C. Burge and S. Karlin, J. Mol. Biol., 268:78-94 (1997), incorporated herein by reference). Method C refers to a polypeptide obtained by using a Hyseq proprietary software program that translates the novel polynucleotide and its complementary strand into six possible amino acid sequences (forward and reverse frames) and chooses the polypeptide with the longest open reading frame.

5.3 EXAMPLE 3

Novel Nucleic Acids

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), full length gene cDNA sequences and their corresponding protein sequences were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank. Other computer programs which may have been used in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide sequences are shown in the Sequence Listing as SEQ ID NO:1-351. The amino acids are SEQ ID NO:985-1335.

Table 1 shows the various tissue sources of SEQ ID NO: 1-351.

The nearest neighbor results for SEQ ID NO: 1-351 were obtained by a BLASTP version 2.0a1 19MP-WashU search against Genpept release 120 and Geneseq October 12, 2000 release 21 (Derwent), using BLAST algorithm. The nearest neighbor result showed the closest homologue for SEQ ID NO: 1-351 from Genpept. The translated amino acid sequences for which the nucleic acid sequence encodes are shown in the Sequence Listing. The homologs with identifiable functions for SEQ ID NO: 1-351 are shown in Table 2 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 3 shows the signature region found in the indicated polypeptide sequences, the description of the signature, the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 4 shows the name of the domain found, the description, the p-value and the pFam score for the identified domain within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determine from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" Protein Engineering, Vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et as reference, was obtained for the polypeptide sequences. Table 7 shows the position of the signal peptide in each of the polypeptides and the maximum score and mean score associated with that signal peptide.

5.4 EXAMPLE 4

Novel Nucleic Acids

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTV and/or BLAST against Genbank (i.e. dbEST version 117, gb pri 117, UniGene version 117, Genpept release 117). Other computer programs which may have been used in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide, including splice variants resulting from these procedures are shown in the Sequence Listing as SEQ ID NOS: 352-766. The corresponding amino acids are SEQ ID NO: 1336-1750.

Table 1 shows the various tissue sources of SEQ ID NO: 352-766.

The nearest neighbor results for SEQ ID NO: 352-766 were obtained by a BLASTP version 2.0a1 19MP-WashU search against Genpept release 120 and Geneseq October 12, 2000 release 21 (Derwent), using BLAST algorithm. The nearest neighbor result showed the closest homologue for SEQ ID NO: 352-766 from Genpept. The translated amino acid sequences for which the nucleic acid sequence encodes are shown in the Sequence Listing. The homologs with identifiable functions for SEQ ID NO: 352-766 are shown in Table 2 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 3 shows the signature region found in the indicated polypeptide sequences, the description of the signature, the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 4 shows the name of the domain found, the description, the p-value and the pFam score for the identified domain within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determine from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" Protein Engineering, Vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et as reference, was obtained for the polypeptide sequences. Table 7 shows the position of the signal peptide in each of the polypeptides and the maximum score and mean score associated with that signal peptide.

5.5 EXAMPLE 5

Novel Nucleic Acids

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank (i.e., dbEST version 118, gb pri 118, UniGene version 118, Genpept release 118). Other computer programs which may have been used in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide, including splice variants resulting from these procedures are shown in the Sequence Listing as SEQ ID NOS: 767-930. The corresponding amino acid sequences are SEQ ID NO:1751-1914.

Table 1 shows the various tissue sources of SEQ ID NO: 767-930.

The homology results for SEQ ID NO: 767-930 were obtained by a BLASTP version 2.0a1 19MP-WashU search against Genpept release 120 and Geneseq October 12, 2000 release 21(Derwent), using BLAST algorithm. The nearest neighbor result showed the homologs for SEQ-ID NO: 767-930 from Genpept. The translated amino acid sequences for which the nucleic acid sequence encodes are shown in the Sequence Listing. The homologues with identifiable functions for SEQ ID NO: 767-930 are shown in Table 2 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 3 shows the signature region found in the indicated polypeptide sequences, the description of the signature, the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 4 shows the name of the domain found, the description, the p-value and the pFam score for the identified domain within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determine from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" Protein Engineering, Vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et al reference, was obtained for the polypeptide sequences. Table 7 shows the position of the signal peptide in each of the polypeptides and the maximum score and mean score associated with that signal peptide.

5.6 EXAMPLE 6

Novel Nucleic Acids

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank (i.e. dbEST version 118, gb pri 118, UniGene version 118, Genpept release 118). Other computer programs which may have been used

in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide, including splice variants resulting from these procedures are shown in the Sequence Listing as SEQ ID NOS: 931-965. The corresponding amino acid sequences are shown in SEQ ID NO:1915-1949.

5 Table 1 shows the various tissue sources of SEQ ID NO: 931-965.

The nearest neighbor results for SEQ ID NO: 931-965 were obtained by a BLASTP version 2.0al 19MP-WashU search against Genpept release 120 and Geneseq October 12, 2000 release (Derwent), using BLAST algorithm. The nearest neighbor result showed the closest homologue for SEQ ID NO: 931-965 from Genpept. The translated amino acid sequences for
10 which the nucleic acid sequence encodes are shown in the Sequence Listing. The homologs with identifiable functions for SEQ ID NO: 931-965 are shown in Table 2 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 3 shows the
15 signature region found in the indicated polypeptide sequences, the description of the signature, the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 4 shows the name of
20 the domain found, the description, the p-value and the pFam score for the identified domain within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determine from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process
25 for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" Protein Engineering, Vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et al as reference,
30 was obtained for the polypeptide sequences. Table 7 shows the position of the signal peptide in each of the polypeptides and the maximum score and mean score associated with that signal peptide.

5.7 EXAMPLE 7

Novel Nucleic Acids

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank (i.e. dbEST version 119, gb pri 119, UniGene version 119, Genpept release 119). Other computer programs which may have been used in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide, including splice variants resulting from these procedures are shown in the Sequence Listing as SEQ ID NOS:966-974. The corresponding amino acid sequences are SEQ ID NO:1950-1958.

Table 1 shows the various tissue sources of SEQ ID NO: 966-974.

The nearest neighbor results for SEQ ID NO: 966-974 were obtained by a BLASTP version 2.0al 19MP-WashU search against Genpept release 120 and Geneseq October 12, 2000 release (Derwent), using BLAST algorithm. The nearest neighbor result showed the closest homologue for SEQ ID NO: 966-974 from Genpept. The translated amino acid sequences for which the nucleic acid sequence encodes are shown in the Sequence Listing. The homologs with identifiable functions for SEQ ID NO: 966-974 are shown in Table 2 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 3 shows the signature region found in the indicated polypeptide sequences, the description of the signature, the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 4 shows the name of the domain found, the description, the p-value and the pFam score for the identified domain within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determine from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" Protein Engineering, Vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et as reference, was obtained for the polypeptide sequences. Table 7 shows the position of the signal peptide in

each of the polypeptides and the maximum score and mean score associated with that signal peptide.

5.8 EXAMPLE 8

Novel Nucleic Acids

5 Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank (i.e. dbEST version 120, gb pri 120, UniGene version 120, Genpept release 120). Other computer programs which may have been used
10 in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide, including splice variants resulting from these procedures are shown in the Sequence Listing as SEQ ID NOS:975-984. The corresponding amino acid sequences are SEQ ID NO:1959-1968.

Table 1 shows the various tissue sources of SEQ ID NO: 975-984.

15 The nearest neighbor results for SEQ ID NO: 975-984 were obtained by a BLASTP version 2.0al 19MP-WashU search against Genpept release 120 and Geneset October 21, 2000 release (Derwent), using BLAST algorithm. The nearest neighbor result showed the closest homologue for SEQ ID NO: 975-984 from Genpept. The translated amino acid sequences for which the nucleic acid sequence encodes are shown in the Sequence Listing. The homologs
20 with identifiable functions for SEQ ID NO: 975-984 are shown in Table 2 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 3 shows the signature region found in the indicated polypeptide sequences, the description of the signature,
25 the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 4 shows the name of the domain found, the description, the p-value and the pFam score for the identified domain
30 within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determine from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also

disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" Protein Engineering, Vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et al. as reference, was obtained for the polypeptide sequences. Table 7 shows the position of the signal peptide in each of the polypeptides and the maximum score and mean score associated with that signal peptide.

5.9 EXAMPLE 9

Novel Nucleic Acids

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank (i.e. dbEST version 120, gb pri 120, UniGene version 120, Genpept release 120). Other computer programs which may have been used in the editing process were phredPhrap and Consed (University of Washington) and ed-ready, ed-ext and gc-zip-2 (Hyseq, Inc.). The full-length nucleotide, including splice variants resulting from these procedures are shown in the Sequence Listing as SEQ ID NOS:3937-3942. The corresponding peptide sequence is SEQ ID NO: 3943-3948.

Table 1 shows the various tissue sources of SEQ ID NO: 3937-3942.

The nearest neighbor results for SEQ ID NO: 3937-3942 were obtained by a BLASTP version 2.0a1 19MP-WashU search against Genpept release 120 and Geneseq October 12, 2000 release 21 (Derwent), using BLAST algorithm. The nearest neighbor result showed the closest homologue for SEQ ID NO: 3937-3942 from Genpept. The translated amino acid sequences for which the nucleic acid sequence encodes are shown in the Sequence Listing. The homologs with identifiable functions for SEQ ID NO: 3937-3942 are shown in Table 9 below.

Using eMatrix software package (Stanford University, Stanford, CA) (Wu et al., J. Comp. Biol., Vol. 6 pp. 219-235 (1999) herein incorporated by reference), all the sequences were examined to determine whether they had identifiable signature regions. Table 10 shows the signature region found in the indicated polypeptide sequences, the description of the signature, the eMatrix p-value(s) and the position(s) of the signature within the polypeptide sequence.

Using the pFam software program (Sonnhammer et al., Nucleic Acids Res., Vol. 26(1) pp. 320-322 (1998) herein incorporated by reference) all the polypeptide sequences were examined for domains with homology to certain peptide domains. Table 11 shows the name of

the domain found, the description, the p-value and the pFam score for the identified domain within the sequence.

The nucleotide sequence within the sequences that codes for signal peptide sequences and their cleavage sites can be determined from using Neural Network SignalP V1.1 program (from
 5 Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their
 10 cleavage sites" Protein Engineering, Vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et al reference, was obtained for the polypeptide sequences. Table 12 shows the position of the signal peptide in each of the polypeptides and the maximum score and mean score associated with that signal peptide.

15 Tables 5 and 13 are correlation tables of all of the sequences and the SEQ ID NOS.

TABLE 1

Tissue Origin	RNA Source	Library Name	SEQ ID NOS:
lung			3 11 25 49 65 75 114 141 156 160 172 190 198 209 217 224 229 234-235 267 269 274 277 282 284 303 308 312 320 334 336 352 372 396 398 412 414 417 453 464 470 481 492-504 508-509 532 539 581 584 617-619 621 628 633 643 688 691 745 752 761 768 794 822 837 848 876 887 953 967 973
adult brain	GIBCO	AB3001	1 3 12-13 16 22-24 28-29 41 48 58 65 78 82 89-90 94 97 103 112 114-115 117 120 122 130-131 168 181 184 186-187 189- 190 198 208 216 247 249 259 270 277 297 301 308 312 314 321 333 348 374 396 403 406 410 412 416-417 420 423 426-427 431 456 474 481 484-485 488 498 500 508-509 530 549 553 558 563- 564 583 596 602-603 608 612 621-622 624 643 650 674 699 711 736 738-739 753 770 779-780 785-786 802-803 816 822 839 842 848 859 861 871 893-894 897 900 903 925 954 958 967 969
adult brain	GIBCO	ABD003	3 19 21-25 28-29 31 33-34 37 39 41 46-48 53 58 63-64 66 72 78 80 99 103 109-110 112 114 118 120-124 126 132-133 135

			139 143 146 148-149 159 163 168 174 176 179-180 184-185 188-190 202 208- 209 216-217 221 223 230 234-235 240 244 249 251 253 255 258-259 263 269- 270 277 282 285-286 290 294-295 297 301-302 304-305 307-308 311-312 314 320 329 333 335-336 342 344 346 349 354 358 365 370 373-374 377 380 382- 383 388 394-396 399 401-402 406 409- 410 413 416 420-421 425 428 430-431 436-437 442 456 462 464 466-467 474 484 486 495-496 500-501 506 508-509 519 530 537 542 549 561-562 564 572 574 577-578 580-583 586-587 589 592- 593 596-597 601 608 610 612-614 617- 624 630-632 635 637 650 658 663-664 668 676 679 681 689-690 693 699 724 726 732 736 742-743 747 767-770 780 784 789 793 799 802-805 813 817-818 822 824 829-831 837 839 845 848 856 859-860 864 871-872 875-876 881 887 896-897 901 903 907 910-911 925 930 933 943-944 947 952-953 958 962-963 965 967 972 977
adult brain	Clontech	ABR001	3 53 66 113 115 126 135 160 172 179 185 204 263 273 305 312 323 358 380 383 395-396 403 420 428-429 431 461 542 583 586 606-607 611 620 645-646 688 690 715 732 736 740 748 754 768 784- 786 790 796 800 878 897 906-907 947 977
adult brain	Clontech	ABR006	19 32 49 53 68 72 91 103 118 125 130- 131 134 184 224 275 338 350 354 361- 363 374 384 390 394 396 431-432 434- 435 445 468 549 621 732 734-736 745 760-761 764 768-769 775 787 806 811 818 887 903 906 918 930 942 947 957 973 977
adult brain	Clontech	ABR008	2-3 9-11 14 17 21 23-25 28-29 31-35 37 41-42 45 47-48 56-57 65-66 69-70 72 75 77-78 88 91-92 97-99 101 103 112-115 118-128 130-131 135 138-140 142 144- 146 148 152 156-157 159-160 163 168 172 174 176 178-180 182-190 194 196- 198 200-201 204 209-214 218 220-225 228-230 232-233 238-240 243-244 246 254-256 260-264 270 272-274 278-279 282-285 289-291 293-294 296-297 301 303-306 312-314 317 321-322 325-328 334 336 338 340-342 344 346 348 350- 352 354 356-358 363 366 369-374 376 379-381 383-386 388-394 398-399 402-

			403 405 409-412 414 418-421 423-424 426-427 430 433-437 443 445-450 452 456-457 460 462 464 471 479 482-483 485 488 490-498 505 507 510 516 519- 522 524 527-532 535 538-539 542-545 548 551 553 555 561-562 566 569 571 574 580-583 588-589 593 597 601-608 611-612 614-615 617-618 621-622 624 630-635 642 644 646-648 650-652 655 657 659-661 664-665 668 672 674 689 693-699 701-702 708 711 715 717 724 728-730 732 734-735 738-740 745 747- 750 753-755 757 761 763-764 766-769 772-773 775 780-781 789-791 793-795 799-800 802-806 809 812 818-819 821- 822 826 829-830 832 834-835 841 843 845 856 858-859 861 864 866 870 872 876 880 883 885 887 893-898 902 906- 916 918 921 925-926 930-931 933 942- 943 946 948 950-951 953-954 958-960 962-965 967 969-970 972 977
adult brain	Clontech	ABR011	57 196 270 304 344 436 834
adult brain	BioChain	ABR012	14 82 121-122 168 691
adult brain	Invitrogen	ABR013	72 108 263 270 336 425 492-494 732 787 790 826 880
adult brain	Invitrogen	ABR014	293 394 399 764 768-769 928 967
adult brain	Invitrogen	ABR015	738-739 764
adult brain	Invitrogen	ABR016	320 374 396 399 405 684 742-743 767 931 947 967
adult brain	Invitrogen	ABT004	21 33-34 37-38 47 52 57-58 69 72 91-93 109 119 122-124 126-127 135 142-143 158 167-168 185-188 194 200 212 232 242 246 255 258 270 277 279 293 301 312-313 319 322-323 331 341 346 348 371 374 388 391 394 399 401 409 411 429 436-437 456 462 477 488 496 498 510 512 515 539 542 545 549 559 563 573 579 587 589 601-605 612 620-621 624 640 643 647 681 715 723 728 732 735-736 740 745 748 753 766 785-786 792-793 797-801 812 822 829-831 853- 856 859 876-877 884 893-894 908-909 918 925 933 950 969 978
cultured preadipocytes	Stratagene	ADP001	4 28-29 69 93 114 121 132-133 135 151- 152 159 167 172 178 181 184 190 194- 195 203-204 209 217 219 240 248 260- 262 267 273-274 277 282 297 301 304 312 314 326-327 361-362 371 374 388 394 401 403 405 411 420 437 453 466- 467 470 474 478 496 507-509 517 530 532-533 584 588 593 602-603 608 610 617-621 630-631 633 639 642-643 661

			693 729 746 761 765 769 834 842 848 887 907 923 947-950 957 967 969
adrenal gland	Clontech	ADR002	1 3 12-13 21 23-24 27-29 67 74 78 103- 105 108-109 113 115 118 120-121 128- 133 149 156 160 172 177 182 214 217 223 232-233 247 254 269-270 273-274 277 283 285 288 298-299 308 317 319 328 338 340 342 361-362 364 372 376- 377 382 384 401-402 405-406 416 420 431 437 444 446 448 457 462 484 500 507 517 524 532-533 539 545 554 561- 562 564 588 597 602-603 606-607 635 642 646 649 658 664 674 693 703 730 740 745 752 759 765 767 775 779 799 809 817-818 839 845 856 859 863 887 890-891 896 948 953 958 961-963 973
adult heart	GIBCO	AHR001	1 3-4 8 10 14 20-21 25 28-29 33-34 37-38 41 48 54-57 65 69-72 75 78 80 82-83 97 99-100 108 112-115 117-121 123-124 128-133 141 144-146 149 152 159 162- 163 168 172 176 179 181 184 186-187 190-191 201 203 208-209 212 216-218 221 223 227 229 233 244 247 249 253- 255 258 263-264 267 269-270 274 278 280-282 285 289 291 295 297-299 301 303-304 308 313 317 321-322 326 328 334 344 348 352 358 361-363 370-371 380 382-383 388 394-396 398 401 403 405-406 410-416 423 425-427 430-431 436 452-453 464-465 470-474 481-484 487-488 490 492-494 496 499-500 505- 506 508-509 514 523 529-530 533 547- 548 553 558 563-565 577-578 586-588 590 593 597 601-603 606-608 610-613 617-619 621-622 626-628 637-638 642- 644 652 658 661 672 682-683 688 691 693 697 699 708 711 713 715 732 737 745 747-748 750-753 759 761 765 768- 770 775 790 802-803 814-815 818-819 830 837 839-840 842 845 848 859 861- 862 867 876-877 887 891-892 896 900- 901 903 905-906 908-909 919-920 922 925 928 936 939-940 946-947 950 953 959 967 970-971 973 977
adult kidney	GIBCO	AKD001	1 3 8 12-14 17 19-25 28-29 33-34 37-39 41 46-48 50 52 55-60 62 65-67 69 71-72 75 77-78 82 84 89-90 93 97 108-110 114- 116 118-121 123-125 128 130-133 135 138 144 146 149 156 159-161 163-164 167-172 176 179 184 186-187 189-190 194 196 200-202 204 209 211-212 216- 217 219 221 223-224 229 232-235 244

			247 250 253 255-256 258 263-264 268- 272 274 277-281 283 286 288-290 292 294-295 297 301 303-309 311-314 316 319-323 325 328-338 342 348-349 352 354-355 358 361-363 365 370-371 373 376-378 380 382-383 388 395-399 401- 403 405-406 409-413 416 418-420 425- 428 430-431 440 442 452-454 462 464- 465 470 472-474 477 479 481 483-485 487-489 492-495 498-500 504 506 510 517 522 525 529-530 532-533 539 542- 543 547 551-552 558 560-564 569-570 573-574 577-578 580-583 585-590 594- 596 601-608 610-613 617-621 624 626- 628 630-631 634-636 639 642-643 648 652 656 658 664-665 676-677 679 681 688-691 693 697 699 708 711 715 717 720-722 724 729-732 738-741 747-748 751-753 761 765 770-778 780 784 789 791 793 797 804 813 817 823-824 834 837 839 842-843 845 848 859 861-862 864 867 870 876-877 887 889 892-894 896-897 900-901 903 907 913-915 918 921 923 925 929-930 932 939 942 946- 947 949-950 953 958-959 961-963 967 969 972 977
adult kidney	Invitrogen	AKT002	1 3 16 21 30 32 35 38-41 46-47 56 77 92 109 123-124 130-131 146 149 161 167- 168 172 176 190 209 212 234-235 258 279 292 301 303 308 314 333 355 363 372 380 383 396 399 402 418-419 426- 427 431 448 454 461 471-474 488 492 495 504 506 508-509 520-521 530 537 539-541 545 547 563 582-583 592 613 617-618 621 623-624 633 655 688 690 693 699 704 713 732 745 752-753 761 766-768 770 784 789 797 837 842 848-849 866-867 877 887 893-894 903 914-915 925 929-930 937 944-945 947- 949 955 961 967 984
adult lung	GIBCO	ALG001	1 3 14 18 28-29 38 54-56 59 92 110 114- 115 130-131 146 149 156 159 164 167 176 184 209 217 234-236 240 255-256 258 263-264 269 271 276 280-281 297 305 308 312 314 322 325 332 336 344 353 361-362 388 401 410 420-421 426- 427 431 465 469 474 484 498 500 506 508-509 517 530 532 573 592 596 613 619-620 623 626-628 638 658 679 681 684 689 717 731 741 771 791 799 817 834 845 861-862 864 875-876 901 921 925 928 932 940 947 949 959 962-963

			967
lymph node	Clontech	ALN001	3 10 110 146 160 168 196 209 221 269 278 301 336 348 394 405 411 420 422 459 464 474 485 503 506-507 532 563 582 619 623 630-631 642 669 684 697 713 715 727 747 767 769 789 825 839 842 849 887 896 913 921 925
young liver	GIBCO	ALV001	3 14 16 37-38 41 51 56 60 97 104-105 108 110 117 119 128 130-131 134 139 149 152 169-172 176 184 189-190 200 209 212 216 218 228 232 255 258 263 270-271 275 285-286 292 295 298-299 301 304 314 341 358 365 368 376 400 410-412 431 474 481-482 485 496 500 504-505 517 520-522 524 530 532-533 547 551 563 581 583 610-611 621 624 635 643 691 708 711 715 720 752 755 761 768 796-797 811 818 830 845-847 852 864-865 867-869 896 899 910-911 949 958 965 969 972-973
adult liver	Invitrogen	ALV002	3 37 42 56 60 71 82 104-105 114-115 117-118 125 130-131 134-135 164 169- 172 176 179 200 203-204 212 217 223 226 232 237 244 263 274-275 292 301 310-312 314 317 349 354 364 368 372 376 398-399 402 426-427 439 442 451 458 465 474 482 485 490 506 515 525 527 545 547 552 568 571 573-575 582 587 594-595 604-605 608 610 621 630- 631 634-635 637 657 664 690 693 699 723 726 745 751 763 767 784 793 811 822 845 848 852 856 861-862 864 892 899 908-909 925 950 958 967 983
adult liver	Clontech	ALV003	60 134 169-171 275
adult ovary	Invitrogen	AOV001	1 3 9-10 12-14 16 18 20 22-25 28-29 33- 35 37 39 41-42 46 48-50 55-57 59 63-67 69 71-72 75 77-80 82 88-89 92 101 103- 106 108-110 113 115 119-121 123-126 128-133 135 138 142-146 149 151-152 159-161 167-168 172 174 176-177 179 181 184-190 194 198 200 203 208-209 211-212 214 217 219 221 224 226 232- 235 240-242 246-247 249 251 254-255 258-259 264 269-271 274 276-277 279- 283 285 288 290 293-294 297 301-304 306-308 311 314 319-322 325-326 328- 329 331-332 335-338 341-342 344 348 354-358 361-363 365 368 370-372 374 376 379-380 382-383 388 394-396 398- 399 401-402 405-406 409-412 416 418- 421 423 425-433 438 442-443 449-452 454 462 464 466-467 469-471 474 479

			482-484 488 490 492-496 498 500-504 506-509 511 515-518 520-524 529-530 532-533 537 539-542 545 551 555 558 560-565 569 571 573 577-578 581-583 585-590 592-593 596-597 600-605 608 610-611 613-614 617-628 633-637 639 642-643 646-648 650 652 654 656 658 664 668-670 672 674 679 681 684 688 691 693 697-699 701-702 713 717 721- 722 724 729-732 738-744 747-750 752- 753 755 759 761 765 767-774 779-780 783-784 789 793 795-797 801 813-818 823-824 828 830-832 834 837 839 841- 842 845 848-851 856 859 862 864 866- 867 870-871 874-878 881-883 887-889 891 893-894 896-897 901 903 906-911 913 919-922 925 928 930 936 939-940 943-944 946-947 949-950 952-953 955 957-958 962-963 965 967 969 971 973 977 981-982
adult placenta	Invitrogen	APL001	41 56 67 253 301 304 334 380 383 451 474 479 500 577-578 643 648 729 767 856 859 866 873 962-963
placenta	Invitrogen	APL002	3 21 31 38 63-64 78 135 143 168 186-187 212 232 244 263 280-281 334 336 344 348 371 374 394 399 461 490 582 588 602-607 610 620 699 745 769 793 817 822 859 897-898 923 928 931 943 949 969 973
adult spleen	GIBCO	ASP001	1 3 21-22 46 52 54-55 57-58 61-62 72 74 78 82 88 118 121 130-131 137 152 159 168 172 189 202 209 217 223 234-235 252 255 263 269 271 274 282 288 290 301 314 322 335 350 363 394 403 405- 406 410-412 415 431 459 464 472-474 482 488 500 506 510 514 517 532 537 542 561-563 589 593 602-603 610 613 619 621 636 642-643 655 658 662 674 676 679 681-682 684 689 691-692 697 699 715 720 723 729 747-748 769-770 782 793 818 830 834 845 856 859 862 877 887 893-894 896 903 906-907 914- 915 918 925 928 930 940 946 965 967 977 982
testis	GIBCO	ATS001	6 22 28-29 33-34 41 48 52 62 65 72 97 106 109 118 132-133 145-146 168 172 176 183 185 189-191 195 209 211-212 214 221 223 230 254-255 258 263 269 283 297 312 314 321 342 352 361-362 365 380 383 388 395 401 405-406 412 430-431 441 469-470 474 479 495-496 500 506 520-521 533 543 545 548 560

			563 574 582 589-590 593 608 616-618 620 623-624 638 642-643 697 699 708 711 745 747-748 765 767-768 779 784 789 812-813 834 837 839 848 859 862 868-869 875-877 887 889 893-894 896 928 944 947 953-955 972 981
Genomic DNA from BAC 63I18	Research Genetics (CITB BAC Library)	BAC001	515
Genomic DNA from BAC 393I6	Research Genetics (CITB BAC Library)	BAC002	640
Genomic DNA from BAC 393I6	Research Genetics (CITB BAC Library)	BAC003	640
adult bladder	Invitrogen	BLD001	50 55 66 71 111 143-144 148 160 201 209 223 255-256 280-281 286 305 315 319 340 394 431 442 488 497 505 518 552 588-589 621 636 664 676 715 738-739 769 790 824 837 845 877 887 936 940 948 962-963 967
bone marrow	Clontech	BMD001	3 10-13 16 18 20-21 25 28-29 31-34 41 45 48 52 54-55 57 59 61 65 67 72-73 75 78 80 82 84 99 103 108 110 114-115 118- 120 123-124 128 130-133 143-144 148 152 159-161 163 168 172 174 176 178 190 192 198 203 209 211 217-218 221 223-224 227 233-236 244 247 249 252 254 258 260-262 267 269 272 278 280- 281 284-285 288 290 294-297 301 304 308 314 317-318 320-321 325 328-330 333-335 349 351-354 358 363 365 367 377 382 388 394-397 400 405 408 410- 412 418-421 425-428 431 433 435 442 449-450 453 455 459 464 468-470 474 478-479 481 484 490 496 504 506 508- 509 511 519-521 530 532 539 553 558- 559 561-563 580 582 586 592 599 608 610 613-614 617-619 623 625-628 635 638 641-643 658 664 672 682 699 711 713 717 731 734 740 742-743 745 761 768-771 774 776-778 784 787 789 813 817-818 822 834 839-840 842 848 862 866 870 876 885-887 891 896-898 900 903 906 913 919 921-922 927-928 939 944 947 950 953 959 961-963 967-968 970 973 977
bone marrow	Clontech	BMD002	3 9-10 15-19 30 33-34 39 45 54 57 63-64 71 82 102 116 119 130-133 148 152 156

			159-160 168 176 182 224 254-255 271-272 282 285 290 297-299 301 305 323 333 340 344 351-355 358 361-362 364 367 370 372 387 394-395 399 403 405 409 411 449-450 459 461 468 474 488-489 524 530 532 580-582 592 602-603 611 617-618 621-622 630-632 642 661 663 694 717 730 734 740 745 752 755 761 767 769-771 775-778 784 787 811 813 818 832 840 842 849 859 878 887 893-894 896-898 903 906 908-909 923 928 944 946-949 953 958-963 965 982
bone marrow	Clontech	BMD004	54
bone marrow	Clontech	BMD007	766 887 928
adult colon	Invitrogen	CLN001	22 37 67 97 117 121 148-149 168 172 190 200 204-205 232 244 263 268 292 301-302 363 377 384 452 455 459 470 530 582 602-603 619 687 723 728 751 761 831 861 887 914-916 934 955 969 984
Mixture of 16 tissues – mRNAs*	Various Vendors*	CTL016	358 740 760
Mixture of 16 tissues - mRNAs*	Various Vendors*	CTL021	468 527 928
adult cervix	BioChain	CVX001	1 3 10 14 22 28-30 37 41 47-48 51-52 54-57 71 82 89-90 92 106 108 110-111 117-118 121 129-131 135 141 143-146 160-161 164 168 172 177 189-190 193 195 200 204 209 211-212 217 226 229-230 232 234-235 240-242 246 254 260-263 268-270 274 277 282 285 292 295 297 305-308 314-316 319 328 343-344 348 354 358 363 368 380 382-384 389 394 396 399 401 405-407 410 416 418-421 428 430-431 437 442 453-454 459 464 469 471-473 476 480 484 492-495 500 504 506-509 516-517 526 530 532 545 550-551 563-565 569 577-578 585-586 590 608 611 613 619 621 623 628 630-631 634-637 641 643 648 656-658 664-665 674 679 682 689-690 693 700 703 708 713 721-722 724 728 732 742-743 747 750 752 755 757 761 763 767-769

* The 16 tissue-mRNAs and their vendor source, are as follows: 1) Normal adult brain mRNA (Invitrogen), 2) normal adult kidney mRNA (Invitrogen), 3) normal adult liver mRNA (Invitrogen), 4) normal fetal brain mRNA (Invitrogen), 5) normal fetal kidney mRNA (Invitrogen), 6) normal fetal liver mRNA (Invitrogen), 7) normal fetal skin mRNA (Invitrogen), 8) human adrenal gland mRNA (Clontech), 9) human bone marrow mRNA (Clontech), 10) human leukemia lymphoblastic mRNA (Clontech), 11) human thymus mRNA (Clontech), 12) human lymph node mRNA (Clontech), 13) human spinal cord mRNA (Clontech), 14) human thyroid mRNA (Clontech), 15) human esophagus mRNA (BioChain), 16) human conceptional umbilical cord mRNA (BioChain).

			779-780 784 788 810-811 813-815 822 834 836-837 839 848 861 866-867 871 874 877 887 891-894 897-898 901 913 916 919 921-922 925 946-947 953 958- 959 967 969 973
diaphragm	BioChain	DIA002	3 39 184 203 431 563 848 967
endothelial cells	Strategene	EDT001	3 6 8-10 14 19-24 28-29 33-34 37 39 41 46 48 52 55-58 62-65 67 69 71-72 75 78 80 82-83 87 101-102 108-109 114-115 117 123-124 128 130-133 135 138 143 145-146 149 156 159-160 167-168 172 174 176-177 179 181 184-187 189-190 194-195 200 203 208-209 212 216-217 219 223-224 226-227 229 234-235 244 248-249 254-256 258 263-264 267 269 271 274 276-282 285 290-291 294 297 301-304 308 311 313-314 316-317 320- 321 323 325-326 328-329 331-332 334- 337 339-341 344 348-349 352 354-355 358 361-363 365 367 371-372 375 379- 380 383 389 394-395 398-403 405-406 409-412 425-428 437 442-443 448 454 464 466-467 474 479 481 490 492-498 500 503 506-509 511 517 520-521 523- 524 530 532 537 540-542 558 561-563 565 569-570 573 581-583 586 588-589 596 602-608 610-611 613 617-622 625 628 630-631 633-637 642-643 646 648 650 652 659 661-662 682 688 690-693 696 698-699 708 712 715 717 720-722 724 727 729 740 745 748-750 752 761 765 767-770 772-773 779 784 789 792 794 796 802-803 811 817-818 821 824 827-828 830 834-835 837 842 845 848 859 861-862 864 866-867 870 876 885 887 891 893-894 897-898 900 903 906- 907 913 916 921 925 939 947 950 953 955 957-958 962-963 967 973 978 984
Genomic clones from the short arm of chromosome 8	Genomic DNA from Genetic Research	EPM001	324 515 640
esophagus	BioChain	ESO002	97 103 128 371 474
fetal brain	Clontech	FBR001	67 129 156 159 232 267 433 446 503 845 952
fetal brain	Clontech	FBR004	28-29 185 213 277 350 384 432 485 501 549 651 747 754 761 780 787 848 870 887 906 958
fetal brain	Clontech	FBR006	10-11 14 21 30 32 47 49 56 65 69 72 77- 78 82 84 97 101 115 118 121 125 128 130-131 138 142 148 152 159-160 179 185 188 194 197 203 210 212 214 219

			222 227-229 243-246 249 252 256 264 270 273 282 285 290-291 293 301-303 305-306 312 321-322 325 327 339-340 344 346 350 354-357 363 367-371 374 388 391 394-395 399 402 405-406 410 414 420 426-427 436-437 442 444 454 456-457 460 462 464 470 480 485 492- 494 507 510 516 524 528 530-532 539- 542 549 553-554 561-562 580-582 588- 589 602-608 611 615 617-619 621-622 624 632 636 641-642 646-647 651-653 661-662 666-669 672 677 691 715-716 730 735 740 752 754 761 767-770 772- 775 780-781 799-801 808 818 822-823 835 843 845 856 859 864 867 876 880 885 887 890 893-894 896 913 918 926 942 946-947 951 957-959 962-963 970- 971
fetal brain	Clontech	FBRs03	130-131 312 517 637 691 738-739
fetal brain	Invitrogen	FBT002	3 22 28-31 47 57 63-64 72 75 77-78 86 94-95 97-98 126-127 135 140 143 156 159-160 167-168 177 185 190 196 201 203-204 214 217 230 254-255 258 267 273-274 277 279 282-283 292 301-302 305 312 314 323 329 346 348 367 374 382 394 399 401 403 412 415 420 432 437 474 482 485 495 507 513 517 527 529-530 539-542 548 552 579 587-588 600 604-605 612 617-618 621-622 624 634 642-643 647-648 650 679 689 693 699 712 715 742-743 745 748-749 753 768-769 793 797 829-831 834 845 848 856 859 893-894 908-909 913 916 931 933 940 950 967 969
fetal heart	Invitrogen	FHR001	19 57 130-131 354 431 642 769 844
fetal kidney	Clontech	FKD001	3 31 33-34 38 48 54 72 160 208-209 211 223 264 269 277 283 290 313 325 341 348 358 396 418-420 474 484 506 508- 509 517 520-521 532 547 553 558 567 569 587 596 608 610 613 619 622 626- 627 642 679 734 745 818 843 887 896 903 916 969 971
fetal kidney	Clontech	FKD002	19 474 726 903
fetal kidney	Invitrogen	FKD007	3 118 186-187 230 244 271 432 887 969
fetal lung	Clontech	FLG001	69 132-133 156 168 208-209 217 267 269 274-275 286 354 394 396 406 462 483- 484 608 619 751 769 771 834 914-915 925
fetal lung	Invitrogen	FLG003	3 8 28-29 32 39 50 66 82 88 92 168 186- 187 200 204 212 226 229 246 274 309 327 332 368 374 382 394 398 426-427 431-432 442 485 536 555-557 587 604-

			605 621 624 636 642-643 661 677-678 724 753 769 848 859 864 877-878 896 902 904 914-915 958
fetal lung	Clontech	FLG004	130-131 394 664 769 942
fetal liver- spleen	Columbia University	FLS001	3 8-10 12-13 16-17 19-25 27-29 33-35 37- 38 41 45-46 48 52 55-58 60-67 69 71-74 77-78 80 82 84 87-90 104-106 108-109 112-121 123-125 128-134 138 141 143- 146 149 151 156 159 163-164 167-172 174 176-179 181 184 186-188 190 194 200-201 203 208-209 211-212 216-217 219 224-227 229-230 232 234-235 237 241 243-244 246-248 254-255 258 260- 263 267 269-270 273-282 284-285 288- 290 292-295 297-299 301-306 308 311- 318 320-323 326 328 332 335 341-344 348 352 354-359 361-365 367-368 371- 374 376-380 382-383 388-389 394-396 398-399 401-411 413-414 416 418-421 425 428-430 432-433 437 439 442-444 449-450 452 456-457 461-470 472-474 478-479 481-482 484-485 487 490-494 497-499 504-507 511 514-515 517-521 523-524 526 529 532 537 540-541 547 555 558-559 563 575 577-578 580-596 598-599 601-603 606-608 610-613 617- 624 626-628 630-631 634-636 639 642- 643 647-648 654-656 663-665 672 674- 675 679 681 684 686 688 691 693-699 711 713 715 717 719-726 729 732-733 738-740 745 748-749 751-753 757 759 761 767-770 776-778 780 784 787 792- 794 799 804 809 811 813 817-819 822- 825 830-831 834 837 840 842 845-848 852 856 859 861-862 865 867-869 871 874-878 887-888 891 893-894 896-900 903 905-911 913 916 918 923 928 930- 931 936 939 942 944 946-950 952 958- 959 961-963 965 967 969-970 972-973 976-977 981-983
fetal liver- spleen	Columbia University	FLS002	3 8-13 15-17 19-20 22 25 28-29 33-35 37 41 45-46 52 54-56 60-61 63-64 66-70 73- 74 78 80 82 92 99 104-106 108-109 112 115-116 118 120-121 123-125 128 132- 135 139 141 143-144 146 149 152 156 159-161 167 169-172 174 176-177 179 181 185 188 190 194 196-197 200 204 212 214 216-218 223-224 226-230 232- 235 237 246-247 252 254-255 258-263 267 270-277 284-286 288 292 294-295 297-299 301 303-305 308 310 314 318 320 323 328 330-332 335-337 340 342-

			344 352 354-355 358 361-365 367-368 371 373-374 376-377 382 388 394-396 398-399 401 405-406 409-411 413 418- 421 429 431 439-440 442-444 451-452 457 462-463 466-468 470 474 477-479 481 483-484 487-488 491 495 499 504 508-509 516 519-521 524 526-528 530 532 537 540-541 543 545-547 550-551 553 555 560 564 568 574-575 577-578 580-592 596-597 600 602-603 608 610- 611 613-614 617-618 621-622 628 630- 631 634 637 639 642 644 647 654 658- 659 665-667 669-675 679 681 684-685 688-690 693 695 697 708 711 713 715 717-719 723-727 729 731-734 738-739 741 745-746 749-750 753 759 761 766- 767 769-770 776-779 782 784 791-792 794 805 808 817-818 822 824-825 830 834 837 842 845-849 852 856 859 864- 865 867 874-878 888 891-892 896-900 903 905-906 908-909 913 916 918 921 923 925 932 936 939-940 942 944 946- 947 949-950 953 955-956 958-959 961- 963 965 968-970 973 977-978 981
fetal liver- spleen	Columbia University	FLS003	19 60 78 224 273 275 370 373-374 401 602-603 639 643 730 732 738-739 748 752 770 782 928 930 947 949
fetal liver	Invitrogen	FLV001	37 55 60 69 72-73 97 104-105 108 113- 114 116-118 121 135 143 152 167-168 186-187 195 200-201 209 217 223 240 244 253 255 275 284 301 311 314 317 336 342 348-349 358 371 374 382 394 402 411-412 418-419 428 430 442 453 517 568-569 580 582 584 587 589 601- 603 606-608 617-618 624 634 639 642- 644 646 664-665 669 679 715 717 720 726 745 748 751 769-770 782 791 794 797 824 830-831 845-847 852 859 870 899 913-916 925 928 948 956 958 969 976 982
fetal liver	Clontech	FLV002	72 418-419 632
fetal liver	Clontech	FLV004	3 160 169-171 355 367 374 376 547 617- 618 621 646 717 741 771 836 878 976
fetal muscle	Invitrogen	FMS001	15 27 32 37 67 72 83 99 112 121 138 167 174 177 186-187 190 203-204 211 215 230 252 259 312 374 403 406 409 457 461 485 505 517 528 530 540-541 544 549 554 558 579-580 583 602-603 608 639 642-643 654 664 699 715 730 737 751 772-773 788 802-803 810 848 856 859 864 868-869 887 893-894 905-906 910-911 923 948 967

fetal muscle	Invitrogen	FMS002	15 99 130-131 223 361-362 431 474 505 581 639 643 666-667 784 790 808 810- 811 874 880 887 903 946 950 958 962- 963 973
fetal skin	Invitrogen	FSK001	3 6 20-22 32-34 41-45 47 49-52 55 63-64 66 69 77 80 88 91 98 101 111-112 115 126 130-131 135 142 144 146 160 163 167 176 188-190 196 201 204 208 213 215 217-218 229 232 244 246 248 255 263 265-269 274 279-281 283 285 288 292 294 297 301 303 308 314 321 341- 342 344 348 354-355 358 361-362 366 369 371-372 374 381-382 384 386 394 401 403 405 413 415 428 431 437 440 460 466-467 472-473 477 481 483 495 499 504 517 522 532 536-537 539-541 545 556-558 569 574 576-578 580 584- 585 587-589 592-593 602-603 606-608 612 617-618 621 624 634 637 639 642- 643 647 664 673-674 676 680-681 689 699 705-707 709-715 724 728-730 738- 740 745 748 752 765 768-769 772-773 793 797 817 823 830 834 842 848 859 861 864 870 874 883 887-888 893-894 901 904 908-909 913-916 923 925 947 950 958 962-964 967 975
fetal skin	Invitrogen	FSK002	3 130-131 146 194 306 354 367 400 405 474 489 520-521 547 558 561-562 585 596 730 740 748 755 767 771 810 840 893-894 946 959
fetal spleen	BioChain	FSP001	276 563 842
umbilical cord	BioChain	FUC001	3 20 33 34 39 48 50 52 55-57 65 67 69 72 77 79 82 92 109 112-113 121 130-133 138-143 156 167-168 172 174 179 184- 185 190 194-196 200 202-203 208-209 229-230 244 269-271 278 284-285 290 297-299 303 305 308 320 331-332 336 338 342-343 363 367 372 374 379-380 383-384 392-394 397 399 402 405-406 410 425-427 429-430 449-450 474 476 484 497 499 501 504-505 510 515 517 532-533 539 549 551 558 563 569 574 577-578 581 586-587 597 602-603 608 610 617-619 621 626-627 634-637 639 642-643 658 663-664 674 690-691 693- 694 699 713 715-717 720 724 726 729 738-739 746-747 749 759 761 765 768- 769 774-775 793 797 807 818 822 837 848-849 856 862 868-869 874 885 887 892-894 903 906-907 916-917 919-920 928 936 939 944 946-947 962-963 967 969

fetal brain	GIBCO	HFB001	3 9-10 12-14 16 21 25 28-30 32-34 37-39 41 47-48 52-53 56 65 67 69 71-72 75 80 84 92 97 103 106 110 114 117-119 123- 124 127 129 132-133 135 138 141-142 144-146 148-149 152 156 159-160 168 172 174 176 179 181 184-185 190 198 208-209 212 214 219 221 223-224 229- 230 233-236 240 244 247 251 253-255 258-259 270 273 276-277 285 297 304- 305 308 312 314 322-323 325 328 332- 333 335-337 339-340 342-344 346 352 354 358 363 365 370-372 374 382 394- 396 398 401 403 405-406 409-412 414 416 425-427 431-432 437 442 445 453 456 462 466-467 469-470 472-474 479 483 488 490 492-497 500-501 504 506- 510 520-521 524 530 537 539 545 549 552 558 560-562 564 569 579 582-583 586-587 596 602-608 610-612 614 617- 624 626-628 630-631 633 635 638 641 643 647-648 656 658 661 676 679 688- 689 693 696-697 711-712 715 724 726 731 735 745 747-749 752 754 761 765 767-770 774 779-781 784-786 789 799- 800 802-803 813 818-819 823-824 831 834-835 837 839 845 848 859 864 866- 867 871 874-875 881 887 891 893-894 896-897 900 906-907 910-911 918 921- 922 925 927-928 930 943-944 946-947 950 953 962-963 965 969 972-973 977
macrophage	Invitrogen	HMP001	86 168 186-187 297 537 608 681 761 845 877
infant brain	Columbia University	IB2002	2-3 9-10 12-14 16 21 25 27-30 32 37-38 46-47 49 55-56 58 63 69 71-72 78-79 82 84-86 91-92 98-99 106 109-110 113-115 118 127-128 130-133 135 138 142 144 151 156 168 173-176 180-181 185-188 192 194 196-201 203 208 210-212 214 217-218 224 229-231 233 236 238 240- 241 244 246 251-256 259 263 270-271 277-279 284-285 287 293-294 296 301- 302 308 312-314 317 322-323 327 330 333 339 342 345-346 351 354 358 361- 362 365-366 368 370-371 373-374 382 388 394-396 402 405-406 411-412 415- 416 420 424-425 428 431 436-437 440- 441 444-445 453 456 460 465 474 479 482-483 488 495-496 498 501 503-504 506-510 515-517 520-521 524-525 529 531-532 534-535 537 539-542 544-545 549 561-562 569 574 577-578 580-583 586-587 589 592 596 600-608 610 612-

			613 616-618 620 622 624 629-632 634-635 637 641 643-644 650-651 653 661 663-664 676-677 689 693 695-698 708 711 720-722 724 730 732 735 740 745-748 754 765-766 768-769 779-781 785-786 789 791 796 798 800-803 807 811-813 818-819 822-824 830-831 834-835 837 839 842-843 845 854 856 858 864 867-869 875-877 879 881 887 892-894 896 903 907-911 913 916 919-920 925 930-932 936 939 943 946-947 953 958 970-973 977-978 982 984
infant brain	Columbia University	IB2003	3 12-13 21 27-29 32 39 49 69 72 82 91 113 116 126 128 132-133 142 144 156 176-177 184-185 188 194 208 212 223-224 228 230 244 255 259 267 270 273 276 293-294 312 320 326-327 337 342 346 354-355 358 361-363 382 388 390 394 396 399 402 420 425 431 442 462 474 482 484 488 495-496 510 520-522 524 529 540-541 549 563 582 586 588-589 596 600-603 606-607 612 617-618 620-621 632 647 650 679 720-722 724 735-736 746 751 754 769 785-786 793 800 807 811-813 818-819 822 824 831 834 838-840 843 856 864 892 896 907 919-920 925 930-931 936 947 950 957 973 982
infant brain	Columbia University	IBM002	16 47 82 84 201 263 302 376 394 421 440 488 537 592 606-607 635 740 769 887 892 906 921 926 971
infant brain	Columbia University	IBS001	84 86 180 185 198 201 203 230 279 312 326 346 354 366 388 488 542 561 588 620 647 664 732 740 785-786 801 807 822 827 910-911 925 931
lung, fibroblast	Stratogene	LFB001	3 11 25 49 65 75 114 141 156 160 172 190 198 209 217 224 229 234-235 267 269 274 277 282 284 303 308 312 320 334 336 352 372 396 398 412 414 437 453 464 470 481 492-494 508-509 532 539 581 584 617-619 621 628 633 643 688 691 745 752 761 768 794 822 837 848 876 887 953 967 973
lung tumor	Invitrogen	LGT002	1 3 9-10 12-13 20 31 38 41 46 48 51-52 56 58 63-64 72 74-75 78 82 88 101 106-107 110 114-115 117-118 120-121 123-124 128-133 135 143-146 149 151 156 159-161 163-164 167-168 172 176 178-179 184-185 189-191 194-196 200 203 209 212 216-217 226 228-229 232 234-236 241 246 248 256 258-259 263-264 269-271 274 282-283 285-286 290 292

			294 297 301 308-309 311 314 317 321 326 328-329 331 333-334 341 348 352 354-355 363 365 371 380 382-383 388 394-395 398-402 405-406 410-411 413 416 418-419 426-427 439 442 452-453 458-459 461-462 464-465 470-471 474 478 483-484 490 495-496 499 510 522 524 528 536-537 540-541 543 548 556- 558 560-565 571-573 580 582 587-588 592 597 602-605 608 610 612-613 617- 622 625-629 633-634 636 642-644 648 661 664 669 679 688-689 691 693 699- 700 708 717 723-724 730 733-734 738- 740 745 747 749 752-753 761 767-768 770 779 782 784-786 789 793-794 797 817-818 820 823-824 834 837 842 845 848 855 857 859 862 864 866 870 875- 877 887 892 896 900-901 907-909 914- 915 919-920 923-925 939 943 947 949 953 958 962-963 965 968 970 972-973 977
lymphocytes	ATCC	LPC001	3 9-11 32 47 50 56 71 75 88 97 99 102 121 125 128-129 135 138 141 149 163 167-168 212-213 217 233 255 290 294 301 305 311 314 342 372 377 388 398- 399 410 437 442 453 470 474 481 495 500 506 510 529 532 537 542 558 571 579 604-605 610 620 628 637 643 658 666-667 676 679 697 708 713 728 730 734 749 765 768 796 807 818 822 834 839 848 859 875 885 887 896 903 906 914-915 928 947 973 981 987
leukocyte	GIBCO	LUC001	1 3 9 11 18-19 21 23-25 27 31-34 39 41- 42 46-48 52 54-58 62-69 71-72 74-75 78- 80 82 89-90 93 99 110 115-121 123-124 128-133 135 138 141 143-146 149 152 156 159-161 163 167-168 176 179 181 186-187 189-190 194 198 200 203-204 209 211-212 218-219 226 232-236 240 244 247 251 253-255 258-259 263-264 269 271 274 278-279 282-283 285 288- 290 294-295 297 301-306 311 313-314 317 320-321 325 328 330-331 335 337 342 344 348 350-351 353-354 358-359 361-365 368 371-372 375 388-389 394- 395 397-401 403 405 407 409-412 421 425-427 432 437 442 448-450 452 457 460-461 468-471 474 476 479-482 484 492-494 496-498 500 506-510 516-517 520-521 524 529-530 532 537 540-544 551 553-554 558 560-565 569 577-578 580-583 586-587 589 592 596-597 602-

			603 606-608 610-624 626-628 630-631 634-635 641-643 654 657-658 661 663- 665 669 672 677 679 684-689 691 696- 697 699 708 711 713 715 717 721-724 728 730 738-740 747-749 755 761 765 767-769 771 774-779 782 784 789 791- 792 794-795 797 807-808 811-815 817- 818 822 824 828 830 832 834 839-840 842 845 848 856 859 862 864 867 871 875-877 887 891 893-894 896-898 903 906-911 913-916 921 923 925 927-928 930 932 935-936 939 943-944 947 949- 950 953 958-959 961-963 965 967 972- 973 982
leukocyte	Clontech	LUC003	1 41 82 106 119 123-124 160 177 184 201 212 221 228 271 279 285 295 321 325 372 394 411-412 443 468-470 530 532 537 551 569 580-581 613 619 623 626- 627 642 655 697 761 767 769 775 789 809 867 887 923 928 950
melanoma from cell line ATCC #CRL 1424	Clontech	MEL004	3 25 55-56 67 71 78 109 121 129 146 167 172-173 176 200 209 212 258-259 263 278 297 301 306 312 335 338 340 352 361-362 367 388 395 402 410 418-419 429 437 454 464-465 481 496 500 503 507 524 532 539 560-562 581-582 587 589 599 612-613 617-621 623 643 657 663-664 672 715 724 748 752 761 767- 768 770 785-786 789 835 848 877 887 896 916 919-920 947 967 978-980
mammary gland	Invitrogen	MMG001	1 14 19 21 28-29 31-37 47 49-51 55 57 63-67 69 71 72 75-78 92 108-109 111 116 121 123-124 126 128 130-133 135 142 144 148-150 156 159 164 168 172 177- 179 184 186-187 190 194 200-204 209 212 217 226 230 232-236 241 244 246- 247 252 255 258-259 263 268 270 275 279-283 285 290 292-293 301 304-305 311 313-314 317 320 322-323 326-327 330 332 338 342-344 348-349 354 360 363 367 371 374 380 382-383 385 388 394-395 398 401-403 407 409 411-412 418-420 426-427 430 435 437 442 449- 453 459 461 465-468 470 474 477-478 480 483 485 488 498 500 503-504 507 515 519 522 524 529-532 538-541 544 547 555 560 563 565 569 573-574 579- 580 582 584 587-589 593 597 601-610 612-613 615-618 620-622 624 634 636- 637 639 642-644 646-647 650 657 663- 664 674 676 679 688-689 691 693 696 701-703 713 715 717 728 730 732 738-

			739 741-743 745 749 751 753 763 767 769 772-773 785-786 793 796-797 812 821-824 830-833 837 848 856 859 861 864 868-870 876-877 887 891 893-894 898 903-904 907-911 913-918 921 923 925-926 930-931 936 942 949-950 958 961 966-967 969 972-973
induced neuron cells	Stratogene	NTD001	9 65 82 92 106 113 142 146 156 172 176 191 208 221 258 277 328 333 346 361- 362 371-372 375 388 410 414 418-419 440 471 484 495 516 524 529-530 592 610 628 642 650 745 748 752 761 793 818 848 851 897
retinoid acid induced neuron cells	Stratogene	NTR001	19 87 184 305 385 440 474 626-627 643 748 799 834 977
neuronal cells	Stratogene	NTU001	19 33-34 42 70 82 87 109 115 126 146 172 185 188 194 212 255 269 274 283 312 317 329 340 361-362 367 379 394 399 401 410 420 426-427 474 479 507 530 579 582-583 610 617-618 636 643 658 732 740 765 769 784 791 793 799 802-803 818 842 851 864 897 907 932
pituitary gland	Clontech	PIT004	3 19 123-124 194 255 354 358 373-374 377 426-427 462 492-494 635 785-786 793 893-894
placenta	Clontech	PLA003	138 176 574 896 972
prostate	Clontech	PRT001	3 9 16 57 65 75 83 108 130-134 138 141 146 149-150 159 182 186-187 190 203 209 234-235 276 283 322 413 415 442 449-450 453 480 484 490 499-500 503 505-506 523 537 542 564 583 602-603 611 619 623 643 650 697 711 729 761 765 770 776-778 784 789 819 822 831 839 862 866 887 904 907 921 935 962- 963 967 973
rectum	Invitrogen	REC001	19 30 33-34 66 108-109 123-124 126 129- 131 143 149 151 156 164 190 201 240 247 250 263 268 274 279 287 295 298- 299 310 314 332 341 354 384 394 401 420 425 442 446 459 483 485 520-521 532 545 559 580-581 584 592 602-607 610 612 615 619 634 637 646 655 664 683-684 741 769 793 822 870 908-911 914-916 934 937-938 942 967 973 982
salivary gland	Clontech	SAL001	16 68 74 84 121 123-124 156 172 190 203 209 232 248 254 269 292 294 363 377 395 398 400 402 405-406 410 430 442 459 462 474 483 485 563-564 579 587- 588 599 602-603 643 658 699 728 730 737 741 748 794 822 867 876 897 903 981

salivary gland	Clontech	SALs03	217 254 270 388 610
skin fibroblast	ATCC	SFB001	517 949
skin fibroblast	ATCC	SFB002	269 688
skin fibroblast	ATCC	SFB003	3 203 897 907
small intestine	Clontech	SIN001	3-4 47 57 68-69 92 99 125-126 130-131 135 149 151-152 156 159 185 204 241 246 291-292 318-319 338 343 348 363 373 375 382 388-389 392-394 397 400 437 466-467 471 484 500 517 520-521 525 547 560 580-581 588 599 602-603 612 624 643 711 731 733-734 757 761 769 774-775 794 824 864 904 906 910- 911 913 948 953 959 976 984
skeletal muscle	Clontech	SKM001	15 75 135 146 172 190 218 267 282 308 410 426-427 474 505 588 620 623 658 692 713 737 779 790 862 874 878 887 952 962-963
skeletal muscle	Clontech	SKMs04	215
spinal cord	Clontech	SPC001	14 20-21 25 28-29 31 39 46 48 59 78 83- 84 91-92 103 112-113 135 160 168 172 176 188 190 205 209 229 232 258 285 301 308 312-314 321 323 329 346 374 377 380 383 388 394 398 406 409-410 431 449-450 453 455 466-467 470-471 484-486 488 495 497 500 503 508-509 524 537 539 558 581 586 604-605 611 619 623 630-631 633 656 663 711 715 729 736 740-741 761 767 769 776-778 780 818 822 831 835-836 840 843 859 861 871 875 887-888 897 906-907 913 919-920 928 931 953 958
adult spleen	Clontech	SPLc01	3 6 12-13 15 156 131 178 365 431 451 461 558 610 715 797 809 876 947 967
stomach	Clontech	STO001	35 114 130-131 144 155 176 189 206-207 249 260-262 336 382 398 425 431 453 461 483 496 500 527 530 580 642 657 663 669 748 765 768 802-803 839 891 942 981
thalamus	Clontech	THA002	30-32 48 66 109 127 130-131 135 142 145 156-158 168 172 174 185 199 224- 225 233 246 277 282 286 293 322 332 334 346 374 384 400 402 420 424 435- 437 446 466-467 485 503 506 527 542 549 572 612 615 622 624 633 643-644 658 676 736 790 794 824 831 835 896 907 950 969
thymus	Clonotech	THM001	10 16 20 28-29 32 37 41 52 57 66-67 74- 75 110 118 121 129-131 141 151 159-160 208 211 218 247 269 289 295 297 320 325 354 358 365 367 372 378 388-389 395 398 411-412 420 423 435 452 500 508-509 517 524 532 537 551 558 560

			569 577-578 582 586 598 608 611 622 643 684 715 721-723 728 740 766 772- 773 795 834 837 849 864 885 900 921 946 948 958 962-963 965 972-973 982
thymus	Clontech	THMc02	1 3 9-11 16 21 27 32-34 38-39 51 55-57 66 72 74 77-78 80 82 89-90 101 112 115 118-119 121 123-124 126 138 144 152 159 168 174 176 178 186-188 197 200 208 212-214 217 225 233 243-244 246 254 256-262 279 282 285 288-289 296- 297 313-314 322 334 343 354-355 358- 359 363-364 367-368 372-373 382 387- 389 395 400 402 411 414 426-427 437 440 442 449-450 454 457 462 464 469 474 479 481 485 490-491 506 508-509 511 517 522 526 528 532 542 551 554 561-562 564 566-570 580-582 585 589 597 599-600 602-608 611 613-614 619- 621 625 628 630-631 644 646 655 669 672 677 684 686-693 697 713 717 720 728 740 746 749 760-762 767 771 775 794 797 804 808 811 816 818-819 837 840 859 880 883 887-888 896-897 903 908-911 913 916 924 936 947-948 950 962-963 965 967 970
thyroid gland	Clontech	THR001	3 8-9 14-15 19-22 28-29 39 41 55-56 66 69 71-72 78-79 97 104-105 109 113 115 119 121 123-124 130-133 135 138 143- 144 146 148 151-152 156 159-163 165 168 172 174 177 183-184 196 199-200 203 209 211 215-218 228-229 232-236 244 254 255 258 273 282 283 282 294 297 303-306 308 311 317-318 322-323 325-326 334-335 340 342 348 354 358 373 377 381-382 387 394 398 401-402 405-406 409-412 416 422 425-427 429- 431 440 449-453 462 466-468 474 478- 479 481-484 490 492-496 500-501 505- 506 517-518 522-525 532 537 540-541 545 551 558 560 563-564 580 583 587- 589 593 597 599 606-607 610 617-621 625-628 633 635 641-643 658-659 664- 669 674 682 686 688-691 696 699 715 724 730 740 742-743 747 750 752 759 761 765-766 768-769 779 789 796 802- 803 813 818-819 822 831 837 843 845 848-849 862 864 868-869 871 874 876- 877 887 893-894 896-897 907-909 912 919-921 923 925 928 936 940-942 944 946-947 950 953 955 958-959 962-963 967 969 973 981
trachea	Clontech	TRC001	33-34 55-56 69 74 163 172 190 209 212

			267 270 297 305 314 352 413 426-427 466-467 500 502 504 580 586 610 613 633 642 688 691 711 724 738-739 774 782 816 820 839 848 862 868-869 914- 915 928 968
uterus	Clontech	UTR001	4 9 18 37 63-64 74 108 114-115 130-131 160 166 179 184 190 209 233 249 269 285 301 314 327 337 348 384 394 399- 400 403 406 411 425 431 434 437 440 462 474 485 490 508-509 526 532 579 617-619 636 642-643 672 761 769 793 837 849 864 887 903 906 928 934 947 967

TABLE 2

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
1	L06175	Homo sapiens	occurs in MHC class I region; ORF	308	98
2	Y70775	Homo sapiens	Follistatin-related protein zfst.	3094	98
3	X15187	Homo sapiens	precursor polypeptide (AA -21 to 782)	4112	100
4	AF110640	Homo sapiens	orphan seven-transmembrane receptor	344	100
5	G03798	Homo sapiens	Human secreted protein, SEQ ID NO: 7879.	158	72
6	W85607	Homo sapiens	Secreted protein clone da228_6.	1477	100
7	Y30162	Homo sapiens	Human dorsal root receptor 4 hDRR4.	884	88
8	Y15227	Homo sapiens	Leu1	391	100
9	Y28817	Homo sapiens	pt326_4 secreted protein.	3338	100
10	X92106	Homo sapiens	bleomycin hydrolase	2445	100
11	Y15228	Homo sapiens	Leu2	445	100
12	U27838	Mus musculus	glycosyl-phosphatidyl-inositol-anchored protein homolog	432	34
13	U27838	Mus musculus	glycosyl-phosphatidyl-inositol-anchored protein homolog	320	27
14	Y71062	Homo sapiens	Human membrane transport protein, MTRP-7.	2323	99
15	U96781	Homo sapiens	Ca2+ ATPase of fast-twitch skeletal muscle sarcoplasmic reticulum, adult isoform	5145	100
16	M16653	Homo sapiens	pancreatic elastase IIB zymogen	1435	99
17	Y13398	Homo sapiens	Amino acid sequence of protein PRO346.	1749	99
18	Y02283	Homo sapiens	Secreted protein clone br342_11 polypeptide sequence.	1399	99
19	Y53030	Homo sapiens	Human secreted protein clone d24_1 protein sequence SEQ ID NO:66.	1371	100
20	AL031320	Homo sapiens	dJ20N2.5 (novel protein similar to fucosidase, alpha-L-1, tissue (EC 3.2.1.51, alpha-l-fucosidase fucosylhydrolase))	2597	99
21	B01384	Homo sapiens	Neuron-associated protein.	1876	100
22	Y68778	Homo sapiens	Amino acid sequence of a human phosphorylation effector PHSP-10.	2470	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
23	Y55935	Homo sapiens	Human KHS2 protein.	4781	99
24	Y55935	Homo sapiens	Human KHS2 protein.	2807	100
25	AC024792	Caenorhabditis elegans	contains similarity to TR:O95029	463	31
26	Y07972	787	Human secreted protein fragment	1540	100
27	X97630	Homo sapiens	serine/threonine protein kinase	3781	98
28	AF150755	Mus musculus	microtubule-actin crosslinking factor	3514	68
29	AF150755	Mus musculus	microtubule-actin crosslinking factor	3725	70
30	Z38011	Mus musculus	DMR-N9	2988	86
31	AJ000522	Homo sapiens	axonemal dynein heavy chain	6058	99
32	AF037256	Mus musculus	ES2 protein	2260	91
33	S62140	Homo sapiens	TLS=nuclear RNA-binding protein	2917	100
34	S62140	Homo sapiens	TLS=nuclear RNA-binding protein	2890	98
36	AB038237	Homo sapiens	G protein-coupled receptor C5L2	1767	100
37	D79994	Homo sapiens	similar to ankyrin of Chromatium vinosum.	6089	99
38	X63380	Homo sapiens	serum response factor-related protein	1966	99
39	AL022072	Schizosaccharomyces pombe	lipoic acid synthetase	1067	61
40	J03930	Homo sapiens	alkaline phosphatase	2751	100
41	AF132968	Homo sapiens	CGI-34 protein	1088	98
42	AL117637	Homo sapiens	hypothetical protein	2208	100
43	AL021393	Homo sapiens	bK747E2.1 (novel protein)	1526	100
44	X68011	Homo sapiens	ZNF81	1886	100
45	AC002464	Homo sapiens	organic cation transporter; 50% similarity to JC4884 (PID:g2143892)	2423	100
46	W78245	Homo sapiens	Fragment of human secreted protein encoded by gene 19.	1949	100
47	Y41765	Homo sapiens	Human PRO1083 protein sequence.	3604	100
48	AF097330	Homo sapiens	H1 chloride channel; p64H1; CLIC4	1305	99
50	U09413	Homo sapiens	zinc finger protein ZNF135	1361	57
51	AF061812	Homo sapiens	keratin 16	2374	100
52	W63681	Homo sapiens	Human secreted protein 1.	1326	99
53	AB035303	Homo sapiens	cadherin-10	4094	100
54	A12022	synthetic construct	MRP-3	403	100
55	AL121897	Homo sapiens	bA392M18.3 (KIAA0180)	1867	100
56	Y73330	Homo sapiens	HTRM clone 397663 protein sequence.	818	96
57	AF151018	Homo sapiens	HSPC184	955	100
58	AF125042	Homo sapiens	bisphosphate 3'-nucleotidase	1586	100
59	AF118670	Homo sapiens	orphan G protein-coupled receptor	1971	100
60	X04494	Homo sapiens	precursor polypeptide	1903	100
61	AF208865	Homo sapiens	EDRF	528	100
62	D15057	Homo sapiens	DAD-1	567	100
63	AF260665	Homo sapiens	histone acetyltransferase	1510	100
64	AF260665	Homo sapiens	histone acetyltransferase	1429	96
65	AJ277145	Homo sapiens	ras-related small GTPase RAB18	1073	100
66	Y94950	Homo sapiens	Human secreted protein clone dh1073_12 protein sequence SEQ ID NO:106.	348	100
67	Y82744	Homo sapiens	DNA replication and repair associated protein (DRASP).	1028	100
68	Y44486	Homo sapiens	Human GPRW receptor polypeptide.	1721	100
69	AL031228	Homo sapiens	dJ1033B10.2 (WD40 protein BING4 (similar to S. cerevisiae YER082C, M. sexta MNG10 and C. elegans F28D1.1))	3196	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
70	AJ276316	Homo sapiens	zinc finger protein 304	1751	52
71	Y18314	Homo sapiens	paraplegin-like protein	4146	99
72	AF157028	Homo sapiens	protein phosphatase methylesterase-1	2017	100
74	Y71082	Homo sapiens	Human B-aggressive lymphoma (BAL) protein.	1765	99
75	AF225420	Homo sapiens	AD025	734	100
76	X95235	Homo sapiens	transcription factor AP2	217	100
77	AF108420	Takifugu rubripes	1-aminocyclopropane-carboxylate synthase	733	56
78	G01349	Homo sapiens	Human secreted protein, SEQ ID NO: 5430.	650	99
79	AL117635	Homo sapiens	hypothetical protein	922	99
81	Z85986	Homo sapiens	dJ108K11.3 (similar to yeast suppressor protein SRP40)	865	77
82	AF183414	Homo sapiens	hemin-sensitive initiation factor 2a kinase	3231	99
83	G01143	Homo sapiens	Human secreted protein, SEQ ID NO: 5224.	495	98
84	U03985	Homo sapiens	N-ethylmaleimide-sensitive factor	3744	99
85	Y17791	Homo sapiens	VAX2 protein	1496	100
87	AF263538	Homo sapiens	growth differentiation factor 3	1944	99
88	Y19757	Homo sapiens	SEQ ID NO 475 from WO9922243.	1361	100
89	AF161493	Homo sapiens	HSPC144	1185	100
90	AF161493	Homo sapiens	HSPC144	856	100
91	B25780	787	Human secreted protein SEQ ID	647	41
92	U57344	Mus musculus	Meis3	1007	89
93	AF172854	Homo sapiens	cardiotrophin-like cytokine CLC	1197	98
94	AL390114	Leishmania major	extremely cysteine/valine rich protein	223	29
95	AB016886	Arabidopsis thaliana	contains similarity to adenylate kinase-gene_id:MCA23.18	287	38
96	AC005525	Homo sapiens	F22162_1	1855	96
97	B20997	Homo sapiens	Human nucleic acid-binding protein, NuABP-1.	3836	99
98	AJ006692	Homo sapiens	high sulfur keratin	507	77
99	AF172264	Homo sapiens	Trat2 and NCK interacting kinase, splice variant 1	694	99
100	L11239	Homo sapiens	homeobox protein	717	100
101	AC004890	Homo sapiens	similar to zinc finger proteins; similar to AAC01956 (PID:g2843171)	2154	98
102	AC003682	Homo sapiens	R28830_2	1287	48
103	AF201839	Rattus norvegicus	dynamamin IIIbb isoform	4270	95
104	Y79510	Homo sapiens	Human carbohydrate-associated protein CRBAP-6.	1394	100
105	Y79510	Homo sapiens	Human carbohydrate-associated protein CRBAP-6.	1209	90
106	AL096748	Homo sapiens	hypothetical protein	1216	100
108	X97260	Homo sapiens	Metallothionein 2	381	100
109	AL034422	Homo sapiens	dJ1141E15.2 (novel protein)	433	100
110	AF191338	Homo sapiens	anaphase-promoting complex subunit 4	683	100
111	AL021712	Arabidopsis thaliana	putative protein	185	26
112	AF250138	Homo sapiens	small stress protein-like protein HSP22	1063	100
113	AL109976	Homo sapiens	dJ79416.1.1 (novel protein)	4176	99
114	Y36151	787	Human secreted protein	668	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
115	AF110399	Homo sapiens	elongation factor Ts	1666	100
116	AF210317	Homo sapiens	facilitative glucose transporter family member GLUT9	2052	99
117	Y73328	Homo sapiens	HTRM clone 082843 protein sequence.	931	100
118	X04085	Homo sapiens	catalase	2846	100
119	AF147717	Homo sapiens	ubiquitin C-terminal hydrolase UCH37	1695	100
120	X73882	Homo sapiens	microtubule associated protein	3801	99
121	AC004882	Homo sapiens	similar to CAA16821 (PID:g3255952)	3223	100
122	M93311	Homo sapiens	metallothionein-III	421	100
123	G03827	Homo sapiens	Human secreted protein, SEQ ID NO: 7908.	557	94
124	G03827	Homo sapiens	Human secreted protein, SEQ ID NO: 7908.	222	53
125	AF232009	Homo sapiens	peroxisomal trans 2-enoyl CoA reductase	1565	99
126	AB004906	Ipomoea purpurea	transposase	146	20
127	M60165	Homo sapiens	guanine nucleotide-binding regulatory protein 2	1832	99
128	Y10319	Homo sapiens	carnitine carrier	1592	100
129	U75467	Drosophila melanogaster	Atu	937	36
130	Z21507	Homo sapiens	human elongation factor-1-delta	494	87
131	Z21507	Homo sapiens	human elongation factor-1-delta	938	100
132	Y58633	Homo sapiens	Protein regulating gene expression PRGE-26.	6745	100
133	Y58633	Homo sapiens	Protein regulating gene expression PRGE-26.	4818	95
134	M13692	Homo sapiens	alpha-1 acid glycoprotein precursor	1064	99
135	U72970	Sus scrofa	calcium/calmodulin-dependent protein kinase II isoform gamma-B	2723	99
136	G32213	Homo sapiens	Human secreted protein, SEQ ID NO: 7294.	450	100
137	AC005102	Homo sapiens	small inducible cytokine subfamily A member 24	627	99
138	AF155343	Homo sapiens	putative zinc finger protein	5855	92
139	AF144638	Homo sapiens	sphingosine-1-phosphate lyase	2977	100
140	AF152318	Homo sapiens	protocadherin gamma A1	4778	100
141	B08517	Homo sapiens	Amino acid sequence of a beta-tubulin antigen.	5841	100
142	X56667	Homo sapiens	calretinin	1410	99
143	X92763	Homo sapiens	tafazzins	1605	100
144	Y95293	Homo sapiens	Human GEF containing NEK-like kinase substrate sGNK.	4092	99
145	AF226046	Homo sapiens	GK003	1198	100
146	M22877	Homo sapiens	cytochrome c	554	98
147	AJ272212	Homo sapiens	protein serine kinase	2196	100
148	AB026491	Homo sapiens	PICK1	2114	98
149	AB018580	Homo sapiens	hluPGFS	1699	100
150	X91868	Homo sapiens	six1	1509	100
151	AF266505	Mus musculus	pseudouridine synthase 3	2135	84
152	U29170	Drosophila melanogaster	ANON-23D	883	43
153	G04075	Homo sapiens	Human secreted protein, SEQ ID NO: 8156.	567	99
154	AY009128	Homo sapiens	ISCU2	138	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
155	AF141315	Homo sapiens	alpha-1,4-N-acetylglucosaminyltransferase	1842	100
156	AF110645	Homo sapiens	candidate tumor suppressor p33 ING1 homolog	1294	99
157	AF159297	Zea mays	extensin-like protein	238	25
158	AL133325	Homo sapiens	dJ984P4.3 (Homeobox protein NKX2B)	1437	100
159	AF073298	Homo sapiens	small EDRK-rich factor 2	294	100
160	AC004858	Homo sapiens	U1 small ribonucleoprotein 1SNRP homolog; match to PID:g4050087	4032	100
161	AB012109	Homo sapiens	APC10	990	100
162	AL162751	Arabidopsis thaliana	putative protein	194	32
163	AJ005698	Homo sapiens	poly(A)-specific ribonuclease	3351	100
164	AF117646	Homo sapiens	long CBL-3 protein	2547	99
165	AC004002	Homo sapiens	similar to ciliary dynein beta heavy chain; 78% Similarity to P23098 (PID:g118965)	5065	100
166	M10942	Homo sapiens	human metallothionein-Ie	381	100
167	AF126484	Homo sapiens	CARD4	4961	100
168	AF161518	Homo sapiens	HSPC169	1604	100
169	M64983	Homo sapiens	fibrinogen beta chain	2482	100
170	M64983	Homo sapiens	fibrinogen beta chain	2679	100
171	M58514	Gallus gallus	fibrinogen beta chain	1059	78
172	AF078845	Homo sapiens	16.7Kd protein	786	100
173	AC004774	Homo sapiens	Dlx-6	923	100
174	Z98974	Schizosaccharomyces pombe	putative vacuolar protein sorting-associated protein	185	31
175	X56203	Plasmodium falciparum	liver stage antigen	283	23
176	W74726	Homo sapiens	Human secreted protein fg949_3.	1879	100
177	AJ222967	Homo sapiens	cystinosin	1920	100
178	AC024796	Caenorhabditis elegans	contains similarity to TR:O76167	221	27
179	Y66632	Homo sapiens	Mechanone-bound protein PRO276.	1370	100
180	AF151803	Homo sapiens	CGI-45 protein	215	28
181	G02694	Homo sapiens	Human secreted protein, SEQ ID NO: 6775.	283	100
182	Y17292	Homo sapiens	Human cell death preventing kinase (DPK-1) protein sequence.	2676	100
183	AF234765	Rattus norvegicus	serine-arginine-rich splicing regulatory protein SRRP86	148	27
184	AF151855	Homo sapiens	CGI-97 protein	1214	96
185	AF289664	Mus musculus	CYLN2	4673	90
186	AL022238	Homo sapiens	dJ1042K10.2 (supported by GENSCAN, FGENES and GENEWISE)	4059	100
187	AL022238	Homo sapiens	dJ1042K10.2 (supported by GENSCAN, FGENES and GENEWISE)	2332	100
188	X83543	Homo sapiens	APXL	8513	99
189	AF059569	Homo sapiens	actin binding protein MAYVEN	3106	99
190	M18135	Rattus norvegicus	smooth-muscle alpha tropomyosin	1306	95
191	AF242194	Drosophila melanogaster	brakeless-B	147	52
192	D30689	Bacillus subtilis	subunit of nitrite reductase	113	29
193	Y44984	Homo sapiens	Human epidermal protein-1.	538	97

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
194	B25679	Homo sapiens	Human secreted protein sequence encoded by gene 15 SEQ ID NO:68.	760	100
195	AB020315	787	homologue of mouse dkk-1 gene:Acc	1466	100
196	U35730	Mus musculus	jerky	2021	75
197	AL136450	Homo sapiens	dJ510O21.1 (novel protein)	632	100
198	X56203	Plasmodium falciparum	liver stage antigen	512	24
199	Y70775	Homo sapiens	Follistatin-related protein zfst.	2027	63
200	X87237	Homo sapiens	a-glucosidase I	4447	99
201	AF101078	Caenorhabditis elegans	CLU-1	1393	46
202	X04571	Homo sapiens	precursor polypeptide (AA -22 to 1185)	6611	100
203	X00474	Homo sapiens	pS2 precursor	466	100
204	AB029333	Halocynthia roretzi	HrPET-1	974	54
205	AF146019	Homo sapiens	hepatocellular carcinoma antigen gene 520	998	100
206	AF071002	Homo sapiens	minK-related peptide 1; MiRP1	632	100
207	AB038162	Homo sapiens	trefoil factor 2	744	100
208	U30521	Homo sapiens	P311 HUM	363	100
209	AB000911	Sus scrofa	ribosomal protein	782	100
210	AB021227	Homo sapiens	membrane-type-5 matrix metalloproteinase	3545	100
211	AF180920	Homo sapiens	cyclin L ania-6a	2722	100
212	AF105365	Homo sapiens	K-CI cotransporter KCC4	5624	100
213	U29244	Caenorhabditis elegans	similar to human (TRE) transforming protein (PIR:S22157)	602	32
214	AL033538	Homo sapiens	dJ477H23.1 (novel protein)	3195	100
215	X52011	Homo sapiens	muscle determination factor	1262	100
216	AF083248	Homo sapiens	ribosomal protein L26 homolog	739	100
217	AF006751	Homo sapiens	ES/130	4793	99
218	AB007859	Homo sapiens	KIAA0399 protein	3559	99
219	AK026291	Homo sapiens	unnamed protein product	826	100
221	Y34045	Homo sapiens	Splice variant of cancer associated polypeptide CH1-9a11-2.	551	97
222	Z67996	Homo sapiens	tenascin-R (restrictin)	7186	100
223	AF134802	Homo sapiens	cofilin isoform 1	345	100
224	Y17711	Homo sapiens	atopy related autoantigen CALC	1611	99
225	AF190051	Gallus gallus	hepatocyte nuclear factor 1a dimerization cofactor isoform	443	81
226	AK026256	Homo sapiens	unnamed protein product	866	98
227	Z69368	Schizosaccharomyces pombe	nuf2-like coiled-coil protein	230	25
228	AF275948	Homo sapiens	ABCA1	11763	99
229	AF161384	Homo sapiens	HSPC266	2006	98
230	Y16270	Homo sapiens	paralemin	1951	100
231	AJ245599	Homo sapiens	putative secreted ligand	2379	99
232	W88499	Homo sapiens	Human stomach carcinoma clone HP10412-encoded protein.	1545	99
233	AF096286	Mus musculus	pecanex 1	3623	93
234	V64619_cd 1	Homo sapiens	30-NOV-1990 Human HE1 cDNA.	796	100
235	V64619_cd 1	Homo sapiens	30-NOV-1990 Human HE1 cDNA.	470	98
236	AF227258	Bos taurus	RPGR-interacting protein-1	1262	38
237	AJ132445	Homo sapiens	claudin-14	1181	100
238	AL034562	Homo sapiens	dJ684O24.2 (prodynorphin (Beta-	1330	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
			Neoeendorphin-Dynorphin precursor, Proenkephalin B precursor))		
239	AF262027	Homo sapiens	eIF-5A2	808	100
240	AL079344	Arabidopsis thaliana	putative protein	194	33
241	AC002394	Homo sapiens	Gene product with similarity to dynein beta subunit	1542	51
242	AJ271361	Takifugu rubripes	FRANK2 protein	303	30
243	AL021918	Homo sapiens	b34I8.1 (Kruppel related Zinc Finger protein 184)	1476	48
244	AF190167	Homo sapiens	membrane associated protein SLP-2	1736	99
245	Y10601	Homo sapiens	ankyrin-like protein	5877	100
246	AL121771	Homo sapiens	dJ548G19.1.1 (novel protein (ortholog of mouse zinc finger protein ZFP64) (translation of cDNA NT2RP3001398 (Em:AK001596)) (isoform 1))	3628	100
247	L25314	Drosophila melanogaster	actin-related protein	984	47
248	X63745	Homo sapiens	KDEL receptor	1095	100
249	AF112208	Homo sapiens	13kDa differentiation-associated protein	816	100
250	AP001707	Homo sapiens	human gene for claudin-8, Accession No. AJ250711	1172	100
251	AL136125	Homo sapiens	dJ304B14.1 (novel protein)	778	100
252	AL031186	Homo sapiens	bK984G1.1 (supported by FGENES)	532	100
253	Y17531	Homo sapiens	Human secreted protein clone BL205 14 protein.	639	100
254	AL049843	Homo sapiens	dJ392M17.3 (KIAA0349 protein)	6741	99
255	AJ242972	Homo sapiens	TOLLIP protein	1424	99
256	Y94873	Homo sapiens	Human protein clone HP02632.	1876	100
257	AF279865	Homo sapiens	kinesin-like protein GAKIN	2903	100
258	AL024498	Homo sapiens	dJ417M14.1 (novel protein)	589	100
259	R66278	Homo sapiens	Therapeutic polypeptide from glioblastoma cell line.	850	100
260	AF101784	Homo sapiens	b-TRCP variant E3RS-IkappaB	3226	99
261	AF101784	Homo sapiens	b-TRCP variant E3RS-IkappaB	2821	100
262	AF101784	Homo sapiens	b-TRCP variant E3RS-IkappaB	3149	99
263	AF197060	Homo sapiens	src homology 3 domain-containing protein HIP-55	2257	100
264	Y86262	Homo sapiens	Human secreted protein HAQAR23, SEQ ID NO:177.	766	100
265	Y56966	Homo sapiens	Human SBPSAPL polypeptide.	2779	100
266	Y56966	Homo sapiens	Human SBPSAPL polypeptide.	1018	99
267	AJ300465	Homo sapiens	putative white family ATP-binding cassette transporter	1557	95
268	AC004030	Homo sapiens	F21856_2	3579	99
269	X55954	Homo sapiens	HL23 ribosomal protein	714	100
270	AB033921	Mus musculus	Ndr1 related protein Ndr2	1855	94
271	AF081886	Homo sapiens	ERO1-like protein	1905	99
272	AF166492	Homo sapiens	small GTPase RAB6B	1060	100
273	AL022238	Homo sapiens	dJ1042K10.4 (novel protein)	2201	100
274	W88667	Homo sapiens	Secreted protein encoded by gene 134 clone HAIBP89.	1530	99
275	X00129	Homo sapiens	precursor RBP	1044	97
276	Z47500_cdl	Homo sapiens	11-MAY-1998 Human RHOH gene sequence.	1161	100
277	AB049188	Equus caballus	ubiquitin C-terminal hydrolase	1118	96

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
278	AF270647	Homo sapiens	GTT1	1564	100
279	AF143956	Mus musculus	coronin-2	2414	94
280	R85151	Homo sapiens	Endothelial cell polypeptide.	911	92
281	R85151	Homo sapiens	Endothelial cell polypeptide.	1031	100
282	D83948	Rattus norvegicus	S1-1 protein	3975	90
283	Y14768	Homo sapiens	I Kappa B-like protein	2037	100
286	AL031316	Homo sapiens	dJ28O10.3(HSD11B1 (hydroxysteroid (11-beta) dehydrogenase 1)	294	100
287	D64109	Homo sapiens	tob family	1773	99
288	AB026043	Homo sapiens	MS4A7	1230	100
289	M61866	Homo sapiens	Krueppel-related DNA-binding protein	209	90
290	AJ001810	Homo sapiens	mRNA cleavage factor I 25 kDa subunit	1217	100
291	Y99454	Homo sapiens	Human PRO1605 (UNQ786) amino acid sequence SEQ ID NO:395.	694	100
292	Y44824	Homo sapiens	Human molecule associated with cell proliferation, MACP-4.	2370	100
293	AJ276101	Homo sapiens	GPRC5B protein	2099	100
294	AF161406	Homo sapiens	HSPC288	719	100
295	Y58628	Homo sapiens	Protein regulating gene expression PRGE-21.	1276	100
296	U91561	Rattus norvegicus	pyridoxine 5'-phosphate oxidase	1239	87
297	L02956	Xenopus laevis	ribonucleoprotein	1624	83
298	AF226730	Homo sapiens	Cyt19	1729	99
299	AF226730	Homo sapiens	Cyt19	906	98
300	Y54324	Homo sapiens	Amino acid sequence of a human gastric cancer antigen protein.	718	89
301	AF125533	Homo sapiens	NADH-cytochrome b5 reductase isoform	1606	100
302	Y32226	Homo sapiens	Human receptor molecule (REC) encoded by Incyte clone 2825826.	1675	87
303	AF247565	Homo sapiens	hepatocellular carcinoma associated ring finger protein	525	100
304	AF208844	Homo sapiens	BM-002	428	100
305	AC004983	Homo sapiens	similar to PID:g3877944	1988	100
306	AL132978	Arabidopsis thaliana	putative protein	210	25
307	Y10530	Homo sapiens	olfactory receptor	1645	100
308	AF180681	Homo sapiens	guanine nucleotide exchange factor	3597	100
309	AF111856	Homo sapiens	sodium dependent phosphate transporter isoform NaPi-3b	3591	99
310	Y13583	Homo sapiens	G-protein coupled receptor	2171	100
311	Z73420	Homo sapiens	cE146D10.2 (mercaptopyruvate sulfurtransferase (EC 2.8.1.2))	1598	100
312	X79535	Homo sapiens	beta tubulin	2348	100
313	AF070658	Homo sapiens	HSPC002	861	100
314	AF078866	Homo sapiens	SURF-4	1395	100
317	Z37986	Homo sapiens	phenylalkylamine binding protein	1258	100
320	AB047892	Macaca fascicularis	hypothetical protein	258	82
321	Y25755	Homo sapiens	Human secreted protein encoded from gene 45.	1440	100
322	AB016531	Homo sapiens	PEX16	1741	100
323	AL391141	Arabidopsis	putative protein	274	49

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
		<i>thaliana</i>			
325	AF140501	<i>Homo sapiens</i>	DNA polymerase iota	3691	99
326	X96698	<i>Homo sapiens</i>	D1075-like	1450	96
327	AF152325	<i>Homo sapiens</i>	protocadherin gamma A5	4769	100
328	AF151803	<i>Homo sapiens</i>	CGI-45 protein	1970	100
329	X74070	<i>Homo sapiens</i>	transcription factor BTF3	639	81
330	AF171102	<i>Homo sapiens</i>	retinal degeneration B beta	1302	95
331	W54040	<i>Homo sapiens</i>	Human interferon-inducible protein, HIF1.	484	98
332	AF024617	<i>Homo sapiens</i>	transcription-associated zinc ribbon protein	691	100
333	U19181	<i>Rattus norvegicus</i>	Rabin3	2129	90
334	G03877	<i>Homo sapiens</i>	Human secreted protein, SEQ ID NO: 7958.	621	100
335	AL008582	<i>Homo sapiens</i>	bK223H9.2 (ortholog of <i>A. thaliana</i> F23F1.8)	626	100
336	AF110774	<i>Homo sapiens</i>	adrenal gland protein AD-001	647	100
337	AB011414	<i>Homo sapiens</i>	Kruppel-type zinc finger protein	1674	58
338	AF207600	<i>Homo sapiens</i>	ethanolamine kinase	129	100
340	AC020579	<i>Arabidopsis thaliana</i>	putative phosphoribosylformylglycinamide synthase; 25509-29950	3283	50
341	Y28576	<i>Homo sapiens</i>	Secreted peptide clone pe503_1.	944	100
342	U32274	<i>Saccharomyces cerevisiae</i>	Ydr386wp; CAI: 0.12	191	37
343	A01771	synthetic construct	vascular anticoagulating protein	1661	99
344	AF220052	<i>Homo sapiens</i>	uncharacterized hematopoietic stem/progenitor cells protein MDS032	1285	100
345	Y70400	<i>Homo sapiens</i>	Human cell-signalling protein-2.	754	100
346	Y50926	<i>Homo sapiens</i>	Human fetal brain cDNA clone vc16_1 derived protein.	962	100
347	AF183423	<i>Homo sapiens</i>	Da protein	1329	100
348	AC006069	<i>Arabidopsis thaliana</i>	putative cleavage and polyadenylation specificity factor	1388	55
349	AL032631	<i>Caenorhabditis elegans</i>	Y106G6H.8	194	39
350	U70669	<i>Homo sapiens</i>	Fas-ligand associated factor 3	167	23
351	Y93468	<i>Homo sapiens</i>	Amino acid sequence of a potassium channel interactor protein.	1182	92
352	AF005856	<i>Drosophila yakuba</i>	anon2A5	111	45
353	AJ271684	<i>Homo sapiens</i>	myeloid DAP12-associating lectin	1013	100
354	AF099100	<i>Homo sapiens</i>	WD-repeat protein 6	2882	99
355	U51730	Murine leukemia virus	reverse transcriptase	316	42
356	D50617	<i>Saccharomyces cerevisiae</i>	YFL042C	279	27
357	D50617	<i>Saccharomyces cerevisiae</i>	YFL042C	279	27
358	AF161432	<i>Homo sapiens</i>	HSPC314	1059	93
359	AB029488	<i>Homo sapiens</i>	C11orf21	758	99
360	AJ251024	<i>Homo sapiens</i>	putative odorant binding protein ag	1239	100
361	U43281	<i>Saccharomyces cerevisiae</i>	Lpg22p	2074	74
362	U43281	<i>Saccharomyces cerevisiae</i>	Lpg22p	2153	74

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
363	AC007153	Arabidopsis thaliana	100632	156	24
364	AF197927	Homo sapiens	AF5q31 protein	3992	99
365	D28500	Homo sapiens	mitochondrial isoleucine tRNA synthetase	4286	98
366	X97868	Homo sapiens	arylsulphatase	3141	98
367	AL162048	Homo sapiens	hypothetical protein	1532	100
368	L36062	Mus musculus	steroidogenic acute regulatory protein	189	25
369	AF113249	Homo sapiens	multiple domain putative nuclear protein	1022	59
370	M15888	Bos taurus	endozepine-related protein precursor	2425	84
371	X66363	Homo sapiens	serine/threonine protein kinase	2562	100
372	W74802	Homo sapiens	Human secreted protein encoded by gene 73 clone HSQEL25.	1532	89
373	AF100772	Homo sapiens	tenascin-M1	11535	99
374	AF090934	Homo sapiens	PRO0518	382	100
375	AB021643	Homo sapiens	gonadotropin inducible transcription repressor-3	2761	99
376	AB049758	Homo sapiens	MAWD binding protein	1331	100
377	AF070666	Homo sapiens	Kruppel-associated box protein	466	97
378	S59342	Mus sp.	nuclear pore complex glycoprotein p62	464	60
379	AF149205	Mus musculus	Su(var)3-9 homolog Suv39h2	1690	88
380	AF227906	Homo sapiens	UDP-glucose:glycoprotein glucosyltransferase 2 precursor	7851	99
381	AF118566	Mus musculus	hematopoietic zinc finger protein	1769	92
382	AK000619	Homo sapiens	unnamed protein product	810	100
383	AF227906	Homo sapiens	UDP-glucose:glycoprotein glucosyltransferase 2 precursor	7851	99
384	AF117946	Homo sapiens	Link guanine nucleotide exchange factor II	2363	100
385	AF125390	Drosophila melanogaster	L82G	139	41
386	Y34037	Homo sapiens	Human secreted protein clone ca106_19x protein sequence SEQ ID NO:20.	1092	50
387	U18795	Saccharomyces cerevisiae	Yel064cp	206	28
388	AF177388	Homo sapiens	cancer-amplified transcriptional coactivator ASC-2	10748	99
389	AJ002744	Homo sapiens	UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 7	3469	96
390	AF097366	Homo sapiens	cone sodium-calcium potassium exchanger	3166	100
391	AF217525	Homo sapiens	Down syndrome cell adhesion molecule	5337	60
392	U81035	Rattus norvegicus	ankyrin binding cell adhesion molecule neurofascin	3967	91
393	X65224	Gallus gallus	neurofascin	4097	78
394	X13916	Homo sapiens	LDL-receptor related precursor (AA -19 to 4525)	4292	99
395	AF151083	Homo sapiens	HSPC249	444	98
396	AB017026	Mus musculus	oxysterol-binding protein	2173	98
397	AL035587	Homo sapiens	dJ475N16.4 (KIAA0240)	2393	100
398	W74813	Homo sapiens	Human secreted protein encoded by gene 85 clone HSDFV29.	722	92
399	Y71110	Homo sapiens	Human Hydrolase protein-8 (HYDRL-8).	1637	99

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
400	AF039718	Caenorhabditis elegans	contains similarity to lupus LA protein homologs	325	43
401	AE000877	Methanothermobacter thermoautotrophicus	conserved protein	231	36
402	Y27795	Homo sapiens	Human secreted protein encoded by gene No. 79.	1539	99
403	Z50853	Homo sapiens	CLPP	615	100
405	X03475	Rattus norvegicus	ribosomal protein L35a (aa 1-110)	576	99
406	AF144237	Homo sapiens	LOMP protein	252	44
407	U20239	Mus musculus	fibrosin	288	76
409	AL033378	Homo sapiens	dJ323M4.1 (KIAA0790 protein)	6026	99
410	X54326	Homo sapiens	glutamyl-tRNA synthetase	7577	99
411	X61585	Bos taurus	polynucleotide adenylyltransferase	3715	97
412	AF217190	Homo sapiens	MLEL1 protein	5271	99
414	G02815	Homo sapiens	Human secreted protein, SEQ ID NO: 6896.	314	95
415	AJ245922	Homo sapiens	alpha-tubulin 8	2370	100
416	AF203032	Homo sapiens	neurofilament protein	220	21
417	Z97653	Homo sapiens	c380A1.2.1 (novel protein (isoform 1))	1567	100
418	AJ404326	Homo sapiens	SR+89	1871	99
419	AJ404326	Homo sapiens	SR+89	902	64
420	AF134726	Homo sapiens	G9A	5334	99
421	L28125	Podospira anserina	beta transducin-like protein	288	39
422	W21733	Homo sapiens	NIP-1 encoded by clone 59.	110	72
423	S67970	Homo sapiens	ZNF75=KRAB zinc finger	951	76
424	L28035	Mus musculus	protein kinase C gamma	3768	98
426	Y73373	Homo sapiens	HTRM clone 921803 protein sequence.	555	56
427	Y73373	Homo sapiens	HTRM clone 921803 protein sequence.	266	49
428	X61118	Homo sapiens	TTC-2a/RBTN-2a	876	100
429	Z96932	Homo sapiens	nuclear autoantigen fo 14 kDa	496	83
430	AJ277291	Homo sapiens	HELG protein	678	72
431	X82157	Homo sapiens	hevin	3525	99
432	AC007192	Homo sapiens	P85B_HUMAN; PTDINS-3-KINASE P85-BETA	3825	99
433	AL021918	Homo sapiens	b34I8.1 (Kruppel related Zinc Finger protein 184)	1713	50
434	AF084464	Rattus norvegicus	GTP-binding protein REM2	141	29
435	AL049795	Homo sapiens	dJ622L5.2 (novel protein)	1756	98
436	M14513	Rattus norvegicus	(Na ⁺ and K ⁺) ATPase, alpha(III) catalytic subunit	4269	99
437	U33460	Homo sapiens	DNA-directed RNA polymerase I, largest subunit	8777	98
438	D87076	Homo sapiens	similar to human bromodomain protein BR140(JC2069)	3067	100
439	L43912	Macaca mulatta	mannose-binding protein A	589	93
440	D31763	Homo sapiens	ha0946 protein is Kruppel-related.	927	49
441	U70976	Homo sapiens	arrestin	2068	99
442	B08069	Homo sapiens	A human beta-alanine-pyruvate aminotransferase (HAPA).	2343	99
443	AF100662	Caenorhabditis	contains similarity to ubiquitin	166	24

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
		elegans	carboxyl-terminal hydrolase (Pfam: UCH-1.hmm, score: 28.46) (Pfam: UCH-2.hmm, score: 47.53)		
444	D78017	Rattus norvegicus	NFI-A1	2667	98
445	AL049569	Homo sapiens	dJ37C10.3 (novel ATPase)	2418	100
448	AJ242540	Volvox carteri f. nagariensis	hydroxyproline-rich glycoprotein DZ-HRGP	165	34
449	AJ133352	Homo sapiens	ZNF237 protein	2006	100
450	AJ133352	Homo sapiens	ZNF237 protein	1025	96
451	AF170708	Homo sapiens	T-box protein TBX3	3700	99
452	AK002080	Homo sapiens	unnamed protein product	1546	99
453	L32977	Homo sapiens	Rieske Fe-S protein	1239	93
454	X51760	Homo sapiens	zinc finger protein (583 AA)	1533	57
455	Y01141	Homo sapiens	Secreted protein encoded by gene 7 clone HTLFA90.	1453	99
456	AB006631	Homo sapiens	The human homolog of mouse Cux-2	6559	100
457	AF067165	Homo sapiens	zinc finger protein 3	977	64
458	AF038169	Homo sapiens	unknown	154	38
459	W75214	Homo sapiens	Human secreted protein encoded by gene 19 clone HRSMC69.	1180	95
460	U97002	Caenorhabditis elegans	similar to acyl-CoA dehydrogenases and epoxide hydrolases; Pfam domain PF00441 (Acyl-CoA dh), Score=57.4, E-value=1.7e-16, N=2; contains similarity to Pfam domain PF00702 (Hydrolase), Score=57.4, E-value=1e-13, N=1	583	37
461	AK023114	Homo sapiens	unnamed protein product	1041	99
462	M93134	Friend murine leukemia virus	pol protein	289	44
463	AF055473	Homo sapiens	GAGE-8	232	47
466	Y51415	Homo sapiens	Human wild type pKe83 protein.	2625	100
467	Y51417	787	Human pKe83 splice variant protein	2433	100
468	V57936	Homo sapiens	Human transmembrane protein HTMPN-60.	1629	96
469	D38552	Homo sapiens	The ha1539 protein is related to cyclophilin.	2995	100
470	Y70013	Homo sapiens	Human Protease and associated protein-7 (PPRG-7).	3530	100
471	AJ224747	Homo sapiens	C-terminal variant of hINADL including 2 amino acid exchanges and an insertion of 28 amino acids in frame.	7969	100
472	W99665	Homo sapiens	Human secreted protein clone du157_12 protein.	1546	100
473	W99665	Homo sapiens	Human secreted protein clone du157_12 protein.	998	98
474	X63526	Homo sapiens	homologue to elongation factor 1-gamma from A.salina	2273	99
475	X15940	Homo sapiens	ribosomal protein L31 (AA 1-125)	644	100
476	M60832	Homo sapiens	alpha-2 type VIII collagen	3581	99
477	AF039697	Homo sapiens	antigen NY-CO-31	1213	97
478	AF156929	Sus scrofa	inflammatory response protein 6	1588	83
479	AF264717	Homo sapiens	FYVE domain-containing dual specificity protein phosphatase FYVE-DSP2	5610	99
480	AF044578	Homo sapiens	putative DNA polymerase; POL4P	2478	94
481	X89750	Homo sapiens	TGIF protein	1413	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
482	M93107	Homo sapiens	(R)-3-hydroxybutyrate dehydrogenase	1663	96
483	U58334	Homo sapiens	Bbp/53BP2	1556	41
484	AF151538	Homo sapiens	deoxycytidyl transferase; Rev1p	4281	99
485	Z98884	Homo sapiens	dJ467L1.1 (KIAA0833)	699	73
486	AJ243874	Homo sapiens	oligophrenin-4	3682	100
487	Z11737	Homo sapiens	flavin-containing monooxygenase 4	2969	100
488	X56123	Mus musculus	talin	4353	77
489	AJ278112	Homo sapiens	putative cell cycle control protein	335	23
490	W74843	Homo sapiens	Human secreted protein encoded by gene 115 clone HOVBA03.	1013	98
491	Y41337	Homo sapiens	Human secreted protein encoded by gene 30 clone HRDDV47.	509	36
492	X90530	Homo sapiens	ragB	1926	99
493	X90530	Homo sapiens	ragB	1405	99
494	X90530	Homo sapiens	ragB	1893	96
495	AL022394	Homo sapiens	dJ511B24.3 (KIAA0395 (probable homeobox protein))	4990	99
496	Y11395	Homo sapiens	lanthionine synthetase C-like protein 1	2168	100
497	AJ010119	Homo sapiens	Ribosomal protein kinase B (RSK-B)	4001	100
498	G01563	Homo sapiens	Human secreted protein, SEQ ID NO: 5644.	330	100
499	X54131	Homo sapiens	protein-tyrosine phosphatase	10465	99
500	G01082	Homo sapiens	Human secreted protein, SEQ ID NO: 5163.	549	100
501	AC004142	Homo sapiens	similar to murine leucine-rich repeat protein; possible role in neural development by protein-protein interactions; 93% similarity to D49802 (PID:g1369906)	3676	100
502	AL117544	Homo sapiens	hypothetical protein	1226	100
503	AF203032	Homo sapiens	neurofilament protein	5115	99
504	AL034417	Homo sapiens	bK215D11.2 (similar to rat gene 33)	2476	100
505	X69090	Homo sapiens	190kD protein	7343	99
506	U58755	Caenorhabditis elegans	coded for by C. elegans cDNA yk34b1.5; coded for by C. elegans cDNA yk13f10.5; coded for by C. elegans cDNA yk46e8.5; coded for by C. elegans cDNA yk46d5.5; coded for by C. elegans cDNA yk43c2.5; coded for by C. elegans cDNA yk46e8.3; coded for by C. elegans cDNA yk43c2.3; coded for by C. elegans cDNA yk46d5.3; coded for by C. elegans cDNA yk13f10.3; coded for by C. elegans cDNA yk34b1.3	782	55
507	AJ293309	Homo sapiens	NHP2 protein	801	100
508	U39045	Rattus norvegicus	cytoplasmic dynein intermediate chain 2B	3241	97
509	AF063231	Mus musculus	cytoplasmic dynein intermediate chain 2	3159	97
510	AF202893	Mus musculus	Kif21b	4336	95
511	Y13115	Homo sapiens	serine/threonine protein kinase	5071	99
512	AB030207	Homo sapiens	G gamma subunit	364	100
513	AF039571	Homo sapiens	peripheral benzodiazepine receptor interacting protein; PBR-IP/PRAX1	495	33
514	AB037883	Homo sapiens	Gb3/CD77 synthase	1916	99

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
515	D90868	Escherichia coli	similar to	1489	100
516	X98834	Homo sapiens	zinc finger protein Hsal2	5290	100
517	AF055668	Mus musculus	apoptosis-linked gene 4, deltaC form	2904	78
518	AF019926	Mus musculus	protein kinase	1694	90
519	M34513	Homo sapiens	omega protein	317	91
520	Y08612	Homo sapiens	88kDa nuclear pore complex protein	2313	99
521	Y08612	Homo sapiens	88kDa nuclear pore complex protein	1561	99
522	AL096766	Homo sapiens	dA59H18.1 (KIAA0767 protein)	2497	100
523	AF186249	Homo sapiens	six transmembrane epithelial antigen of prostate	1790	100
524	AB029012	Homo sapiens	KIAA1089 protein	4933	100
525	AB026893	Homo sapiens	vascular cadherin-2	5962	100
526	X74331	Homo sapiens	DNA primase (p58 subunit)	1720	100
528	AC007228	Homo sapiens	R31665_2	1488	47
529	X14830	Homo sapiens	acetylcholine receptor beta-subunit preprotein	2639	100
530	U80446	Caenorhabditis elegans	coded for by C. elegans cDNA yk172e6.3; coded for by C. elegans cDNA yk158f7.3; coded for by C. elegans cDNA yk158f7.5; coded for by C. elegans cDNA yk172e6.5	420	39
531	S76838	Mus sp.	Dbs	4821	88
532	Z82215	Homo sapiens	dJ68O2.2 (myosin, heavy polypeptide 9, non-muscle)	9828	100
533	AF245505	Homo sapiens	adlcan	277	31
534	AF300612	Homo sapiens	N-acetylgalactosamine-4-O-sulfotransferase	993	59
535	AL121928	Homo sapiens	bA18I14.3 (pleckstrin and Sec7 domain protein)	3333	99
536	AJ271055	Mus musculus	iroquois homeobox protein 6	1724	76
537	AF180473	Homo sapiens	Not2p	2267	100
538	AF071059	Mus musculus	zinc finger RNA binding protein	1089	51
539	AF023453	Homo sapiens	actin-related protein 3-beta	2219	100
540	AC005030	Homo sapiens	R29828_1	1461	70
541	AC003030	Homo sapiens	R29828_1	2294	100
542	AL121889	Homo sapiens	dJ1076E17.1 (KIAA0823 protein (continues in AL023803))	2152	100
543	AB006135	Rattus norvegicus	db83	1238	98
544	G02650	Homo sapiens	Human secreted protein, SEQ ID NO: 6731.	644	97
545	Y07595	Homo sapiens	transcription factor TFIIH	2373	100
546	AL133545	Homo sapiens	bA386N14.1 (novel protein similar to a dual specificity phosphatase)	964	99
547	X83618	Homo sapiens	hydroxymethylglutaryl-CoA synthase	2647	100
548	AF134726	Homo sapiens	NG37	4359	99
549	AB035356	Homo sapiens	neurexin I-alpha protein	6948	99
551	AB037901	Homo sapiens	gene amplified in squamous cell carcinoma-1	5215	99
552	AB043634	Homo sapiens	PAR-6A	885	100
553	AP000693	Homo sapiens	partial CDS	4875	99
554	AF002223	Homo sapiens	myotubularin related 1	3490	100
555	AC004893	Homo sapiens	similar to NEDD-4 (KIA0093); similar to P46934 (PID:g1171682)	1611	100
556	AJ404468	Homo sapiens	axonemal dynein heavy chain	8328	100
557	AJ404468	Homo sapiens	axonemal dynein heavy chain	11137	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
558	X65873	Homo sapiens	kinesin heavy chain	4860	100
559	AJ277365	Homo sapiens	polyglutamine-containing protein	592	36
560	AF205600	Homo sapiens	transposase-like protein	407	27
561	X71125	Homo sapiens	glutaminyl-peptide cyclotransferase	1914	100
562	X71125	Homo sapiens	glutaminyl-peptide cyclotransferase	1456	97
563	X54304	Homo sapiens	myosin regulatory light chain	897	100
564	AF250842	Drosophila melanogaster	multiple asters	130	23
565	Y58608	Homo sapiens	Protein regulating gene expression PRGE-1.	1619	99
566	AL121893	Homo sapiens	bA189K21.5 (novel protein similar to retinoblastoma binding protein (RBBP9))	1012	100
567	AL117352	Homo sapiens	dJ876B10.2 (novel protein (ortholog of rat EXO84))	3713	99
568	AF228603	Homo sapiens	pleckstrin 2	1841	100
569	AF239243	Homo sapiens	histone deacetylase 7	3244	86
570	AF087695	Mus musculus	veli 3	989	100
571	AB046381	Homo sapiens	testis-abundant finger protein	1346	99
572	AC005551	Homo sapiens	R26529_2, partial CDS	1020	100
573	Y90290	Homo sapiens	Human peptidase, HPEP-7 protein sequence.	274	52
574	W76734	Homo sapiens	Human mDia Rho targeting protein.	712	32
575	AL121935	Homo sapiens	bA517H2.3 (t-complex 10 (a murine tcp homolog))	853	78
576	Y86217	Homo sapiens	Human secreted protein HWHGU54, SEQ ID NO:132.	2123	99
577	AL121716	Homo sapiens	dJ202D23.2 (novel protein)	6329	99
578	AL121716	Homo sapiens	dJ202D23.2 (novel protein)	6329	99
579	X92715	Homo sapiens	KRAB/C2H2 zinc finger protein	3102	97
580	X54637	Homo sapiens	protein tyrosine kinase	5564	98
581	X78817	Homo sapiens	p115	1148	44
582	AJ251245	Rattus norvegicus	SECIS binding protein 2	3086	71
583	AF113125	Homo sapiens	E-1 enzyme	591	100
584	M19529	Sus scrofa	folliculin A	1936	98
585	AF169677	Homo sapiens	leucine-rich repeat transmembrane protein FLRT3	3403	100
586	D87635	Homo sapiens	similar to human transcription factor TFIIS (S34159).	8083	99
587	Y00876	Homo sapiens	Human LAPH-1 protein sequence.	2110	100
588	Y99674	Homo sapiens	Human GTPase associated protein-25.	2111	99
589	D86973	Homo sapiens	similar to Yeast translation activator GCN1 (P1:A48126)	12033	99
590	AL034452	Homo sapiens	dJ682J15.1 (novel Collagen triple helix repeat containing protein)	1979	100
591	Y57396	Homo sapiens	Human lysoenzyme LYC4 polypeptide.	814	100
592	AJ297743	Mus musculus	torsinB protein	1448	85
593	AF164796	Homo sapiens	NADH:ubiquinone oxidoreductase MLRQ subunit homolog	469	100
594	Y41312	Homo sapiens	Human secreted protein encoded by gene 5 clone HLDRM43.	749	94
595	Y41312	Homo sapiens	Human secreted protein encoded by gene 5 clone HLDRM43.	824	100
596	Y77123	Homo sapiens	Human neurotransmission-associated protein (NTAP) 998868.	2102	98
597	AF215703	Drosophila	KISMET-L long isoform	1880	65

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
		melanogaster			
598	AF070447	Homo sapiens	barrier-to-autointegration factor	290	90
599	X56203	Plasmodium falciparum	liver stage antigen	372	22
600	X79828	Mus musculus	NK10	202	53
601	AB004109	Cricetus griseus	phosphatidylserine synthase II	2262	92
602	U94988	Mus musculus	Nulp1	2912	89
603	U94988	Mus musculus	Nulp1	2800	86
604	AF006264	Homo sapiens	recombination and sister chromatid cohesion protein homolog	2850	100
605	AF006264	Homo sapiens	recombination and sister chromatid cohesion protein homolog	2530	100
606	X82260	Homo sapiens	RanGAP1	2929	100
607	X82260	Homo sapiens	RanGAP1	1843	97
608	AF160909	Drosophila melanogaster	BcDNA.LD03471	943	58
610	X74801	Homo sapiens	gamma subunit of CCT chaperonin	2745	99
611	AL031427	Homo sapiens	dJ167A19.1 (novel protein)	1608	100
612	Y71072	Homo sapiens	Human membrane transport protein, MTRP-17.	445	100
613	X16396	Homo sapiens	precursor polypeptide (AA -29 to 315)	1749	100
614	AK000281	Homo sapiens	unnamed protein product	1814	99
615	AB011128	Homo sapiens	KIAA0556 protein	5761	99
616	U19361	Petromyzon marinus	NF-180	205	21
617	AF045555	Homo sapiens	wbscr1	1208	100
618	AF045555	Homo sapiens	wbscr1 alternative spliced product	1318	100
619	U22229	Felis catus	ribosomal protein L41	128	100
620	Y17169	Homo sapiens	A6 related protein	1819	100
621	Y12065	Homo sapiens	hNop56	2956	99
622	AF177758	Homo sapiens	ubiquitin specific protease 16	2998	100
623	AF317425	Homo sapiens	GAC-1	3866	100
624	AL050297	Homo sapiens	hypothetical protein	1227	99
625	AC007204	Homo sapiens	BC273239	3398	99
626	Z68747	Homo sapiens	imogen 38	2024	99
627	Z68747	Homo sapiens	imogen 38	1958	97
628	Y70229	Homo sapiens	Human RNA-associated protein-10 (RNAAP-10).	3424	99
629	AF191492	Homo sapiens	nasopharyngeal carcinoma associated gene protein-8	613	100
630	AF119664	Homo sapiens	transcriptional regulator protein HCNGP	1574	100
631	AF119664	Homo sapiens	transcriptional regulator protein HCNGP	1150	89
632	Y17849	Homo sapiens	ganglioside-induced differentiation associated protein 1	1839	98
633	X55740	Homo sapiens	5'-nucleotidase	3012	100
634	AF039688	Homo sapiens	antigen NY-CO-3	931	100
635	AF119662	Homo sapiens	E46 protein	2424	100
636	AB007836	Homo sapiens	Hic-5	2544	100
637	AF077818	Mus musculus	syntrophin-associated serine-threonine protein kinase	2027	44
638	AL035455	Homo sapiens	dJ1018E9.1 (VAMP (vesicle-associated membrane protein)-associated protein B and C)	150	26
639	AF078844	Homo sapiens	hqp0376 protein	416	81

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
640	U28377	Escherichia coli	ORF_f239; was ORF_f191 and ORF_f194 before splice	1198	100
641	AK024442	Homo sapiens	FLJ00032 protein	1677	56
642	U58682	Homo sapiens	ribosomal protein S28	340	100
643	X57432	Rattus rattus	ribosomal protein S2	1520	98
644	AB002348	Homo sapiens	KIAA0350 protein	5186	99
646	Y96202	Homo sapiens	IkappaB kinase (IKK) binding protein, Y2H56.	1178	98
647	AB029482	Mus musculus	JNK-binding protein JNKBP1	4609	81
648	AB009053	Arabidopsis thaliana	contains similarity to isoamyl acetate-hydrolyzing esterase-gene_id:MQB2.25	407	44
650	AC002550	Homo sapiens	Unknown gene product	858	99
651	U26592	Homo sapiens	diabetes mellitus type I autoantigen	253	66
652	X60155	Homo sapiens	zinc finger 41	4349	100
653	X53330	Platynereis dumerilii	H4 protein (AA 1 - 103)	523	100
654	AC003682	Homo sapiens	R27945_2	2558	100
655	X80473	Mus musculus	rab19	596	56
656	J02649	Rattus norvegicus	unknown protein	201	95
657	AC006014	Homo sapiens	similar to RFP transforming protein; similar to P14373 (PID:g132517)	1331	99
658	X92972	Homo sapiens	protein phosphatase 6	1666	100
659	L35269	Homo sapiens	zinc finger protein	2803	99
660	AC003682	Homo sapiens	F18547_1	3184	96
661	X79204	Homo sapiens	ataxin-1	4195	99
662	X17620	Homo sapiens	Nm23 protein	965	99
663	AB015617	Homo sapiens	ELKS	1501	80
664	Z56281	Homo sapiens	interferon regulatory factor 3	2331	100
665	AJ248283	Pyrococcus abyssi	LACTOYLGLUTATHIONE LYASE (EC 4.4.1.5) METHYLGLYOXALASE) (ALDOKETOMUTASE) (GLYOXALASE D).	254	40
666	Z70200	Homo sapiens	U5 snRNP-specific 200kD protein	8819	99
667	Z70200	Homo sapiens	U5 snRNP-specific 200kD protein	8589	97
668	AF153450	Manduca sexta	juvenile hormone esterase binding protein	225	32
669	AF227198	Homo sapiens	CrkRS	7231	99
670	X99586	Homo sapiens	SMT3C protein	441	87
671	Z61589_cd1	Homo sapiens	17-AUG-1998 DNA encoding a human OC-2 protein.	2593	100
672	AJ132702	Mus musculus	ATFa-associated factor	3240	88
673	AF204159	Homo sapiens	potassium large conductance calcium-activated channel beta 3a subunit	1486	100
674	G02061	Homo sapiens	Human secreted protein, SEQ ID NO: 6142.	558	99
675	G01246	Homo sapiens	Human secreted protein, SEQ ID NO: 5327.	141	77
676	AB016839	Homo sapiens	mob1	419	42
677	D86970	Homo sapiens	similar to myosin heavy chain: Containing ATP/GTP-binding site motif A(P-loop)	161	28
678	U83115	Homo sapiens	non-lens beta gamma-crystallin like protein	8569	99
679	AF203687	Homo sapiens	prolactin regulatory element-binding protein	2181	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
680	M27685	Mus musculus	ultra-high sulphur keratin	650	58
681	U04968	Cricetulus griseus	nucleotide excision repair protein	3712	97
682	AF119663	Homo sapiens	G-protein gamma-12 subunit	356	100
683	G03733	Homo sapiens	Human secreted protein, SEQ ID NO: 7814.	342	100
684	X67699	Homo sapiens	CDw52 antigen	297	100
685	AF022789	Homo sapiens	ubiquitin hydrolyzing enzyme I	1892	100
686	AJ001006	Mus musculus	EMeg32 protein	938	96
687	W03516	Homo sapiens	Prostaglandin DP receptor.	1864	100
688	AF019661	Mus musculus	zeta proteasome chain; PSMA5	1214	100
689	AF156557	Homo sapiens	stomatin related protein	2036	100
690	G03960	Homo sapiens	Human secreted protein, SEQ ID NO: 8041.	593	100
691	AF161512	Homo sapiens	HSPC163	738	100
692	AL031115	Homo sapiens	ZXDA, ZXDB (zinc finger X-linked protein)	4298	100
693	L40410	Homo sapiens	thyroid receptor interactor	806	100
694	AC004542	Homo sapiens	OXYSTEROL-BINDING PROTEIN-like; similar to P22059 (PID:g129308)	2533	99
695	AF169411	Rattus norvegicus	PAPIN	4144	52
696	Y58168	Homo sapiens	Human hydrolase homologue HHH-4.	2144	100
697	AF271994	Homo sapiens	dopamine responsive protein DRG-1	1613	100
698	Y41741	Homo sapiens	Human PRO704 protein sequence.	1323	100
699	AL133506	Unknown	/prediction=(method:"genscan", version:"1.0", score:"109.13"); /prediction=(method:	825	48
700	Y96870	Homo sapiens	Human goose-type lysozyme (GOLY).	1032	100
701	AC003034	Homo sapiens	Gene with similarity to rat kidney-specific (KS) gene	1190	100
702	AC003034	Homo sapiens	Gene with similarity to rat kidney-specific (KS) gene	337	95
703	AJ142832	Homo sapiens	calpain	3756	100
704	S52624	Homo sapiens	unknown	185	100
705	AF005081	Homo sapiens	skin-specific protein	652	100
706	Y16793	Homo sapiens	keratin, type I	2232	100
707	Y44985	Homo sapiens	Human epidermal protein-2.	455	69
708	AF113220	Homo sapiens	MSTP040	686	100
709	Y44985	Homo sapiens	Human epidermal protein-2.	408	65
710	Y16132	Homo sapiens	CDT6	1874	100
711	Y68775	Homo sapiens	Amino acid sequence of a human phosphorylation effector PHSP-7.	2407	100
712	X63422	Homo sapiens	H(+)-transporting ATP synthase	209	100
713	AF169968	Mus musculus	DNA binding protein DESRT	1467	79
714	X52563	Bos taurus	permeability increasing protein	383	29
715	AJ277739	Homo sapiens	RPB11b1alpha protein	480	98
716	AL135791	Homo sapiens	bA162G10.3 (zinc finger protein)	401	98
717	AF223466	Homo sapiens	HT015 protein	1311	97
719	AF117383	Homo sapiens	placental protein 13; PP13	746	100
720	Z98743	Homo sapiens	dJ181C9.2 (Rho GTPase activating protein 8 (RhoGAP, p50RhoGAP))	324	100
721	AL163815	Arabidopsis thaliana	putative protein	653	61
722	G01436	Homo sapiens	Human secreted protein, SEQ ID	418	96

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
			NO: 5517.		
723	AF282919	Mus musculus	Zfp228	349	49
724	AB023191	Homo sapiens	KIAA0974 protein	2953	100
725	AL031778	Homo sapiens	dJ34B21.1 (novel BZRP (benzodiazapine receptor (peripheral) (MBR, PBR, PBKS, IBP, Isoquinoline-binding protein)) LIKE protein)	920	100
726	AL021939	Homo sapiens	dJ352A20.2 (aldehyde dehydrogenase family protein)	1764	100
727	AF182426	Rattus norvegicus	arylacetamide deacetylase	791	42
728	Y08565	Homo sapiens	UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase	3331	99
729	AF155135	Homo sapiens	novel retinal pigment epithelial cell protein	1652	99
730	AL078606	Arabidopsis thaliana	putative protein	277	55
731	Y73352	Homo sapiens	HTRM clone 1732368 protein sequence.	1720	100
732	AF178432	Homo sapiens	SH3 protein	3302	100
733	Y17832	Human endogenous retrovirus K	env protein	223	34
734	Y28859	Homo sapiens	Human mesoderm induction early response protein ER1.	2067	98
735	U09355	Oryctolagus cuniculus	protein phosphatase 2A1 B gamma subunit	2352	99
736	Y94922	Homo sapiens	Human secreted protein clone pv6_1 protein sequence SEQ ID NO:50.	724	99
737	AB027003	Mus musculus	protein phosphatase	378	84
738	AF112200	Homo sapiens	NADH-oxidoreductase B18 subunit	739	100
739	AF112200	Homo sapiens	NADH-oxidoreductase B18 subunit	613	88
740	AF302154	Homo sapiens	SPG protein	6556	100
741	B256F1	Homo sapiens	Human secreted protein sequence encoded by gene 17 SEQ ID NO:70.	1410	99
742	L27479	Homo sapiens	X123	1237	99
743	L27479	Homo sapiens	X123	1206	97
744	Y66745	Homo sapiens	Membrane-bound protein PRO1135.	588	99
745	AJ001019	Homo sapiens	ring finger protein	1292	99
746	X68453	Sus scrofa	tubulin-tyrosine ligase	1882	94
747	Y57897	Homo sapiens	Human transmembrane protein HTMPN-21.	1173	100
748	AF151069	Homo sapiens	HSPC235	1694	96
749	AF182404	Homo sapiens	mitochondrial uncoupling protein 1	1674	100
750	AL121993	Homo sapiens	dJ776P7.1 (Novel protein)	2500	99
751	AF149825	Homo sapiens	PACSN3	2253	100
752	AL008635	Homo sapiens	dJ510H16.2 (high-mobility group protein 2-like 1)	3026	99
753	Y57914	Homo sapiens	Human transmembrane protein HTMPN-38.	1124	100
754	AF285109	Homo sapiens	septin 3 isoform B	1766	100
755	AF004161	Oryctolagus cuniculus	peroxisomal Ca-dependent solute carrier	2371	95
756	Z19585	Homo sapiens	thrombospondin-4	4239	100
757	AP001745	Homo sapiens	similar to zinc finger 5 protein	1857	100
758	AF190664	Mus musculus	LMBR2	555	72
759	AF090326	Mus musculus	AE-1 binding protein AEBP2	1540	97
760	AL096677	Homo sapiens	dJ322G13.3 (novel protein similar to	999	94

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
			bovine and mouse beta-soluble NSF attachment protein (SNAP-beta))		
761	AC003007	Homo sapiens	Unknown gene product (partial)	649	96
762	U66372	Bos taurus	ribosomal protein S29	230	73
764	Y90899	Homo sapiens	D1-like dopamine receptor activity modifying protein SEQ ID NO:1.	1152	100
765	U88169	Caenorhabditis elegans	similar to molybdopterin biosynthesis MOEB proteins	1204	65
766	AL118506	Homo sapiens	dJ591C20.3.1 (novel DnaJ domain protein, similar to mouse and bovine cysteine string protein)	1091	100
767	AK024693	Homo sapiens	unnamed protein product	3767	100
768	Z11518	Homo sapiens	histidyl-tRNA synthetase	2582	100
769	X13916	Homo sapiens	LDL-receptor related precursor (AA -19 to 4525)	25529	100
770	AC009360	Arabidopsis thaliana	Contains 3 PF00400 WD40, G-beta repeat domains.	333	33
771	AB037685	Mus musculus	LANP-like protein	1246	91
772	AL161578	Arabidopsis thaliana	putative protein	335	46
773	AL161578	Arabidopsis thaliana	putative protein	333	47
774	AY008271	Homo sapiens	helicase SMARCAD1	5264	99
775	Y21591	Homo sapiens	Human secreted protein (clone CC332-33).	1127	96
776	W88853	Homo sapiens	Polypeptide fragment encoded by gene 89.	752	100
777	W88853	Homo sapiens	Polypeptide fragment encoded by gene 89.	752	100
778	W88853	Homo sapiens	Polypeptide fragment encoded by gene 89.	752	100
779	AF196481	Homo sapiens	RING finger protein; FXY2	3644	100
780	AL035427	Homo sapiens	dJ769N13.1 (KIAA0443 protein.)	1609	54
781	AB026187	Homo sapiens	protocadherin-Xa	5244	100
782	U24458	Homo sapiens	Human secreted protein sequence encoded by gene 22 SEQ ID NO:83.	1002	100
783	AB027289	Homo sapiens	cyclin-E binding protein 1	5421	100
784	G02916	Homo sapiens	Human secreted protein, SEQ ID NO: 6997.	627	100
785	AJ245822	Homo sapiens	type I transmembrane receptor	4560	100
786	AJ245820	Homo sapiens	type I transmembrane receptor	4624	100
787	Z48042	Homo sapiens	GPI-anchored protein p137	3340	99
788	AL031782	Homo sapiens	dJ708F5.1 (PUTATIVE novel Collagen alpha 1 LIKE protein)	2739	100
789	AJ131245	Homo sapiens	Sec24B protein	6602	100
790	AF107203	Homo sapiens	ataxin 2-binding protein	2008	100
791	Y14690	Homo sapiens	procollagen alpha 2(V)	600	34
792	AL031055	Homo sapiens	dJ28H20.2 (novel protein)	1267	100
793	Y36194	787	Human secreted protein	2051	99
794	AB028127	Homo sapiens	mannosyltransferase	2138	96
795	AC007228	Homo sapiens	R31665_2	2738	79
796	AL049482	Arabidopsis thaliana	putative protein	436	47
797	AC004528	Homo sapiens	R32184_3	891	91
798	AB037830	Homo sapiens	KIAA1409 protein	7532	100
799	X53793	Homo sapiens	5' half of the product is homologues to Bacillus subtilis SAICAR synthetase, 3' half corresponds to the catalytic subunit of AIR carboxylase	2232	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
800	Y99350	Homo sapiens	Human PRO1378 (UNQ715) amino acid sequence SEQ ID NO:33.	1343	100
801	AB042636	Homo sapiens	junctophilin type3	1225	47
802	AB029324	Rattus norvegicus	TIP120-family protein TIP120B	3916	90
803	AB029324	Rattus norvegicus	TIP120-family protein TIP120B	4961	90
804	AF251040	Homo sapiens	putative nuclear protein	2119	100
805	AB033281	Homo sapiens	F-box and WD-repeats protein beta-TRCP2 isoform C	2879	100
806	U87305	Rattus norvegicus	transmembrane receptor UNC5H1	3257	90
807	AF118889	Rattus norvegicus	b-tomoxin isoform	3155	97
808	AF226993	Rattus norvegicus	selective LIM binding factor	8793	95
809	W19919	Homo sapiens	Human Ksr-1 (kinase suppressor of Ras).	3939	99
810	AL031782	Homo sapiens	dJ708F5.1 (PUTATIVE novel Collagen alpha 1 LIKE protein)	1546	100
811	AC002542	Homo sapiens	similar to C. elegans F11A10.5; 80% similarity to Z68297 (PID:g1130619)	2294	100
812	U83246	Homo sapiens	copine I	606	52
813	AF242552	Gallus gallus	retinovin	945	34
814	X52332	Homo sapiens	zinc finger protein 10	1651	93
815	X52332	Homo sapiens	zinc finger protein 10	2423	99
816	Y09631	Homo sapiens	PIBF1 protein	2935	99
817	X71997	Rattus norvegicus	myosin I	3883	98
818	AY004877	Mus musculus	cytoplasmic dynein heavy chain	11105	98
819	Y27196	Homo sapiens	Human cyclic nucleotide phosphodiester PDE8B(E) amino acid sequence.	3790	100
820	AF081947	Mus musculus	tektin	1134	81
821	AL035106	Homo sapiens	dJ998C11.1 (continues in Embodiment as ba269H4.1)	871	100
822	AF022795	Homo sapiens	TGF beta receptor associated protein-1	385	24
823	AF015770	Mus musculus	radical fringe	1422	82
824	U82695	Homo sapiens	expressed-Xq28STS protein	1444	99
825	X77371	Mesocricetus auratus	COR1	641	78
826	AB014576	Homo sapiens	KIAA0676 protein	296	79
827	AL049733	Homo sapiens	dJ875H3.1 (APK1 antigen)	1584	72
828	AF222980	Homo sapiens	disrupted in Schizophrenia 1 protein	4418	100
829	Z31560	Homo sapiens	sox-2	1683	100
830	AF295773	Homo sapiens	ral guanine nucleotide dissociation stimulator	4717	99
831	AB041926	Homo sapiens	GCK family kinase MINK-2	6866	100
832	L04948	Saccharomyces cerevisiae	mitochondrial transporter protein	338	35
833	AJ007012	Mus musculus	Fish protein	704	94
834	Z34289	Homo sapiens	nucleolar phosphoprotein p130	3455	99
835	U10991	Homo sapiens	G2	8436	98
836	AF230877	Homo sapiens	MIP-T3	2945	99
837	X58288	Homo sapiens	protein-tyrosine phosphatase	7734	99
838	X56958	Homo sapiens	ankyrin (brank-2)	9631	100
839	AC024791	Caenorhabditis elegans	contains similarity to beta-lactamases	370	24

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
840	D83197	Homo sapiens	ankyrin repeat protein	802	99
841	AF053711	Serinus canaria	neurofilament medium subunit	192	31
842	AF283772	Homo sapiens	similar to Homo sapiens ribosomal protein L10 encoded by GenBank Accession Number L25899	990	96
843	U76343	Homo sapiens	GABA transport protein	2992	98
844	Y13645	Homo sapiens	uroplakin II	897	100
845	D21064	Homo sapiens	similar to rat general mitochondrial matrix processing protease mRNA (RATMPP).	2710	99
846	AF192522	Homo sapiens	Niemann-Pick C3 protein; NPC3	7047	100
847	AF192522	Homo sapiens	Niemann-Pick C3 protein; NPC3	5472	100
848	X60489	Homo sapiens	elongation factor-1-beta	1162	100
849	AC007204	Homo sapiens	BC273239_1	2277	67
850	AC003682	Homo sapiens	R28830_1	2401	100
851	AL121583	Homo sapiens	bA358N2.1 (novel protein)	353	61
852	Z48475	Homo sapiens	glucokinase regulator	3155	99
853	Z83844	Homo sapiens	dJ37E16.2 (SH3-domain binding protein 1)	1884	98
854	AF233323	Homo sapiens	Fas-associated phosphatase-1	390	36
855	AF062741	Rattus norvegicus	pyruvate dehydrogenase phosphatase isoenzyme 2	447	80
856	Y11411	Homo sapiens	pristanoyl-CoA oxidase	3595	98
857	M97188	Strongylocentrotus purpuratus	tektin A1	290	46
858	AB001105	Homo sapiens	hippocalcin-like protein 4	995	100
859	AF164791	Homo sapiens	putative 38.3kDa protein	1795	100
860	AF298117	Homo sapiens	homeobox protein OTX2	1477	93
861	AF015264	Rattus norvegicus	golgi peripheral membrane protein p65	1820	81
862	X16901	Homo sapiens	30kb subunit of RAB30 /74	1284	100
863	M12140	Homo sapiens	envelope protein	202	81
864	AF151459	Homo sapiens	HSPC109	315	98
865	AL109983	Homo sapiens	dJ718P11.1.1 (novel class II aminotransferase similar to serine palmitoyltransferase (isoform 1))	444	100
866	M77183	Rattus norvegicus	alpha-1-macroglobulin	227	45
867	AF272663	Homo sapiens	gephyrin	3785	100
868	X75285	Mus musculus	fibulin-2	3258	87
869	X82494	Homo sapiens	fibulin-2	3407	99
870	AJ297743	Mus musculus	torsinB protein	169	43
871	AJ278313	Homo sapiens	phospholipase C-beta-1a	6258	99
872	AF073344	Homo sapiens	ubiquitin-specific protease 3	256	43
873	Y91955	Homo sapiens	Human cytoskeleton associated protein 10 (CYSKP-10).	535	100
874	AJ000414	Homo sapiens	Cdc42-interacting protein 4	1136	53
875	AF265555	Homo sapiens	ubiquitin-conjugating BIR-domain enzyme APOLLON	627	100
876	Y48586	Homo sapiens	Human breast tumour-associated protein 47.	2537	98
877	AF182198	Homo sapiens	intersectin 2 long isoform	8764	99
878	L17308	Gossypium hirsutum	proline-rich cell wall protein	192	35
879	AF177169	Homo sapiens	tropomodulin 2	1769	100
880	W03627	Homo sapiens	Human follicle stimulating hormone GPR N-terminal sequence.	210	23

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
881	AL021068	Homo sapiens	dJ206D15.3	2615	99
882	AC005498	Homo sapiens	R31665_2	318	82
883	AF165518	Homo sapiens	MAGOH isoform	182	94
884	D21211	Homo sapiens	protein tyrosine phosphatase (PTP-BAS, type 3)	368	43
885	U13045	Homo sapiens	nuclear respiratory factor-2 subunit beta 1	869	62
886	X52836	Homo sapiens	tryptophan hydroxylase (AA 1 - 444)	2320	98
887	X51466	Homo sapiens	elongation factor 2	4460	100
888	AB039903	Homo sapiens	interferon-responsive finger protein 1 long form	1096	98
889	X51760	Homo sapiens	zinc finger protein (583 AA)	3130	100
890	AJ243396	Homo sapiens	voltage-gated sodium channel beta-3 subunit	1024	100
891	W67928	Homo sapiens	Fragment of human secreted protein encoded by gene 4.	391	100
892	AB020598	Homo sapiens	peptide transporter 3	3017	100
893	Y66648	Homo sapiens	Membrane-bound protein PRO1120.	4722	99
894	Y66648	Homo sapiens	Membrane-bound protein PRO1120.	3606	96
895	A29218_cd 1	Homo sapiens	19-NOV-1998 DNA encoding G-protein coupled 7 TM receptor with AXOR15 activity.	2178	100
896	AJ000332	Homo sapiens	Glucosidase II	5063	99
897	X98259	Homo sapiens	M-phase phosphoprotein 8	1085	100
898	X57110	Homo sapiens	c-cbl protein	4849	99
899	X63652	Homo sapiens	inter-alpha-trypsin inhibitor heavy chain ITIH1	3376	98
900	X85134	Homo sapiens	RB protein binding protein	2816	99
901	L11672	Homo sapiens	zinc finger protein	2047	58
902	Y85565	Homo sapiens	Human homologue of UNC-53 (Hs-UNC-53/2) sequence.	369	83
903	X54871	Homo sapiens	ras related protein Rab5b	1094	100
904	Z98265	Homo sapiens	plakophilin 3	4065	100
905	AL035295	Homo sapiens	hypothetical protein	959	99
906	AF051782	Homo sapiens	hypothetical protein	531	35
907	AF208536	Homo sapiens	nucleotide binding protein; NBP	1372	100
908	U79240	Homo sapiens	serine/threonine protein kinase	2365	98
909	U79240	Homo sapiens	serine/threonine protein kinase	2386	99
910	AJ132545	Homo sapiens	protein kinase	2921	100
911	AJ132545	Homo sapiens	protein kinase	1637	99
912	AL121733	Homo sapiens	hypothetical protein	1344	99
913	Y67579	Homo sapiens	Human death inducer-obliterator 1 (DIO-1) polypeptide.	1586	100
914	X87342	Homo sapiens	Human giant larvae homologue	5317	99
915	X87342	Homo sapiens	Human giant larvae homologue	3495	96
916	M94362	Homo sapiens	lamin B2	2357	93
917	AJ011654	Homo sapiens	triple LIM domain protein	3432	100
918	AJ131899	Rattus norvegicus	proline rich synapse associated protein 1	5776	88
919	AF054986	Homo sapiens	putative transmembrane GTPase	1816	100
920	U95822	Homo sapiens	putative transmembrane GTPase	1237	100
921	Y11588	Homo sapiens	apoptosis specific protein	1492	100
922	X84195	Homo sapiens	acylphosphatase	510	100
923	U72882	Homo sapiens	interferon-induced leucine zipper protein	1409	99
924	AE000660	Homo sapiens	hADV36S1	573	100
925	AF126245	Homo sapiens	acyl-Coenzyme A dehydrogenase-8 precursor	2162	100

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
926	AE001968	Deinococcus radiodurans	hypothetical protein	147	27
927	W81576	Homo sapiens	EBV-induced G-protein coupled receptor (EBI-2) polypeptide.	1778	100
928	U01317	Homo sapiens	beta-globin	687	94
929	X98333	Homo sapiens	organic cation transporter	2933	100
930	Y91444	Homo sapiens	Human secreted protein sequence encoded by gene 42 SEQ ID NO:165.	1401	100
931	Y91644	Homo sapiens	Human secreted protein sequence encoded by gene 43 SEQ ID NO:317.	1243	100
932	D90279	Homo sapiens	collagen alpha 1(V) chain precursor	569	39
933	Z31560	Homo sapiens	sox-2	1587	96
934	AF147790	Homo sapiens	transmembrane mucin 12	3047	99
935	Z85996	Homo sapiens	match: multiple proteins; match: Q08151 P28185 Q01111 Q43554; match: Q08150 Q40195 P20340 Q39222; match: Q40368 P36412 P40393 Q40723; match: CE01798 Q38923 Q40191 Q41022; match: Q39433 Q40177 Q40218 Q08146; match: P10949 P11023 Q16948 Q20337; match: Q25389 P25228 P20336 P05713; match: P35276 Q08147 P17609 P22128; match: Q15771 P36410 P35291; GTP-binding	726	94
936	AB041533	Homo sapiens	sperm antigen	1054	38
937	X91906	Homo sapiens	voltage-gated chloride ion channel	3914	100
938	AB032481	Homo sapiens	homeobox transcription factor	1744	100
939	AF111106	Homo sapiens	protein serine/threonine phosphatase 4 regulatory subunit 1	4682	99
940	Y17999	Homo sapiens	Dyrk1B protein kinase	3331	99
941	AF305872	Homo sapiens	thyroglobulin	455	99
942	AF33462	Homo sapiens	cingulin	5939	99
943	AK024442	Homo sapiens	FLJ00032 protein	1616	61
944	Y35911	Homo sapiens	Extended human secreted protein sequence, SEQ ID NO. 160.	262	55
945	AB015320	Homo sapiens	sigma1B subunit of AP-1 clathrin adaptor complex	599	71
946	Z82287	Caenorhabditis elegans	ZK550.2	229	35
947	D84223	Homo sapiens	leucyl tRNA synthetase	6207	99
948	U49057	Rattus norvegicus	rA9	3846	62
949	AK000568	Homo sapiens	unnamed protein product	1659	100
950	AL021578	Homo sapiens	dJ453C12.6.1 (uncharacterized hypothalamus protein (isoform 1))	257	42
951	AB032435	Homo sapiens	differentiation-associated Na-dependent inorganic phosphate cotransporter	3063	99
952	AF110532	Homo sapiens	uncoupling protein UCP-4	1561	100
953	X83587	Mus musculus	1A13 protein	1420	59
954	AL031665	Homo sapiens	dJ545L17.5.1 (novel protein)	386	53
955	Y87600	Homo sapiens	Human fatty acid synthase-like protein (HFASLP).	2377	100
956	Y99421	Homo sapiens	Human PRO1433 (UNQ738) amino acid sequence SEQ ID NO:292.	522	55

SEQ ID NO:	ACCESSION NUMBER	SPECIES	DESCRIPTION	SMITH-WATERMAN SCORE	% IDENTITY
957	U68535	Mus musculus	aldo-keto reductase	451	73
958	AC007067	Arabidopsis thaliana	T10O24.10	1594	57
959	U72194	Mus musculus	muskelin	3947	99
960	AB003661	Drosophila melanogaster	CG15168 gene product	277	54
961	X80332	Mus musculus	rab20	983	82
962	Y67315	Homo sapiens	Human secreted protein BL89_13 amino acid sequence.	3916	99
963	Y67315	Homo sapiens	Human secreted protein BL89_13 amino acid sequence.	3916	99
964	L32602	Rattus norvegicus	homeodomain 159..341	1821	96
965	Z97832	Homo sapiens	dJ329A5.3 (KIAA06460 protein)	3581	99
966	W88995	Homo sapiens	Polypeptide fragment encoded by gene 146.	176	39
967	U12465	Homo sapiens	ribosomal protein L35	604	100
968	AF151803	Homo sapiens	CGI-45 protein	1101	78
969	W74865	Homo sapiens	Human secreted protein encoded by gene 137 clone HMWIF35.	1348	98
970	L21936	Homo sapiens	succinate dehydrogenase flavoprotein subunit	703	100
971	AJ133521	Drosophila buzzatii	protease, reverse transcriptase, ribonuclease H, integrase	194	23
972	AC006017	Homo sapiens	N-acetylgalactosaminyltransferase; similar to Q10473 (PID:g1709559)	3271	100
973	Z81317	Schizosaccharomyces pombe	DNA2-NAM7 helicase family protein	685	31
974	M17885	Homo sapiens	acidic ribosomal phosphoprotein (P0)	792	100
975	U22829	Mus musculus	P2Y purinoceptor	399	40
976	AL132772	Homo sapiens	dJ1013A22.1 (hepatic nuclear factor 4, alpha)	2466	99
977	AC003973	Homo sapiens	ZNF91L	1550	43
978	J04031	Homo sapiens	MDMCSF (EC 1.5.1.5; EC 3.5.4.9; EC 3.3.4.3)	2824	63
979	AF136715	Homo sapiens	taxol resistant associated protein	217	76
980	AF136715	Homo sapiens	taxol resistant associated protein	306	95
981	Z92822	Caenorhabditis elegans	ZK520.1	1109	44
982	AJ295149	Homo sapiens	putative dipeptidase	1564	99
983	AL021331	Homo sapiens	dJ366N23.3 (KIAA0173 and Tubulin-Tyrosine Ligase LIKE)	1492	100
984	AL161501	Arabidopsis thaliana	putative adenosine deaminase	370	38

TABLE 3

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
2	BL00282	Kazal serine protease inhibitors family proteins.	BL00282 16.88 4.259e-14 97-120
3	BL00298	Heat shock hsp90 proteins family proteins.	BL00298A 10.97 1.000e-40 74-119 BL00298E 27.30 1.000e-40 321-376 BL00298F 11.21 1.000e-40 409-464 BL00298H 20.50 1.000e-40 553-607 BL00298C 16.40 2.286e-40 186-230

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			BL00298B 15.64 1.290e-39 134-181 BL00298G 24.57 5.345e-39 465-520 BL00298I 30.07 7.818e-34 661-715 BL00298D 17.97 6.226e-33 242-282
4	PR00237	RHODOPSIN-LIKE GPCR SUPERFAMILY SIGNATURE	PR00237A 11.48 4.316e-13 57-82
5	PD02454	!!!! PROTEIN ALU SUBFAMILY WARNING ENTRY NUCLEAR PHOSPHO.	PD02454B 11.61 4.309e-17 75-103
6	DM00864	EGF-LIKE DOMAIN.	DM00864A 15.21 7.429e-09 98-119
7	PR00237	RHODOPSIN-LIKE GPCR SUPERFAMILY SIGNATURE	PR00237A 11.48 1.750e-11 29-54 PR00237D 8.94 7.000e-09 138-160 PR00237B 13.50 8.250e-09 61-83
9	PF00855	PWWP domain proteins.	PF00855 13.75 5.667e-15 272-289
10	BL00139	Eukaryotic thiol (cysteine) proteases cysteine proteins.	BL00139D 9.24 4.400e-11 391-408 BL00139A 10.29 7.511e-09 67-77
12	BL01113	C1q domain proteins.	BL01113B 18.26 9.294e-19 689-725 BL01113C 13.18 4.857e-11 757-777 BL01113D 7.47 2.161e-10 790-800
13	BL01113	C1q domain proteins.	BL01113B 18.26 3.813e-14 599-635 BL01113C 13.18 4.857e-11 667-687 BL01113D 7.47 2.161e-10 700-710
14	BL00594	Aromatic amino acids permeases proteins.	BL00594A 16.75 6.531e-10 50-94
15	BL01047	Heavy-metal-associated domain proteins.	BL01047B 19.73 4.913e-13 707-728
16	PR00625	DNAJ PROTEIN FAMILY SIGNATURE	PR00625A 12.84 7.462e-18 310-330 PR00625B 13.48 3.939e-15 340-351
18	BL00615	C-type lectin domain proteins.	BL00615A 5.68 3.700e-09 144-162
20	PR00741	GLYCOSYL HYDROLASE FAMILY 29 SIGNATURE	PR00741D 16.11 9.982e-21 175-195 PR00741F 14.66 9.262e-21 243-265 PR00741B 14.23 1.947e-18 128-145 PR00741G 9.29 2.180e-17 318-340 PR00741C 9.16 7.328e-17 147-166 PR00741H 10.32 2.141e-13 351-374 PR00741A 9.24 3.596e-13 89-105 PR00741E 13.39 3.535e-12 215-232
22	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 3.647e-20 117-148 BL00107B 13.31 1.000e-16 182-198
23	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 1.600e-23 126-157
24	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 1.600e-23 126-157
27	BL00239	Receptor tyrosine kinase class II proteins.	BL00239B 25.15 2.324e-16 91-139
28	BL00018	EF-hand calcium-binding domain proteins.	BL00018 7.41 3.250e-10 681-694 BL00018 7.41 6.400e-10 717-730
29	BL00018	EF-hand calcium-binding domain	BL00018 7.41 3.250e-10 681-694

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		proteins.	BL00018 7.41 6.400e-10 717-730
30	BL01113	Clq domain proteins.	BL01113A 17.99 9.308e-09 54-81
33	PD01168	SYNTHETASE LIGASE PROTEIN ALANYL.	PD01168L 9.47 1.667e-09 401-416
34	PD01168	SYNTHETASE LIGASE PROTEIN ALANYL.	PD01168L 9.47 1.667e-09 411-426
36	PR00426	CSA-ANAPHYLATOXIN RECEPTOR SIGNATURE	PR00426D 10.59 3.618e-12 110-122
37	PF00791	Domain present in ZO-1 and Unc5-like netrin receptors.	PF00791B 28.49 2.049e-10 1080-1135
38	BL00350	MADS-box domain proteins.	BL00350 20.79 1.000e-40 1-55
40	BL00123	Alkaline phosphatase proteins.	BL00123B 19.31 1.000e-40 90-133 BL00123C 24.61 1.000e-40 145-195 BL00123E 22.25 1.000e-40 304-358 BL00123G 26.01 1.000e-40 438-488 BL00123F 19.03 8.714e-35 364-399 BL00123A 10.80 9.000e-24 52-77 BL00123D 12.73 1.000e-17 216-229
44	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 2.800e-14 346-359 PD00066 13.92 4.600e-14 486-499 PD00066 13.92 1.000e-13 374-387 PD00066 13.92 6.000e-13 458-471 PD00066 13.92 2.714e-12 234-247 PD00066 13.92 3.143e-12 430-443 PD00066 13.92 8.714e-12 514-527 PD00066 13.92 3.739e-11 402-415 PD00066 13.92 2.038e-10 318-331
45	DM00973	3 kw RESISTANCE BENOMYL YLL028W CYCLOHEXIMIDE.	DM00973A 21.17 2.946e-10 180-217
47	BL00649	G-protein coupled receptors family 2 proteins.	BL00649C 17.82 1.682e-10 475-501 BL00649B 20.68 7.387e-09 417-463
50	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 8.200e-16 415-458 PD00066 13.92 5.846e-15 305-311 PD00066 13.92 1.000e-14 221-234 PD00066 13.92 1.000e-14 417-430 PD00066 13.92 2.800e-14 249-262 PD00066 13.92 2.800e-14 277-290 PD00066 13.92 8.800e-14 333-346 PD00066 13.92 9.400e-14 361-374 PD00066 13.92 4.000e-13 389-402 PD00066 13.92 6.571e-12 473-486
51	BL00226	Intermediate filaments proteins.	BL00226D 19.10 1.000e-40 417-464 BL00226B 23.86 3.348e-35 251-299 BL00226C 13.23 1.429e-24 316-347 BL00226A 12.77 1.857e-15 151-166
52	PR00217	43 KD POSTSYNAPTIC PROTEIN SIGNATURE	PR00217C 10.91 5.648e-09 133-149
53	BL00232	Cadherins extracellular repeat proteins domain proteins.	BL00232B 32.79 1.000e-40 143-191 BL00232A 27.72 2.350e-28 49-82 BL00232B 32.79 7.052e-21 252-300 BL00232C 10.65 6.625e-20 250-268 BL00232B 32.79 1.314e-11 367-415 BL00232C 10.65 9.308e-10 470-488
54	BL00303	S-100/CaBP type calcium binding	BL00303B 26.15 8.759e-23 125-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		protein.	162 BL00303A 21.77 1.000e-21 82-119
58	PR00378	INOSITOL PHOSPHATASE SIGNATURE	PR00378D 16.86 1.000e-15 242-261 PR00378B 13.80 9.250e-13 109-129
59	PR00425	BRADYKININ RECEPTOR SIGNATURE	PR00425C 13.23 9.040e-12 120-140
60	BL00280	Pancreatic trypsin inhibitor (Kunitz) family proteins.	BL00280 24.61 6.727e-38 238-282 BL00280 24.61 1.514e-30 294-338
65	BL01019	ADP-ribosylation factors family proteins.	BL01019A 13.20 1.222e-11 43-83
68	PR00237	RHODOPSIN-LIKE GPCR SUPERFAMILY SIGNATURE	PR00237E 13.03 5.091e-13 188-212 PR00237G 19.63 7.207e-13 268-295 PR00237A 11.48 4.375e-11 24-49 PR00237C 15.69 3.057e-10 101-124 PR00237D 8.94 4.750e-10 137-159 PR00237F 13.57 5.364e-10 230-255 PR00237B 13.50 9.438e-10 57-79
70	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 7.938e-28 31-70
71	PR00830	ENDOPEPTIDASE LA (LON) SERINE PROTEASE (S16) SIGNATURE	PR00830A 8.41 8.759e-12 348-368
72	BL00120	Lipases, serine proteins.	BL00120B 11.37 2.149e-10 148-163
77	PR00753	1-AMINOCYCLOPROPANE-1-CARBOXYLATE SYNTHASE SIGNATURE	PR00753E 8.01 3.552e-11 191-216 PR00753D 6.85 2.778e-09 131-153
78	PR00506	D21 CLASS N6 ADENINE-SPECIFIC DNA METHYLTRANSFERASE SIGNATURE	PR00506C 19.40 8.017e-09 96-119
82	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 3.571e-16 436-467
84	BL00675	Sigma-54 interaction domain proteins ATP-binding region A proteins.	BL00675A 24.86 8.800e-10 256-300
85	BL00027	'Homeobox' domain proteins.	BL00027 26.43 2.286e-30 117-130
87	BL00250	TGF-beta family proteins.	BL00250A 21.24 6.786e-36 264-300 BL00250B 27.37 1.450e-26 328-364
91	BL00215	Mitochondrial energy transfer proteins.	BL00215A 15.82 9.250e-17 10-35 BL00215A 15.82 6.000e-16 221-246 BL00215A 15.82 7.857e-12 108-133 BL00215B 10.44 9.526e-11 168-181
92	BL00027	'Homeobox' domain proteins.	BL00027 26.43 9.526e-24 324-367
95	PR00094	ADENYLATE KINASE SIGNATURE	PR00094C 12.94 1.000e-08 119-136
96	PD02327	GLYCOPROTEIN ANTIGEN PRECURSOR IMMUNOGLO.	PD02327B 19.84 2.091e-09 143-165
97	BL00752	XPA protein.	BL00752B 19.17 7.309e-09 28-72
98	PR00876	NEMATODE METALLOTHIONEIN SIGNATURE	PR00876B 7.66 2.268e-10 135-149
99	PR00109	TYROSINE KINASE CATALYTIC DOMAIN SIGNATURE	PR00109B 12.27 9.824e-12 122-141
100	BL00027	'Homeobox' domain proteins.	BL00027 26.43 7.429e-31 118-161
101	BL00028	Zinc finger, C2H2 type, domain proteins.	BL00028 16.07 6.870e-12 370-387 BL00028 16.07 6.885e-11 398-415 BL00028 16.07 8.269e-11 342-359 BL00028 16.07 4.300e-10 229-246

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			BL00028 16.07 6.100e-10 258-275
102	PR00048	C2H2-TYPE ZINC FINGER SIGNATURE	PR00048A 10.52 7.750e-14 665-679 PR00048A 10.52 8.500e-14 581-595 PR00048A 10.52 9.250e-14 637-651 PR00048A 10.52 2.059e-12 609-623 PR00048A 10.52 2.588e-12 469-483 PR00048A 10.52 7.353e-12 553-567 PR00048A 10.52 2.895e-11 525-539 PR00048A 10.52 4.316e-11 441-455 PR00048A 10.52 5.263e-11 413-427 PR00048B 6.02 2.125e-10 569-579 PR00048B 6.02 4.938e-10 513-523 PR00048A 10.52 5.696e-10 497-511 PR00048B 6.02 8.875e-10 429-439 PR00048B 6.02 1.000e-09 457-467 PR00048B 6.02 6.684e-09 485-495
103	PR00195	DYNAMIN SIGNATURE	PR00195A 11.94 5.364e-22 31-50 PR00195B 9.47 1.783e-21 56-74 PR00195C 11.50 3.455e-21 126-144 PR00195D 11.76 8.714e-21 175-194 PR00195F 16.20 8.500e-20 217-237 PR00195E 9.82 8.650e-20 194-211
104	BL01113	Clq domain proteins.	BL01113A 17.99 1.865e-09 121-148 BL01113A 17.99 5.846e-09 82-109
105	BL00420	Speract receptor repeat proteins domain proteins.	BL00420A 20.42 6.400e-11 70-99 BL00420A 20.42 8.525e-10 73-102 BL00420A 20.42 5.708e-09 85-114
108	PR00860	VERTEBRATE METALLOTHIONEIN SIGNATURE	PR00860B 7.04 2.929e-20 27-41 PR00860A 5.46 5.500e-16 5-18 PR00860C 9.61 1.47e-14 41-51
112	BL01031	Heat shock hsp26 proteins family profile.	BL01031C 17.68 6.400e-10 122-147
114	DM01840	kw SPAC24B11.09 RD7E5.13.	DM01840B 22.04 2.688e-40 59-103 DM01840A 10.95 9.571e-13 31-43
115	BL01126	Elongation factor Ts proteins.	BL01126A 18.48 2.317e-30 46-89 BL01126B 13.15 7.387e-19 116-135 BL01126C 9.20 9.735e-11 190-203
116	BL00216	Sugar transport proteins.	BL00216B 27.64 4.375e-21 35-85
118	BL00437	Catalase proximal heme-ligand proteins.	BL00437A 18.82 1.000e-40 49-101 BL00437B 16.28 1.000e-40 114-168 BL00437C 21.86 1.000e-40 190-239 BL00437D 25.72 1.000e-40 248-301 BL00437E 23.95 1.000e-40 327-379
119	BL00140	Ubiquitin carboxyl-terminal hydrolase family 1 cysteine activ.	BL00140D 22.64 8.274e-14 164-208 BL00140C 11.80 5.444e-10 77-102
120	BL00224	Clathrin light chain proteins.	BL00224B 16.94 6.712e-10 95-148
122	BL00203	Vertebrate metallothioneins proteins.	BL00203 13.94 1.000e-40 16-62
123	PR00041	CAMP RESPONSE ELEMENT	PR00041D 7.95 2.906e-09 24-41

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		BINDING (CREB) PROTEIN SIGNATURE	
124	PR00041	CAMP RESPONSE ELEMENT BINDING (CREB) PROTEIN SIGNATURE	PR00041D 7.95 2.906e-09 24-41
125	BL00061	Short-chain dehydrogenases/reductases family proteins.	BL00061C 7.86 3.250e-10 212-222
126	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 6.400e-25 251-290
127	PR00318	ALPHA G-PROTEIN (TRANSDUCIN) SIGNATURE	PR00318D 16.28 1.900e-34 219-248 PR00318B 14.79 3.455e-27 168-191 PR00318C 12.09 7.000e-23 197-215 PR00318A 7.84 1.600e-19 35-51 PR00318E 7.23 2.500e-12 265-275
128	PR00927	ADENINE NUCLEOTIDE TRANSLOCATOR 1 SIGNATURE	PR00927E 14.93 9.743e-10 67-89 PR00927B 14.66 4.575e-09 69-91
130	BL00824	Elongation factor 1 beta/beta'/delta chain proteins.	BL00824B 9.21 7.750e-22 133-153
131	BL00824	Elongation factor 1 beta/beta'/delta chain proteins.	BL00824C 14.58 1.000e-40 166-204 BL00824D 14.04 1.621e-38 204-239 BL00824B 9.21 7.750e-22 133-153 BL00824E 12.49 1.000e-19 247-263
132	PR00209	ALPHA/BETA GLIADIN FAMILY SIGNATURE	PR00209B 4.88 9.222e-13 1209-1228
133	PR00209	ALPHA/BETA GLIADIN FAMILY SIGNATURE	PR00209B 4.88 9.222e-13 1168-1187
134	PR00708	ALPHA-I-ACID GLYCOPROTEIN SIGNATURE	PR00708D 14.67 1.000e-27 141-168 PR00708C 11.77 1.643e-25 98-120 PR00708B 15.15 2.174e-24 73-95 PR00708E 13.33 1.600e-21 189-207 PR00708A 14.40 2.636e-21 51-70
135	PR00145	TYROSINE KINASE CATALYTIC DOMAIN SIGNATURE	PR00145 12.27 8.168e-10 145
136	PF00023	Ank repeat proteins.	PF00023A 16.03 3.250e-10 211-217
137	BL00471	Small cytokines (intercrine/chemokine) C-x-C subfamily signat.	BL00471 23.92 7.480e-10 42-90
140	PR00205	CADHERIN SIGNATURE	PR00205B 11.39 5.582e-10 328-346 PR00205B 11.39 9.018e-10 543-561
141	BL00412	Neuromodulin (GAP-43) proteins.	BL00412D 16.54 7.704e-09 976-1027
143	PR00979	TAFAZZIN SIGNATURE	PR00979E 10.83 5.950e-26 192-214 PR00979A 11.91 8.773e-25 63-83 PR00979C 12.16 6.400e-19 108-124 PR00979D 12.38 7.955e-19 170-185 PR00979F 10.14 3.382e-15 230-244 PR00979B 15.59 5.636e-15 94-106
145	DM00686	kw REPLICATION REP 28K 17.7K.	DM00686C 14.14 7.720e-09 111-131
146	PR00604	CLASS IA AND IB CYTOCHROME C SIGNATURE	PR00604D 15.86 1.000e-17 87-104 PR00604B 12.73 9.591e-16 57-73 PR00604C 10.21 8.200e-12 73-84 PR00604E 10.13 1.000e-11 106-117 PR00604A 11.13 8.800e-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			11 44-52 PR00604F 8.60 1.000e-10 123-132
147	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 3.864e-15 266-297 BL00107B 13.31 6.143e-11 335-351
148	PD00289	PROTEIN SH3 DOMAIN REPEAT PRESYN.	PD00289 9.97 8.448e-09 67-81
149	PR00069	ALDO-KETO REDUCTASE SIGNATURE	PR00069D 19.36 1.857e-30 187-217 PR00069A 16.01 7.429e-25 41-66 PR00069E 18.14 3.100e-22 235-260 PR00069C 16.03 7.000e-20 151-169 PR00069B 11.33 8.071e-19 101-120
150	BL00027	'Homeobox' domain proteins.	BL00027 26.43 2.688e-27 139-182
151	PD02906	SYNTHASE I PSEUDOURIDYLATE PSEUDOURIDINE LYASE TR.	PD02906C 24.17 7.070e-22 165-200 PD02906B 15.35 8.393e-15 114-127 PD02906A 10.84 6.500e-09 71-84
153	BL00479	Phorbol esters / diacylglycerol binding domain proteins.	BL00479A 19.86 5.091e-12 891-914 BL00479B 12.57 1.837e-11 915-931
158	BL00027	'Homeobox' domain proteins.	BL00027 26.43 6.786e-31 143-186
160	BL00422	Granins proteins.	BL00422C 16.18 7.750e-12 420-448
162	PR00625	DNAJ PROTEIN FAMILY SIGNATURE	PR00625A 12.84 9.297e-11 62-82
164	BL01282	BIR repeat proteins.	BL01282B 30.49 6.182e-10 347-386
166	PR00860	VERTEBRATE METALLOTHIONEIN SIGNATURE	PR00860B 7.04 2.929e-20 83-97 PR00860A 5.46 1.000e-18 61-74 PR00860C 9.61 1.900e-15 97-107
167	PR00449	TRANSFORMING PROTEIN P21 RAS SIGNATURE	PR00449A 13.20 7.052e-09 196-218
169	BL00514	Fibrinogen beta and gamma chains C-terminal domain proteins.	BL00514C 17.41 1.346e-39 316-355 BL00514G 15.98 2.241e-34 471-501 BL00514H 14.95 6.571e-27 510-535 BL00514E 14.28 1.273e-16 388-405 BL00514D 15.35 9.100e-15 369-382 BL00514B 16.42 4.857e-14 260-276 BL00514F 11.65 9.690e-14 416-431 BL00514A 11.68 8.200e-11 149-159
170	BL00514	Fibrinogen beta and gamma chains C-terminal domain proteins.	BL00514C 17.41 1.346e-39 268-305 BL00514G 15.98 2.241e-34 423-453 BL00514H 14.95 6.571e-27 462-487 BL00514E 14.28 1.273e-16 340-357 BL00514D 15.35 9.100e-15 321-334 BL00514B 16.42 4.857e-14 212-228 BL00514F 11.65 9.690e-14 368-383 BL00514A 11.68 8.200e-11 101-111
171	BL00514	Fibrinogen beta and gamma chains C-terminal domain proteins.	BL00514G 15.98 2.241e-34 385-415 BL00514H 14.95 6.571e-27 424-449 BL00514C 17.41 4.632e-24 230-267 BL00514E 14.28 1.273e-16 302-319 BL00514D 15.35 9.100e-15 283-296

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			BL00514B 16.42 4.857e-14 212-228 BL00514F 11.65 9.690e-14 330-345 BL00514A 11.68 8.200e-11 101-111
173	BL00027	'Homeobox' domain proteins.	BL00027 26.43 9.400e-29 119-162
174	DM01970	0 kw ZK632.12 YDR313C ENDOSOMAL III.	DM01970B 8.60 5.119e-15 1391-1404
176	BL00773	Chitinases family 19 proteins.	BL00773C 9.42 8.000e-09 2-16
182	PR00109	TYROSINE KINASE CATALYTIC DOMAIN SIGNATURE	PR00109B 12.27 9.163e-14 141-160
183	PD01937	DNA PROTEIN POLYMERASE ENDONUCLEASE DNA-	PD01937A 6.68 3.475e-09 221-232
185	BL00845	CAP-Gly domain proteins.	BL00845 16.43 2.946e-23 247-272 BL00845 16.43 1.628e-21 107-132
186	PR00452	SH3 DOMAIN SIGNATURE	PR00452B 11.65 6.538e-11 525-541
187	PR00452	SH3 DOMAIN SIGNATURE	PR00452B 11.65 6.538e-11 497-513
188	DM01803	1 HERPESVIRUS GLYCOPROTEIN H.	DM01803A 10.51 1.000e-09 1081-1102
189	PF00651	BTB (also known as BR-C/Ttk) domain proteins.	PF00651 15.00 5.091e-15 69-82
190	PR00194	TROPOMYOSIN SIGNATURE	PR00194C 6.38 1.900e-35 145-174 PR00194E 8.74 3.250e-30 231-257 PR00194D 9.57 1.500e-26 175-199 PR00194B 10.24 5.200e-24 120-141 PR00194A 7.86 4.857e-21 84-102
192	PD02042	IRON-SULFUR ELECTRON TRANSPORT AROMATIC HYDROCARB.	PD02042B 16.75 5.154e-09 131-146 PD02042A 21.13 5.909e-09 94-121
193	PR00021	SMALL PROLINE-RICH PROTEIN SIGNATURE	PR00021A 4.31 2.200e-10 2-15
195	BL00463	Fungal Zn(2)-Cys(6) binuclear cluster domain proteins.	BL00463 8.22 5.071e-09 111-123
196	PR00118	BETA-LACTAMASE CLASS A SIGNATURE	PR00118F 16.42 2.386e-09 165-181
197	DM00215	PROLINE-RICH PROTEIN 3.	DM00215 19.43 5.424e-09 234-267
198	BL00660	Band 4.1 family domain proteins.	BL00660A 31.50 5.500e-11 714-767
199	BL00282	Kazal serine protease inhibitors family proteins.	BL00282 16.88 8.820e-13 70-93
202	PR00009	TYPE I EGF SIGNATURE	PR00009A 14.15 5.345e-15 971-987 PR00009C 14.11 8.773e-13 996-1008 PR00009D 16.83 8.000e-11 1008-1018 PR00009C 14.11 1.882e-09 892-904
203	BL00025	P-type 'Trefoil' domain proteins.	BL00025 17.17 4.536e-19 38-59
205	BL00018	EF-hand calcium-binding domain proteins.	BL00018 7.41 7.300e-10 165-178
206	PR00168	SLOW VOLTAGE-GATED POTASSIUM CHANNEL SIGNATURE	PR00168D 12.88 6.865e-11 67-86
207	BL00025	P-type 'Trefoil' domain proteins.	BL00025 17.17 3.423e-20 39-60 BL00025 17.17 8.750e-16 88-109
209	BL00646	Ribosomal protein S13 proteins.	BL00646B 21.42 6.100e-30 110-143 BL00646A 25.82 6.192e-29 14-62
210	PR00138	MATRIXIN SIGNATURE	PR00138D 16.56 3.605e-25 279-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			305 PR00138C 16.41 3.000e-24 218-247 PR00138E 6.01 8.714e-13 314-328 PR00138A 15.14 9.538e-13 134-148 PR00138B 15.82 4.522e-12 188-204
211	DM01206	CORONAVIRUS NUCLEOCAPSID PROTEIN.	DM01206B 10.69 8.429e-12 386-406 DM01206B 10.69 1.247e-10 384-404 DM01206B 10.69 5.068e-10 388-408
212	PD01941	TRANSMEMBRANE COTRANSPORTER SYMP.	PD01941A 14.81 1.000e-40 163-217 PD01941B 15.02 9.705e-30 420-467 PD01941E 15.92 8.714e-23 837-884 PD01941C 19.96 8.200e-20 508-563 PD01941D 27.18 1.600e-16 661-710 PD01941F 28.52 9.645e-15 1005-1060
213	BL00362	Ribosomal protein S15 proteins.	BL00362 24.67 8.313e-09 330-373
214	BL00115	Eukaryotic RNA polymerase II heptapeptide repeat proteins.	BL00115Z 3.12 2.125e-09 1178-1227 BL00115Z 3.12 6.096e-09 1164-1213
215	BL00038	Myc-type, 'helix-loop-helix' dimerization domain proteins.	BL00038B 16.97 7.600e-18 125-146 BL00038A 13.61 1.474e-13 102-118
216	BL01108	Ribosomal protein L24 proteins.	BL01108A 20.33 2.241e-22 49-82 BL01108B 11.40 8.457e-10 96-107
217	PR00381	KINESIN LIGHT CHAIN SIGNATURE	PR00381A 9.55 1.321e-10 360-378
222	BL00514	Fibrinogen beta and gamma chains C-terminal domain proteins.	BL00514C 17.41 2.358e-26 1166-1203 BL00514G 15.98 9.000e-15 1289-1319 BL00514D 15.35 6.936e-12 1207-1220 BL00514F 11.65 4.288e-10 1253-1268 BL00514E 14.95 8.636e-10 1210-1343
223	BL00325	Actin-depolymerizing proteins.	BL00325B 21.66 1.000e-40 93-139 BL00325A 24.83 9.333e-24 61-93
224	BL00018	EF-hand calcium-binding domain proteins.	BL00018 7.41 1.450e-10 231-244
225	PF01329	Pterin 4 alpha carbinolamine dhydratase.	PF01329B 18.52 1.692e-18 67-92
228	BL00211	ABC transporters family proteins.	BL00211B 13.37 6.250e-18 1033-1065 BL00211B 13.37 8.875e-18 2045-2077 BL00211A 12.23 1.900e-09 931-943
230	PR00761	BINDIN PRECURSOR SIGNATURE	PR00761A 5.81 9.366e-09 275-292
231	PR00049	WILM'S TUMOUR PROTEIN SIGNATURE	PR00049D 0.00 3.500e-10 54-69
232	BL00412	Neuromodulin (GAP-43) proteins.	BL00412D 16.54 1.978e-10 109-160 BL00412D 16.54 4.122e-09 133-184
233	BL01210	Caveolins proteins.	BL01210B 13.92 8.129e-09 106-156
236	BL00939	Ribosomal protein L1e proteins.	BL00939F 17.27 5.393e-09 861-891
238	BL01252	Endogenous opioids neuropeptides precursors proteins.	BL01252D 18.25 3.571e-28 205-233 BL01252B 19.09 5.034e-27

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			37-67 BL01252C 18.10 1.621e-21 164-190 BL01252A 14.22 7.107e-18 14-34
239	BL00302	Eukaryotic initiation factor 5A hypusine proteins.	BL00302 14.81 1.000e-40 25-79
240	PR00420	AROMATIC-RING HYDROXYLASE (FLAVOPROTEIN MONOOXYGENASE) SIGNATURE	PR00420A 14.78 8.851e-13 26-49
241	PD02929	ADHESION GLYCOPROTEIN PRECURSOR I.	PD02929A 28.27 4.529e-09 235-289
243	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 8.527e-25 11-50
244	BL01270	Band 7 protein family proteins.	BL01270C 16.91 6.745e-17 115-144 BL01270B 18.74 6.857e-17 76-115 BL01270E 13.03 6.016e-15 182-211 BL01270D 20.87 9.160e-13 144-182
245	PF00791	Domain present in ZO-1 and Unc5-like netrin receptors.	PF00791B 28.49 6.305e-12 253-308 PF00791B 28.49 1.909e-11 427-482 PF00791B 28.49 2.651e-09 179-234 PF00791B 28.49 3.890e-09 112-167
246	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 2.500e-13 277-290 PD00066 13.92 9.143e-12 193-206 PD00066 13.92 5.304e-11 165-178 PD00066 13.92 6.478e-11 249-262 PD00066 13.92 3.423e-10 221-234
247	BL00406	Actins proteins.	BL00406D 12.58 6.400e-20 465-520 BL00406B 5.47 4.857e-14 249-304 BL00406E 8.44 1.000e-11 522-572 BL00406C 6.75 5.449e-11 313-368
248	BL00951	ER lumen protein retaining receptor proteins.	BL00951C 19.35 1.000e-40 112-161 BL00951A 15.10 7.750e-39 21-57 BL00951D 13.00 2.000e-38 161-196 BL00951B 14.23 3.100e-31 57-88
252	BL01113	Clq domain proteins.	BL01113A 17.99 9.129e-15 200-227 BL01113A 17.99 4.818e-14 194-221 BL01113A 17.99 7.818e-14 182-209 BL01113A 17.99 1.730e-13 185-212 BL01113A 17.99 6.595e-13 191-218 BL01113A 17.99 6.077e-12 203-230 BL01113A 17.99 9.182e-11 179-206 BL01113A 17.99 2.532e-10 176-203 BL01113A 17.99 9.043e-10 218-245 BL01113A 17.99 9.426e-10 209-236 BL01113A 17.99 4.115e-09 137-164
257	BL00845	CAP-Gly domain proteins.	BL00845 16.43 1.837e-21 466-491
259	PR00248	METABOTROPIC GLUTAMATE GPCR SIGNATURE	PR00248G 12.67 2.688e-09 53-78
260	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 3.400e-10 441-452 BL00678 9.67 5.800e-10 481-492 BL00678 9.67 8.800e-10 358-369
261	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 3.400e-10 415-426 BL00678 9.67 5.800e-10 455-466

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			BL00678 9.67 8.800e-10 332-343
262	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 3.400e-10 468-479 BL00678 9.67 5.800e-10 508-519 BL00678 9.67 8.800e-10 385-396
263	BL50002	Src homology 3 (SH3) domain proteins profile.	BL50002B 15.18 2.200e-10 415-429
264	BL00049	Ribosomal protein L14 proteins.	BL00049C 17.38 3.040e-12 94-130
265	PD01469	GLYCOPROTEIN PROTEIN PRECURSOR SA.	PD01469 20.59 2.091e-14 438-470
266	PD01469	GLYCOPROTEIN PROTEIN PRECURSOR SA.	PD01469 20.59 2.091e-14 279-311
267	BL00567	Phosphoribulokinase proteins.	BL00567A 10.66 1.161e-12 36-55
269	BL00049	Ribosomal protein L14 proteins.	BL00049C 17.38 2.688e-12 92-128 BL00049B 18.42 6.806e-24 54-86 BL00049A 13.86 8.333e-19 19-42 BL00049D 13.47 5.765e-12 129-140
272	BL01115	GTP-binding nuclear protein ran proteins.	BL01115A 10.22 9.735e-12 14-58
273	PR00021	SMALL PROLINE-RICH PROTEIN SIGNATURE	PR00021A 4.31 1.911e-09 819-832
275	PR00179	LIPOCALIN SIGNATURE	PR00179B 9.56 2.895e-13 124-137 PR00179A 13.78 3.250e-11 36-49 PR00179C 19.02 6.040e-11 154-170
276	PR00449	TRANSFORMING PROTEIN P21 RAS SIGNATURE	PR00449A 13.20 8.364e-17 22-44 PR00449C 17.27 1.000e-13 62-85 PR00449E 13.50 4.000e-12 172-195 PR00449B 14.34 5.680e-10 45-62
277	BL00140	Ubiquitin carboxyl-terminal hydrolase family I cysteine activ.	BL00140D 22.64 1.000e-40 161-205 BL00140C 11.80 9.053e-30 79-104 BL00140A 15.96 9.400e-28 5-35 BL00140B 12.29 4.649e-17 37-53
278	PD02712	ELEMENT TRANSPOSASE FOR TRANSPOSON TRANSPOSABLE.	PD02712A 23.07 8.015e-09 47-83
279	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 1.474e-10 100-111
282	DM00892	3 RETROVIRAL PROTEINASE.	DM00892C 23.55 4.767e-21 864-898
283	BL00048	Protamine P1 proteins.	BL00048 6.39 9.550e-09 56-83
286	PR00081	GLUCOSE/RIBITOL DEHYDROGENASE FAMILY SIGNATURE	PR00081A 10.53 1.878e-11 36-54
287	PR00310	ANTI-PROLIFERATIVE PROTEIN BTG1 FAMILY SIGNATURE	PR00310B 10.59 4.231e-17 29-59 PR00310D 9.10 6.679e-16 89-119
289	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 7.000e-36 37-76
293	BL00979	G-protein coupled receptors family 3 proteins.	BL00979L 20.63 3.800e-12 111-152
295	PD02411	PROTEIN TRANSCRIPTION REGULATION NUCLEAR.	PD02411 21.89 7.000e-16 195-229
296	BL01064	Pyridoxamine 5'-phosphate oxidase proteins.	BL01064A 27.84 8.313e-28 77-129 BL01064C 15.22 7.136e-25 202-235
297	BL00030	Eukaryotic RNA-binding region RNP-1 proteins.	BL00030A 14.39 2.929e-13 37-56 BL00030B 7.03 1.900e-11 167-177 BL00030A 14.39 2.000e-10 128-147

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
298	BL01183	ubiE/COQ5 methyltransferase family proteins.	BL01183B 21.31 6.660e-12 143-188
299	BL01279	Protein-L-isoaspartate(D-aspartate) O-methyltransferase signa.	BL01279A 24.27 5.862e-11 57-105
301	BL00191	Cytochrome b5 family, heme-binding domain proteins.	BL00191K 17.38 4.951e-27 184-228 BL00191J 11.37 6.447e-17 128-150
302	DM00892	3 RETROVIRAL PROTEINASE.	DM00892C 23.55 3.893e-16 33-67
306	PF01140	Matrix protein (MA), p15.	PF01140D 15.54 2.988e-09 416-451
307	PR00245	OLFACTORY RECEPTOR SIGNATURE	PR00245A 18.03 4.818e-21 59-81 PR00245C 7.84 5.154e-20 238-254 PR00245D 10.47 4.000e-15 274-286 PR00245B 10.38 8.200e-15 177-192 PR00245E 12.40 5.714e-12 291-306
309	BL00203	Vertebrate metallothioneins proteins.	BL00203 13.94 2.245e-10 612-658
310	BL00237	G-protein coupled receptors proteins.	BL00237A 27.68 7.632e-23 119-159 BL00237C 13.19 3.864e-15 251-278 BL00237D 11.23 3.739e-12 312-329
311	BL00380	Rhodanese proteins.	BL00380D 15.90 8.200e-28 110-136 BL00380G 11.26 5.800e-16 267-280 BL00380B 14.77 7.000e-14 49-62 BL00380F 9.76 5.886e-13 203-214 BL00380C 15.67 7.387e-13 82-98 BL00380E 12.44 7.000e-11 181-193 BL00380A 10.48 1.000e-09 10-20
312	BL00227	Tubulin subunits alpha, beta, and gamma proteins.	BL00227B 19.29 1.000e-40 50-105 BL00227C 25.48 1.000e-40 111-163 BL00227D 18.46 1.000e-40 220-274 BL00227F 21.16 1.000e-40 372-426 BL00227A 24.55 3.259e-39 1-35 BL00227E 24.15 8.500e-34 324-359
327	BL00232	Cadherins extracellular repeat proteins domain proteins.	BL00232B 32.79 7.362e-21 225-273 BL00232B 32.79 2.588e-17 435-483 BL00232B 32.79 6.301e-15 116-164 BL00232B 32.79 6.769e-13 330-378 BL00232C 10.65 9.341e-12 223-241 BL00232C 10.65 5.696e-11 328-346 BL00232C 10.65 3.942e-10 433-451
329	PD02749	TRANSCRIPTION PROTEIN FACTOR BTF3 REGULATION NUCL.	PD02749B 12.75 2.241e-37 35-71 PD02749C 13.96 4.892e-28 87-121 PD02749A 9.56 6.000e-15 2-15
330	PR00391	PHOSPHATIDYLINOSITOL TRANSFER PROTEIN SIGNATURE	PR00391E 12.50 7.785e-15 211-231 PR00391B 8.39 1.000e-13 83-104 PR00391D 12.21 9.328e-13 191-207 PR00391A 7.83 5.390e-11 16-36
332	BL01030	RNA polymerases M / 15 Kd subunits proteins.	BL01030 23.44 1.818e-23 87-125
337	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 2.929e-32 6-45
340	PD02711	SYNTHASE	PD02711B 14.26 1.973e-20 944-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		PHOSPHORIBOSYLFORMYLGLY.	968
343	BL00223	Annexins repeat proteins domain proteins.	BL00223C 24.79 1.000e-40 245-300 BL00223B 28.47 8.714e-38 168-218 BL00223A 15.59 8.250e-27 98-132 BL00223A 15.59 8.750e-27 26-60 BL00223C 24.79 9.438e-16 13-68 BL00223C 24.79 2.735e-15 85-140 BL00223A 15.59 2.253e-11 258-292
346	PR00345	STATHMIN FAMILY SIGNATURE	PR00345B 7.12 2.800e-28 81-110 PR00345E 8.54 7.652e-28 158-183 PR00345C 4.54 9.100e-28 110-134 PR00345D 10.97 1.964e-24 134-158 PR00345A 13.46 5.645e-16 52-71
347	BL00586	Ribosomal protein L16 proteins.	BL00586B 17.00 3.215e-15 184-221
348	PR00388	3',5'-CYCLIC NUCLEOTIDE CLASS II PHOSPHODIESTERASE SIGNATURE	PR00388A 10.45 2.778e-09 86-105
351	BL00018	EF-hand calcium-binding domain proteins.	BL00018 7.41 3.118e-11 160-173 BL00018 7.41 2.350e-10 244-257
354	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 1.947e-09 256-267
358	DM01206	CORONAVIRUS NUCLEOCAPSID PROTEIN.	DM01206B 10.69 3.278e-09 175-195 DM01206B 10.69 6.696e-09 183-203 DM01206B 10.69 8.633e-09 132-152 DM01206B 10.69 8.861e-09 181-201 DM01206B 10.69 9.316e-09 177-197
361	PD01498	OXIDASE BIOSYNTHESIS OXIDOREDUCTASE PORP.	PD01498C 24.90 6.880e-14 219-263
362	PD01498	OXIDASE BIOSYNTHESIS OXIDOREDUCTASE PORP.	PD01498C 24.90 6.880e-14 219-263
365	BL00178	Aminoacyl-transfer RNA synthetases class I proteins.	BL00178B 7.11 1.000e-11 589-600 BL00178A 14.23 9.100e-09 46-56
366	BL00523	Sulfatases proteins.	BL00523E 19.27 1.000e-23 318-348 BL00523A 13.36 5.500e-16 30-47 BL00523B 8.64 1.964e-13 78-90 BL00523C 12.64 9.625e-13 129-140 BL00523G 9.46 5.500e-10 506-516
369	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 4.818e-09 21-52
370	BL00880	Acyl-CoA-binding protein.	BL00880 17.52 1.000e-40 75-125
371	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 1.000e-23 276-307 BL00107B 13.31 1.692e-12 342-358
372	PR00211	GLUTELIN SIGNATURE	PR00211B 0.86 6.602e-11 326-347 PR00211B 0.86 6.106e-10 320-341 PR00211B 0.86 3.167e-09 333-354
373	BL00279	Membrane attack complex components / perforin proteins.	BL00279E 37.11 9.349e-10 749-797
375	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 1.231e-33 10-49
377	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 7.563e-28 10-49
379	BL00598	Chromo domain proteins.	BL00598 14.45 5.781e-16 3-25

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
380	PR00413	HALOACID DEHALOGENASE/EPOXIDE HYDROLASE FAMILY SIGNATURE	PR00413D 11.28 8.941e-09 864-878
383	PR00413	HALOACID DEHALOGENASE/EPOXIDE HYDROLASE FAMILY SIGNATURE	PR00413D 11.28 8.941e-09 864-878
387	BL01060	Flagella transport protein flp family proteins.	BL01060A 15.65 1.535e-09 131-174
388	PR00209	ALPHA/BETA GLIADIN FAMILY SIGNATURE	PR00209B 4.88 6.318e-11 1009-1028
389	PR00837	ALLERGEN V5/TPX-1 FAMILY SIGNATURE	PR00837B 11.64 1.000e-10 469-483
391	BL00240	Receptor tyrosine kinase class III proteins.	BL00240B 24.70 7.907e-10 118-142
392	PR00014	FIBRONECTIN TYPE III REPEAT SIGNATURE	PR00014D 12.04 8.412e-10 691-706
393	PR00014	FIBRONECTIN TYPE III REPEAT SIGNATURE	PR00014D 12.04 8.412e-10 706-721
394	BL01209	LDL-receptor class A (LDLRA) domain proteins.	BL01209 9.31 3.368e-15 47-60 BL01209 9.31 5.500e-13 92-105
395	BL00634	Ribosomal protein L30 proteins.	BL00634 34.38 4.090e-13 70-121
396	BL01013	Oxysterol-binding protein family proteins.	BL01013D 26.81 8.000e-26 358-402 BL01013A 25.14 7.231e-21 45-81 BL01013C 9.97 1.000e-13 132-142 BL01013B 11.33 1.000e-11 110-121
397	BL00930	Peripherin / rom-1 proteins.	BL00930E 17.80 1.000e-40 56-92 BL00930D 9.12 4.632e-37 12-56 BL00930F 16.91 2.800e-36 92-133
400	PR00780	LEUSERPIN 2 SIGNATURE	PR00780B 4.89 4.491e-09 262-285
401	PR00819	CBXX/CFQX SUPERFAMILY SIGNATURE	PR00819B 10.83 7.158e-11 4-20
403	BL00381	Endopeptidase Clp serine proteins.	BL00381C 2.05 1.2 32 1 194 BL00381A 15.48 2.286e-22 74-111 BL00381B 21.42 8.326e-14 78-130
405	BL01105	Ribosomal protein L35Ae proteins.	BL01105A 17.37 1.000e-40 4-49 BL01105B 12.95 1.000e-40 68-108
406	BL00344	GATA-type zinc finger domain proteins.	BL00344 17.99 7.000e-12 814-852
407	PR00211	GLUTELIN SIGNATURE	PR00211B 0.86 9.750e-09 73-94
409	PR00910	LUTEOVIRUS ORF6 PROTEIN SIGNATURE	PR00910A 2.51 4.321e-09 9-22
410	BL00762	WHEP-TRS domain proteins.	BL00762A 23.43 1.000e-28 752-789 BL00762A 23.43 4.400e-21 903-940 BL00762A 23.43 5.415e-18 825-862 BL00762B 16.14 8.759e-12 1154-1168
412	BL00690	DEAH-box subfamily ATP-dependent helicases proteins.	BL00690B 13.38 5.320e-15 262-280 BL00690A 6.87 1.818e-13 230-240
415	BL00227	Tubulin subunits alpha, beta, and gamma proteins.	BL00227B 19.29 1.000e-40 52-107 BL00227C 25.48 1.000e-40 113-165 BL00227D 18.46 1.000e-40 222-276 BL00227F 21.16 1.000e-40 382-436 BL00227E 24.15 1.750e-34 326-361

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			BL00227A 24.55 1.000e-33 1-35
416	PF00992	Troponin.	PF00992A 16.67 1.711e-09 557-592
418	BL00541	Nuclear transition protein 1 proteins.	BL00541 8.44 9.875e-09 256-310
419	BL00541	Nuclear transition protein 1 proteins.	BL00541 8.44 9.875e-09 197-251
420	PF00856	SET domain proteins.	PF00856A 26.14 9.074e-13 901-938 PF00856B 16.42 2.397e-12 951-973
421	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 8.200e-12 33-44
423	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 8.600e-30 130-169
424	PF00564	Octicosapeptide repeat proteins.	PF00564B 24.74 1.305e-17 421-472
426	PR00988	URIDINE KINASE SIGNATURE	PR00988A 6.39 4.569e-12 3-21
427	PR00988	URIDINE KINASE SIGNATURE	PR00988A 6.39 4.569e-12 3-21
428	BL00478	LIM domain proteins.	BL00478B 14.79 3.250e-13 115-130 BL00478B 14.79 9.036e-13 50-65
431	BL00282	Kazal serine protease inhibitors family proteins.	BL00282 16.88 8.875e-12 464-487
432	PD00930	PROTEIN GTPASE DOMAIN ACTIVATION.	PD00930B 33.72 7.800e-18 316-357 PD00930A 25.62 9.617e-12 125-151 PD00930B 33.72 2.521e-10 214-255
433	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 4.649e-34 34-73
434	PR00449	TRANSFORMING PROTEIN P21 RAS SIGNATURE	PR00449A 13.20 7.563e-11 56-78
436	PR00120	H ⁺ -TRANSPORTING ATPASE (PROTON PUMP) SIGNATURE	PR00120C 9.90 5.800e-19 705-722
437	BL00115	Eukaryotic RNA polymerase II heptapeptide repeat proteins.	BL00115T 8.45 7.273e-29 1208-1242 BL00115Q 18.08 2.776e-21 953-983 BL00115Y 11.86 8.000e-17 1604-1650 BL00115M 19.19 3.130e-15 731-774 BL00115A 14.34 9.392e-16 463-496 BL00115A 15.44 7.414e-15 43-82 BL00115R 6.50 6.128e-14 983-1010 BL00115J 16.71 9.289e-14 591-617 BL00115I 8.33 4.336e-13 535-590 BL00115L 12.25 5.939e-13 662-694 BL00115G 11.65 6.011e-13 435-463 BL00115K 15.03 3.417e-10 617-659 BL00115O 16.76 5.805e-10 863-913 BL00115P 11.54 7.538e-10 913-953 BL00115S 18.24 7.968e-10 1010-1052 BL00115U 10.34 4.475e-09 1242-1265
438	PF00628	PHD-finger.	PF00628 15.84 4.536e-10 219-234
440	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 6.351e-34 10-49
441	PR00309	ARRESTIN SIGNATURE	PR00309A 9.68 5.250e-24 32-55 PR00309D 7.09 4.938e-23 290-309 PR00309B 7.81 2.800e-21 69-88 PR00309C 8.22 1.621e-19 165-183 PR00309E 9.82 9.438e-15 374-389
442	BL00600	Aminotransferases class-III pyridoxal-	BL00600B 19.60 7.324e-14 103-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		phosphate attachment si.	129 BL00600G 12.43 2.125e-12 306-325 BL00600F 8.77 8.105e-12 271-284 BL00600E 16.43 3.167e-11 228-257 BL00600D 8.71 8.650e-09 207-221
443	BL00972	Ubiquitin carboxyl-terminal hydrolases family 2 proteins.	BL00972A 11.93 3.160e-18 69-87
444	BL00349	CTF/NF-I proteins.	BL00349A 10.07 1.000e-40 8-54 BL00349C 9.33 1.000e-40 82-125 BL00349E 10.79 1.000e-40 152-195 BL00349F 11.81 1.000e-40 213-255 BL00349H 15.70 7.387e-36 361-399 BL00349B 10.51 2.227e-34 54-82 BL00349D 11.70 9.100e-34 125-152 BL00349G 19.72 5.781e-30 323-356
445	BL00154	E1-E2 ATPases phosphorylation site proteins.	BL00154F 8.23 8.941e-21 271-295 BL00154E 20.37 2.620e-15 124-165
448	DM00215	PROLINE-RICH PROTEIN 3.	DM00215 19.43 4.882e-11 82-115 DM00215 19.43 6.492e-09 87-120
451	BL01283	T-box domain proteins.	BL01283A 24.15 3.100e-40 112-160 BL01283D 11.70 6.000e-39 253-286 BL01283B 23.17 6.538e-38 170-212 BL01283C 13.05 7.750e-19 222-236
452	PR00420	AROMATIC-RING HYDROXYLASE (FLAVOPROTEIN MONOOXYGENASE) SIGNATURE	PR00420A 14.78 2.579e-11 3-26
453	PR00162	RIESKE 2FE-2S SUBUNIT SIGNATURE	PR00162B 12.77 7.429e-17 215-228 PR00162A 9.35 2.324e-14 193-205 PR00162C 8.10 7.120e-14 227-240
454	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 7.000e-30 87-126
456	BL00027	'Homeobox' domain proteins.	BL00027 25.43 9.333e-18 1149-1192
457	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 2.737e-24 16-55
459	BL00290	Immunoglobulins and major histocompatibility complex proteins.	BL00290A 20.89 1.529e-14 154-177 BL00290B 13.17 9.000e-12 214-232
460	PR00413	HALOACID DEHALOGENASE/EPOXIDE HYDROLASE FAMILY SIGNATURE	PR00413F 14.91 7.333e-11 193-214 PR00413E 15.78 5.714e-09 175-192
463	PR00759	BASIC PROTEASE (KUNITZ-TYPE) INHIBITOR FAMILY SIGNATURE	PR00759B 11.26 8.385e-09 74-85
466	BL00019	Actinin-type actin-binding domain proteins.	BL00019D 15.33 4.200e-19 300-330
467	BL00019	Actinin-type actin-binding domain proteins.	BL00019D 15.33 4.200e-19 300-330
469	PR00153	CYCLOPHILIN PEPTIDYL-PROLYL CIS-TRANS ISOMERASE SIGNATURE	PR00153D 11.99 3.250e-15 510-523 PR00153C 11.01 4.682e-14 495-511 PR00153E 9.10 8.548e-14 523-539 PR00153B 11.57 1.720e-13 452-465
470	BL00491	Aminopeptidase P and proline dipeptidase proteins.	BL00491C 12.15 3.912e-09 557-572
471	PD00289	PROTEIN SH3 DOMAIN REPEAT	PD00289 9.97 1.000e-14 1482-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		PRESYNA.	1496 PD00289 9.97 8.650e-11 1122-1136
474	BL50040	Elongation factor 1 gamma chain profile.	BL50040D 17.41 1.000e-40 279-329 BL50040E 18.79 1.000e-40 333-388 BL50040F 18.99 5.320e-40 390-428 BL50040C 22.62 3.739e-38 141-184 BL50040B 13.65 7.000e-30 59-85 BL50040A 12.98 1.450e-14 10-22
475	BL01144	Ribosomal protein L31e proteins.	BL01144 25.07 1.000e-40 22-74
476	PR00007	COMPLEMENT C1Q DOMAIN SIGNATURE	PR00007C 15.60 2.421e-21 589-611 PR00007B 14.16 3.500e-21 544-564 PR00007A 19.33 6.897e-20 517-544 PR00007D 9.64 6.571e-12 623-634
477	BL50002	Src homology 3 (SH3) domain proteins profile.	BL50002A 14.19 5.846e-10 170-189
479	DM01970	0 kw ZK632.12 YDR313C ENDOSOMAL III.	DM01970B 8.60 9.500e-17 967-980
480	PR00868	DNA-POLYMERASE FAMILY A (POL I) SIGNATURE	PR00868C 13.76 5.688e-17 284-308 PR00868A 16.33 3.186e-13 224-247 PR00868H 12.51 3.388e-13 431-448 PR00868I 10.87 7.938e-11 462-476 PR00868E 13.19 1.608e-10 340-366
481	BL00027	'Homeobox' domain proteins.	BL00027 26.43 9.182e-22 53-96
482	BL00061	Short-chain dehydrogenases/reductases family proteins.	BL00061B 25.79 3.647e-21 188-226
483	BL50002	Src homology 3 (SH3) domain proteins profile.	BL50002A 14.19 1.750e-12 1032-1051
485	PF00023	Ank repeat proteins.	PF00023A 16.03 9.625e-10 760-776 PF00023A 16.03 3.571e-09 715-731
486	PD02870	RECEPTOR INTERLEUKIN-1 PRECURSOR.	PD02870B 18.83 9.262e-20 103-131 PD02870A 18.74 9.476e-02 201-236
487	PR00370	FLAVIN-CONTAINING MONOOXYGENASE (FMO) SIGNATURE	PR00370G 15.45 3.769e-28 471-493 PR00370E 10.91 1.000e-24 27-46 PR00370C 12.72 4.000e-21 140-157 PR00370E 11.96 9.229e-21 320-339 PR00370D 16.33 1.750e-20 185-204 PR00370F 17.75 7.395e-20 375-395 PR00370A 3.35 2.038e-18 4-20
489	PD01675	GLYCOPROTEIN MAJOR ENVELOPE PROBABLE U3.	PD01675C 19.89 2.330e-10 55-89
492	BL00211	ABC transporters family proteins.	BL00211A 12.23 5.050e-09 45-57
493	BL00211	ABC transporters family proteins.	BL00211A 12.23 5.050e-09 45-57
494	BL00211	ABC transporters family proteins.	BL00211A 12.23 5.050e-09 58-70
495	BL00027	'Homeobox' domain proteins.	BL00027 26.43 6.786e-12 509-552 BL00027 26.43 9.143e-12 319-362 BL00027 26.43 2.600e-11 627-670 BL00027 26.43 3.625e-10 779-822
497	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 5.800e-22 214-245 BL00107B 13.31 1.000e-13 281-297 BL00107A 18.39 3.520e-13 583-614 BL00107B 13.31 8.615e-12 652-668
499	BL00383	Tyrosine specific protein phosphatases	BL00383E 10.35 1.000e-14 1902-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		proteins.	1913 BL00383D 11.92 3.077e-14 1862-1875 BL00383A 13.34 5.500e-14 1730-1745 BL00383C 10.10 2.000e-13 1785-1796 BL00383F 15.51 9.069e-12 1940- 1956 BL00383B 7.61 1.692e-11 1755-1764
501	PR00019	LEUCINE-RICH REPEAT SIGNATURE	PR00019B 11.36 1.360e-09 136- 150 PR00019A 11.19 1.667e-09 91-105 PR00019B 11.36 4.600e- 09 160-174
503	BL00226	Intermediate filaments proteins.	BL00226D 19.10 1.000e-40 367- 414 BL00226B 23.86 6.143e-27 195-243 BL00226A 12.77 7.840e- 14 96-111 BL00226C 13.23 2.600e-13 309-340 BL00226C 13.23 6.143e-12 266-297 BL00226B 23.86 1.209e-09 146- 194
505	PD02407	3-BISPHOSPHOGLYCERATE-INDEPENDENT PHOSPHOGLYCER.	PD02407F 7.61 6.739e-09 916- 930
506	PF00632	HECT-domain (ubiquitin-transferase).	PF00632C 20.66 9.830e-19 991- 1023 PF00632B 18.45 1.155e-11 940-968
507	BL01082	Ribosomal protein L7Ae proteins.	BL01082 20.37 4.273e-20 76-116
508	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 2.421e-09 493-504
509	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 2.421e-09 473-484
510	PR00320	G-PROTEIN BETA WD-40 REPEAT SIGNATURE	PR00320B 12.19 4.774e-11 567- 582 PR00320B 12.19 5.886e-10 763-778 PR00320C 13.01 6.760e- 10 567-582 PR00320A 16.74 7.618e-10 846-861 PR00320A 16.74 3.415e-09 763-778 PR00320A 16.74 6.268e-09 567- 582
511	BL00479	Phorbol esters / diacylglycerol binding domain proteins.	BL00479C 11.11 3.250e-12 170- 183
512	BL50058	G-protein gamma subunit profile.	BL50058 27.23 7.494e-09 10-58
513	BL00524	Somatomedin B domain proteins.	BL00524A 9.65 8.925e-14 80-101
515	BL00041	Bacterial regulatory proteins, araC family proteins.	BL00041 23.99 1.964e-19 492-524
516	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 8.500e-13 391-404
517	BL00415	Synapsins proteins.	BL00415E 4.82 9.291e-09 959- 996
518	PR00109	TYROSINE KINASE CATALYTIC DOMAIN SIGNATURE	PR00109B 12.27 9.471e-12 126- 145
519	BL00290	Immunoglobulins and major histocompatibility complex proteins.	BL00290B 13.17 4.750e-09 47-65
522	PR00505	D12 CLASS N6 ADENINE-SPECIFIC DNA METHYLTRANSFERASE SIGNATURE	PR00505A 14.15 7.128e-09 364- 381
525	BL00312	Glycophorin A proteins.	BL00312B 9.22 5.781e-10 891- 920
528	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 2.500e-32 16-55
529	PR00254	NICOTINIC ACETYLCHOLINE RECEPTOR SIGNATURE	PR00254D 15.50 4.000e-17 131- 150 PR00254A 11.23 4.706e-14 61-78 PR00254C 11.36 4.000e-12

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			113-126 PR00254B 12.97 1.486e-11 95-110
531	BL00741	Guanine-nucleotide dissociation stimulators CDC24 family sign.	BL00741B 14.27 6.870e-16 787-810
532	PR00193	MYOSIN HEAVY CHAIN SIGNATURE	PR00193D 14.36 3.143e-34 447-476 PR00193C 12.60 7.632e-32 216-244 PR00193B 11.69 7.750e-29 167-193 PR00193A 15.41 2.588e-22 111-131 PR00193E 19.47 2.200e-21 501-530
533	PD02870	RECEPTOR INTERLEUKIN-1 PRECURSOR.	PD02870B 18.83 5.596e-09 348-381
535	PR00683	SPECTRIN PLECKSTRIN HOMOLOGY DOMAIN SIGNATURE	PR00683D 15.87 2.452e-10 465-484
536	BL00027	'Homeobox' domain proteins.	BL00027 26.43 6.684e-24 164-207
538	PR00239	MOLLUSCAN RHODOPSIN C-TERMINAL TAIL SIGNATURE	PR00239E 1.58 2.739e-09 225-237
539	BL00406	Actins proteins.	BL00406C 6.75 1.000e-40 157-212 BL00406B 5.47 6.143e-37 90-145 BL00406D 12.58 4.600e-36 291-346 BL00406E 8.44 2.200e-33 364-414 BL00406A 9.95 4.441e-23 7-42
540	PR00456	RIBOSOMAL PROTEIN P2 SIGNATURE	PR00456E 3.06 9.625e-10 44-59
541	PR00456	RIBOSOMAL PROTEIN P2 SIGNATURE	PR00456E 3.06 9.625e-10 44-59
542	PF00023	Ank repeat proteins.	PF00023A 16.03 7.857e-11 138-154
544	PF00642	Zinc finger C-x8-C-x5-C-x3-H type (and similar).	PF00642 11.59 9.082e-10 838-849
546	BL00383	Tyrosine specific protein phosphatases proteins.	BL00383E 10.35 4.115e-10 104-115
547	BL01226	Hydroxymethylglutaryl-coenzyme A lyase proteins.	BL01226A 13.79 1.000e-40 50-83 BL01226C 13.51 1.000e-40 127-167 BL01226D 13.79 1.000e-40 174-210 BL01226E 13.74 1.000e-40 212-253 BL01226H 13.74 1.000e-40 386-434 BL01226F 25.06 1.000e-40 460-508 BL01226G 15.76 3.483e-32 292-321 BL01226B 13.35 1.818e-31 95-127 BL01226F 9.78 8.714e-23 253-271
549	BL00964	Syndecans proteins.	BL00964B 12.05 2.426e-10 1246-1289
551	DM01930	2 kw FINGER SMCX SMCY YDR096W.	DM01930B 15.41 1.367e-37 170-215 DM01930F 14.16 8.232e-28 267-303 DM01930B 19.86 9.163e-10 37-71
552	BL00195	Glutaredoxin proteins.	BL00195B 15.31 7.158e-09 9-29
554	BL00383	Tyrosine specific protein phosphatases proteins.	BL00383E 10.35 2.756e-12 436-447
555	PR00403	WW DOMAIN SIGNATURE	PR00403B 12.19 7.612e-11 122-137 PR00403A 16.82 3.912e-10 107-121 PR00403B 12.19 2.068e-09 76-91
558	PR00380	KINESIN HEAVY CHAIN SIGNATURE	PR00380A 14.18 2.714e-26 76-98 PR00380D 9.93 3.000e-24 275-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			297 PR00380C 13.18 5.154e-20 226-245 PR00380B 12.64 9.400e-20 195-213
559	BL00518	Zinc finger, C3HC4 type (RING finger), proteins.	BL00518 12.23 5.333e-09 522-531
561	PD01795	PROTEIN AMINOPEPTIDASE PRECURSOR HYDROLASE SIGNA.	PD01795B 11.56 2.333e-12 159-172 PD01795A 10.27 1.000e-09 135-144
562	PD01795	PROTEIN AMINOPEPTIDASE PRECURSOR HYDROLASE SIGNA.	PD01795B 11.56 2.333e-12 110-123 PD01795A 10.27 1.000e-09 86-95
563	BL00018	EF-hand calcium-binding domain proteins.	BL00018 7.41 1.391e-09 41-54
565	BL00348	p53 tumor antigen proteins.	BL00348F 23.19 4.143e-09 188-231
567	PD00301	PROTEIN REPEAT MUSCLE CALCIUM-BL.	PD00301B 5.49 4.115e-09 284-295
569	PF00850	Histone deacetylase family.	PF00850E 8.88 6.553e-21 756-782 PF00850D 14.76 1.519e-16 722-746 PF00850F 15.70 1.118e-11 794-827 PF00850G 22.75 8.375e-11 833-875
570	PD00289	PROTEIN SH3 DOMAIN REPEAT PRESYN.	PD00289 9.97 4.960e-10 137-151
571	BL00518	Zinc finger, C3HC4 type (RING finger), proteins.	BL00518 12.23 8.800e-11 44-53
573	BL00299	Ubiquitin domain proteins.	BL00299 28.84 1.123e-11 123-175
574	PF01140	Matrix protein (MA), p15.	PF01140D 15.54 3.700e-10 986-1021
576	BL00284	Serpins proteins.	BL00284C 28.56 5.200e-26 200-242 BL00284A 15.64 4.913e-18 71-95 BL00284B 17.99 7.261e-15 173-194 BL00284D 16.34 5.846e-13 306-333 BL00284E 19.15 2.28e-11 41-54
579	PD01066	PROTEIN ZINC FINGER METAL-BINDING NUCLEIC ACID-BINDING	PD01066 19.43 6.553e-29 15
580	BL50001	Src homology 2 (SH2) domain proteins profile.	BL50001B 17.40 4.500e-12 1010-1031
581	PD00930	PROTEIN GTPASE DOMAIN ACTIVATION.	PD00930B 33.72 3.189e-22 608-649 PD00930A 25.62 6.806e-17 505-531
584	BL00612	Osteonectin domain proteins.	BL00612B 11.35 2.034e-11 93-126
585	DM01551	kw OSTEOINDUCTIVE YOPM MEMBRANE OUTER.	DM01551C 14.62 8.859e-10 102-122
586	PF00628	PHD-finger.	PF00628 15.84 3.455e-12 235-250
587	BL00027	'Homeobox' domain proteins.	BL00027 26.43 6.063e-10 85-128
588	PR00326	GTP1/OBG GTP-BINDING PROTEIN FAMILY SIGNATURE	PR00326A 8.75 7.525e-16 227-248 PR00326C 9.79 6.760e-15 276-292 PR00326D 19.09 6.657e-13 293-312 PR00326B 16.74 9.229e-13 248-267
589	BL00422	Granins proteins.	BL00422A 28.34 7.429e-09 2349-2378
590	BL00415	Synapsins proteins.	BL00415N 4.29 9.794e-10 295-339
591	BL00128	Alpha-lactalbumin / lysozyme C proteins.	BL00128A 20.76 3.423e-13 35-65 BL00128C 19.34 2.980e-11 110-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			132
596	PR00049	WILM'S TUMOUR PROTEIN SIGNATURE	PR00049D 0.00 3.136e-09 31-46
597	DM00547	1 kw CHROMO BROMODOMAIN SHADOW GLOBAL.	DM00547C 17.30 1.667e-19 207-229 DM00547E 13.94 6.200e-18 319-342 DM00547B 11.28 1.000e-17 179-193 DM00547D 11.60 9.250e-13 289-303 DM00547F 23.43 6.727e-12 679-726 DM00547A 12.38 4.818e-11 158-170
600	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 1.882e-27 13-52
601	BL00192	Cytochrome b/b6 heme-ligand proteins.	BL00192A 11.90 6.400e-09 390-430
602	BL00936	Ribosomal protein L35 proteins.	BL00936B 27.27 8.615e-09 118-157
603	BL00936	Ribosomal protein L35 proteins.	BL00936B 27.27 8.615e-09 118-157
606	PR00019	LEUCINE-RICH REPEAT SIGNATURE	PR00019B 11.36 7.300e-10 292-306 PR00019A 11.19 5.667e-09 323-337
607	PR00019	LEUCINE-RICH REPEAT SIGNATURE	PR00019B 11.36 7.300e-10 292-306 PR00019A 11.19 5.667e-09 323-337
608	PR00320	G-PROTEIN BETA WD-40 REPEAT SIGNATURE	PR00320C 13.01 9.500e-12 168-183 PR00320A 16.74 2.853e-10 60-75 PR00320A 16.74 4.706e-10 14-29 PR00320C 13.01 5.320e-10 60-75 PR00320C 13.01 5.680e-10 14-29 PR00320A 16.74 6.049e-09 217-232 PR00320B 12.19 8.875e-09 168-183
610	BL00750	Chaperonins TCP-1 proteins.	BL00750B 16.17 1.000e-40 70-120 BL00750A 20.17 6.211e-31 26-69 BL00750G 20.12 8.000e-31 431-471 BL00750F 18.40 5.100e-30 370-411 BL00750E 24.59 8.650e-29 295-332 BL00750H 21.44 1.000e-27 489-524 BL00750C 25.65 5.345e-17 149-181 BL00750D 16.16 6.318e-14 203-222
613	BL00766	Tetrahydrofolate dehydrogenase/cyclohydrolase proteins.	BL00766B 24.49 1.000e-40 142-190 BL00766E 13.78 1.000e-40 322-359 BL00766C 25.86 5.500e-39 208-256 BL00766D 17.05 4.536e-26 283-313 BL00766A 21.48 6.063e-24 102-132
615	BL00256	Adipokinetic hormone family proteins.	BL00256 12.28 3.298e-10 746-755
616	BL00319	Amyloidogenic glycoprotein extracellular domain proteins.	BL00319C 17.12 9.053e-09 419-453
617	BL00030	Eukaryotic RNA-binding region RNP-1 proteins.	BL00030A 14.39 4.429e-09 44-63
618	BL00030	Eukaryotic RNA-binding region RNP-1 proteins.	BL00030A 14.39 4.429e-09 44-63
620	BL00325	Actin-depolymerizing proteins.	BL00325B 21.66 5.817e-16 77-123
622	BL00972	Ubiquitin carboxyl-terminal hydrolases	BL00972A 11.93 5.500e-19 213-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		family 2 proteins.	231 BL00972D 22.55 2.742e-16 501-526 BL00972B 9.45 1.000e-11 297-307 BL00972C 16.48 3.160e-11 370-385 BL00972E 20.72 7.517e-10 526-548
625	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 6.333e-39 6-45
628	BL00039	DEAD-box subfamily ATP-dependent helicases proteins.	BL00039D 21.67 7.750e-31 478-524 BL00039A 18.44 2.000e-25 198-237 BL00039C 15.63 1.844e-15 327-351 BL00039B 19.19 5.636e-14 242-268
630	PD00306	PROTEIN GLYCOPROTEIN PRECURSOR RE.	PD00306A 10.26 7.000e-12 232-246
631	PD00306	PROTEIN GLYCOPROTEIN PRECURSOR RE.	PD00306A 10.26 7.000e-12 290-304
633	BL00785	5'-nucleotidase proteins.	BL00785C 9.45 3.625e-16 108-122 BL00785E 15.85 4.000e-16 279-295 BL00785A 9.73 6.500e-14 29-40 BL00785B 10.65 5.500e-13 72-86 BL00785D 9.89 4.000e-12 135-145
636	PR00832	PAXILLIN SIGNATURE	PR00832E 14.43 9.901e-14 85-108
637	PR00109	TYROSINE KINASE CATALYTIC DOMAIN SIGNATURE	PR00109B 12.27 6.362e-13 221-240
638	PF00635	MSP (Major sperm protein) domain proteins.	PF00635B 15.84 4.900e-11 463-502
639	PR00860	VERTEBRATE METALLOTHIONEIN SIGNATURE	PR00860B 7.04 1.900e-18 85-99 PR00860C 9.61 1.474e-14 99-109 PR00860A 5.46 1.720e-14 63-76
641	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 4.462e-15 271-284 PD00066 13.92 4.462e-15 299-312 PD00066 13.92 2.800e-14 327-340 PD00066 13.92 2.800e-14 383-396 PD00066 13.92 2.800e-14 407-424 PD00066 13.92 7.000e-14 355-368 PD00066 13.92 8.800e-14 439-452 PD00066 13.92 8.800e-14 495-508 PD00066 13.92 1.500e-13 551-564 PD00066 13.92 7.000e-13 467-480 PD00066 13.92 7.000e-13 523-536 PD00066 13.92 9.500e-13 215-228 PD00066 13.92 9.500e-13 243-256 PD00066 13.92 9.500e-13 579-592 PD00066 13.92 8.615e-10 607-620 PD00066 13.92 1.600e-09 187-200
642	BL00961	Ribosomal protein S28e proteins.	BL00961B 11.24 7.429e-37 67-100 BL00961A 9.90 4.079e-26 42-66
643	BL00585	Ribosomal protein S5 proteins.	BL00585A 28.43 1.391e-40 103-155 BL00585B 18.78 3.250e-30 193-230
647	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 9.400e-10 181-192
648	PR00876	NEMATODE METALLOTHIONEIN SIGNATURE	PR00876C 6.15 9.229e-09 112-126
652	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 5.941e-27 29-68
653	BL00047	Histone H4 proteins.	BL00047A 13.53 1.000e-40 2-41

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			BL00047B 6.51 1.429e-40 41-74 BL00047C 12.18 1.310e-38 74-104
654	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 4.109e-25 30-69
655	BL01115	GTP-binding nuclear protein ran proteins.	BL01115A 10.22 3.483e-17 19-63
657	BL00518	Zinc finger, C3HC4 type (RING finger), proteins.	BL00518 12.23 8.286e-10 31-40
658	BL00125	Serine/threonine specific protein phosphatases proteins.	BL00125B 21.48 1.000e-40 89-135 BL00125C 19.97 1.000e-40 153-200 BL00125D 33.11 1.000e-40 213-268 BL00125A 14.83 8.941e-38 47-84
659	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 8.200e-16 492-505 PD00066 13.92 9.308e-15 380-393 PD00066 13.92 6.000e-13 352-365 PD00066 13.92 7.000e-13 240-253 PD00066 13.92 7.500e-13 268-281 PD00066 13.92 7.500e-13 408-421 PD00066 13.92 2.174e-11 464-477 PD00066 13.92 1.000e-10 436-449
660	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 2.189e-26 29-68
661	BL00795	Involucrin proteins.	BL00795C 17.06 7.882e-15 193-238 BL00795C 17.06 3.797e-13 187-232 BL00795C 17.06 5.014e-13 188-233 BL00795C 17.06 4.506e-12 196-241 BL00795C 17.06 7.896e-12 191-236 BL00795C 17.06 1.667e-11 185-230 BL00795C 17.06 2.000e-11 198-243 BL00795C 17.06 3.778e-11 171-216 BL00795C 17.06 6.111e-11 197-242 BL00795C 17.06 6.44e-11 194-237 BL00795C 17.06 8.000e-11 189-234 BL00795C 17.06 8.556e-11 192-237 BL00795C 17.06 1.733e-10 195-240 BL00795C 17.06 2.779e-10 184-229 BL00795C 17.06 4.035e-10 199-244 BL00795C 17.06 5.081e-10 186-231 BL00795C 17.06 6.965e-10 190-235 BL00795C 17.06 2.700e-09 200-245 BL00795C 17.06 5.800e-09 175-220 BL00795C 17.06 6.500e-09 182-227 BL00795C 17.06 6.600e-09 201-246 BL00795C 17.06 6.600e-09 202-247 BL00795C 17.06 6.600e-09 208-253
662	BL00469	Nucleoside diphosphate kinases proteins.	BL00469 22.22 1.000e-40 149-204
663	BL01160	Kinesin light chain repeat proteins.	BL01160B 19.54 9.411e-11 331-385
664	BL00601	Tryptophan pentad repeat proteins (TRF family) proteins.	BL00601A 20.29 5.500e-23 7-46 BL00601B 20.92 3.631e-13 69-98
665	BL00082	Extradiol ring-cleavage dioxygenases proteins.	BL00082A 19.07 8.615e-12 49-72
666	DM01537	kw SKI2W SKI2 NUCLEOLAR	DM01537B 21.63 4.073e-37 834-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		HELICASE.	881 DM01537B 21.63 9.750e-21 1669-1716 DM01537A 15.14 8.650e-18 698-718 DM01537A 15.14 6.766e-12 1537-1557
667	DM01537	kw SKI2W SKI2 NUCLEOLAR HELICASE.	DM01537B 21.63 7.923e-38 820- 867 DM01537B 21.63 9.750e-21 1655-1702 DM01537A 15.14 8.650e-18 684-704 DM01537A 15.14 6.766e-12 1523-1543
669	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 6.786e-24 849- 880 BL00107B 13.31 6.727e-13 916-932
670	BL00299	Ubiquitin domain proteins.	BL00299 28.84 9.735e-27 37-89
671	BL00027	'Homeobox' domain proteins.	BL00027 26.43 6.571e-12 432-475
676	PR00861	ALPHA-LYTIC ENDOPEPTIDASE SERINE PROTEASE (S2A) SIGNATURE	PR00861E 9.88 2.385e-09 206- 221
678	BL00225	Crystallins beta and gamma 'Greek key' motif proteins.	BL00225B 18.06 7.517e-24 1805- 1840 BL00225B 18.06 8.297e-20 1987-2022 BL00225B 18.06 2.575e-19 1896-1931 BL00225B 18.06 8.200e-19 175-210 BL00225B 18.06 8.200e-19 1698- 1733 BL00225B 18.06 4.808e-14 73-108 BL00225B 18.06 4.808e- 14 1596-1631 BL00225B 18.06 5.500e-14 2077-2112 BL00225A 13.82 5.829e-12 2043-2064 BL00225A 13.82 3.127e-09 1759- 1780
679	PR00320	G-PROTEIN BETA WD-40 REPEAT SIGNATURE	PR00320C 13.01 4.240e-10 169- 184 PR00320A 16.74 6.294e-10 169-184
680	BL00243	Integrins beta chain cysteine-rich domain proteins.	BL00243I 31.77 1.143e-11 172- 212
681	PR00852	XERODERMA PIGMENTOSUM GROUP D PROTEIN SIGNATURE	PR00852E 15.90 1.000e-29 612- 634 PR00852E 8.14 3.769e-27 348-371 PR00852D 11.38 8.875e- 27 309-331 PR00852B 11.08 2.800e-25 249-269 PR00852I 17.26 3.500e-25 683-704 PR00852F 11.85 5.909e-24 379- 398 PR00852G 16.19 4.462e-23 468-486 PR00852C 8.81 9.143e- 23 284-303
682	BL50058	G-protein gamma subunit profile.	BL50058 27.23 1.375e-35 15-63
685	BL00972	Ubiquitin carboxyl-terminal hydrolases family 2 proteins.	BL00972A 11.93 7.500e-20 40-58 BL00972D 22.55 3.903e-16 300- 325 BL00972B 9.45 1.000e-13 120-130 BL00972E 20.72 5.500e- 11 325-347
687	BL00237	G-protein coupled receptors proteins.	BL00237A 27.68 4.273e-14 98- 138
688	BL00388	Proteasome A-type subunits proteins.	BL00388A 23.14 1.000e-40 8-54 BL00388B 31.38 3.864e-33 66- 108 BL00388D 20.71 1.000e-21 153-184 BL00388C 18.79 8.147e- 16 126-148
689	PD02796	PROTEIN STEROL CARRIER LIPID-	PD02796B 20.92 1.105e-15 347-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		TRAN.	394
691	PD01572	PHOTOSYSTEM II REACTION CENTRE T PROTEIN PHOTOS.	PD01572 8.77 4.083e-09 1-31
692	BL00028	Zinc finger, C2H2 type, domain proteins.	BL00028 16.07 7.600e-10 488-505
694	BL01013	Oxysterol-binding protein family proteins.	BL01013A 25.14 9.357e-33 527-563 BL01013D 26.81 8.235e-23 814-858 BL01013C 9.97 6.211e-14 615-625 BL01013B 11.33 3.605e-13 592-603
695	PD00289	PROTEIN SH3 DOMAIN REPEAT PRESYN.	PD00289 9.97 3.571e-13 164-178 PD00289 9.97 8.650e-11 2147-2161 PD00289 9.97 2.552e-09 23-37
698	PR00161	NICKEL-DEPENDENT HYDROGENASE/B-TYPE CYTOCHROME SIGNATURE	PR00161C 9.51 4.930e-09 282-302
700	PR00749	LYSOZYME G SIGNATURE	PR00749F 13.63 8.636e-13 139-156 PR00749H 8.22 3.681e-12 173-194 PR00749B 16.54 1.419e-11 48-70 PR00749C 7.26 3.060e-11 72-91 PR00749A 10.33 4.815e-10 24-45
703	PR00704	CALPAIN CYSTEINE PROTEASE (C2) FAMILY SIGNATURE	PR00704I 9.52 1.000e-29 476-505 PR00704D 11.05 2.500e-27 132-158 PR00704E 12.55 5.500e-27 162-186 PR00704F 13.61 1.000e-22 187-215 PR00704G 13.87 1.237e-21 317-339 PR00704H 13.38 8.138e-21 367-385 PR00704A 14.68 2.125e-19 27-51 PR00704C 11.88 1.257e-17 96-113 PR00704B 17.94 1.833e-15 72-95
705	PR00859	PROKARYOTE METALLOTHIONEIN SIGNATURE	PR00859C 7.06 2.776e-09 94-111
706	BL00225	Intermediate filament proteins.	BL00226D 19.10 3.31e-26 369-416 BL00226B 23.86 3.250e-24 203-251 BL00226C 13.23 8.269e-21 268-299 BL00226A 12.77 8.200e-14 103-118
707	PR00021	SMALL PROLINE-RICH PROTEIN SIGNATURE	PR00021A 4.31 2.440e-10 2-15
708	BL00361	Ribosomal protein S10 proteins.	BL00361B 18.34 5.101e-10 82-105
709	PR00021	SMALL PROLINE-RICH PROTEIN SIGNATURE	PR00021A 4.31 2.200e-10 2-15
710	BL00514	Fibrinogen beta and gamma chains C-terminal domain proteins.	BL00514C 17.41 8.412e-27 160-197 BL00514E 14.28 8.909e-16 219-236 BL00514H 14.95 1.551e-15 317-342 BL00514G 15.98 7.750e-15 284-314 BL00514D 15.35 4.789e-10 201-214
711	PD00930	PROTEIN GTPASE DOMAIN ACTIVATION.	PD00930B 33.72 8.714e-12 49-90
714	BL00400	LBP / BPI / CETP family proteins.	BL00400C 24.53 6.029e-17 158-202 BL00400D 23.26 2.080e-14 222-259 BL00400A 21.59 1.600e-10 27-59
715	BL01154	RNA polymerases L / 13 to 16 Kd	BL01154B 24.55 5.500e-36 40-76

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		subunits proteins.	BL01154A 18.70 3.000e-22 19-40
716	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 9.786e-32 10-49
717	BL00215	Mitochondrial energy transfer proteins.	BL00215A 15.82 9.206e-14 77-102 BL00215A 15.82 8.412e-10 175-200
719	BL00309	Vertebrate galactoside-binding lectin proteins.	BL00309C 18.65 2.241e-09 62-87
726	BL00687	Aldehyde dehydrogenases glutamic acid proteins.	BL00687E 25.37 7.136e-33 266-316 BL00687D 26.00 5.333e-28 151-198 BL00687B 17.54 3.647e-26 39-81 BL00687C 24.13 6.087e-22 96-133 BL00687F 9.55 2.500e-11 352-363
727	DM01354	kw TRANSCRIPTASE REVERSE II ORF2.	DM01354N 13.17 1.000e-40 129-174 DM01354O 8.73 6.605e-15 180-226
734	PD00301	PROTEIN REPEAT MUSCLE CALCIUM-BL.	PD00301A 10.24 6.400e-09 101-112
735	BL01024	Protein phosphatase 2A regulatory subunit PR55 proteins.	BL01024A 10.26 1.000e-40 22-69 BL01024B 8.91 1.000e-40 86-127 BL01024C 7.80 1.000e-40 146-185 BL01024D 13.22 1.000e-40 185-222 BL01024E 11.96 1.000e-40 222-266 BL01024F 9.42 1.000e-40 266-317 BL01024G 11.09 1.000e-40 317-349 BL01024H 13.88 1.000e-40 389-442
736	PF00913	Trypanosome variant surface glycoprotein.	PF00913D 11.90 7.130e-10 24-51
737	PR00700	PROTEIN TYROSINE PHOSPHATASE SIGNATURE	PR00700D 12.47 2.200e-09 82-101
740	PR00320	G-PROTEIN BETA WD-40 REPEAT SIGNATURE	PR00320C 13.01 1.600e-09 68-83 PR00320A 10.74 7.366e-09 62-82
743	PR00871	DNA NUCLEOTIDYLEXOTRANSFERASE (TDT) SIGNATURE	PR00871G 14.48 8.000e-09 115-201
745	BL00518	Zinc finger, C3HC4 type (RING finger), proteins.	BL00518 12.23 2.286e-10 33-42
749	BL00215	Mitochondrial energy transfer proteins.	BL00215A 15.82 5.200e-15 221-246 BL00215A 15.82 7.618e-14 20-45 BL00215A 15.82 8.851e-11 123-148 BL00215B 10.44 9.526e-11 69-82 BL00215B 10.44 7.300e-09 272-285 BL00215B 10.44 8.500e-09 165-178
751	BL50002	Src homology 3 (SH3) domain proteins profile.	BL50002A 14.19 1.000e-14 370-389 BL50002B 15.18 2.200e-10 408-422
752	BL00353	HMG1/2 proteins.	BL00353B 11.47 3.089e-12 390-440
753	PF00622	Domain in SPla and the RYanodine Receptor.	PF00622B 21.00 4.214e-14 47-69
754	BL00211	ABC transporters family proteins.	BL00211A 12.23 8.941e-10 66-78
755	PR00926	MITOCHONDRIAL CARRIER PROTEIN SIGNATURE	PR00926F 17.75 7.750e-19 392-415 PR00926C 16.07 5.935e-17 253-274 PR00926D 10.53 2.059e-15 301-320 PR00926E 11.70

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			4.971e-15 344-363 PR00926B 16.07 9.526e-13 210-225 PR00926A 10.41 1.514e-12 197-211
756	BL01187	Calcium-binding EGF-like domain proteins pattern proteins.	BL01187A 9.98 2.125e-12 324-336 BL01187A 9.98 4.789e-11 377-389 BL01187B 12.04 3.057e-10 439-455
757	PF00651	BTB (also known as BR-C/Ttk) domain proteins.	PF00651 15.00 4.429e-10 43-56
758	PR00055	HIV TAT DOMAIN SIGNATURE	PR00055A 8.13 8.855e-09 144-156
759	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 5.304e-11 110-123
760	PR00448	NSF ATTACHMENT PROTEIN SIGNATURE ^o	PR00448D 12.42 3.455e-27 162-186 PR00448A 10.74 1.273e-22 37-57 PR00448B 16.01 9.379e-21 100-118 PR00448C 11.46 1.000e-20 129-147
765	BL01042	Homoserine dehydrogenase proteins.	BL01042A 13.29 5.909e-11 74-95
766	PR00625	DNAJ PROTEIN FAMILY SIGNATURE	PR00625A 12.84 2.154e-18 26-46 PR00625B 13.48 9.000e-16 57-78
768	BL00762	WHEP-TRS domain proteins.	BL00762A 23.43 8.500e-28 112-149 BL00762B 16.14 3.793e-12 64-78 BL00762A 23.43 6.625e-12 6-43 BL00762C 15.58 4.176e-09 459-472 BL00762D 11.15 9.667e-09 210-220
769	PR00709	AVIDIN SIGNATURE	PR00709A 4.60 1.934e-09 1-20
770	PR00320	G-PROTEIN BETA WD-40 REPEAT SIGNATURE	PR00320C 13.01 1.720e-10 262-277 PR00320A 16.74 2.853e-10 262-277 PR00320C 13.01 4.300e-09 96-111 PR00320B 12.19 5.500e-09 262-277 PR00320A 16.74 6.278e-09 35-70
771	PR00019	LEUCINE-RICH REPEAT SIGNATURE	PR00019B 11.50 1.714e-12 87-101 PR00019A 11.19 1.000e-10 90-104
772	PD02807	APOLIPOPROTEIN E PRECURSOR APO-E GLYCOPROTEIN PLAS.	PD02807C 8.91 6.308e-10 110-159
773	PD02807	APOLIPOPROTEIN E PRECURSOR APO-E GLYCOPROTEIN PLAS.	PD02807C 8.91 6.308e-10 155-204
774	DM00547	1 kw CHROMO BROMODOMAIN SHADOW GLOBAL.	DM00547F 23.43 3.942e-28 943-990 DM00547E 13.94 9.750e-21 652-675 DM00547B 11.28 1.818e-18 518-532 DM00547C 17.30 3.531e-17 546-568 DM00547A 12.38 1.273e-11 497-509 DM00547D 11.60 9.200e-11 622-636
776	PR00779	INOSITOL 1,4,5-TRISPHOSPHATE-BINDING PROTEIN RECEPTOR SIGNATURE	PR00779F 14.51 5.147e-09 769-792
777	PR00779	INOSITOL 1,4,5-TRISPHOSPHATE-BINDING PROTEIN RECEPTOR SIGNATURE	PR00779F 14.51 5.147e-09 742-765
778	PR00779	INOSITOL 1,4,5-TRISPHOSPHATE-BINDING PROTEIN RECEPTOR SIGNATURE	PR00779F 14.51 5.147e-09 742-765

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
779	BL01282	BIR repeat proteins.	BL01282B 30.49 2.543e-09 6-45
781	PR00205	CADHERIN SIGNATURE	PR00205B 11.39 3.118e-11 654-672 PR00205B 11.39 8.588e-11 230-248 PR00205B 11.39 8.527e-10 551-569 PR00205B 11.39 4.203e-09 336-354
783	BL00625	Regulator of chromosome condensation (RCC1) proteins.	BL00625B 17.69 2.167e-19 193-227 BL00625A 16.21 5.500e-17 199-228 BL00625B 17.69 1.885e-16 140-174 BL00625B 17.69 2.770e-16 245-279 BL00625A 16.21 9.115e-16 251-280 BL00625A 16.21 6.507e-14 146-175
785	PF00084	Sushi domain proteins (SCR repeat proteins).	PF00084B 9.45 7.188e-10 595-607 PF00084B 9.45 6.400e-09 656-668
786	PF00084	Sushi domain proteins (SCR repeat proteins).	PF00084B 9.45 7.188e-10 595-607 PF00084B 9.45 6.400e-09 656-668
787	BL00826	MARCKS family proteins.	BL00826C 7.63 6.738e-09 203-230
788	PR00453	VON WILLEBRAND FACTOR TYPE A DOMAIN SIGNATURE	PR00453A 12.79 1.310e-14 36-54 PR00453B 14.65 8.568e-10 75-90
789	PR00102	ORNITHINE CARBAMOYLTRANSFERASE SIGNATURE	PR00102B 14.82 5.418e-09 963-977
790	BL00030	Eukaryotic RNA-binding region RNP-1 proteins.	BL00030B 7.03 5.500e-11 199-209
791	BL00415	Synapsins proteins.	BL00415N 4.29 9.519e-10 393-437 BL00415N 4.29 2.117e-09 103-147 BL00415N 4.29 3.628e-09 97-141 BL00415N 4.29 5.664e-09 387-431
795	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 2.091e-36 105-144
799	PF00731	AIN carbonyl	PF00731C 23.16 7.333e-35 337-380 PF00731B 19.47 7.429e-28 299-336 PF00731A 19.32 6.333e-24 268-297
804	BL00170	Cyclophilin-type peptidyl-prolyl cis-trans isomerase signatur.	BL00170B 20.97 8.071e-09 297-337
805	BL00678	Trp-Asp (WD) repeat proteins proteins.	BL00678 9.67 3.400e-10 378-389 BL00678 9.67 5.800e-10 418-429 BL00678 9.67 8.800e-10 295-306
806	PD01719	PRECURSOR GLYCOPROTEIN SIGNAL RE.	PD01719A 12.89 7.571e-14 290-318
807	PR00320	G-PROTEIN BETA WD-40 REPEAT SIGNATURE	PR00320B 12.19 9.100e-09 451-466
809	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 4.462e-12 564-595
810	PR00453	VON WILLEBRAND FACTOR TYPE A DOMAIN SIGNATURE	PR00453A 12.79 1.310e-14 36-54 PR00453B 14.65 8.568e-10 75-90
814	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 2.047e-31 16-55
815	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 2.047e-31 16-55
817	PR00193	MYOSIN HEAVY CHAIN SIGNATURE	PR00193D 14.36 5.154e-36 125-154 PR00193E 19.47 3.919e-18 179-208
818	PR00830	ENDOPEPTIDASE LA (LON) SERINE	PR00830A 8.41 9.571e-11 115-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		PROTEASE (S16) SIGNATURE	135
819	BL00126	3'5'-cyclic nucleotide phosphodiesterases proteins.	BL00126C 22.07 7.857e-24 528-569 BL00126B 35.22 3.714e-15 669-724 BL00126D 25.50 1.173e-14 584-623 BL00126B 15.20 1.000e-12 502-514 BL00126A 27.56 3.361e-09 461-498
820	PR00511	TEKTIN SIGNATURE	PR00511B 12.25 8.826e-22 174-195 PR00511A 13.59 7.723e-11 155-172
821	BL00741	Guanine-nucleotide dissociation stimulators CDC24 family sign.	BL00741B 14.27 2.800e-15 13-36
822	PF00780	Domain found in NIK1-like kinases, mouse citron and yeast ROM.	PF00780I 14.69 4.825e-09 231-261
827	BL00030	Eukaryotic RNA-binding region RNP-1 proteins.	BL00030A 14.39 5.235e-11 144-163
828	BL00326	Tropomyosins proteins.	BL00326D 8.76 9.357e-11 545-586
829	PD02448	TRANSCRIPTION PROTEIN DNA-BINDIN.	PD02448A 9.37 1.000e-40 46-85 PD02448B 10.17 1.000e-40 85-133 PD02448C 13.62 1.000e-40 152-189 PD02448E 11.33 9.000e-30 235-261 PD02448F 14.22 9.654e-25 279-303 PD02448D 11.48 3.659e-18 197-211 PD02448G 10.73 7.857e-16 305-318
830	BL00720	Guanine-nucleotide dissociation stimulators CDC25 family sign.	BL00720B 16.57 4.500e-23 483-507
831	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 6.625e-21 143-174 BL00107B 13.31 4.214e-10 213-229
832	BL00215	Mitochondrial energy transfer proteins.	BL00215A 15.82 5.787e-11 32-57
833	PR00497	NEUTROPHIL CYTOSOL FACTOR P40 SIGNATURE	PR00497A 6.92 4.375e-09 41-59
834	BL00229	Tau and MAP proteins tubulin-binding domain proteins.	BL00229A 23.57 9.565e-10 99-138
835	BL00421	Transmembrane 4 family proteins.	BL00421E 20.87 2.216e-09 1053-1083
836	BL00795	Involucrin proteins.	BL00795B 12.41 7.931e-09 405-445
837	PR00020	MAM DOMAIN SIGNATURE	PR00020A 18.17 1.000e-17 34-53 PR00020B 15.52 5.846e-16 68-85 PR00020D 12.70 2.543e-15 147-162 PR00020C 13.66 3.483e-13 95-107 PR00020E 8.64 6.586e-13 165-179
838	BL50017	Death domain proteins profile.	BL50017B 17.60 6.897e-13 1499-1515
839	PF00850	Histone deacetylase family.	PF00850C 14.55 9.542e-09 1352-1369
840	PF00023	Ank repeat proteins.	PF00023A 16.03 4.500e-12 44-60 PF00023B 14.20 7.923e-11 73-83 PF00023B 14.20 9.000e-10 139-149 PF00023B 14.20 5.500e-09 40-50
842	BL01194	Ribosomal protein L15e proteins.	BL01194B 13.66 1.000e-40 37-85 BL01194C 12.35 9.250e-40 103-138 BL01194A 18.70 7.632e-38

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			2-37 BL01194D 19.02 2.658e-36 139-178
843	BL00610	Sodium:neurotransmitter symporter family proteins.	BL00610A 17.73 1.000e-40 40-90 BL00610B 23.65 1.000e-40 104-154 BL00610C 12.94 1.000e-40 206-258 BL00610E 20.34 1.000e-40 355-398 BL00610F 29.02 1.000e-40 454-509 BL00610D 20.97 6.063e-35 272-325 BL00610G 12.89 8.588e-13 514-537
845	BL00143	Insulinase family, zinc-binding region proteins.	BL00143A 20.91 4.300e-20 94-121 BL00143C 14.16 5.500e-13 245-258 BL00143B 14.41 9.053e-10 141-156
846	PR00543	OESTROGEN RECEPTOR SIGNATURE	PR00543D 10.87 1.355e-09 898-914
847	PR00543	OESTROGEN RECEPTOR SIGNATURE	PR00543D 10.87 1.355e-09 898-914
848	BL00824	Elongation factor 1 beta/beta'/delta chain proteins.	BL00824C 14.58 1.000e-40 129-167 BL00824D 14.04 6.192e-39 167-202 BL00824B 9.21 2.080e-21 96-116 BL00824E 12.49 3.333e-19 210-226 BL00824A 13.78 8.650e-14 19-34
849	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 1.000e-40 12-51
850	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 7.316e-24 10-49
852	BL01272	Glucokinase regulatory protein family proteins.	BL01272B 19.61 6.870e-30 136-171 BL01272C 11.68 3.314e-25 249-274 BL01272A 6.49 1.231e-18 99-117
853	PD00930	PROTEIN GTPASE DOMAIN ACTIVATION	PD00930B 33.72 9.341e-20 65-105
854	PD00289	PROTEIN SH3 DOMAIN REPEAT PRESYN.	PD00289 9.97 6.850e-11 140-154
858	PR00450	RECOVERIN FAMILY SIGNATURE	PR00450C 12.22 3.250e-25 68-90 PR00450B 11.76 8.125e-23 22-42 PR00450D 16.58 8.920e-22 92-112 PR00450E 12.14 1.581e-19 114-133 PR00450G 15.33 5.500e-19 166-187 PR00450F 12.30 4.375e-15 140-156 PR00450A 13.58 1.857e-14 8-23
860	BL00027	'Homeobox' domain proteins.	BL00027 26.43 7.188e-27 74-117
866	BL00477	Alpha-2-macroglobulin family thiolester region proteins.	BL00477L 23.51 7.480e-20 54-87
867	BL01078	Molybdenum cofactor biosynthesis proteins.	BL01078B 14.20 1.621e-20 408-429 BL01078A 10.16 2.000e-13 366-379 BL01078D 5.99 3.455e-11 566-576 BL01078C 10.52 3.793e-11 501-513
868	BL01177	Anaphylatoxin domain proteins.	BL01177E 20.64 5.800e-24 462-489 BL01177C 17.39 5.333e-19 416-435 BL01177B 13.61 7.840e-16 122-138 BL01177D 17.50 1.900e-15 441-459
869	BL01177	Anaphylatoxin domain proteins.	BL01177E 20.64 5.800e-24 415-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			442 BL01177C 17.39 5.333e-19 369-388 BL01177B 13.61 7.840e-16 122-138 BL01177D 17.50 1.900e-15 394-412
871	BL50007	Phosphatidylinositol-specific phospholipase X-box domain proteins prof.	BL50007A 19.61 1.000e-40 322-368 BL50007D 19.54 1.000e-40 589-631 BL50007B 20.90 6.700e-36 383-421 BL50007E 25.63 9.053e-33 748-785 BL50007C 8.97 5.200e-19 452-469
872	BL00972	Ubiquitin carboxyl-terminal hydrolases family 2 proteins.	BL00972D 22.55 3.250e-17 90-115
874	PR00452	SH3 DOMAIN SIGNATURE	PR00452B 11.65 4.250e-09 370-386
877	BL00741	Guanine-nucleotide dissociation stimulators CDC24 family sign.	BL00741B 14.27 5.500e-13 1343-1366
878	DM00215	PROLINE-RICH PROTEIN 3.	DM00215 19.43 2.525e-09 52-85
881	PD02807	APOLIPOPROTEIN E PRECURSOR APO-E GLYCOPROTEIN PLAS.	PD02807E 10.90 4.702e-09 358-407
882	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 7.188e-37 8-47
885	PF00023	Ank repeat proteins.	PF00023A 16.03 8.071e-09 10-26
886	PR00372	BIOPTERIN-DEPENDENT AROMATIC AMINO ACID HYDROXYLASE SIGNATURE	PR00372B 10.30 9.308e-27 225-248 PR00372A 13.39 7.000e-24 134-154 PR00372E 12.62 2.125e-23 360-380 PR00372C 7.90 3.025e-22 289-309 PR00372F 13.09 6.333e-21 395-414 PR00372D 10.22 1.000e-19 329-348
887	BL00301	GTP-binding elongation factors proteins.	BL00301B 20.09 2.800e-24 103-135 BL00301A 12.41 4.316e-13 21-33
888	BL00518	Zinc finger, C3HC4 type (RING finger), proteins.	BL00518 12.23 1.667e-09 30-39
889	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 4.906e-26 6-45
890	DM00179	w KINASE ALPHA ADHESION T-CELL.	DM00179 13.97 7.652e-09 113-123
892	BL01022	PTR2 family proton/oligopeptide symporters proteins.	BL01022B 22.19 6.016e-14 72-118 BL01022E 23.51 1.173e-12 472-508 BL01022A 11.58 9.135e-12 42-61 BL01022D 9.42 3.455e-11 199-212
893	PD02407	3-BISPHOSPHOGLYCERATE-INDEPENDENT PHOSPHOGLYCER.	PD02407K 12.59 6.529e-10 360-383
894	PD02407	3-BISPHOSPHOGLYCERATE-INDEPENDENT PHOSPHOGLYCER.	PD02407K 12.59 6.529e-10 360-383
895	PR00237	RHODOPSIN-LIKE GPCR SUPERFAMILY SIGNATURE	PR00237B 13.50 9.100e-14 116-138 PR00237F 13.57 1.360e-13 312-337 PR00237G 19.63 9.069e-13 353-380 PR00237E 13.03 7.120e-12 243-267 PR00237D 8.94 4.150e-11 194-216 PR00237A 11.48 4.375e-11 83-108
896	BL00129	Glycosyl hydrolases family 31 proteins.	BL00129D 16.76 8.258e-26 634-678 BL00129A 26.21 1.720e-25 384-430 BL00129E 22.60 4.857e-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			23 698-734 BL00129C 15.12 1.750e-22 596-624 BL00129B 19.19 5.891e-18 495-522 BL00129F 26.19 7.545e-15 814-852
897	BL00598	Chromo domain proteins.	BL00598 14.45 1.220e-13 9-31
898	BL00518	Zinc finger, C3HC4 type (RING finger), proteins.	BL00518 12.23 6.000e-09 396-405
899	PD01101	INHIBITOR HEAVY CHAIN CHANNEL IN.	PD01101B 21.53 1.000e-40 274-327 PD01101D 24.45 1.000e-40 457-512 PD01101A 18.25 6.268e-23 83-117 PD01101C 12.69 1.237e-16 366-386 PD01101E 6.73 7.750e-12 566-576
900	PR00600	PROTEIN PHOSPHATASE PP2A 55KD REGULATORY SUBUNIT SIGNATURE	PR00600A 11.61 5.979e-09 31-52
901	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 8.116e-31 24-63
903	BL01115	GTP-binding nuclear protein ran proteins.	BL01115A 10.22 1.509e-11 21-65
906	DM00215	PROLINE-RICH PROTEIN 3.	DM00215 19.43 2.174e-13 539-572 DM00215 19.43 4.750e-12 549-582 DM00215 19.43 9.824e-11 551-584 DM00215 19.43 2.929e-10 548-581 DM00215 19.43 4.054e-10 550-583 DM00215 19.43 5.339e-10 552-585 DM00215 19.43 7.107e-10 544-577
907	PR00988	URIDINE KINASE SIGNATURE	PR00988A 6.39 6.276e-12 314-332
908	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 5.950e-17 1125-1156
909	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 5.950e-17 1118-1140
910	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 8.560e-13 150-181
911	BL00107	Protein kinases ATP-binding region proteins.	BL00107A 18.39 8.560e-13 150-181
912	PF00856	SET domain proteins.	PF00856A 26.14 4.553e-11 243-280
913	PF00628	PHD-finger.	PF00628 15.84 6.400e-13 197-212
914	PR00962	LETHAL(2) GIANT LARVAE PROTEIN SIGNATURE	PR00962D 10.40 1.000e-27 435-459 PR00962G 15.71 4.086e-26 593-618 PR00962B 11.98 9.122e-26 296-319 PR00962A 13.28 6.143e-22 15-34 PR00962C 8.00 4.000e-21 348-369 PR00962F 12.39 9.769e-21 552-572 PR00962H 13.32 2.636e-20 623-643 PR00962I 11.68 9.786e-20 692-712 PR00962E 8.81 2.915e-18 515-534
915	PR00962	LETHAL(2) GIANT LARVAE PROTEIN SIGNATURE	PR00962D 10.40 1.000e-27 365-389 PR00962G 15.71 4.086e-26 523-548 PR00962A 13.28 6.143e-22 15-34 PR00962C 8.00 4.000e-21 278-299 PR00962F 12.39 9.769e-21 482-502 PR00962H

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			13.32 2.636e-20 553-573 PR00962I 11.68 9.786e-20 622-642 PR00962E 8.81 2.915e-18 445-464
916	BL00134	Serine proteases, trypsin family, histidine proteins.	BL00134A 11.96 5.886e-14 90-107
917	BL00478	LIM domain proteins.	BL00478B 14.79 8.393e-13 211-226 BL00478B 14.79 6.712e-10 271-286
918	PR00049	WILM'S TUMOUR PROTEIN SIGNATURE	PR00049D 0.00 5.729e-09 973-988
922	BL00150	Acylphosphatase proteins.	BL00150 25.33 1.000e-40 37-84
924	DM00031	IMMUNOGLOBULIN V REGION.	DM00031B 15.41 8.063e-09 79-113
925	BL00072	Acyl-CoA dehydrogenases proteins.	BL00072D 30.08 2.837e-24 280-331 BL00072E 24.12 8.200e-24 368-411 BL00072C 25.30 7.873e-20 226-267 BL00072B 9.48 6.049e-12 183-196
927	BL00237	G-protein coupled receptors proteins.	BL00237C 13.19 1.692e-13 229-256 BL00237A 27.68 6.657e-13 90-130 BL00237D 11.23 9.571e-13 290-307
928	BL01033	Globins profile.	BL01033A 16.94 7.923e-18 25-47 BL01033B 13.81 1.000e-15 93-105
929	BL00216	Sugar transport proteins.	BL00216B 27.64 8.714e-13 203-253
932	BL00415	Synapsins proteins.	BL00415N 4.29 9.519e-10 353-397 BL00415N 4.29 2.117e-09 63-107 BL00415N 4.29 3.628e-09 57-101 BL00415N 4.29 5.664e-09 347-391
933	PD02448	TRANSCRIPTION PROTEIN DNA-BINDIN.	PD02448A 9.37 1.000e-40 46-85 PD02448E 12.17 1.000e-40 25-153 PD02448C 13.62 1.000e-40 152-189 PD02448E 11.33 9.000e-30 223-249 PD02448F 14.22 9.654e-25 267-291 PD02448D 11.48 3.659e-18 197-211 PD02448G 10.73 7.857e-16 293-306
934	DM00191	w SPAC8A4.04C RESISTANCE SPAC8A4.05C DAUNORUBICIN.	DM00191D 13.94 9.083e-10 136-175
935	BL01115	GTP-binding nuclear protein ran proteins.	BL01115A 10.22 4.696e-10 67-111
936	BL00019	Actinin-type actin-binding domain proteins.	BL00019D 15.33 8.138e-14 865-895
937	PR00762	CHLORIDE CHANNEL SIGNATURE	PR00762A 14.22 4.000e-22 183-201 PR00762C 9.29 1.000e-21 268-288 PR00762E 12.07 3.250e-20 520-537 PR00762D 11.29 1.000e-19 470-491 PR00762F 15.12 1.429e-19 538-558 PR00762B 12.12 1.818e-18 214-234 PR00762G 14.13 3.455e-17 577-592
938	BL00027	'Homeobox' domain proteins.	BL00027 26.43 9.500e-25 291-334
939	DM01111	4 kw PHOSPHATASE	DM01111E 17.28 1.568e-10 248-

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
		TRANSFORMING 61K PDF1.	297 DM01111E 17.28 5.168e-10 659-708 DM01111D 16.76 5.263e-09 279-325 DM01111M 10.67 8.674e-09 911-935
940	BL00107	Protein kinases ATP-binding region proteins.	BL00107B 13.31 1.000e-14 293-309 BL00107A 18.39 6.760e-13 229-260
942	BL01160	Kinesin light chain repeat proteins.	BL01160B 19.54 9.832e-11 543-597
943	PD01066	PROTEIN ZINC FINGER ZINC-FINGER METAL-BINDING NU.	PD01066 19.43 3.500e-35 8-47
945	BL00989	Clathrin adaptor complexes small chain proteins.	BL00989B 26.51 1.000e-40 66-117 BL00989A 11.66 1.000e-13 5-19
946	PR00178	FATTY ACID-BINDING PROTEIN SIGNATURE	PR00178D 13.52 9.571e-09 450-469
947	BL00178	Aminoacyl-transfer RNA synthetases class-I proteins.	BL00178B 7.11 4.857e-09 713-724
948	PF00628	PHD-finger.	PF00628 15.84 8.412e-14 201-216
951	BL00216	Sugar transport proteins.	BL00216B 27.64 2.050e-10 180-230
952	PR00926	MITOCHONDRIAL CARRIER PROTEIN SIGNATURE	PR00926F 17.75 4.300e-11 26-49 PR00926F 17.75 6.348e-09 134-157
955	PF00109	Beta-ketoacyl synthase.	PF00109 13.08 2.846e-12 342-357
957	PR00069	ALDO-KETO REDUCTASE SIGNATURE	PR00069A 16.01 8.826e-24 26-51 PR00069B 11.33 1.514e-17 86-105 PR00069C 16.03 8.816e-14 155-173
958	PF00583	Acetyltransferase (GNAT) family.	PF00583A 12.53 5.500e-10 631-642
961	PR00328	GTP-BINDING SAR1 PROTEIN SIGNATURE	PR00328A 10.62 8.740e-10 7-31
962	BL00354	HMG-I and HMG-Y DNA-binding domain proteins (A+T-hoc).	BL00354A 3.83 9.438e-10 1489-1499
963	BL00354	HMG-I and HMG-Y DNA-binding domain proteins (A+T-hoc).	BL00354A 3.83 9.438e-10 1489-1499
964	BL00027	'Homeobox' domain proteins.	BL00027 26.43 7.188e-27 53-96
965	PF00992	Troponin.	PF00992A 16.67 2.421e-09 581-616
966	PR00515	5-HYDROXYTRYPTAMINE 1F RECEPTOR SIGNATURE	PR00515D 7.91 5.741e-09 13-33
967	BL00579	Ribosomal protein L29 proteins.	BL00579B 21.99 5.065e-21 164-194
970	BL00504	Fumarate reductase / succinate dehydrogenase FAD-binding site proteins.	BL00504C 18.68 2.227e-24 34-59 BL00504D 10.43 7.261e-21 75-93
973	PF00580	UvrD/REP helicase.	PF00580A 13.37 4.720e-09 249-271
974	PR00456	RIBOSOMAL PROTEIN P2 SIGNATURE	PR00456F 5.86 1.000e-10 242-254
975	BL00237	G-protein coupled receptors proteins.	BL00237A 27.68 4.429e-22 99-139
976	BL00031	Nuclear hormones receptors DNA-binding region proteins.	BL00031A 19.55 7.158e-33 60-93 BL00031B 22.25 5.500e-28 94-126
977	PD00066	PROTEIN ZINC-FINGER METAL-BINDI.	PD00066 13.92 8.200e-16 196-209 PD00066 13.92 8.200e-16 336-349 PD00066 13.92 2.385e-15 476-489

SEQ ID NO:	ACCESSION NO.	DESCRIPTION	RESULTS*
			PD00066 13.92 9.308e-15 252-265 PD00066 13.92 2.800e-14 448-461 PD00066 13.92 4.600e-14 392-405 PD00066 13.92 5.200e-14 280-293 PD00066 13.92 4.000e-13 224-237 PD00066 13.92 4.429e-12 308-321 PD00066 13.92 9.571e-12 420-433 PD00066 13.92 6.870e-11 168-181
978	BL00721	Formate--tetrahydrofolate ligase proteins.	BL00721B 13.21 1.000e-40 346-401 BL00721D 13.90 1.000e-40 538-592 BL00721E 13.46 1.000e-40 597-646 BL00721I 18.79 2.500e-40 814-860 BL00721H 21.20 8.239e-39 763-814 BL00721A 15.31 9.719e-32 287-321 BL00721C 16.92 4.000e-30 498-535 BL00721F 15.96 8.232e-27 660-702 BL00721G 7.97 3.017e-10 721-734
981	PD00126	PROTEIN REPEAT DOMAIN TPR NUCLEA.	PD00126A 22.53 2.552e-09 180-201
982	BL00869	Renal dipeptidase proteins.	BL00869C 12.58 3.172e-19 59-95 BL00869E 13.12 9.129e-18 120-157 BL00869J 15.60 6.032e-17 270-310 BL00869H 11.08 1.840e-16 219-242 BL00869G 13.55 2.543e-16 192-214 BL00869F 12.77 7.031e-14 157-192 BL00869I 12.92 3.274e-12 242-270 BL00869D 14.02 5.282e-10 95-124 BL00869B 15.55 9.382e-10 31-61
983	PR00196	ANNEXIN FAMILY SIGNATURE	PR00196F 13.89 2.125e-09 92-108
984	BL00485	Adenosine and AMP deaminase proteins.	BL00485D 30.82 2.427e-10 154-209

* Results include in order: accession number subtype; raw score; p-value; position of signature in amino acid sequence

5

TABLE 4

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
2	ig	Immunoglobulin domain	3.9e-17	60.3
3	HSP90	Hsp90 protein	0	1548.4
6	tsp_1	Thrombospondin type 1 domain	0.002	22.1
7	7tm_1	7 transmembrane receptor (rhodopsin family)	6.7e-08	27.3
9	PWWP	PWWP domain	8.1e-16	66.0
12	C1q	C1q domain	1.7e-26	101.5
13	C1q	C1q domain	2e-20	81.3
14	Aa_trans	Transmembrane amino acid transporter protein	2.7e-42	153.9
15	E1-E2_ATPase	E1-E2 ATPase	6.3e-124	412.2
16	trypsin	Trypsin	1.2e-87	278.6
17	ig	Immunoglobulin domain	7.6e-12	43.2
18	lectin_c	Lectin C-type domain	0.0003	21.2
20	Alpha_L_fucos	Alpha-L-fucosidase	1.2e-217	736.5

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
22	pkinase	Eukaryotic protein kinase domain	3.3e-87	303.1
23	pkinase	Eukaryotic protein kinase domain	2.7e-85	296.8
24	pkinase	Eukaryotic protein kinase domain	2.7e-85	296.8
25	ank	Ank repeat	5.5e-14	59.9
27	pkinase	Eukaryotic protein kinase domain	1.5e-100	347.4
28	spectrin	Spectrin repeat	4e-57	203.2
29	spectrin	Spectrin repeat	4e-57	203.2
30	WD40	WD domain, G-beta repeat	1.2e-07	38.8
33	rrm	RNA recognition motif.	1.1e-17	72.2
34	rrm	RNA recognition motif.	1.1e-17	72.2
36	7tm_1	7 transmembrane receptor (rhodopsin family)	3e-36	117.3
37	ank	Ank repeat	5.9e-25	96.3
38	SRF-TF	SRF-type transcription factor	1.4e-36	133.9
40	alk_phosphatase	Alkaline phosphatase	0	1034.9
44	zf-C2H2	Zinc finger, C2H2 type	8.6e-103	354.9
45	sugar_tr	Sugar (and other) transporter	3.1e-08	40.3
47	7tm_2	7 transmembrane receptor (Secretin family)	6.4e-79	275.6
50	zf-C2H2	Zinc finger, C2H2 type	1.3e-98	341.0
51	filament	Intermediate filament proteins	1.2e-176	600.3
52	zf-C3HC4	Zinc finger, C3HC4 type (RING finger)	2.7e-10	37.7
53	Cadherin_C_term	Cadherin cytoplasmic region	1.9e-94	327.2
54	S_100	S-100/ICaBP type calcium binding domain	5.2e-18	73.3
58	inositol_P	Inositol monophosphatase family	5e-13	49.8
59	7tm_1	7 transmembrane receptor (rhodopsin family)	8.8e-46	147.6
60	Kunitz_BPTI	Kunitz/Bovine pancreatic trypsin inhibitor	3.7e-47	148.6
62	DAD	DAD family	2.5e-74	260.3
63	MOZ_SAS	MOZ/SAS family	5.9e-133	455.1
64	MOZ_SAS	MOZ/SAS family	1.7e-123	425.6
65	ras	Ras family	9.5e-27	308.5
67	Ham1p_like	Ham1 family	3.7e-49	176.7
68	7tm_1	7 transmembrane receptor (rhodopsin family)	5.2e-39	126.1
70	zf-C2H2	Zinc finger, C2H2 type	1.5e-112	387.3
71	Peptidase_M41	Peptidase family M41	1.2e-110	381.0
72	abhydrolase	alpha/beta hydrolase fold	9.8e-05	26.5
81	K_tetra	K+ channel tetramerisation domain	0.022	-16.8
82	pkinase	Eukaryotic protein kinase domain	5e-49	176.3
84	AAA	ATPases associated with various cellular act	1.3e-77	271.3
85	homeobox	Homeobox domain	1.4e-28	108.3
87	TGF-beta	Transforming growth factor beta like	6.7e-68	210.2
91	mito_carr	Mitochondrial carrier proteins	4.6e-57	198.5
95	adenylatekinase	Adenylate kinase	1.1e-15	60.0
96	ig	Immunoglobulin domain	4.1e-20	69.8
99	CNH	CNH domain	3.4e-120	412.7
100	homeobox	Homeobox domain	7.4e-32	119.3
101	zf-C2H2	Zinc finger, C2H2 type	2.2e-47	170.8
102	zf-C2H2	Zinc finger, C2H2 type	4.4e-89	309.4
103	dynamin	Dynamin family	1.4e-150	513.6
104	lectin_c	Lectin C-type domain	4.2e-15	63.6
105	lectin_c	Lectin C-type domain	4.2e-15	63.6
108	metalthio	Metallothionein	2e-25	97.9

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
112	HSP20	Hsp20/alpha crystallin family	2.6e-20	77.7
115	EF_TS	Elongation factor TS	3.8e-63	221.1
116	sugar_tr	Sugar (and other) transporter	4e-63	223.1
118	catalase	Catalase	0	1158.9
119	UCH	Ubiquitin carboxyl-terminal hydrolase, famil	1e-10	24.4
122	metalthio	Metallothionein	2.8e-25	97.4
125	adh_short	short chain dehydrogenase	1.6e-45	164.6
126	KRAB	KRAB box	7.9e-25	95.9
127	G-alpha	G-protein alpha subunit	1e-249	843.0
128	mito_carr	Mitochondrial carrier proteins	2e-65	227.2
131	EF1BD	EF-1 guanine nucleotide exchange domain	4.9e-53	189.6
132	GYF	GYF domain	4.9e-28	106.6
133	GYF	GYF domain	4.9e-28	106.6
134	lipocalin	Lipocalin / cytosolic fatty-acid binding pr	2.1e-33	119.1
135	pkinase	Eukaryotic protein kinase domain	3.3e-86	299.8
136	ank	Ank repeat	2.2e-29	111.1
137	IL8	Small cytokines (intecrine/chemokine), inter	3.1e-18	65.2
139	pyridoxal_deC	Pyridoxal-dependent decarboxylase conse	0.00011	19.0
140	cadherin	Cadherin domain	1.3e-88	307.8
142	efhand	EF hand	5.7e-33	123.0
143	Acyltransferase	Acyltransferase	2e-29	111.2
146	cytochrome_c	Cytochrome c	1.7e-33	124.7
147	pkinase	Eukaryotic protein kinase domain	2.3e-86	300.3
148	PDZ	PDZ domain (Also known as DHR or GLGF).	1.7e-09	45.0
149	aldo_ket_red	Aldo/keto reductase family	7.4e-189	640.8
150	homeobox	Homeobox domain	3.2e-08	38.7
151	PseudoU_synth_1	tRNA pseudouridine synthase	4.7e-57	203.0
152	abhdrolase	alpha/beta hydrolase fold	1.7e-31	118.0
153	PDZ	PDZ domain (Also known as DHR or GLGF).	1.1e-09	45.0
156	PHD	PHD-finger	7.6e-15	62.8
157	fn3	Fibronectin type III domain	0.015	21.9
158	homeobox	Homeobox domain	2.7e-27	104.1
160	PWI	PWI domain	3.9e-24	93.6
162	DnaJ	DnaJ domain	2e-06	34.8
164	Cbl_N	CBL proto-oncogene N-terminal domain	8e-117	401.5
166	metalthio	Metallothionein	3.1e-26	100.6
167	LRR	Leucine Rich Repeat	0.00069	26.3
169	fibrinogen_C	Fibrinogen beta and gamma chains, C-term	5.3e-180	611.4
170	fibrinogen_C	Fibrinogen beta and gamma chains, C-term	5.3e-180	611.4
171	fibrinogen_C	Fibrinogen beta and gamma chains, C-term	1e-149	510.8
173	homeobox	Homeobox domain	1.5e-29	111.6
174	FYVE	FYVE zinc finger	7.4e-28	103.8
175	GRIP	GRIP domain	3.9e-08	40.5
182	pkinase	Eukaryotic protein kinase domain	3.4e-71	250.0
185	CAP_GLY	CAP-Gly domain	5.6e-51	182.8
186	TBC	TBC domain	2.2e-50	180.8
187	TBC	TBC domain	2.2e-50	180.8

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
188	PDZ	PDZ domain (Also known as DHR or GLGF).	4e-13	57.0
189	Kelch	Kelch motif	5.2e-106	365.6
190	Tropomyosin	Tropomyosins	3.8e-171	535.4
192	Rieske	Rieske [2Fe-2S] domain	0.0016	18.5
199	ig	Immunoglobulin domain	5.9e-19	66.1
202	EGF	EGF-like domain	3.4e-54	193.5
203	trefoil	Trefoil (P-type) domain	1e-24	95.5
204	TBC	TBC domain	8.5e-38	139.0
205	efhand	EF hand	0.0096	22.6
206	ISK_Channel	Slow voltage-gated potassium channel	0.0031	8.1
207	trefoil	Trefoil (P-type) domain	2.9e-48	173.7
209	Ribosomal_S13	Ribosomal protein S13/S18	1.2e-78	274.7
210	hemopexin	Hemopexin	1.3e-62	221.5
213	TBC	TBC domain	2.5e-48	174.0
215	Basic	Myogenic Basic domain	4.3e-50	179.8
216	Ribosomal_L24	KOW motif	8.2e-23	89.2
222	fn3	Fibronectin type III domain	7.3e-141	481.4
223	cofilin_ADF	Cofilin/tropomyosin-type actin-binding pr	9.3e-47	168.8
224	efhand	EF hand	6.1e-06	33.2
225	Pterin_4a	Pterin 4 alpha carbinolamine dehydratase	9.3e-42	152.1
228	ABC_tran	ABC transporter	4.1e-110	379.2
234	E1_DerP2_DerF2	E1 family	3.7e-90	312.9
235	E1_DerP2_DerF2	E1 family	1.6e-48	174.6
237	PMP22_Claudin	PMP-22/EMP/MP20/Claudin family	1.7e-25	98.1
238	Opioids_neurope p	Vertebrate endogenous opioids neurope	1.8e-159	543.2
239	eIF-5a	Eukaryotic initiation factor 5A hypusine	5.9e-104	358.8
240	Amino oxidase	Flavin containing amine oxidase	2.5e-11	37.8
243	zf-C2H2	Zinc finger, C2H2 type	2.1e-99	343.6
244	Band_7	SPFH domain / Band 7 family	2.3e-53	190.7
245	ank	Ank repeat	1.6e-88	307.5
246	zf-C2H2	Zinc finger, C2H2 type	6.7e-49	175.9
247	actin	Actin	2.3e-42	140.3
248	ER_lumen_recept t	ER lumen protein retaining receptor	2.4e-155	529.5
250	PMP22_Claudin	PMP-22/EMP/MP20/Claudin family	2.2e-38	140.9
252	Collagen	Collagen triple helix repeat (20 copies)	1.4e-13	58.6
255	C2	C2 domain	0.052	7.8
257	CAP_GLY	CAP-Gly domain	1.4e-20	81.8
260	WD40	WD domain, G-beta repeat	9.9e-62	218.5
261	WD40	WD domain, G-beta repeat	9.9e-62	218.5
262	WD40	WD domain, G-beta repeat	9.9e-62	218.5
263	cofilin_ADF	Cofilin/tropomyosin-type actin-binding pr	7.8e-21	82.6
264	Ribosomal_L14	Ribosomal protein L14p/L23e	9.2e-10	40.6
265	SAPA	Saposin A-type domain	4.4e-27	103.4
266	SAPA	Saposin A-type domain	4.4e-27	103.4
267	ABC_tran	ABC transporter	9.5e-39	142.2
269	Ribosomal_L14	Ribosomal protein L14p/L23e	6.2e-62	219.2
270	abhydrolase	alpha/beta hydrolase fold	0.042	-3.3
272	ras	Ras family	4.3e-87	302.8

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
273	rrm	RNA recognition motif.	0.074	14.6
275	lipocalin	Lipocalin / cytosolic fatty-acid binding pr	2.5e-41	146.4
276	ras	Ras family	1.1e-67	238.3
277	UCH	Ubiquitin carboxyl-terminal hydrolase, famil	1.2e-147	503.9
278	START	START domain	3.2e-09	44.1
279	WD40	WD domain, G-beta repeat	1.8e-27	104.7
282	G-patch	G-patch domain	7.8e-22	86.0
287	Anti_proliferat	BTG1 family	1.2e-101	351.0
289	KRAB	KRAB box	7.1e-21	82.8
293	7tm_3	7 transmembrane receptor	3.3e-73	256.6
295	SET	SET domain	5e-30	113.2
296	Pyridox_oxidase	Pyridoxamine 5'-phosphate oxidase	1.3e-76	268.0
297	rrm	RNA recognition motif.	5.4e-45	162.9
298	Ubie_methyltran	ubiE/COQ5 methyltransferase family	6.3e-05	-96.3
299	Ubie_methyltran	ubiE/COQ5 methyltransferase family	0.0024	-118.1
301	Cyt_reductase	FAD/NAD-binding Cytochrome reductase	7.7e-61	215.5
302	G-patch	G-patch domain	3.1e-14	60.7
307	7tm_1	7 transmembrane receptor (rhodopsin family)	7.7e-43	138.2
308	PH	PH domain	0.0015	17.8
310	7tm_1	7 transmembrane receptor (rhodopsin family)	1.4e-84	270.8
311	Rhodanese	Rhodanese-like domain	3.3e-64	226.7
312	tubulin	Tubulin/FtsZ family	4.9e-286	963.6
314	SURF4	SURF4 family	1.2e-199	676.6
325	IMS	impB/mucB/samB family	2e-58	207.5
327	cadherin	Cadherin domain	4.3e-91	316.0
329	NAC	NAC domain	2.1e-28	107.8
330	IP_trans	Phosphatidylinositol transfer protein	6.5e-98	338.7
332	TFIIS	Transcription factor S-II (TFIIS)	8.8e-05	29.3
337	zf-C2H2	Zinc finger, C2H2 type	3.6e-61	216.6
340	AIRS	AIR synthase related protein	4e-32	120.2
341	annexin	Annexin	4.6e-80	152.4
345	Stathmin	Stathmin family	1.8e-90	514.0
347	Ribosomal_L16	Ribosomal protein L16	4.6e-09	34.5
348	lactamase_B	Metallo-beta-lactamase superfamily	0.012	-6.0
351	efhand	EF hand	2.5e-14	61.0
353	lectin_c	Lectin C-type domain	1.3e-05	32.1
354	WD40	WD domain, G-beta repeat	2.2e-18	74.5
360	lipocalin	Lipocalin / cytosolic fatty-acid binding pr	6.3e-10	38.3
362	Acetyltransf	Acetyltransferase (GNAT) family	0.0019	24.9
365	tRNA-synt_1	tRNA synthetases class I (I, L, M and V)	4.6e-185	628.2
366	Sulfatase	Sulfatase	6.1e-228	770.6
368	START	START domain	3.8e-11	50.5
369	pkinase	Eukaryotic protein kinase domain	2.4e-10	41.3
370	ACBP	Acyl CoA binding protein	4.4e-56	199.7
371	pkinase	Eukaryotic protein kinase domain	1.6e-94	327.5
373	EGF	EGF-like domain	2.6e-12	54.3
375	zf-C2H2	Zinc finger, C2H2 type	8.2e-64	225.4
377	KRAB	KRAB box	3.7e-27	103.7
379	SET	SET domain	7.3e-61	215.6
380	Glyco_transf_8	Glycosyl transferase family 8	0.0028	-40.1
381	zf-C2H2	Zinc finger, C2H2 type	4.3e-06	33.7
383	Glyco_transf_8	Glycosyl transferase family 8	0.0028	-40.1

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
384	RasGEF	RasGEF domain	8.1e-43	155.7
385	TBC	TBC domain	0.017	-66.6
389	Glycosyl_transf_2	Glycosyl transferases	1.3e-15	65.3
390	Na_Ca_Ex	Sodium/calcium exchanger protein	3.9e-105	362.7
391	fn3	Fibronectin type III domain	4.1e-102	352.6
392	fn3	Fibronectin type III domain	3.4e-45	163.6
393	fn3	Fibronectin type III domain	3.4e-45	163.6
394	ldl_recept_b	Low-density lipoprotein receptor repeat	7.1e-49	175.8
395	Ribosomal_L30	Ribosomal protein L30p/L7e	0.0023	16.0
396	Oxysterol_BP	Oxysterol-binding protein	1.5e-94	327.5
397	RDS_ROM1	Peripherin/rom-1	2.9e-33	123.9
399	lactamase_B	Metallo-beta-lactamase superfamily	3.4e-39	143.6
402	F-box	F-box domain.	0.0002	28.1
403	CLP_protease	Clp protease	4.8e-64	226.2
405	Ribosomal_L35_Ae	Ribosomal protein L35Ae	6e-77	269.0
406	LIM	LIM domain containing proteins	0.00021	20.7
410	tRNA-synt_1c	tRNA synthetases class I (E and Q)	1e-236	799.8
411	NTP_transf_2	Nucleotidyltransferase domain	3.9e-16	67.0
412	DEAD	DEAD/DEAH box helicase	0.00016	17.2
414	DUF94	Domain of unknown function DUF94	0.00011	26.9
415	tubulin	Tubulin/FtsZ family	4.5e-289	973.7
420	SET	SET domain	3.3e-57	203.5
421	WD40	WD domain, G-beta repeat	6.1e-29	109.6
423	zf-C2H2	Zinc finger, C2H2 type	1.5e-39	144.9
424	pkinese	Eukaryotic protein kinase domain	8.9e-75	261.8
428	LIM	LIM domain containing proteins	1.8e-34	126.7
431	kazal	Kazal-type serine protease inhibitor domain	3.7e-18	73.8
432	SH2	Src homology domain 2	1.4e-67	198.4
433	zf-C2H2	Zinc finger, C2H2 type	2.8e-144	492.7
434	ras	Ras family	0.012	-106.8
436	E1-E2_ATPase	E1-E2 ATPase	1.6e-117	391.0
437	RNA_pol_A	RNA polymerase alpha subunit	0	1077.7
438	PHD	PHD finger	1.6e-11	51.7
439	lectin_c	Lectin C-type domain	4.7e-30	113.3
440	zf-C2H2	Zinc finger, C2H2 type	1.1e-65	231.6
441	arrestin	Arrestin (or S-antigen)	2.9e-254	858.1
442	aminotran_3	Aminotransferases class-III pyridoxal-pho	8.2e-80	231.1
443	UCH-1	Ubiquitin carboxyl-terminal hydrolases famil	8.5e-12	52.6
444	CTF_NFI	CTF/NF-I family	2.6e-277	934.6
451	T-box	T-box	3.8e-117	402.6
453	Rieske	Rieske [2Fe-2S] domain	2.6e-13	57.7
454	zf-C2H2	Zinc finger, C2H2 type	3.9e-64	226.5
456	homeobox	Homeobox domain	2.8e-08	38.9
459	ig	Immunoglobulin domain	2.6e-20	70.5
460	Hydrolase	haloacid dehalogenase-like hydrolase	4e-25	96.9
462	rve	Integrase core domain	1.6e-13	50.7
466	CH	Calponin homology (CH) domain	2.4e-17	71.1
467	CH	Calponin homology (CH) domain	2.4e-17	71.1
468	Sterol_desat	Sterol desaturase	7.5e-38	139.2
469	pro_isomerase	Cyclophilin type peptidyl-prolyl cis-tr	2.6e-63	220.9
470	Peptidase_M24	metallopeptidase family M24	6e-08	28.1
471	PDZ	PDZ domain (Also known as DHR or GLGF).	5.4e-129	441.9

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
472	myb_DNA-binding	Myb-like DNA-binding domain	3.6e-06	33.9
473	ZZ	Zinc finger present in dystrophin, CB	0.012	20.0
474	EF1G_domain	Elongation factor 1 gamma, conserved doma	6.3e-88	305.5
475	Ribosomal_L31e	Ribosomal protein L31e	6.1e-66	232.5
476	Clq	Clq domain	2.5e-75	263.7
477	SH3	SH3 domain	1.1e-12	55.6
478	MoaA_NifB_Pq qE	moaA / nifB / pqqE family	0.002	-17.7
479	FYVE	FYVE zinc finger	9.3e-21	78.6
480	DNA_pol_A	DNA polymerase family A	2.3e-46	167.4
482	adh_short	short chain dehydrogenase	1.2e-62	221.6
483	ank	Ank repeat	1.3e-17	71.9
484	IMS	impB/mucB/samB family	2.2e-83	290.5
486	TIR	TIR domain	3.2e-19	67.8
487	FMO-like	Flavin-binding monooxygenase-like	0	1425.5
488	I_LWEQ	I/LWEQ domain	9.5e-101	341.0
495	homeobox	Homeobox domain	3.6e-06	30.8
497	pkinese	Eukaryotic protein kinase domain	2.3e-166	566.1
499	fn3	Fibronectin type III domain	2.5e-237	801.8
501	LRR	Leucine Rich Repeat	9.3e-31	115.6
502	RGS	Regulator of G protein signaling domain	0.041	11.9
503	filament	Intermediate filament proteins	1e-142	487.5
505	fn3	Fibronectin type III domain	1.3e-100	347.7
506	HECT	HECT-domain (ubiquitin-transferase).	1e-13	59.0
507	Ribosomal_L7A e	Ribosomal protein L7Ae	5.7e-26	99.7
508	WD40	WD domain, G-beta repeat	0.063	19.8
509	WD40	WD domain, G-beta repeat	0.063	19.8
510	WD40	WD domain, G-beta repeat	2.1e-42	154.3
511	pkinese	Eukaryotic protein kinase domain	2.3e-86	300.4
512	G-gamma	GGL domain	1.9e-08	34.3
513	SH3	SH3 domain	1e-06	54.2
515	HTH_AraC	Bacterial regulatory helix-turn-helix protei	3.9e-27	103.6
516	zf-C2H2	Zinc finger, C2H2 type	1.7e-34	128.0
517	S1	S1 RNA binding domain	6.1e-58	205.9
518	pkinese	Eukaryotic protein kinase domain	1.8e-75	264.2
525	cadherin	Cadherin domain	2e-80	280.6
528	zf-C2H2	Zinc finger, C2H2 type	4e-70	246.4
529	neur_chan	Neurotransmitter-gated ion-channel	5.8e-222	750.8
531	RhoGEF	RhoGEF domain	3.5e-44	160.2
532	myosin_head	Myosin head (motor domain)	0	1494.5
533	LRR	Leucine Rich Repeat	8.3e-15	62.6
535	Sec7	Sec7 domain	5.1e-92	319.1
536	homeobox	Homeobox domain	4.8e-05	26.4
539	actin	Actin	2.4e-100	330.6
542	ank	Ank repeat	1.9e-35	131.2
544	zf-CCCH	Zinc finger C-x8-C-x5-C-x3-H type	2.8e-10	41.7
546	DSPc	Dual specificity phosphatase, catalytic doma	2.4e-40	147.4
547	HMG_CoA_synt	Hydroxymethylglutaryl-coenzyme A synthas	0	1250.8
549	laminin_G	Laminin G domain	3.3e-76	266.6
551	PHD	PHD-finger	0.008	9.3
552	PDZ	PDZ domain (Also known as DHR or	0.0017	25.0

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
		GLGF).		
555	WW	WW domain	1.3e-24	95.3
558	kinesin	Kinesin motor domain	1.8e-176	599.7
559	zf-C3HC4	Zinc finger, C3HC4 type (RING finger)	0.00085	16.5
563	efhand	EF hand	7.9e-11	49.4
567	PH	PH domain	7.8e-06	25.9
568	PH	PH domain	3.1e-39	143.8
569	Hist_deacetyl	Histone deacetylase family	5.2e-106	365.6
570	PDZ	PDZ domain (Also known as DHR or GLGF).	3.4e-20	80.5
571	zf-C3HC4	Zinc finger, C3HC4 type (RING finger)	1e-16	58.5
573	ubiquitin	Ubiquitin family	1.4e-08	31.1
574	FH2	Formin Homology 2 Domain	1.3e-110	380.9
576	serpin	Serpins (serine protease inhibitors)	4.3e-146	496.4
579	zf-C2H2	Zinc finger, C2H2 type	5.7e-76	265.8
580	pkinase	Eukaryotic protein kinase domain	6.9e-79	275.5
581	RhoGAP	RhoGAP domain	4.4e-53	189.8
582	Ribosomal_L7Ae	Ribosomal protein L7Ae	0.028	1.0
584	kazal	Kazal-type serine protease inhibitor domain	2.2e-52	187.4
585	LRR	Leucine Rich Repeat	4.4e-28	106.7
586	PHD	PHD-finger	3.8e-12	53.8
588	GTP1_OBG	GTP1/OBG family	1.1e-62	215.2
590	Collagen	Collagen triple helix repeat (20 copies)	8e-42	152.4
591	lys	C-type lysozyme/alpha-lactalbumin family	1.6e-31	116.4
596	ACBP	Acyl CoA binding protein	0.0022	-9.4
597	SNF2_N	SNF2 and others N-terminal domain	3.7e-98	339.5
600	KRAB	KRAB box	1.3e-29	111.8
606	LRR	Leucine Rich Repeat	1e-05	32.5
607	LRR	Leucine Rich Repeat	1e-05	32.5
608	WD40	WD domain, G-beta repeat	5.5e-23	39.8
610	cpn60_TCP1	TCP-1/cpn60 chaperonin family	1.7e-237	802.4
613	THF_DHG_CYH	Tetrahydrofolate dehydrogenase/cyclohydro	4.9e-173	588.3
617	rrm	RNA recognition motif.	4e-14	60.4
618	rrm	RNA recognition motif.	4e-14	60.4
620	cofilin_ADF	Cofilin/tropomyosin-type actin-binding pr	3e-06	34.2
621	Nop	Putative snoRNA binding domain	6.1e-95	328.8
622	UCH-2	Ubiquitin carboxyl-terminal hydrolase family	5.8e-21	83.1
625	zf-C2H2	Zinc finger, C2H2 type	2.5e-124	426.4
628	DEAD	DEAD/DEAH box helicase	2.5e-68	219.0
632	GST	Glutathione S-transferases.	4.8e-26	89.0
633	5_nucleotidase	5'-nucleotidase	6.6e-248	837.0
636	LIM	LIM domain containing proteins	1.6e-88	307.5
637	pkinase	Eukaryotic protein kinase domain	1.5e-73	257.8
638	MSP_domain	MSP (Major sperm protein) domain	8.4e-09	42.7
639	metalthio	Metallothionein	2e-24	94.6
641	zf-C2H2	Zinc finger, C2H2 type	6.1e-114	391.9
642	Ribosomal_S28e	Ribosomal protein S28e	9.3e-48	172.1
643	Ribosomal_S5	Ribosomal protein S5	8.3e-87	301.8
646	PHD	PHD-finger	0.00025	23.1
647	WD40	WD domain, G-beta repeat	1.5e-22	88.4

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
648	Lipase_GDSL	Lipase/Acylhydrolase with GDSL-like motif	0.015	2.2
652	zf-C2H2	Zinc finger, C2H2 type	4.1e-146	498.8
653	histone	Core histone H2A/H2B/H3/H4	1.2e-10	48.8
654	zf-C2H2	Zinc finger, C2H2 type	1.9e-87	303.9
655	ras	Ras family	6.4e-77	269.0
657	zf-C3HC4	Zinc finger, C3HC4 type (RING finger)	5.3e-13	46.4
658	STphosphatase	Ser/Thr protein phosphatase	2.6e-182	619.1
659	zf-C2H2	Zinc finger, C2H2 type	1.3e-92	321.1
660	zf-C2H2	Zinc finger, C2H2 type	1.5e-85	297.6
662	NDK	Nucleoside diphosphate kinases	1.4e-119	410.7
664	IRF	Interferon regulatory factor transcription f	7e-20	79.5
665	4HPPD_C	4-hydroxyphenylpyruvate dioxygenase C term	1.4e-16	68.5
666	DEAD	DEAD/DEAH box helicase	4.8e-74	237.1
667	DEAD	DEAD/DEAH box helicase	2.9e-70	225.1
669	pkinese	Eukaryotic protein kinase domain	6.1e-93	322.2
671	homeobox	Homeobox domain	0.018	16.5
678	crystall	Beta/Gamma crystallin	4.7e-106	365.8
679	WD40	WD domain, G-beta repeat	1.9e-06	34.9
680	Keratin_B2	Keratin, high sulfur B2 protein	4.1e-06	15.9
682	G-gamma	GGL domain	8.5e-33	117.9
685	UCH-2	Ubiquitin carboxyl-terminal hydrolase family	1.4e-29	111.7
686	Acetyltransf	Acetyltransferase (GNAT) family	6.6e-10	46.4
687	7tm_1	7 transmembrane receptor (rhodopsin family)	4.6e-15	50.0
688	proteasome	Proteasome A-type and B-type	6.5e-64	225.7
689	SCP2	SCP-2 sterol transfer family	6.2e-37	136.1
690	TS-N	TS-N domain	0.041	20.1
692	zf-C2H2	Zinc finger, C2H2 type	9.9e-60	211.9
693	zf-MYND	MYND finger	0.038	5.5
694	Oxysterol BP	Oxysterol-binding protein	3.9e-133	455.7
695	PDZ	PDZ domain (Also known as DHR or GLGF)	3e-30	115.1
703	Peptidase_C2	Calpain family cysteine protease	2.3e-175	596.0
706	filament	Intermediate filament proteins	7.2e-107	368.5
710	fibrinogen_C	Fibrinogen beta and gamma chains, C-term	7e-80	278.0
711	SH2	Src homology domain 2	2.3e-65	192.1
712	ATP-synt_DE	ATP synthase, Delta/Epsilon chain	0.00062	19.0
713	ARID	ARID DNA binding domain	2e-17	71.3
714	LBP_BPI_CETP	LBP / BPI / CETP family	8.6e-34	125.7
715	RNA_pol_L	RNA polymerases L / 13 to 16 kDa subunit	4.8e-49	176.3
716	KRAB	KRAB box	1.3e-42	155.0
717	mito_carr	Mitochondrial carrier proteins	4.8e-38	133.3
719	Gal-bind_lectin	Vertebrate galactoside-binding lectin	1.5e-25	90.2
726	aldehydh	Aldehyde dehydrogenase family	1.3e-119	410.8
728	Glycos_transf_2	Glycosyl transferases	4e-21	83.6
734	ELM2	ELM2 domain	2e-34	127.8
735	PR55	Protein phosphatase 2A regulatory subunit PR	0	1038.2
737	DSPc	Dual specificity phosphatase, catalytic doma	4e-14	60.4
740	WD40	WD domain, G-beta repeat	5.6e-14	59.9
745	zf-C3HC4	Zinc finger, C3HC4 type (RING	3.8e-13	46.9

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
		finger)		
749	mito_carr	Mitochondrial carrier proteins	4.5e-67	232.8
750	DUF27	Domain of unknown function DUF27	4.5e-12	53.5
751	SH3	SH3 domain	3.6e-17	70.5
752	HMG_box	HMG (high mobility group) box	8.6e-13	55.9
753	SPRY	SPRY domain	5.9e-05	23.3
754	GTP_CDC	Cell division protein	7.5e-153	521.2
755	mito_carr	Mitochondrial carrier proteins	3e-88	305.4
756	TSPN	Thrombospondin N-terminal -like domains	8.1e-58	205.5
757	BTB	BTB/POZ domain	5.7e-23	89.7
759	zf-C2H2	Zinc finger, C2H2 type	1.2e-12	55.4
760	NSF	NSF attachment protein	6.4e-127	435.1
762	Ribosomal_S14	Ribosomal protein S14p/S29e	2.1e-06	24.8
765	ThiF_family	ThiF family	1.7e-39	144.6
766	DnaJ	DnaJ domain	3.9e-36	133.5
768	tRNA-synt_2b	tRNA synthetase class II	9.1e-81	281.7
769	ldl_recept_a	Low-density lipoprotein receptor domain	0	1404.5
770	WD40	WD domain, G-beta repeat	2e-21	84.6
771	LRR	Leucine Rich Repeat	3.8e-06	33.9
774	SNF2_N	SNF2 and others N-terminal domain	5.5e-99	342.3
776	VPS9	Vacuolar sorting protein 9 (VPS9) domain	1.1e-30	115.4
777	VPS9	Vacuolar sorting protein 9 (VPS9) domain	1.1e-30	115.4
778	VPS9	Vacuolar sorting protein 9 (VPS9) domain	1.1e-30	115.4
779	zf-C3HC4	Zinc finger, C3HC4 type (RING finger)	3.1e-08	31.0
781	cadherin	Cadherin domain	5.6e-113	388.7
783	HECT	HECT-domain (ubiquitin-transferase).	4.2e-31	116.8
785	sushi	Sushi domain (SCR repeat)	1.8e-60	214.3
786	sushi	Sushi domain (SCR repeat)	1.8e-60	214.3
788	vwa	von Willebrand factor type A domain	9e-52	187.7
790	rrm	RNA recognition motif.	2.8e-20	80.8
791	Collagen	Collagen triple helix repeat (20 copies)	0.00097	9.7
792	pkinase	Eukaryotic protein kinase domain	0.023	12.4
795	zf-C2H2	Zinc finger, C2H2 type	6.5e-95	328.7
796	adh_short	short chain dehydrogenase	4.1e-05	-7.3
799	SAICAR_synt	SAICAR synthetase	6e-125	428.5
805	WD40	WD domain, G-beta repeat	4e-65	229.8
806	ZU5	ZU5 domain	4.7e-37	136.5
807	WD40	WD domain, G-beta repeat	0.016	21.8
808	WD40	WD domain, G-beta repeat	0.0041	23.8
809	pkinase	Eukaryotic protein kinase domain	2e-31	117.2
810	vwa	von Willebrand factor type A domain	1.9e-52	187.7
814	zf-C2H2	Zinc finger, C2H2 type	4.5e-83	289.4
815	zf-C2H2	Zinc finger, C2H2 type	6e-74	259.1
817	myosin_head	Myosin head (motor domain)	1.5e-176	599.9
818	GSPII_E	Bacterial type II secretion system protein	0.012	11.5
819	PDEase	3'5'-cyclic nucleotide phosphodiesterase	1.1e-74	215.5
821	PH	PH domain	0.00025	20.5
822	CNH	CNH domain	0.00015	-24.7
827	rrm	RNA recognition motif.	1.5e-06	35.2

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
829	HMG_box	HMG (high mobility group) box	7.8e-34	125.8
830	RasGEF	RasGEF domain	2.2e-102	353.5
831	CNH	CNH domain	3e-118	406.2
832	mito_carr	Mitochondrial carrier proteins	3.7e-37	130.3
833	PX	PX domain	2.7e-19	77.5
837	Y_phosphatase	Protein-tyrosine phosphatase	1.6e-263	888.8
838	ank	Ank repeat	2.4e-270	911.5
840	ank	Ank repeat	5.8e-38	139.6
842	Ribosomal_L15e	Ribosomal L15	4.8e-131	448.8
843	SNF	Sodium:neurotransmitter symporter family	0	1201.8
845	Peptidase_M16	Insulinase (Peptidase family M16)	4.7e-67	236.2
848	EF1BD	EF-1 guanine nucleotide exchange domain	2.2e-56	200.7
849	zf-C2H2	Zinc finger, C2H2 type	1.5e-122	420.5
850	zf-C2H2	Zinc finger, C2H2 type	2e-67	237.4
852	SIS	SIS domain	3.8e-30	113.6
853	RhoGAP	RhoGAP domain	1.1e-37	138.6
854	PDZ	PDZ domain (Also known as DHR or GLGF).	5.1e-10	46.7
856	ACOX	Acyl-CoA oxidase	9.1e-263	886.3
858	efhand	EF hand	2.4e-18	74.4
860	homeobox	Homeobox domain	4e-22	86.9
862	TFIIF_beta	Transcription initiation factor IIF, beta	2.2e-134	459.8
866	A2M	Alpha-2-macroglobulin family	4.9e-21	70.9
867	MoCF_biosynth	Molybdenum cofactor biosynthesis protei	5.8e-205	694.3
868	EGF	EGF-like domain	4.1e-22	86.9
869	EGF	EGF-like domain	1.1e-22	88.8
871	PI-PLC-X	Phosphatidylinositol-specific phospholipase	7.2e-95	328.6
872	UCH-2	Ubiquitin carboxyl-terminal hydrolase family	1.1e-20	82.1
874	SH3	SH3 domain	2.2e-14	61.1
877	SH3	SH3 domain	8.6e-14	311.7
882	KRAB	KRAB box	6.9e-42	162.6
885	ank	Ank repeat	7.1e-07	36.3
886	biopterin_H	Biopterin-dependent aromatic amino acid h	0	988.3
887	GTP_EFTU	Elongation factor Tu family	4.9e-129	437.5
888	zf-C3HC4	Zinc finger, C3HC4 type (RING finger)	1.6e-14	51.4
889	zf-C2H2	Zinc finger, C2H2 type	3.7e-92	319.6
890	ig	Immunoglobulin domain	3.8e-06	24.8
892	PTR2	POT family	9.5e-48	163.0
893	Sulfatase	Sulfatase	3.5e-78	273.2
894	Sulfatase	Sulfatase	3.5e-78	273.2
895	7tm_1	7 transmembrane receptor (rhodopsin family)	4.5e-51	164.4
896	Glyco_hydro_31	Glycosyl hydrolases family 31	0	1277.3
897	chromo	'chromo' (CHRromatin Organization MODifier)	3.9e-06	26.0
898	Cbl_N	CBL proto-oncogene N-terminal domain	1.2e-273	922.4
899	vwa	von Willebrand factor type A domain	5.5e-32	119.7
900	WD40	WD domain, G-beta repeat	2.7e-07	37.7
901	zf-C2H2	Zinc finger, C2H2 type	4e-156	532.1
903	ras	Ras family	6.6e-101	348.6

SEQ ID NO:	PFAM NAME	DESCRIPTION	p-value	PFAM SCORE
904	Armadillo_seg	Armadillo/beta-catenin-like repeats	1.1e-06	35.6
906	FH2	Formin Homology 2 Domain	4.5e-112	385.7
907	Cytidylyltransf	Cytidylyltransferase	1.4e-05	29.3
908	pkinase	Eukaryotic protein kinase domain	1.2e-64	228.2
909	pkinase	Eukaryotic protein kinase domain	8.5e-70	245.3
910	pkinase	Eukaryotic protein kinase domain	2.9e-42	153.8
911	pkinase	Eukaryotic protein kinase domain	1.2e-35	131.8
912	PHD	PHD-finger	5.1e-06	33.4
913	PHD	PHD-finger	5.5e-16	66.5
916	filament	Intermediate filament proteins	9.7e-121	414.5
917	LIM	LIM domain containing proteins	5.9e-15	57.9
918	SAM	SAM domain (Sterile alpha motif)	4.3e-16	66.9
922	Acylphosphatase	Acylphosphatase	2.9e-63	223.6
924	ig	Immunoglobulin domain	1.3e-08	32.8
925	Acyl-CoA_dh	Acyl-CoA dehydrogenase	2.4e-131	449.8
927	7tm_1	7 transmembrane receptor (rhodopsin family)	2.9e-45	145.9
928	globin	Globin	2.4e-52	186.9
929	sugar_tr	Sugar (and other) transporter	1.2e-16	68.8
932	Collagen	Collagen triple helix repeat (20 copies)	0.00097	9.7
933	HMG_box	HMG (high mobility group) box	7.8e-34	125.8
934	SEA	SEA domain	0.0021	24.7
935	ras	Ras family	6.4e-59	209.2
936	CH	Calponin homology (CH) domain	3.8e-21	83.7
937	voltage_CLC	Voltage gated chloride channels	1.9e-199	676.0
938	homeobox	Homeobox domain	1.9e-25	98.0
940	pkinase	Eukaryotic protein kinase domain	9.9e-58	205.2
942	Myosin tail	Myosin tail	3.7e-09	38.2
943	zf-C2H2	Zinc finger, C2H2 type	2.2e-92	320.3
945	Clat_adaptor_s	Clathrin adaptor complex small chain	1.3e-76	268.0
946	sugar_tr	Sugar (and other) transporter	0.017	-122.8
947	tRNA-synt_1e	tRNA synthetases class I (C)	0.00097	15.6
948	PHD	PHD-finger	2.2e-17	71.2
951	sugar_tr	Sugar (and other) transporter	0.0082	-113.9
952	mito_carr	Mitochondrial carrier proteins	1.7e-22	189.7
953	myb_DNA-binding	Myb-like DNA-binding domain	4.5e-23	80.1
955	ketoacyl-synt	Beta-ketoacyl synthase	7.1e-133	454.8
957	aldo_ket_red	Aldo/keto reductase family	1.5e-98	340.8
959	Kelch	Kelch motif	0.02	20.8
961	ras	Ras family	2.2e-29	111.1
964	homeobox	Homeobox domain	5.4e-22	86.5
965	PH	PH domain	3e-21	80.9
966	zf-C3HC4	Zinc finger, C3HC4 type (RING finger)	2.2e-09	34.7
967	Ribosomal_L29	Ribosomal L29 protein	1.6e-15	65.0
970	FAD_binding_2	FAD binding domain	8.9e-47	166.6
971	rve	Integrase core domain	0.00015	19.8
972	Glycos_transf_2	Glycosyl transferases	2.1e-21	84.5
974	Ribosomal_L10	Ribosomal protein L10	3.3e-48	173.6
975	7tm_1	7 transmembrane receptor (rhodopsin family)	1.6e-37	121.3
976	zf-C4	Zinc finger, C4 type (two domains)	2.1e-52	178.5
977	zf-C2H2	Zinc finger, C2H2 type	6.6e-150	511.4
978	FTHFS	Formate-tetrahydrofolate ligase	0	1367.2
982	Renal_dipeptase	Renal dipeptidase	1.3e-73	258.0
984	A_deaminase	Adenosine/AMP deaminase	2.6e-05	-48.6

TABLE 5

SEQ ID NO: of full-length nucleotide sequence	SEQ ID NO: of full-length peptide sequence	SEQ ID NO: of contig nucleotide sequence	SEQ ID NO: of contig. peptide sequence	Priority docket number correspondin g SEQ ID NO: in priority application	SEQ ID NO: in U.S.S.N. 09/496,914
1	985	1969	2953	787CIP2_1	150
2	986	1970	2954	787CIP2_2	223
3	987	1971	2955	787CIP2_3	1884
4	988	1972	2956	787CIP2_4	2123
5	989	1973	2957	787CIP2_5	2313
6	990	1974	2958	787CIP2_6	3284
7	991	1975	2959	787CIP2_7	3324
8	992	1976	2960	787CIP2_8	6182
9	993	1977	2961	787CIP2_9	6210
10	994	1978	2962	787CIP2_10	6213
11	995	1979	2963	787CIP2_11	6257
12	996	1980	2964	787CIP2_12	6294
13	997	1981	2965	787CIP2_13	6294
14	998	1982	2966	787CIP2_14	6330
15	999	1983	2967	787CIP2_15	6364
16	1000	1984	2968	787CIP2_16	6455
17	1001	1985	2969	787CIP2_17	6486
18	1002	1986	2970	787CIP2_18	6503
19	1003	1987	2971	787CIP2_19	6528
20	1004	1988	2972	787CIP2_20	6572
21	1005	1989	2973	787CIP2_21	6578
22	1006	1990	2974	787CIP2_22	6593
23	1007	1991	2975	787CIP2_23	6603
24	1008	1992	2976	787CIP2_24	6603
25	1009	1993	2977	787CIP2_25	6679
26	1010	1994	2978	787CIP2_26	6744
27	1011	1995	2979	787CIP2_27	6762
28	1012	1996	2980	787CIP2_28	6770
29	1013	1997	2981	787CIP2_29	6770
30	1014	1998	2982	787CIP2_30	6787
31	1015	1999	2983	787CIP2_31	6858
32	1016	2000	2984	787CIP2_32	6866
33	1017	2001	2985	787CIP2_33	6938
34	1018	2002	2986	787CIP2_34	6938
35	1019	2003	2987	787CIP2_35	6977
36	1020	2004	2988	787CIP2_36	7001
37	1021	2005	2989	787CIP2_37	7002
38	1022	2006	2990	787CIP2_38	7004
39	1023	2007	2991	787CIP2_39	7005
40	1024	2008	2992	787CIP2_40	7006
41	1025	2009	2993	787CIP2_41	7008
42	1026	2010	2994	787CIP2_42	7014
43	1027	2011	2995	787CIP2_43	7021
44	1028	2012	2996	787CIP2_44	7022
45	1029	2013	2997	787CIP2_46	7057
46	1030	2014	2998	787CIP2_47	7058
47	1031	2015	2999	787CIP2_49	7088
48	1032	2016	3000	787CIP2_50	7089
49	1033	2017	3001	787CIP2_51	7182
50	1034	2018	3002	787CIP2_52	7489
51	1035	2019	3003	787CIP2_53	7564
52	1036	2020	3004	787CIP2_54	7566
53	1037	2021	3005	787CIP2_55	7587

54	1038	2022	3006	787CIP2_56	7591
55	1039	2023	3007	787CIP2_57	7600
56	1040	2024	3008	787CIP2_58	7604
57	1041	2025	3009	787CIP2_59	7612
58	1042	2026	3010	787CIP2_60	7613
59	1043	2027	3011	787CIP2_61	7615
60	1044	2028	3012	787CIP2_62	7616
61	1045	2029	3013	787CIP2_63	7617
62	1046	2030	3014	787CIP2_64	7623
63	1047	2031	3015	787CIP2_65	7625
64	1048	2032	3016	787CIP2_66	7625
65	1049	2033	3017	787CIP2_67	7630
66	1050	2034	3018	787CIP2_68	7638
67	1051	2035	3019	787CIP2_69	7640
68	1052	2036	3020	787CIP2_70	7670
69	1053	2037	3021	787CIP2_71	7676
70	1054	2038	3022	787CIP2_72	7688
71	1055	2039	3023	787CIP2_73	7690
72	1056	2040	3024	787CIP2_74	7700
73	1057	2041	3025	787CIP2_75	7774
74	1058	2042	3026	787CIP2_76	7784
75	1059	2043	3027	787CIP2_77	7785
76	1060	2044	3028	787CIP2_78	7792
77	1061	2045	3029	787CIP2_79	7798
78	1062	2046	3030	787CIP2_80	7807
79	1063	2047	3031	787CIP2_81	7810
80	1064	2048	3032	787CIP2_82	7812
81	1065	2049	3033	787CIP2_83	7816
82	1066	2050	3034	787CIP2_84	7826
83	1067	2051	3035	787CIP2_85	7842
84	1068	2052	3036	787CIP2_86	7850
85	1069	2053	3037	787CIP2_87	7865
86	1070	2054	3038	787CIP2_88	7882
87	1071	2055	3039	787CIP2_89	7891
88	1072	2056	3040	787CIP2_90	7892
89	1073	2057	3041	787CIP2_91	7896
90	1074	2058	3042	787CIP2_92	7896
91	1075	2059	3043	787CIP2_93	7907
92	1076	2060	3044	787CIP2_94	7913
93	1077	2061	3045	787CIP2_95	7914
94	1078	2062	3046	787CIP2_96	7915
95	1079	2063	3047	787CIP2_97	7920
96	1080	2064	3048	787CIP2_98	7921
97	1081	2065	3049	787CIP2_99	7924
98	1082	2066	3050	787CIP2_100	7927
99	1083	2067	3051	787CIP2_101	7929
100	1084	2068	3052	787CIP2_102	7937
101	1085	2069	3053	787CIP2_103	7940
102	1086	2070	3054	787CIP2_104	7942
103	1087	2071	3055	787CIP2_105	7944
104	1088	2072	3056	787CIP2_106	7951
105	1089	2073	3057	787CIP2_107	7951
106	1090	2074	3058	787CIP2_108	7962
107	1091	2075	3059	787CIP2_109	7964
108	1092	2076	3060	787CIP2_110	7977
109	1093	2077	3061	787CIP2_111	7978
110	1094	2078	3062	787CIP2_112	7980
111	1095	2079	3063	787CIP2_113	7982
112	1096	2080	3064	787CIP2_114	8000
113	1097	2081	3065	787CIP2_115	8003

114	1098	2082	3066	787CIP2_116	8004
115	1099	2083	3067	787CIP2_117	8007
116	1100	2084	3068	787CIP2_118	8008
117	1101	2085	3069	787CIP2_119	8009
118	1102	2086	3070	787CIP2_120	8013
119	1103	2087	3071	787CIP2_121	8017
120	1104	2088	3072	787CIP2_122	8018
121	1105	2089	3073	787CIP2_123	8021
122	1106	2090	3074	787CIP2_124	8022
123	1107	2091	3075	787CIP2_125	8023
124	1108	2092	3076	787CIP2_126	8023
125	1109	2093	3077	787CIP2_127	8024
126	1110	2094	3078	787CIP2_128	8026
127	1111	2095	3079	787CIP2_129	8028
128	1112	2096	3080	787CIP2_130	8036
129	1113	2097	3081	787CIP2_131	8038
130	1114	2098	3082	787CIP2_132	8045
131	1115	2099	3083	787CIP2_133	8045
132	1116	2100	3084	787CIP2_134	8048
133	1117	2101	3085	787CIP2_135	8048
134	1118	2102	3086	787CIP2_136	8052
135	1119	2103	3087	787CIP2_137	8053
136	1120	2104	3088	787CIP2_138	8055
137	1121	2105	3089	787CIP2_139	8059
138	1122	2106	3090	787CIP2_140	8061
139	1123	2107	3091	787CIP2_141	8062
140	1124	2108	3092	787CIP2_142	8063
141	1125	2109	3093	787CIP2_143	8064
142	1126	2110	3094	787CIP2_144	8065
143	1127	2111	3095	787CIP2_145	8068
144	1128	2112	3096	787CIP2_146	8069
145	1129	2113	3097	787CIP2_147	8070
146	1130	2114	3098	787CIP2_148	8074
147	1131	2115	3099	787CIP2_149	8076
148	1132	2116	3100	787CIP2_150	8077
149	1133	2117	3101	787CIP2_151	8078
150	1134	2118	3102	787CIP2_152	8079
151	1135	2119	3103	787CIP2_153	8087
152	1136	2120	3104	787CIP2_154	8091
153	1137	2121	3105	787CIP2_155	8100
154	1138	2122	3106	787CIP2_156	8105
155	1139	2123	3107	787CIP2_157	8106
156	1140	2124	3108	787CIP2_158	8108
157	1141	2125	3109	787CIP2_159	8109
158	1142	2126	3110	787CIP2_160	8110
159	1143	2127	3111	787CIP2_161	8112
160	1144	2128	3112	787CIP2_162	8116
161	1145	2129	3113	787CIP2_163	8118
162	1146	2130	3114	787CIP2_164	8124
163	1147	2131	3115	787CIP2_165	8125
164	1148	2132	3116	787CIP2_166	8127
165	1149	2133	3117	787CIP2_167	8132
166	1150	2134	3118	787CIP2_168	8135
167	1151	2135	3119	787CIP2_169	8137
168	1152	2136	3120	787CIP2_170	8139
169	1153	2137	3121	787CIP2_171	8140
170	1154	2138	3122	787CIP2_172	8140
171	1155	2139	3123	787CIP2_173	8140
172	1156	2140	3124	787CIP2_174	8141
173	1157	2141	3125	787CIP2_175	8147

174	1158	2142	3126	787CIP2_176	8149
175	1159	2143	3127	787CIP2_177	8150
176	1160	2144	3128	787CIP2_178	8157
177	1161	2145	3129	787CIP2_179	8161
178	1162	2146	3130	787CIP2_180	8162
179	1163	2147	3131	787CIP2_181	8165
180	1164	2148	3132	787CIP2_182	8166
181	1165	2149	3133	787CIP2_183	8167
182	1166	2150	3134	787CIP2_184	8169
183	1167	2151	3135	787CIP2_185	8170
184	1168	2152	3136	787CIP2_186	8172
185	1169	2153	3137	787CIP2_187	8173
186	1170	2154	3138	787CIP2_188	8174
187	1171	2155	3139	787CIP2_189	8174
188	1172	2156	3140	787CIP2_191	8182
189	1173	2157	3141	787CIP2_192	8186
190	1174	2158	3142	787CIP2_193	8188
191	1175	2159	3143	787CIP2_194	8191
192	1176	2160	3144	787CIP2_195	8192
193	1177	2161	3145	787CIP2_196	8193
194	1178	2162	3146	787CIP2_197	8194
195	1179	2163	3147	787CIP2_198	8195
196	1180	2164	3148	787CIP2_199	8196
197	1181	2165	3149	787CIP2_200	8200
198	1182	2166	3150	787CIP2_201	8201
199	1183	2167	3151	787CIP2_202	8202
200	1184	2168	3152	787CIP2_203	8205
201	1185	2169	3153	787CIP2_204	8206
202	1186	2170	3154	787CIP2_205	8207
203	1187	2171	3155	787CIP2_206	8208
204	1188	2172	3156	787CIP2_207	8209
205	1189	2173	3157	787CIP2_208	8210
206	1190	2174	3158	787CIP2_209	8211
207	1191	2175	3159	787CIP2_210	8212
208	1192	2176	3160	787CIP2_211	8213
209	1193	2177	3161	787CIP2_212	8214
210	1194	2178	3162	787CIP2_213	8215
211	1195	2179	3163	787CIP2_214	8215
212	1196	2180	3164	787CIP2_215	8217
213	1197	2181	3165	787CIP2_217	8221
214	1198	2182	3166	787CIP2_218	8222
215	1199	2183	3167	787CIP2_219	8223
216	1200	2184	3168	787CIP2_220	8224
217	1201	2185	3169	787CIP2_221	8225
218	1202	2186	3170	787CIP2_222	8227
219	1203	2187	3171	787CIP2_223	8232
220	1204	2188	3172	787CIP2_224	8235
221	1205	2189	3173	787CIP2_225	8236
222	1206	2190	3174	787CIP2_227	8238
223	1207	2191	3175	787CIP2_228	8239
224	1208	2192	3176	787CIP2_229	8240
225	1209	2193	3177	787CIP2_230	8242
226	1210	2194	3178	787CIP2_231	8246
227	1211	2195	3179	787CIP2_232	8252
228	1212	2196	3180	787CIP2_233	8257
229	1213	2197	3181	787CIP2_234	8288
230	1214	2198	3182	787CIP2_235	8310
231	1215	2199	3183	787CIP2_236	8311
232	1216	2200	3184	787CIP2_237	8315
233	1217	2201	3185	787CIP2_238	8318

234	1218	2202	3186	787CIP2_239	8326
235	1219	2203	3187	787CIP2_240	8326
236	1220	2204	3188	787CIP2_241	8336
237	1221	2205	3189	787CIP2_242	8351
238	1222	2206	3190	787CIP2_243	8364
239	1223	2207	3191	787CIP2_244	8372
240	1224	2208	3192	787CIP2_245	8376
241	1225	2209	3193	787CIP2_246	8377
242	1226	2210	3194	787CIP2_247	8382
243	1227	2211	3195	787CIP2_248	8404
244	1228	2212	3196	787CIP2_249	8410
245	1229	2213	3197	787CIP2_250	8419
246	1230	2214	3198	787CIP2_251	8430
247	1231	2215	3199	787CIP2_252	8448
248	1232	2216	3200	787CIP2_253	8458
249	1233	2217	3201	787CIP2_254	8461
250	1234	2218	3202	787CIP2_255	8466
251	1235	2219	3203	787CIP2_256	8468
252	1236	2220	3204	787CIP2_257	8477
253	1237	2221	3205	787CIP2_258	8481
254	1238	2222	3206	787CIP2_259	8491
255	1239	2223	3207	787CIP2_260	8503
256	1240	2224	3208	787CIP2_261	8513
257	1241	2225	3209	787CIP2_262	8514
258	1242	2226	3210	787CIP2_263	8518
259	1243	2227	3211	787CIP2_264	8547
260	1244	2228	3212	787CIP2_265	8549
261	1245	2229	3213	787CIP2_266	8549
262	1246	2230	3214	787CIP2_267	8549
263	1247	2231	3215	787CIP2_268	8550
264	1248	2232	3216	787CIP2_269	8603
265	1249	2233	3217	787CIP2_270	8625
266	1250	2234	3218	787CIP2_271	8625
267	1251	2235	3219	787CIP2_272	8633
268	1252	2236	3220	787CIP2_273	8648
269	1253	2237	3221	787CIP2_274	8654
270	1254	2238	3222	787CIP2_275	8671
271	1255	2239	3223	787CIP2_276	8733
272	1256	2240	3224	787CIP2_277	8735
273	1257	2241	3225	787CIP2_278	8747
274	1258	2242	3226	787CIP2_279	8748
275	1259	2243	3227	787CIP2_280	8753
276	1260	2244	3228	787CIP2_281	8770
277	1261	2245	3229	787CIP2_282	8777
278	1262	2246	3230	787CIP2_283	8828
279	1263	2247	3231	787CIP2_284	8836
280	1264	2248	3232	787CIP2_285	8842
281	1265	2249	3233	787CIP2_286	8842
282	1266	2250	3234	787CIP2_287	8850
283	1267	2251	3235	787CIP2_288	8851
284	1268	2252	3236	787CIP2_289	8852
285	1269	2253	3237	787CIP2_290	8853
286	1270	2254	3238	787CIP2_291	8854
287	1271	2255	3239	787CIP2_292	9084
288	1272	2256	3240	787CIP2_293	9099
289	1273	2257	3241	787CIP2_294	9691
290	1274	2258	3242	787CIP2_295	9699
291	1275	2259	3243	787CIP2_296	9883
292	1276	2260	3244	787CIP2_297	9886
293	1277	2261	3245	787CIP2_298	10334

294	1278	2262	3246	787CIP2_299	10335
295	1279	2263	3247	787CIP2_300	10336
296	1280	2264	3248	787CIP2_301	10338
297	1281	2265	3249	787CIP2_302	10339
298	1282	2266	3250	787CIP2_304	10342
299	1283	2267	3251	787CIP2_305	10342
300	1284	2268	3252	787CIP2_306	10343
301	1285	2269	3253	787CIP2_307	10344
302	1286	2270	3254	787CIP2_308	10345
303	1287	2271	3255	787CIP2_309	10346
304	1288	2272	3256	787CIP2_310	10347
305	1289	2273	3257	787CIP2_311	10348
306	1290	2274	3258	787CIP2_312	10349
307	1291	2275	3259	787CIP2_314	10351
308	1292	2276	3260	787CIP2_315	10352
309	1293	2277	3261	787CIP2_316	10353
310	1294	2278	3262	787CIP2_317	10354
311	1295	2279	3263	787CIP2_318	10355
312	1296	2280	3264	787CIP2_319	10356
313	1297	2281	3265	787CIP2_320	10357
314	1298	2282	3266	787CIP2_321	10358
315	1299	2283	3267	787CIP2_322	10360
316	1300	2284	3268	787CIP2_323	10361
317	1301	2285	3269	787CIP2_324	10362
318	1302	2286	3270	787CIP2_325	10363
319	1303	2287	3271	787CIP2_326	10365
320	1304	2288	3272	787CIP2_327	10366
321	1305	2289	3273	787CIP2_328	10367
322	1306	2290	3274	787CIP2_329	10369
323	1307	2291	3275	787CIP2_330	10370
324	1308	2292	3276	787CIP2_331	10371
325	1309	2293	3277	787CIP2_332	10372
326	1310	2294	3278	787CIP2_333	10373
327	1311	2295	3279	787CIP2_334	10375
328	1312	2296	3280	787CIP2_335	10377
329	1313	2297	3281	787CIP2_336	10379
330	1314	2298	3282	787CIP2_337	10381
331	1315	2299	3283	787CIP2_338	10382
332	1316	2300	3284	787CIP2_339	10383
333	1317	2301	3285	787CIP2_340	10384
334	1318	2302	3286	787CIP2_341	10385
335	1319	2303	3287	787CIP2_342	10386
336	1320	2304	3288	787CIP2_343	10387
337	1321	2305	3289	787CIP2_346	10391
338	1322	2306	3290	787CIP2_348	10393
339	1323	2307	3291	787CIP2_349	10394
340	1324	2308	3292	787CIP2_350	10395
341	1325	2309	3293	787CIP2_351	10396
342	1326	2310	3294	787CIP2_352	10397
343	1327	2311	3295	787CIP2_353	10399
344	1328	2312	3296	787CIP2_354	10400
345	1329	2313	3297	787CIP2_355	10401
346	1330	2314	3298	787CIP2_357	10403
347	1331	2315	3299	787CIP2_358	10404
348	1332	2316	3300	787CIP2_359	10407
349	1333	2317	3301	787CIP2_360	10408
350	1334	2318	3302	787CIP2_361	10409
351	1335	2319	3303	787CIP2_362	10410
352	1336	2320	3304	787CIP2B_1	44
353	1337	2321	3305	787CIP2B_2	50

354	1338	2322	3306	787CIP2B_3	93
355	1339	2323	3307	787CIP2B_4	224
356	1340	2324	3308	787CIP2B_5	318
357	1341	2325	3309	787CIP2B_6	318
358	1342	2326	3310	787CIP2B_7	795
359	1343	2327	3311	787CIP2B_8	857
360	1344	2328	3312	787CIP2B_9	924
361	1345	2329	3313	787CIP2B_10	944
362	1346	2330	3314	787CIP2B_11	944
363	1347	2331	3315	787CIP2B_12	967
364	1348	2332	3316	787CIP2B_13	1055
365	1349	2333	3317	787CIP2B_14	1091
366	1350	2334	3318	787CIP2B_15	1225
367	1351	2335	3319	787CIP2B_16	1257
368	1352	2336	3320	787CIP2B_17	1289
369	1353	2337	3321	787CIP2B_18	1292
370	1354	2338	3322	787CIP2B_19	1455
371	1355	2339	3323	787CIP2B_20	1488
372	1356	2340	3324	787CIP2B_21	1666
373	1357	2341	3325	787CIP2B_22	1811
374	1358	2342	3326	787CIP2B_23	1885
375	1359	2343	3327	787CIP2B_24	1911
376	1360	2344	3328	787CIP2B_25	1935
377	1361	2345	3329	787CIP2B_26	1971
378	1362	2346	3330	787CIP2B_27	1989
379	1363	2347	3331	787CIP2B_28	2041
380	1364	2348	3332	787CIP2B_29	2178
381	1365	2349	3333	787CIP2B_30	2237
382	1366	2350	3334	787CIP2B_31	2279
383	1367	2351	3335	787CIP2B_32	2338
384	1368	2352	3336	787CIP2B_33	2351
385	1369	2353	3337	787CIP2B_34	2405
386	1370	2354	3338	787CIP2B_35	2531
387	1371	2355	3339	787CIP2B_36	2584
388	1372	2356	3340	787CIP2B_37	2608
389	1373	2357	3341	787CIP2B_38	2655
390	1374	2358	3342	787CIP2B_39	2655
391	1375	2359	3343	787CIP2B_40	2866
392	1376	2360	3344	787CIP2B_41	3015
393	1377	2361	3345	787CIP2B_42	3015
394	1378	2362	3346	787CIP2B_43	3043
395	1379	2363	3347	787CIP2B_44	3986
396	1380	2364	3348	787CIP2B_45	4647
397	1381	2365	3349	787CIP2B_46	4659
398	1382	2366	3350	787CIP2B_47	5032
399	1383	2367	3351	787CIP2B_48	5244
400	1384	2368	3352	787CIP2B_49	5268
401	1385	2369	3353	787CIP2B_50	5281
402	1386	2370	3354	787CIP2B_51	5282
403	1387	2371	3355	787CIP2B_52	6147
404	1388	2372	3356	787CIP2B_53	6178
405	1389	2373	3357	787CIP2B_54	6184
406	1390	2374	3358	787CIP2B_55	6187
407	1391	2375	3359	787CIP2B_56	6190
408	1392	2376	3360	787CIP2B_57	6191
409	1393	2377	3361	787CIP2B_58	6194
410	1394	2378	3362	787CIP2B_59	6196
411	1395	2379	3363	787CIP2B_60	6201
412	1396	2380	3364	787CIP2B_61	6208
413	1397	2381	3365	787CIP2B_62	6214

414	1398	2382	3366	787CIP2B_63	6217
415	1399	2383	3367	787CIP2B_64	6220
416	1400	2384	3368	787CIP2B_65	6221
417	1401	2385	3369	787CIP2B_66	6222
418	1402	2386	3370	787CIP2B_67	6223
419	1403	2387	3371	787CIP2B_68	6223
420	1404	2388	3372	787CIP2B_69	6226
421	1405	2389	3373	787CIP2B_70	6227
422	1406	2390	3374	787CIP2B_71	6229
423	1407	2391	3375	787CIP2B_72	6248
424	1408	2392	3376	787CIP2B_73	6260
425	1409	2393	3377	787CIP2B_74	6264
426	1410	2394	3378	787CIP2B_75	6269
427	1411	2395	3379	787CIP2B_76	6269
428	1412	2396	3380	787CIP2B_77	6275
429	1413	2397	3381	787CIP2B_78	6276
430	1414	2398	3382	787CIP2B_79	6280
431	1415	2399	3383	787CIP2B_80	6287
432	1416	2400	3384	787CIP2B_81	6290
433	1417	2401	3385	787CIP2B_82	6293
434	1418	2402	3386	787CIP2B_83	6305
435	1419	2403	3387	787CIP2B_84	6308
436	1420	2404	3388	787CIP2B_85	6309
437	1421	2405	3389	787CIP2B_86	6312
438	1422	2406	3390	787CIP2B_87	6314
439	1423	2407	3391	787CIP2B_88	6316
440	1424	2408	3392	787CIP2B_89	6336
441	1425	2409	3393	787CIP2B_90	6341
442	1426	2410	3394	787CIP2B_91	6343
443	1427	2411	3395	787CIP2B_92	6346
444	1428	2412	3396	787CIP2B_93	6357
445	1429	2413	3397	787CIP2B_94	6359
446	1430	2414	3398	787CIP2B_95	6367
447	1431	2415	3399	787CIP2B_96	6383
448	1432	2416	3400	787CIP2B_97	6385
449	1433	2417	3401	787CIP2B_98	6396
450	1434	2418	3402	787CIP2B_99	6396
451	1435	2419	3403	787CIP2B_100	6403
452	1436	2420	3404	787CIP2B_101	6405
453	1437	2421	3405	787CIP2B_102	6414
454	1438	2422	3406	787CIP2B_103	6418
455	1439	2423	3407	787CIP2B_104	6422
456	1440	2424	3408	787CIP2B_105	6425
457	1441	2425	3409	787CIP2B_106	6436
458	1442	2426	3410	787CIP2B_107	6471
459	1443	2427	3411	787CIP2B_108	6474
460	1444	2428	3412	787CIP2B_109	6482
461	1445	2429	3413	787CIP2B_110	6504
462	1446	2430	3414	787CIP2B_111	6510
463	1447	2431	3415	787CIP2B_112	6515
464	1448	2432	3416	787CIP2B_113	6529
465	1449	2433	3417	787CIP2B_114	6535
466	1450	2434	3418	787CIP2B_115	6536
467	1451	2435	3419	787CIP2B_116	6536
468	1452	2436	3420	787CIP2B_117	6541
469	1453	2437	3421	787CIP2B_118	6542
470	1454	2438	3422	787CIP2B_119	6547
471	1455	2439	3423	787CIP2B_120	6548
472	1456	2440	3424	787CIP2B_121	6552
473	1457	2441	3425	787CIP2B_122	6552

474	1458	2442	3426	787CIP2B_123	6555
475	1459	2443	3427	787CIP2B_124	6560
476	1460	2444	3428	787CIP2B_125	6566
477	1461	2445	3429	787CIP2B_126	6576
478	1462	2446	3430	787CIP2B_127	6584
479	1463	2447	3431	787CIP2B_128	6588
480	1464	2448	3432	787CIP2B_129	6589
481	1465	2449	3433	787CIP2B_130	6590
482	1466	2450	3434	787CIP2B_131	6597
483	1467	2451	3435	787CIP2B_132	6600
484	1468	2452	3436	787CIP2B_133	6602
485	1469	2453	3437	787CIP2B_134	6604
486	1470	2454	3438	787CIP2B_135	6605
487	1471	2455	3439	787CIP2B_136	6608
488	1472	2456	3440	787CIP2B_137	6610
489	1473	2457	3441	787CIP2B_138	6614
490	1474	2458	3442	787CIP2B_139	6623
491	1475	2459	3443	787CIP2B_140	6629
492	1476	2460	3444	787CIP2B_141	6631
493	1477	2461	3445	787CIP2B_142	6631
494	1478	2462	3446	787CIP2B_143	6631
495	1479	2463	3447	787CIP2B_144	6632
496	1480	2464	3448	787CIP2B_145	6633
497	1481	2465	3449	787CIP2B_146	6634
498	1482	2466	3450	787CIP2B_147	6635
499	1483	2467	3451	787CIP2B_148	6639
500	1484	2468	3452	787CIP2B_149	6649
501	1485	2469	3453	787CIP2B_150	6651
502	1486	2470	3454	787CIP2B_151	6655
503	1487	2471	3455	787CIP2B_152	6658
504	1488	2472	3456	787CIP2B_153	6667
505	1489	2473	3457	787CIP2B_154	6672
506	1490	2474	3458	787CIP2B_155	6682
507	1491	2475	3459	787CIP2B_156	6683
508	1492	2476	3460	787CIP2B_157	6687
509	1493	2477	3461	787CIP2B_158	6687
510	1494	2478	3462	787CIP2B_159	6693
511	1495	2479	3463	787CIP2B_160	6696
512	1496	2480	3464	787CIP2B_161	6701
513	1497	2481	3465	787CIP2B_162	6707
514	1498	2482	3466	787CIP2B_163	6712
515	1499	2483	3467	787CIP2B_164	6714
516	1500	2484	3468	787CIP2B_165	6720
517	1501	2485	3469	787CIP2B_166	6721
518	1502	2486	3470	787CIP2B_167	6722
519	1503	2487	3471	787CIP2B_168	6736
520	1504	2488	3472	787CIP2B_169	6740
521	1505	2489	3473	787CIP2B_170	6740
522	1506	2490	3474	787CIP2B_171	6760
523	1507	2491	3475	787CIP2B_172	6775
524	1508	2492	3476	787CIP2B_173	6784
525	1509	2493	3477	787CIP2B_174	6793
526	1510	2494	3478	787CIP2B_175	6795
527	1511	2495	3479	787CIP2B_176	6796
528	1512	2496	3480	787CIP2B_177	6807
529	1513	2497	3481	787CIP2B_178	6808
530	1514	2498	3482	787CIP2B_179	6810
531	1515	2499	3483	787CIP2B_180	6815
532	1516	2500	3484	787CIP2B_181	6819
533	1517	2501	3485	787CIP2B_182	6821

534	1518	2502	3486	787CIP2B_183	6827
535	1519	2503	3487	787CIP2B_184	6829
536	1520	2504	3488	787CIP2B_185	6830
537	1521	2505	3489	787CIP2B_186	6835
538	1522	2506	3490	787CIP2B_187	6848
539	1523	2507	3491	787CIP2B_188	6849
540	1524	2508	3492	787CIP2B_189	6851
541	1525	2509	3493	787CIP2B_190	6851
542	1526	2510	3494	787CIP2B_191	6863
543	1527	2511	3495	787CIP2B_192	6869
544	1528	2512	3496	787CIP2B_193	6874
545	1529	2513	3497	787CIP2B_194	6887
546	1530	2514	3498	787CIP2B_195	6890
547	1531	2515	3499	787CIP2B_196	6894
548	1532	2516	3500	787CIP2B_197	6899
549	1533	2517	3501	787CIP2B_198	6900
550	1534	2518	3502	787CIP2B_199	6903
551	1535	2519	3503	787CIP2B_200	6910
552	1536	2520	3504	787CIP2B_201	6913
553	1537	2521	3505	787CIP2B_202	6918
554	1538	2522	3506	787CIP2B_203	6923
555	1539	2523	3507	787CIP2B_204	6926
556	1540	2524	3508	787CIP2B_205	6929
557	1541	2525	3509	787CIP2B_206	6929
558	1542	2526	3510	787CIP2B_207	6932
559	1543	2527	3511	787CIP2B_208	6941
560	1544	2528	3512	787CIP2B_209	6951
561	1545	2529	3513	787CIP2B_210	6954
562	1546	2530	3514	787CIP2B_211	6954
563	1547	2531	3515	787CIP2B_212	6956
564	1548	2532	3516	787CIP2B_213	6957
565	1549	2533	3517	787CIP2B_214	6960
566	1550	2534	3518	787CIP2B_215	6966
567	1551	2535	3519	787CIP2B_216	6968
568	1552	2536	3520	787CIP2B_217	6969
569	1553	2537	3521	787CIP2B_218	6970
570	1554	2538	3522	787CIP2B_219	6971
571	1555	2539	3523	787CIP2B_220	6989
572	1556	2540	3524	787CIP2B_221	6990
573	1557	2541	3525	787CIP2B_223	6996
574	1558	2542	3526	787CIP2B_224	6997
575	1559	2543	3527	787CIP2B_225	7009
576	1560	2544	3528	787CIP2B_226	7016
577	1561	2545	3529	787CIP2B_227	7023
578	1562	2546	3530	787CIP2B_228	7023
579	1563	2547	3531	787CIP2B_229	7035
580	1564	2548	3532	787CIP2B_230	7038
581	1565	2549	3533	787CIP2B_231	7039
582	1566	2550	3534	787CIP2B_232	7040
583	1567	2551	3535	787CIP2B_233	7041
584	1568	2552	3536	787CIP2B_234	7044
585	1569	2553	3537	787CIP2B_235	7059
586	1570	2554	3538	787CIP2B_236	7060
587	1571	2555	3539	787CIP2B_237	7063
588	1572	2556	3540	787CIP2B_238	7067
589	1573	2557	3541	787CIP2B_239	7070
590	1574	2558	3542	787CIP2B_240	7071
591	1575	2559	3543	787CIP2B_241	7079
592	1576	2560	3544	787CIP2B_242	7085
593	1577	2561	3545	787CIP2B_243	7148

594	1578	2562	3546	787CIP2B_244	7156
595	1579	2563	3547	787CIP2B_245	7156
596	1580	2564	3548	787CIP2B_246	7171
597	1581	2565	3549	787CIP2B_248	7265
598	1582	2566	3550	787CIP2B_249	7268
599	1583	2567	3551	787CIP2B_250	7308
600	1584	2568	3552	787CIP2B_251	7336
601	1585	2569	3553	787CIP2B_252	7347
602	1586	2570	3554	787CIP2B_253	7405
603	1587	2571	3555	787CIP2B_254	7405
604	1588	2572	3556	787CIP2B_255	7412
605	1589	2573	3557	787CIP2B_256	7412
606	1590	2574	3558	787CIP2B_257	7436
607	1591	2575	3559	787CIP2B_258	7436
608	1592	2576	3560	787CIP2B_259	7454
609	1593	2577	3561	787CIP2B_260	7476
610	1594	2578	3562	787CIP2B_261	7598
611	1595	2579	3563	787CIP2B_262	7619
612	1596	2580	3564	787CIP2B_263	7644
613	1597	2581	3565	787CIP2B_264	7648
614	1598	2582	3566	787CIP2B_265	7659
615	1599	2583	3567	787CIP2B_266	7661
616	1600	2584	3568	787CIP2B_267	7669
617	1601	2585	3569	787CIP2B_268	7686
618	1602	2586	3570	787CIP2B_269	7686
619	1603	2587	3571	787CIP2B_270	7694
620	1604	2588	3572	787CIP2B_271	7697
621	1605	2589	3573	787CIP2B_272	7733
622	1606	2590	3574	787CIP2B_273	7734
623	1607	2591	3575	787CIP2B_274	7744
624	1608	2592	3576	787CIP2B_275	7751
625	1609	2593	3577	787CIP2B_276	7756
626	1610	2594	3578	787CIP2B_277	7761
627	1611	2595	3579	787CIP2B_278	7761
628	1612	2596	3580	787CIP2B_279	7776
629	1613	2597	3581	787CIP2B_280	7783
630	1614	2598	3582	787CIP2B_281	7800
631	1615	2599	3583	787CIP2B_282	7800
632	1616	2600	3584	787CIP2B_283	7801
633	1617	2601	3585	787CIP2B_284	7811
634	1618	2602	3586	787CIP2B_285	7817
635	1619	2603	3587	787CIP2B_286	7821
636	1620	2604	3588	787CIP2B_287	7822
637	1621	2605	3589	787CIP2B_288	7841
638	1622	2606	3590	787CIP2B_289	7847
639	1623	2607	3591	787CIP2B_290	7880
640	1624	2608	3592	787CIP2B_291	7910
641	1625	2609	3593	787CIP2B_293	7936
642	1626	2610	3594	787CIP2B_294	7945
643	1627	2611	3595	787CIP2B_295	7948
644	1628	2612	3596	787CIP2B_296	7963
645	1629	2613	3597	787CIP2B_297	7984
646	1630	2614	3598	787CIP2B_298	7985
647	1631	2615	3599	787CIP2B_299	8014
648	1632	2616	3600	787CIP2B_301	8029
649	1633	2617	3601	787CIP2B_302	8043
650	1634	2618	3602	787CIP2B_303	8164
651	1635	2619	3603	787CIP2B_304	8175
652	1636	2620	3604	787CIP2B_305	8250
653	1637	2621	3605	787CIP2B_306	8253

654	1638	2622	3606	787CIP2B_307	8255
655	1639	2623	3607	787CIP2B_308	8258
656	1640	2624	3608	787CIP2B_309	8270
657	1641	2625	3609	787CIP2B_310	8271
658	1642	2626	3610	787CIP2B_311	8272
659	1643	2627	3611	787CIP2B_312	8279
660	1644	2628	3612	787CIP2B_313	8284
661	1645	2629	3613	787CIP2B_314	8285
662	1646	2630	3614	787CIP2B_315	8304
663	1647	2631	3615	787CIP2B_316	8309
664	1648	2632	3616	787CIP2B_317	8320
665	1649	2633	3617	787CIP2B_318	8331
666	1650	2634	3618	787CIP2B_319	8332
667	1651	2635	3619	787CIP2B_320	8332
668	1652	2636	3620	787CIP2B_321	8335
669	1653	2637	3621	787CIP2B_322	8337
670	1654	2638	3622	787CIP2B_323	8353
671	1655	2639	3623	787CIP2B_324	8355
672	1656	2640	3624	787CIP2B_325	8358
673	1657	2641	3625	787CIP2B_326	8361
674	1658	2642	3626	787CIP2B_327	8369
675	1659	2643	3627	787CIP2B_328	8385
676	1660	2644	3628	787CIP2B_329	8397
677	1661	2645	3629	787CIP2B_330	8414
678	1662	2646	3630	787CIP2B_331	8431
679	1663	2647	3631	787CIP2B_332	8433
680	1664	2648	3632	787CIP2B_333	8444
681	1665	2649	3633	787CIP2B_334	8446
682	1666	2650	3634	787CIP2B_335	8460
683	1667	2651	3635	787CIP2B_336	8478
684	1668	2652	3636	787CIP2B_337	8490
685	1669	2653	3637	787CIP2B_338	8505
686	1670	2654	3638	787CIP2B_339	8523
687	1671	2655	3639	787CIP2B_340	8530
688	1672	2656	3640	787CIP2B_341	8533
689	1673	2657	3641	787CIP2B_342	8534
690	1674	2658	3642	787CIP2B_343	8537
691	1675	2659	3643	787CIP2B_344	8537
692	1676	2660	3644	787CIP2B_345	8543
693	1677	2661	3645	787CIP2B_346	8546
694	1678	2662	3646	787CIP2B_347	8553
695	1679	2663	3647	787CIP2B_348	8556
696	1680	2664	3648	787CIP2B_349	8561
697	1681	2665	3649	787CIP2B_350	8562
698	1682	2666	3650	787CIP2B_351	8569
699	1683	2667	3651	787CIP2B_352	8587
700	1684	2668	3652	787CIP2B_353	8597
701	1685	2669	3653	787CIP2B_354	8610
702	1686	2670	3654	787CIP2B_355	8610
703	1687	2671	3655	787CIP2B_356	8615
704	1688	2672	3656	787CIP2B_357	8622
705	1689	2673	3657	787CIP2B_358	8626
706	1690	2674	3658	787CIP2B_359	8628
707	1691	2675	3659	787CIP2B_360	8629
708	1692	2676	3660	787CIP2B_361	8630
709	1693	2677	3661	787CIP2B_362	8632
710	1694	2678	3662	787CIP2B_363	8634
711	1695	2679	3663	787CIP2B_364	8643
712	1696	2680	3664	787CIP2B_365	8644
713	1697	2681	3665	787CIP2B_366	8645

714	1698	2682	3666	787CIP2B_367	8646
715	1699	2683	3667	787CIP2B_368	8657
716	1700	2684	3668	787CIP2B_369	8661
717	1701	2685	3669	787CIP2B_370	8670
718	1702	2686	3670	787CIP2B_371	8692
719	1703	2687	3671	787CIP2B_372	8698
720	1704	2688	3672	787CIP2B_373	8762
721	1705	2689	3673	787CIP2B_374	8768
722	1706	2690	3674	787CIP2B_375	8768
723	1707	2691	3675	787CIP2B_376	8799
724	1708	2692	3676	787CIP2B_377	8806
725	1709	2693	3677	787CIP2B_378	8809
726	1710	2694	3678	787CIP2B_379	8814
727	1711	2695	3679	787CIP2B_380	8822
728	1712	2696	3680	787CIP2B_381	8833
729	1713	2697	3681	787CIP2B_382	8835
730	1714	2698	3682	787CIP2B_383	8877
731	1715	2699	3683	787CIP2B_384	8886
732	1716	2700	3684	787CIP2B_385	9003
733	1717	2701	3685	787CIP2B_386	9157
734	1718	2702	3686	787CIP2B_387	9175
735	1719	2703	3687	787CIP2B_388	9205
736	1720	2704	3688	787CIP2B_389	9260
737	1721	2705	3689	787CIP2B_390	9295
738	1722	2706	3690	787CIP2B_391	9307
739	1723	2707	3691	787CIP2B_392	9307
740	1724	2708	3692	787CIP2B_393	9312
741	1725	2709	3693	787CIP2B_394	9347
742	1726	2710	3694	787CIP2B_395	9370
743	1727	2711	3695	787CIP2B_396	9370
744	1728	2712	3696	787CIP2B_397	9382
745	1729	2713	3697	787CIP2B_398	9591
746	1730	2714	3698	787CIP2B_399	9650
747	1731	2715	3699	787CIP2B_400	9655
748	1732	2716	3700	787CIP2B_401	9663
749	1733	2717	3701	787CIP2B_402	9715
750	1734	2718	3702	787CIP2B_403	9735
751	1735	2719	3703	787CIP2B_404	9753
752	1736	2720	3704	787CIP2B_405	9771
753	1737	2721	3705	787CIP2B_406	9784
754	1738	2722	3706	787CIP2B_407	9925
755	1739	2723	3707	787CIP2B_408	9970
756	1740	2724	3708	787CIP2B_409	9997
757	1741	2725	3709	787CIP2B_410	10008
758	1742	2726	3710	787CIP2B_411	10010
759	1743	2727	3711	787CIP2B_412	10023
760	1744	2728	3712	787CIP2B_413	10043
761	1745	2729	3713	787CIP2B_414	10093
762	1746	2730	3714	787CIP2B_415	10172
763	1747	2731	3715	787CIP2B_416	10184
764	1748	2732	3716	787CIP2B_417	10205
765	1749	2733	3717	787CIP2B_418	10246
766	1750	2734	3718	787CIP2B_419	10298
767	1751	2735	3719	787CIP2C_1	886
768	1752	2736	3720	787CIP2C_2	1028
769	1753	2737	3721	787CIP2C_3	1916
770	1754	2738	3722	787CIP2C_4	2072
771	1755	2739	3723	787CIP2C_5	2424
772	1756	2740	3724	787CIP2C_6	2474
773	1757	2741	3725	787CIP2C_7	2474

774	1758	2742	3726	787CIP2C_8	2887
775	1759	2743	3727	787CIP2C_9	3001
776	1760	2744	3728	787CIP2C_10	3182
777	1761	2745	3729	787CIP2C_11	3182
778	1762	2746	3730	787CIP2C_12	3182
779	1763	2747	3731	787CIP2C_13	3193
780	1764	2748	3732	787CIP2C_14	3196
781	1765	2749	3733	787CIP2C_15	3224
782	1766	2750	3734	787CIP2C_16	3225
783	1767	2751	3735	787CIP2C_17	3234
784	1768	2752	3736	787CIP2C_18	3241
785	1769	2753	3737	787CIP2C_19	3243
786	1770	2754	3738	787CIP2C_20	3243
787	1771	2755	3739	787CIP2C_21	3259
788	1772	2756	3740	787CIP2C_22	3272
789	1773	2757	3741	787CIP2C_23	3278
790	1774	2758	3742	787CIP2C_24	3296
791	1775	2759	3743	787CIP2C_25	3327
792	1776	2760	3744	787CIP2C_26	3334
793	1777	2761	3745	787CIP2C_27	3339
794	1778	2762	3746	787CIP2C_28	3347
795	1779	2763	3747	787CIP2C_29	3387
796	1780	2764	3748	787CIP2C_30	3392
797	1781	2765	3749	787CIP2C_31	3411
798	1782	2766	3750	787CIP2C_32	3427
799	1783	2767	3751	787CIP2C_33	3432
800	1784	2768	3752	787CIP2C_34	3441
801	1785	2769	3753	787CIP2C_35	3479
802	1786	2770	3754	787CIP2C_36	3488
803	1787	2771	3755	787CIP2C_37	3488
804	1788	2772	3756	787CIP2C_38	3553
805	1789	2773	3757	787CIP2C_39	3560
806	1790	2774	3758	787CIP2C_40	3618
807	1791	2775	3759	787CIP2C_41	3642
808	1792	2776	3760	787CIP2C_42	3649
809	1793	2777	3761	787CIP2C_43	3676
810	1794	2778	3762	787CIP2C_44	3727
811	1795	2779	3763	787CIP2C_45	3911
812	1796	2780	3764	787CIP2C_46	4218
813	1797	2781	3765	787CIP2C_47	4219
814	1798	2782	3766	787CIP2C_48	4222
815	1799	2783	3767	787CIP2C_49	4222
816	1800	2784	3768	787CIP2C_50	4229
817	1801	2785	3769	787CIP2C_51	4230
818	1802	2786	3770	787CIP2C_52	4240
819	1803	2787	3771	787CIP2C_53	4241
820	1804	2788	3772	787CIP2C_54	4249
821	1805	2789	3773	787CIP2C_55	4252
822	1806	2790	3774	787CIP2C_56	4267
823	1807	2791	3775	787CIP2C_57	4272
824	1808	2792	3776	787CIP2C_58	4273
825	1809	2793	3777	787CIP2C_59	4275
826	1810	2794	3778	787CIP2C_60	4283
827	1811	2795	3779	787CIP2C_61	4290
828	1812	2796	3780	787CIP2C_62	4292
829	1813	2797	3781	787CIP2C_63	4305
830	1814	2798	3782	787CIP2C_64	4306
831	1815	2799	3783	787CIP2C_65	4308
832	1816	2800	3784	787CIP2C_66	4322
833	1817	2801	3785	787CIP2C_67	4351

834	1818	2802	3786	787CIP2C_68	4356
835	1819	2803	3787	787CIP2C_69	4399
836	1820	2804	3788	787CIP2C_70	4400
837	1821	2805	3789	787CIP2C_71	4520
838	1822	2806	3790	787CIP2C_72	4598
839	1823	2807	3791	787CIP2C_73	4599
840	1824	2808	3792	787CIP2C_74	4600
841	1825	2809	3793	787CIP2C_75	4670
842	1826	2810	3794	787CIP2C_76	4708
843	1827	2811	3795	787CIP2C_77	4734
844	1828	2812	3796	787CIP2C_78	4738
845	1829	2813	3797	787CIP2C_79	4749
846	1830	2814	3798	787CIP2C_80	4752
847	1831	2815	3799	787CIP2C_81	4752
848	1832	2816	3800	787CIP2C_82	4770
849	1833	2817	3801	787CIP2C_83	4784
850	1834	2818	3802	787CIP2C_84	4785
851	1835	2819	3803	787CIP2C_85	4792
852	1836	2820	3804	787CIP2C_86	4803
853	1837	2821	3805	787CIP2C_87	4811
854	1838	2822	3806	787CIP2C_88	4817
855	1839	2823	3807	787CIP2C_89	4818
856	1840	2824	3808	787CIP2C_90	4820
857	1841	2825	3809	787CIP2C_91	4831
858	1842	2826	3810	787CIP2C_92	4841
859	1843	2827	3811	787CIP2C_93	4869
860	1844	2828	3812	787CIP2C_94	4876
861	1845	2829	3813	787CIP2C_95	4902
862	1846	2830	3814	787CIP2C_96	4910
863	1847	2831	3815	787CIP2C_97	4931
864	1848	2832	3816	787CIP2C_98	5303
865	1849	2833	3817	787CIP2C_99	5317
866	1850	2834	3818	787CIP2C_100	5322
867	1851	2835	3819	787CIP2C_101	5330
868	1852	2836	3820	787CIP2C_102	5333
869	1853	2837	3821	787CIP2C_103	5333
870	1854	2838	3822	787CIP2C_104	5356
871	1855	2839	3823	787CIP2C_105	5363
872	1856	2840	3824	787CIP2C_106	5364
873	1857	2841	3825	787CIP2C_107	5379
874	1858	2842	3826	787CIP2C_108	5386
875	1859	2843	3827	787CIP2C_109	5397
876	1860	2844	3828	787CIP2C_110	5401
877	1861	2845	3829	787CIP2C_111	5419
878	1862	2846	3830	787CIP2C_112	5420
879	1863	2847	3831	787CIP2C_113	5452
880	1864	2848	3832	787CIP2C_114	5467
881	1865	2849	3833	787CIP2C_115	5482
882	1866	2850	3834	787CIP2C_116	5483
883	1867	2851	3835	787CIP2C_117	5492
884	1868	2852	3836	787CIP2C_118	5499
885	1869	2853	3837	787CIP2C_119	5525
886	1870	2854	3838	787CIP2C_120	5538
887	1871	2855	3839	787CIP2C_121	5539
888	1872	2856	3840	787CIP2C_122	5558
889	1873	2857	3841	787CIP2C_123	5559
890	1874	2858	3842	787CIP2C_124	5586
891	1875	2859	3843	787CIP2C_125	5619
892	1876	2860	3844	787CIP2C_126	5628
893	1877	2861	3845	787CIP2C_127	5640

894	1878	2862	3846	787CIP2C_128	5640
895	1879	2863	3847	787CIP2C_129	5827
896	1880	2864	3848	787CIP2C_130	6094
897	1881	2865	3849	787CIP2C_131	6195
898	1882	2866	3850	787CIP2C_132	6206
899	1883	2867	3851	787CIP2C_133	6355
900	1884	2868	3852	787CIP2C_134	6362
901	1885	2869	3853	787CIP2C_135	6386
902	1886	2870	3854	787CIP2C_136	6431
903	1887	2871	3855	787CIP2C_137	6457
904	1888	2872	3856	787CIP2C_138	6480
905	1889	2873	3857	787CIP2C_139	6497
906	1890	2874	3858	787CIP2C_140	6532
907	1891	2875	3859	787CIP2C_141	6598
908	1892	2876	3860	787CIP2C_142	6644
909	1893	2877	3861	787CIP2C_143	6644
910	1894	2878	3862	787CIP2C_144	6645
911	1895	2879	3863	787CIP2C_145	6645
912	1896	2880	3864	787CIP2C_146	6761
913	1897	2881	3865	787CIP2C_147	6782
914	1898	2882	3866	787CIP2C_148	6981
915	1899	2883	3867	787CIP2C_149	6981
916	1900	2884	3868	787CIP2C_150	7000
917	1901	2885	3869	787CIP2C_151	7029
918	1902	2886	3870	787CIP2C_152	7885
919	1903	2887	3871	787CIP2C_153	8143
920	1904	2888	3872	787CIP2C_154	8143
921	1905	2889	3873	787CIP2C_155	8234
922	1906	2890	3874	787CIP2C_156	8463
923	1907	2891	3875	787CIP2C_157	8467
924	1908	2892	3876	787CIP2C_158	8540
925	1909	2893	3877	787CIP2C_159	8600
926	1910	2894	3878	787CIP2C_160	9656
927	1911	2895	3879	787CIP2C_161	9669
928	1912	2896	3880	787CIP2C_162	9695
929	1913	2897	3881	787CIP2C_163	9744
930	1914	2898	3882	787CIP2C_164	9849
931	1915	2899	3883	787CIP2D_1	4180
932	1916	2900	3884	787CIP2D_2	4181
933	1917	2901	3885	787CIP2D_3	4314
934	1918	2902	3886	787CIP2D_4	4500
935	1919	2903	3887	787CIP2D_5	5651
936	1920	2904	3888	787CIP2D_6	5691
937	1921	2905	3889	787CIP2D_7	5881
938	1922	2906	3890	787CIP2D_8	5882
939	1923	2907	3891	787CIP2D_9	6209
940	1924	2908	3892	787CIP2D_10	6719
941	1925	2909	3893	787CIP2D_11	8130
942	1926	2910	3894	787CIP2D_12	8863
943	1927	2911	3895	787CIP2D_13	8902
944	1928	2912	3896	787CIP2D_14	9162
945	1929	2913	3897	787CIP2D_15	9197
946	1930	2914	3898	787CIP2D_16	9215
947	1931	2915	3899	787CIP2D_17	9232
948	1932	2916	3900	787CIP2D_18	9262
949	1933	2917	3901	787CIP2D_19	9369
950	1934	2918	3902	787CIP2D_20	9371
951	1935	2919	3903	787CIP2D_21	9516
952	1936	2920	3904	787CIP2D_22	9601
953	1937	2921	3905	787CIP2D_23	9731

954	1938	2922	3906	787CIP2D_24	9733
955	1939	2923	3907	787CIP2D_25	9769
956	1940	2924	3908	787CIP2D_26	9804
957	1941	2925	3909	787CIP2D_27	9816
958	1942	2926	3910	787CIP2D_28	9844
959	1943	2927	3911	787CIP2D_29	9924
960	1944	2928	3912	787CIP2D_30	9936
961	1945	2929	3913	787CIP2D_31	10163
962	1946	2930	3914	787CIP2D_32	10165
963	1947	2931	3915	787CIP2D_33	10165
964	1948	2932	3916	787CIP2D_34	10244
965	1949	2933	3917	787CIP2D_35	10278
966	1950	2934	3918	787CIP2E_1	4251
967	1951	2935	3919	787CIP2E_2	5310
968	1952	2936	3920	787CIP2E_3	5697
969	1953	2937	3921	787CIP2E_4	5731
970	1954	2938	3922	787CIP2E_5	5733
971	1955	2939	3923	787CIP2E_6	5734
972	1956	2940	3924	787CIP2E_7	5740
973	1957	2941	3925	787CIP2E_8	7657
974	1958	2942	3926	787CIP2E_9	9572
975	1959	2943	3927	787CIP2F_1	1363
976	1960	2944	3928	787CIP2F_2	4303
977	1961	2945	3929	787CIP2F_3	5760
978	1962	2946	3930	787CIP2F_4	5766
979	1963	2947	3931	787CIP2F_5	5767
980	1964	2948	3932	787CIP2F_6	5767
981	1965	2949	3933	787CIP2F_7	5770
982	1966	2950	3934	787CIP2F_8	6855
983	1967	2951	3935	787CIP2F_9	10026
984	1968	2952	3936	787CIP2F_10	10227

TABLE 6

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
2953	A	3	324	ISEHRIEASGNYLAQRLTSSFLRGLSSWKSNNPLMLCGWTILLTLTMVQGE*GPKGIPGFHTNSSYPHWGTVAKPAGD*DLLPAPGQEGTPLFTR*SLCTYCPID
2954	A	18	467	REELGKDLFDCTLYVLLKYDDFNADKHLALEEFYRAFQVIQLSLPEDQKLSITAATVGQSAVLSCAIQGTLRPPIWKRNNIILNNLDLEDINDFGDDGSLYITKVTTHVGNITCYADGYBQVYQTHIFQVNVPPVIRVYPESQARRAG
2955	A	3	23	FYSAFLVADKGIVTSKHNNDTQHIWESDSNEFSVIADPRGNTLGRGTTIT*VSIPPSL
2956	A	1	493	RTKTDVYILNLAVADLLLLFTLPFWAVNAVHGWVLGKIMCKITSALYTLNLFVSGMQFLACISIDRYVAVTKVPSQSGVGKPCWICFCVWMAAILLSIPQLVFYTVNDNARCIPFPRYLGTSMKALIQMLEICIGFVVPFLIMGVCYFITARTLMKMPNIKIS
2957	A	703	302	EETGVREKRRERMKEKMWQNVLCCCTLQTAVILKLFQNKVLNLIKNNFLSPDTRKNKVFKKWAGGPGAVAHACNPSTLGGRGGRITKSGDRDHPGQHG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ETRSLPACWAQWKSALPVS RAPGRQGS LVFP LP
2958	A	575	1054	CTKCKADCDTCFNKNFCTKCKSGFYHLHGKCLD NCPEGLEANNHTMECVSIHVCEVSEWNPWSPCT KKGKTCGFKRGTTETRVREIIQHPSAKGNLCPPTN ETRKCTVQRKKCKQGERGKKGRERKRKKPNKG ESKEAIPDSKSLESSKEIPEQRENKQQQ
2959	A	1	426	LSMLSTISTEHLRLSVLWPIWYCCCHCPHLSAVMC VLLWALSLLQSILEWMFCSFLSDVSDSNWCQIL DFLTAVWLIFLNLVLCGFTLVLLVRIICGSQKMPL TRLVVTILLTGLVFLFCSLPLSIQ*FLLYWIEKDLD DL
2960	A	1194	852	EKRKTSYSQCLNSKQRNVSMRPSIWIHVHLKPPC RLVELLPFSSALQGLSHLSLGTTLV/*GHLRFRL RNL PQSLRTVILPERNEEQNLQELSHNADKYQM GDCCKEEIDDSIFY
2961	A	274	2250	EKGKVKDAGAEQWISLSLCKGSWETQFSNHLN SLTPPTSVRMPLITTVTLKMMVARHHMKLLCSK AFSTQLQQKIFLHSQMGIHQSVCMKLKPNPTSHII SILMGQPMALVQLETLAPLTIIQKFQTDHMKF WKNLPLHSHHLTPSPQTVIPKKTGSPEIKLKITK TIQNGRELFESSLCGDLLNEVQASE/*NQSIESRK EKRRKSNKHDSSRSEERKSHKIPKLEPEEQNRPN ERVDTVSEKPREEPVLKEGSPSSANTIFCSNNGSV HWFKFQVGDVWSKVGTYPWWPCMVSSDPQL EVHTKINTRGAREYHVQFFSNQPERAWVHEKRV REYKGHKQYEELLA EATKQASNHSEKQKIRKPR PQRERAQWDIGIAHA EKALKMTREBRIEQYTFIYI DKQPEEALSQAKKSVASKTEVKKTRRPRSVLNT QPEQTNAGEVASSLSSTEIRRHSSQRRHTSABEEEP PPVKIAWKTAAR*EASTTM*KGSL* MSPV*EQVFALQ*ATGDGKFIDQ* NKTEISVRGQDRLLISTPNQRNEKPTQSV*PEATS GSTGSVEKKQRRSIRTRSESEKSTEVVPAK* KEQVETVPQATVKTGLQKGSADRGVQGSVRFS* SSVSAAIEETVD
2962	A	2408	836	SASPPPPPPPPSRFPFSGAPGARDRSGPLGSEPQR NPGARPTLEATVTPPGSVGAMSSSGLNSEKVA ALIQKLNSDPQFVLAQNVGTHDLLDICLKRA TV QRAQHVFQHAVPQEGKPTITNQKSSGRCWIFSLN VMRLPFMKKLNIEEFESQSYLFFWDKVERCYFF LSAFVDTAQRKEPEDGRLVQFLMNPANDGGQ WDMLVNIVEKYGVIPKKCFESYTTAATRRMND ILNHKMREFCIRLRLNVHSGATKGEISATQDVM MEEIFRVVCICLGNPPETFTWEYRDKDKNNKKIG PITPLEFNRE/QHVKPLFNMEDKICLVNDPRPQH KYNKLYTVAYL*SNMVWRGEKLFYNNQPIDFLK KMVAASIKDGEAVWFGCDVGKHFNSKLGALSD MNLYDHLEVFGVSLKNMNAERLTFGESLMT HTMTFTA V/SQSRDDSGMVLFTKWRVGEFQWG EDHGHKG YLCMTD*VGSLEYVVEVV/VWDRKH VPAEVLAVLGAGNPFVLPWDFPMGALAE
2963	A	90	543	RHYDSAGKITLKIKNYLEQRAVGGASPRLAQS VLTCREPILENSLTSLIEYLHNALEHDMRLRFNN DRMKTITKETST*LSNSYL VFPLM*SLTYLMKMS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				FERCTARNKMFVNSPFTKVDNYCTSS\WKKFYL KCYFSLNTIKKEKKMT
2964	A	3	2454	FDTYRGLPSISNGNYSQLOFQAREYSGAPYSQRIS AITTVSVAWKVLSGKIGEGAEGNCKCVISEGAW AVCPTQPCGKAKPDKHLKDLLSKLLNSGYFESIP VPKNAKEKEVPLEEEMLIQSEKKTQLSKTESVKE SESLMEFAQPEIQPEFLNRRYMTEVDYSNKQGE EQPWEADYARKPNLPKRWDMLTEPDGQEKKQE SFKSWEASGKHQEVSKPAVSLEQRKQDTSKLRS TLPEEQKKQEISKSKSPSQWKQDTPKSKAGYVQ EEHKKQETPKLWPVQLQKEQDPKKQTPKSWTPS MQSEQNTTKSWTTPMCEEQDSKQPETPKSWENN VESQKHSLTSQSQISPKSWG VATASLIPNDQLLPR KLNTEPKDVP/IACASA*GFLPLQPPFRRI/HVLRK EKLQDLMTQIQGTCNFMQESVLD FDKPSSAIPTS QPPSATPG*PRRHLKEQNLS\VKVIFQGA VTIVF NVNAPLPPRKEQEIKESPYSPGYNQSFTTASTQTP PQCQLPSIHVEQTVHSQETANYHPDGTIQVSNGS LAFYPAQTNVFRPTQPFVNSRGSVRGCTRGGR ITNSYRSPGGYKGFDTYRGLPSISNGNYSQLOFQ AREYSGAPYSQRDNFQCCYKRGGTSGGPRANSR AGWSDSSQVSSPERDNETFNSGDSGQGDSRSMT PVDVPVTNPAATILPVHVYPLPQQMRVAFSAAR TSNLAPGTLDPQIVFDLLNNLGETFDLQLGRFN CPVNGTYVFIFHMLKLA VNVPLYVNL MKNEEVL VSA YANDGAPDHETASNHAILQLFQGDQIWLRL HRGAIYGSSW
2965	A	3	2454	FDTYRGLPSISNGNYSQLOFQAREYSGAPYSQRIS AITTVSVAWKVLSGKIGEGAEGNCKCVISEGAW AVCPTQPCGKAKPDKHLKDLLSKLLNSGYFESIP VP*NAKPKKEVPLEEEMLIQSEKKTQLSKTESVKE SESLMEFAQPEIQPEFLNRRYMTEVDYSNKQGE EQPWEADYARKPNLPKRWDMLTEPDGQEKKQE SFKSWEASGKHQEVSKPAVSLEQRKQDTSKLRS TLPEEQKKQEISKSKSPSQWKQDTPKSKAGYVQ EEHKKQETPKLWPVQLQKEQDPKKQTPKSWTPS MQSEQNTTKSWTTPMCEEQDSKQPETPKSWENN VESQKHSLTSQSQISPKSWG VATASLIPNDQLLPR KLNTEPKDVP/IACASA*GFLPLQPPFRRI/HVLRK EKLQDLMTQIQGTCNFMQESVLD FDKPSSAIPTS QPPSATPG*PRRHLKEQNLS\VKVIFQGA VTIVF NVNAPLPPRKEQEIKESPYSPGYNQSFTTASTQTP PQCQLPSIHVEQTVHSQETANYHPDGTIQVSNGS LAFYPAQTNVFRPTQPFVNSRGSVRGCTRGGR ITNSYRSPGGYKGFDTYRGLPSISNGNYSQLOFQ AREYSGAPYSQRDNFQCCYKRGGTSGGPRANSR AGWSDSSQVSSPERDNETFNSGDSGQGDSRSMT PVDVPVTNPAATILPVHVYPLPQQMRVAFSAAR TSNLAPGTLDPQIVFDLLNNLGETFDLQLGRFN CPVNGTYVFIFHMLKLA VNVPLYVNL MKNEEVL VSA YANDGAPDHETASNHAILQLFQGDQIWLRL HRGAIYGSSW
2966	A	1693	227	DYVLTAE LHRQSPGVSFGLSVFNL MNAIMGSGI LGLAYVMANTGVFGFSFLLLT VALLASYSVHLL LSMCIQTAYLGP*TN YFMVLP AH*LTCLPLIEFLQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				SL*NSL*AVTSYEDLGLFAFGLPGKLVVAGTHIQ NIGAMSSYLLIKTELPAIAEFLTGDYSRYWYLD GQTLIIICVGIVFPLALLPKIGFLGYTSSLFFFM MFFALVVIKKWSIPCPLTLNIVEKGQISNVTDD CKPKLFHFSKESAYALPTMAFSFLCHTSILPIYCE LQSPSKKRMQNVNTNTAIALSFLIYFISALFGYLT YD/GTTKAQRGEVTCHRIKDKVESELLKG***IP* SHDVVVMNVKLCILFAVLLATVPLIHFPARKAVT MMFFSNFPFSWIRHFLITLALNITVLLAIYVPDIRN VFGVVGASTSTCLIFIFPGLFYKLKSREDFLSWKK LGVGCFC/LLSFKTSILRNSLSVYIILPASRKSIFYK I
2967	A	3	3222	SGIVVRALWREKKPGGGRRVKKRNPNRQAVGH TEEDPPRVGTPWKEHTGPGPQEGSTMEAAHAKT TEECLEYGVSETTGLTPDQVKRNLEKYGLNELP AEEGKTLWELVIEQFEDLLVRILLAAACISFVLA WFEEGEETITAFVEPFVILLILIANAIVGVWQERN AENAIEALKKEYPEMCKVYRADRSVQRIKARD IVPGDIVEVAVGDKVPADIRILAIKSTTLRVQDQIL TGEYVSVIKHTEPVDPRAVNQDKKNMFLSGTNI AAGKALGIVATTGVGTEIGKIRDQMAATEQDKT PLQQKLDEFGEQLSKVISLICVAVWLNIGHFNDF VHGGSWFRGAIYYFKIAVALAVAAIPEGLPAVIT TCLALGTRRMAKKNIVRSLPSVETLGCSTVICS DKTGTLTTNQMSVCCKMFIDKVDGDIKLLNEFSIT GSTYAPEGEVLKNDKPVPRGQYDGLVELATICA LCNDSSLDFNEAKGVYEVGEATETALTTLVEK MNVFNTDVRSLSKVERANACNSVIRQLMKKEFT LEFSRDRKSMSVYCSPAKSSRAAVGNKMFVKGA PEGVIDRCNYVRVGTTTRVPLTGPVKEKIMAVIKE WGTIPETLRCLALATRI:TPKPFEMVLDDSFARF LEYETDLTFVGVGMLDPFRKEVTGSLURDA GIRVIMITGDNKGTAIAICRRIGIFGENEEVADRA YTGREFDDLPLABQREACRRACCFARVEPSHK SEIVEYLQSYDEITAMTGDGVNDAPALKKAEIGI AMGSGTAVAKTASEMVLADDNFSTTVAAVEEGR AIYNNMKQFIRYLISNVGEVVCIFLTAALGLPEA LIPVQLLWVNLVTDGLPATALGFNPPDLDIMDRP PRSPKEPLASGWLFFRYMAIGGYVGAATVGAAA WWFLYAEDGPHVNYSQLTTHFMQCTEDNTHFEGI DCEVFEAPEPMTMALSVLVTIEMCNALNSLSEN QSLLRMPWVNIWLLGSICLSMSLHFLILYVDPLP MIFKLRLDLTQWLMVLKISLPVIGLDEILKFVA RNYLEG*LFPLLHL*ARVTDPEDERRK
2968	A	3	2414	GARSCSRLGRCTFPLWKGREMEVRKLSISWQFLI VLVLILQILSALDFDPYRVLGVSRTASQADIKKA YKKLAREWHPDKNDKPGAEDKFIQISKAYEILSN EEKRSNYDQYGDAGENQGYKQKQQQREYRFRH FHENFYFDESFFHFPNSERRDSIDEKYLLHFSHY VNEVAPDSFKKPYLIKITSWCFSCHIEPVWKEV IQELEELGVGIGVVHAGYERRLAHHLGAHSTPSI LGIINGKISFFHNAVRENLRQFVESLLPGNLVEK VTNKNYVRFLSGWQENKPHVLLFDQTPIVPLL YKLTAFAYKDYLSFGYVYVGLRGTEEMTRRYNI NIYAPTLVFKEHINRPADVIQARGMKKQIIDDFI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TRNKYLLAARLTSQKLFHELCPVKRSHRQRKYC VLLTAETTKLSKPFEAFLSFALANTQDTRFVH VYSNRQEFADTLLPDSEAFQKSAVSILERRNT AGRVVYKTLEDPWIGSESDKFILLGYLDQLRKDP ALLSSEAVLPDLTDELAPVFLLRWFYSASDYISD CWDSIFHNNWREMMPLLSLIFSALFILFGTVIVQ AFSDSNDERESSPPEKEEAQEKTKGTEPSFTKENS SKIPKKGFEVTELTDTVYTSNLVRLRPGHMNV VLILSNSTKTSLLQKFALEVYFTTGSSCLHFSFLSL DKHREWLEYLLEFAQDAAPIPNQYDKHFMERDY TGYVLALNGHKYFCLFKPQKTVEEGGKP*GSC SDVDSSLYLGESRGKPSGLGSRPIKGLSKLSL WMERLLEGLQRFYIPSWPELD
2969	A	48	1117	KGLSPDQVLSAFAPLDCMWLVKFTTFLSFATG ACSLKVTVPSTVHGVGRQALYLPVHYGFHTP ASDIQIWLFRPHTMPKYLLGSVNKSVVPD/YGI P/YTSSP*CHPMASLLINPLQFPDEGNYIVKVNIG NGTILSASQKIQVTVDPTKPVVQIHPPSGAVEY VGNMILTCHVEGGTRLAYQWLKNGRPVHTSST YSFSPQNNLTHIAPVTKEDIGNYSCLVRNPVSEM ESDIIMPITYGPGYGLQVNSDKGLKVGCVFTVDL GEAILFDCSADSHPNYTSWIRRTDNTTYIKHGP RLEVASEKVAQKTM DYVCCAYNNITGRQDETHF TVITSVGMCDIQGRDPNKT
2970	A	68	936	HSALLTHSSFCVFTLCQDFFTYSSMSEEVYADL QFQNSSEMEKIPEIGKFGEKAPPAPSHVWRPAAL FLTLLCLLLLIGLGVLASMFHVTLKIEMKKMNKL QNISEELQRNISLQLMSNMNISKIRNLSTTLQTI ATKLCRELYSKEQEHKCKPCRRWIWHKDCYF LSDDVQVTWQESKMACAAONASLLKINNKNALF FISQSRSDYWLGLSPEEENSWVEG*YNO SAWVIRNAPDEYCGTNRLYVQYYHCT QRMICEKMANFVQLGSTYFREA
2971	A	912	2287	VPNYLPSVSSAIGGEVPQRYVWRFICGLHSAPRI LVAFAYWNHYLSCTSPCSCYRPLCRLNFGNLV ENLALLVLTYSSEDF/TWVPG*GRSGEVFEGT GLPLPHSDLPTSWCGHSLQCGSQSSFPPIHENAF IVFIASSLGHMLLTCLWRLTKKHTVSQEDGLSL AGAPRQPRRKSRTSVLRIRVMVRWELSSNGNPG RGVLGLGLGNKL RVVGQNLGL*HCVWVWVE TGE*KRWRLQMGIE*GVASRRQ*VRNSVRGLVC HNSSAPPMYMGFFSPTVFGGVGG*LVHTFILHP PEVEAAGIPLLLGPSLPQRQGREHIVILAAPACA PFHDR*WEPREIRPSP*ELGLRGEPTLSYPASCRVI RQPIP*DRKSYSWKQRLFIINFISFSAVYFRHN MYCEAGVYTIFAILEYTVVLTNMAFHMTAWWD FGNKELLITSQPEEKRF
2972	A	1734	246	GGILSGRDGRTALPRPREPAERTAGLRDRMRPQE LPRLAFPLLLLLLLLLPPPCPAHSATRFDPWES LDARQLPAWFDQAKFGIFHWGVFSVPSFGSEWF WWYWQKEKIPKYVEFMKDNYPSPFKYEDFGPL FTAKFFNANQ\WADIFQASGAKYIVLTSKHHEGF TLWG\SEYSWNWNAIDEGPKRDIVKELEVAINR TDLRFGLYYSLEWFHPLFLEDESSSFHKRQFPVS KTLPELYELVNNYQPEVLWSDGDGGAPDQYWN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				STGFLAWLYNESPVRGTVVTNDRWGAGSICKHG GFYTCSDRYNPGHLLPHKWENCMIDKLSWGY RREAGISDYLTIIEELVKQLVETVSCGGNLLMNIG PTLDGTISVVFEERLRQMGSWLKVNGEAIYETHT WRSQNDTVTPDVWYTSKPKEKLVYAIFLKWPTS GQLFLGHPKAILGATEVKLLGHGQPLNWISLEQN GIMVELPQLTIHQMPCKWGWALALTNVI
2973	A	24	1133	SVPRAGGDMETGAAELYDQALLGILQHVGNVQ DFLRVLFGLYRKTDYFRLLRHPSDRMGFPFGAA QALVLQVFKTFDHMARQDDEKRRQEELEEKIRRK EBEEAKTVSAAAAEKEPVPVPVQEIIEIDSTTELDG HQVEVKVQPPGPVKEMAHGSQEAEPGAVAGA AEVPR\EPILPRIQEQQFQKNPDSYNGAVRENYTW SQDYTDLEVRVPVKHVVKGKQVSVALSSSSIRV AMLENGERVLMEGKLTHTKINTESSLWSLEPGK CVLVNLSKVGEYWWNAILEGEEPIDDKINKERS MATVDEEEQAVLDRLTFDYHQKLQGKPKQSHEL KVHEMLKKGWDAEGSPFRGQRFDPAMFNISPGA VQF
2974	A	271	1854	MQFGRAHGDCVSGAQLCGCPSMDDYMLVLRMIG EGSFGRALLVQHESNQMFMKEIRLPKSFSNTQ NSRKEAVLLAKMKHPNIVAFKESFEAEGHLYTV MEYCDGGDLMOQIKQKQKGLFPEDMILNWFTQ MCLGVNHHKRRVLHRDIKSKNIFLTQNGKGKL GDFGSARLLSNPMAFACTYVGTPTYVPPEIWEN LPYNNKSDIWSLGCILYELCTLKHPFQANSWKNL ILKVCQGCISPLPSHYSYELQFLVKQMFKRNP RPSATTLRSRGIVARLVQKCLPPEIMEYGEEVLE EIKNSKHNTPRKKTNP SRIRIALGNEASTVQEEEQ DRKGSHTDLESINENLVESALRRVNRBKGKNSV HLRKASSPNIHPRQWENVPNTAILENA3LT SSLTAEDMAGCCYKYSKNITRQWLKETPDILL NILKNADLSLQFYTYTYRPGS\EGFLKGPLSEETE ASDSVDGGHDSVLDPERLEPGLDEEDTDFEED DNDPWWSELKKRAGWQGLCDR
2975	A	32	2833	PPGEPGAGRGALSPCGPLSGPPPLPGREAGGTG QPVNPVFDLSRRNPQEDFELIQRIGSGTYGDVYK ARNVNTGELAAIKVIKLEPGEDFAVVQEIIMMK DCKHPDIVAYFGSYLARRDKLWICMEF\CGSGS \QDIYHVTGPLSELQIAYVSRETLOGLYYLHSGK KMRDIKGANILLTDNGHVKLADFGVSAQITATI AKRKSFIGTPYWMAPVAAVERKGGYNQLCDL WAVGITAIELAELOPPMFDLHPMRALFLMTKSNF QPPKLKDKMKWSNSFHHFVKMALTKNPKKRPT AEKLLQHPFVTQHLTRSLAIELLDKVNNDHSTY HDFDDDDPEPLVAVPHRIHSTSRNVREEKTRSEIT FGQVKFDPPLRKETEPHHELPSDGF\LDSSSEIYY TARSNLDLQLEYGQGHQGYFLGANKSLLKSV EELHQRGHVAHLEDEGDDDESKHSTLKAKIP PPLPPKPKSIFIPQEMHSTEDENQGTIKRCPMSGSP \AKPSQVPPRPPPPRLPPHKPVALGNGMSSFQLNG ERDGS\LCQQQNEHRGENLSRKEKKDVPKPSNG LPPTPKVHMGACFSKVFNGCPLKIH\CASSWINPD TRDQYLIFGAEEGIYTLNLNELHETSMEQLFPRR CTWLYVMNCLLSISGKASQLYSHNLPGLFDYA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RQMOKLPVAIPAHLKLPDRILPRKFSVSAKIPETK WCQKCCVVRNPYTGHKYLCGALQTSIVLLEWV EPMQKFMLIKHIDFPIPCPLKMFEMLVVPEQEYP LVCVGVSRRDFNQVVRFETVNPNTSSWFTES DTPQTNVTHVTQLERDTILVCLDCCIKIVNLQGR LKSSRKLSSSELTDFRIESIVCLQDSVLAFWKHG MQGRSFRSNEVTQEISDSTRIFRLGSDRVVVLES RPTDNPTANSNLYLAGHENS
2976	A	32	2833	PPGEPGAGRGALSPCGPLSGPPPLPGREAGGTG QPVNPVFDLSRRNPQEDFELIQRIGSGTYGDVYK ARNVNTGELAAIKVIKLEPGEDFAVVQEEIMMK DCKHPDIVAYFGSYLRRDKLWTCMEFCGSGS LQDIYHVTGPLSELQIAYVSRETLQGLYYLHSGK KMHRIKGANILLTDNGHVKLADFGVSAQITATI AKRKSFIGTPYWMAPEVAAVERKGGYNQLCDL WAVGITAIELAELOPPMFDLHPMRALFLMTKSNF QPPKLKDKMKWSNSFHHFVKMALTKNPKKRPT AEKLLQHPFVTQHLTRSLAIELLDKVNPNPDHSTY HDFDDDDPEPLVAVPHRIHSTSRNVREEKTRSEIT FGQVKFDPPLRKETEPHHELPSDGLDSSEIYY TARNLDLQLEYGQGHQGYFLGANKSLLKSV EELHQRGHVAHLEDEGDDDESKHSTLKAKIP PPLPPKPKSIFIPQEMHSTEDENQGTIKRCPMSGSP VAKPSQVPPRPPPPRLPPHKPVALGNGMSSFQNLG ERDGLCQQQNEHNGENLSRKEKKDVPKPISSNG LPPTPKVHMGACFSKVFNGCPLKIHCASSWINPD TRDQYLIFGAEEGIYTLNLNELHETSMEQLFPRR CTWLYVMNNCLLSISGKASQLYSHNLPGLFDYA RQMOKLPVAIPAHLKLPDRILPRKFSVSAKIPETK WCQKCCVVRNPYTGHKYLCGALQTSIVLLEWV EPMQKFMLIKHIDFPIPCPLKMFEMLVVPEQEYP LVCVGVSRRDFNQVVRFETVNPNTSSWFTES DTPQTNVTHVTQLERDTILVCLDCCIKIVNLQGR LKSSRKLSSSELTDFRIESIVCLQDSVLAFWKHG MQGRSFRSNEVTQEISDSTRIFRLGSDRVVVLES RPTDNPTANSNLYLAGHENS
2977	A	174	1543	YSLRKGITFKLAGAMVHIKKGELTQEEKELLEVI GKGTVQEAGTLLSSKNVRVNCLENGMTPLMH AAYKGKLDCKLLLRHGDVNCHEHGYTA LMFAALSGNKDITWVMLEAGAETDVVNSVGR AAQMAAFVGQHDCTIINNFFPRERLDYYTKPQ GLDKEPKLPPKLAGPLHKIITTNLHPVKIVMLV NENPLLTEEAALNKCVRVMDLICEKCMKQRDM NEVLAMKMHYISCFQKCNFLKDGENKLDTLIK SLLKGRASDGFVYPEKILRESIRKFPYCEATLL QQLVRSIAPVEIGSDPTAFSVLTQAITGQVGFVDV EFCTTCGEKGASKRCSVCKMVIYCDQTCQKTHW FTHKKICKNLKDIYEKQLEAAKEKROEENHGK LDVNSNCVNEEQPEAEVGISQKDSNPEDSGEGK KESLESEAELEGLQDAPAGPQVSEE
2978	A	3	5177	SDDLRTGLFQDVQDAESLKLPGVYEVLFYNETE DCPGMMLWRYPEPRGLTLVRITPVFNTTEDPDI STADLGDVLQDPCSLEYWDELQKVFAFRETNL SESKVCELQLPDINLVNDQKKLVSSDLWRIVLNS SQNGADDQSSASESGSQSTCDPLVTPTALAACTR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				<p> VDSCTPWFPVPSLCVSFQFAHLEFHLCHHLDQLG TAAPQYLQPFVSDRNPSELEYMIVSFREPHMYL RQWNNGSVQCIEIQLAQADCKLLECRNVTMQS VVKPFSIFGQMAVSSDVVEKLLDCTVIVDSVFN LGQHVVHSLNTAIAQAWQONKCPEVEELVFSHFV ICNDTQETLRFQVQDVTDENILLASLHSHQYSWRS HKSPQLLHICIEGWGNWRWSEFSDVHAGTFIRT IQYRGRTASLIKVQQLNGVQKQIICGRQIICSYL SQSIELKVQHYIGQDQGA VVREHFDCLTAKQK LPSYILENNELTELCVKAKGDEDWSRDVCLESK APEYSIVIQVPSSNSSIYVWCTVLTLEPNSQVQQ RMIVFSPLFIMRSHLPDPIIHLEKRSGLSETQIIP GKGEKPLQNIPEDLVHHLTFQAREEYDPSDCA VPISTSLIKQIATKVHPGGTVNQILDEFYGPESKL QPIWYPYNKSDSDRNEQLSQWDSMPRVKLSIWKP YVRTLLIELLPWALLINESKWDLWLFEGEKIVLQ VPAGKIIPPNFQEAQIGIYWANTNTVHKSVAIK LVHNL TSPKWKDGGNGEVVTLDEEAFVDTEIRL GAFFGHQKLCQFCISSMVQQGIQIIEQDKTTINN TPYQIFYKPLSVCNPHSGKEYFRVPDSATFSICP GGEQAMKSSSLPCWDLMPDISQSVLDASLLQK' QIMLGFSAPGADSSQCWSLPAIVRPEFPRQSV VPLGNFRENGFCTRAIVLTYQEH LGVTYLTSED PSPRVIIHNRCPVKMLIKENIKDIPKFEVYCKKIPS ECSIHHEL YHQISSYPDCKTKDLLPSLLLRVEPLD EVTTEWSDAIDINSQGTQVVFLTGFQYVYVDVV HQCGTVFITVAPEGKAGPILTNTNRAPEKIVTF/K MFITQLSLAVFDDLTHHKASAELLRLTDNIFLC VAPGAGPLPGEEPVAALFELYCVEICCGDLQLDN QLYNKS NFHFAVLVCQGEKAEPICSKMQSLLIS NFELEBYKKEKCFIKLC:TLNFGKSLCDINEFS'FL KPARLYNFTYIKTLFDTYLPNSRLAGHSTH LSGGKQVLMQVTQHARALVNPVKLRKLVIQPV NLLVSIHASLKIYASDHTPLSFSVFERGPIFTAR QLVHALAMHYAAGALFRAGWVVGSLDILGSPA SLVRSIGNGVADFFRLPYEGLTRGPGAFVSGVSR GTTSFVKHISGTLTSITNLATSLARNMDRLSLDE EHYNRQEEWRRQLPESLGEGLRQGLSRLGISLLG AIAGIVDQPMQNFQKTSEAQASAGHKAKGVISG VGKGIMGVFTKPIGGAAELVSQTGYGILHGAGLS QLPKQRHQPSDVHADQAPNSHVKYVWKMLQS LGRPEVHMALDVVLVRGSGQEHEGCLLTSEVL FVVSVSEDTQQQAFPVTEIDCAQDSKQNNLLTV QLKQPRVACDVEVDGVRERLSEQQYNRLVDYIT KTSCHLAPSCSSMQIPCVVAAEPPSTVKTYHY LVDPHFAQVFLSKFTMVKNKALRKGF </p>
2979	A	255	2673	<p> AWLFPASVLCPRCLTGS AVGSAEWKSLVVLPF SRPTLGHLD SKPSSKSNMIRGRNSATSADQPHIG NYRLKLTIGKGNFAVKLARHILTGKEVAVKID KTQLNSSSLQKLFREVRIMKVLNHPNIVKLFEB TEKTLYL VMEYASGGEVFDYLVAHGRMKEKEA RAKFRQIVSAVQYCHQKFIVHRDLKAENLLDA DMNIAIDFGFSNEFTFGNKLDTF CGSPPYAAPEL FQGKKYDGPEDVWSLGVILYTLVSGSLPFDGQ NLKELRERVLRGKYRIPFYMSTDCENLLKKFLIL </p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * =Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				NPSKRGTLEQIMKDRWMNVGHEDELKPYGEP LPDYKDPRTTELMVSMGYTREEIQDSL VGQRYN EVMATYLLGYKSSELEGDITTLKPRPSADLTNS SAPSPSHKVQSVSANPKQRRFSDQAGPAIPTSNS YSKKTQSNNAENKRPEEDRESGRKASSTAKVPA SPLPGLERKKTTPTSTNSVLSTSTNRSRNSPLLAE RASL\GQGFHPEWAKTALTMPGSRASASASAA VSAARPRQHQKMSASVHPNKASGLPPTESNCE VPRFRQVCWGCTAPQRVPVAPSAHNISSSGGA PDRTNFPRGVSSRSTFHAGQLRQVRDQQLPYG VTPASPSGHSQGRRGASGSIFSFTSKFVRRNLNE PESKDRIVETLRPHVVNSGGNDKEEFREAKPR SLRFTWSMKTTSMEPNEMMREIRKVLNANSCQ SELHEKYMLLCMHGTGPHGDFVQWEMEVCKLP RLSLNGVRFKRISGTSMFAKNIAKIANELKL
2980	A	120	3433	NCLLLQAKGFHGEIEDLQQWLTDTTERHLLASKP LGGLPETAKEQLNVHMEVCAAFEAKEETYKSLM QKGQQLARCPKSAETNIDQDINNLEK WESVE TKLNER\KT\KLEALNLA\MEFHNSL\QDFINWLT QAEQTLNVASRPSLILDTVLFQIDEHKVFANEVN SHREQUIELDKTGTHLKYPFSQKQDVVLKNNLISV QSRWEKVVQRLVERGRSLDDARKRAKQFHEAW SKLMEWLEESEKSLDSELEIANDPDKIQTQLAQH KEFQKSLGAKHSVYDTTNRTRGRSLKEKTSLADD NLKLDMLSEL RDKWDTICGKSVERQNKLEEA\ LLFSGQFTDALQALIDWL YRVEPQLAEDQPVHG DIDLVMNLIDNHKAFQKELGKRTSSVQALKRSA RELIEGSRDDSSWVKVQMQLSTRWETVCALSIS KQTRLEAALRQAEFFHSVVHALLLEWLAEAEQTL RFHGVLPPDEDALRTLIDQHKEFMKKLEBKRAE ENKATTMGDT\LAICH\PSITTIKHWIT\... VLA WAKQHQQRLASALAG...QELLEALLAW LQWAETITLTDKDKVIPQEIIEVKALIAEHQTFM EEMTRKQPDVDKVTKYKRAADPSSLQSHIPV LDKGRAGRKRFPASSLYPSGSQTQIETKNPRVN LVSKWQQVWLLALERRRKLNDALDRLEELREF ANFDFDIWRKKYMRWMNHKKSVMDFFRRIDK DQDGKITRQEFIDGILSSKFPTSRLMSAVADIFD RDGDGYIDYYEFAALHPNKDAYKPIITDADKIE DEVTRQVAKCKAKRFQVEQIGDNKYRFFLGNQ FGDSQQLRLVRLRSTVMVRVGGGWALDEFL VKNDPCRAKGRTNMELREKFIADGASQGMAA FRPRGRSRPSSRGASPNRSTSVSSQAAQAASQ VPATTTPKILHPLTRNYGKPWLTNSKMSTPCKAA ECSDFPVPSAEGTPIQGSKLRLPGYLSGKGFHSGE DSGLITTAARVRTQFADSKKTPSRPGSRAGSKA GSRASSRRGSDASDFDISEIQSVCSDEVTPQTHR PTPRAGSRPSTAKPSKIPTPQRKSPASKLDKSSKR
2981	A	120	3433	NCLLLQAKGFHGEIEDLQQWLTDTTERHLLASKP LGGLPETAKEQLNVHMEVCAAFEAKEETYKSLM QKGQQLARCPKSAETNIDQDINNLEK WESVE TKLNER\KT\KLEALNLA\MEFHNSL\QDFINWLT QAEQTLNVASRPSLILDTVLFQIDEHKVFANEVN SHREQUIELDKTGTHLKYPFSQKQDVVLKNNLISV QSRWEKVVQRLVERGRSLDDARKRAKQFHEAW

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SKLMEWLEESEKSLDSELEIANDPDKIKTQLAQH KEFQKSLGAKHSVYDTTNRTRGRSLKEKTSLADD NLKLDLMLSELRDKWDTICGKSVERQNKLEEA\ LLFSGQFTDALQALIDWLYRVEPQLAEDQPVHG DIDLVMNLIDNHKAFQKELGKRTSSVQALKRSA RELIEGSRDDSSWVKVQMQLSTRWETVCALSIS KQTRLEAALRQAEFHSVVHALLLEWLAEAEQTL RFHGVLPDDEDALRTLIDQHKEFMKKLEEKRAE LNKATTMGDTVLAICHDPDSITTIKHWITIRARFEE VLAWAKQHQQRLASALAGLIAKQELLEALLAW LQWAETTLTDKKEVIPQEIIEVKALIAEHQTFM EEMTRKQPDVDKVTKYKRRAADPSSLQSHIPV LDKGRAGRKRFPASSLYPSGSQTQIETKNPRVNL LVSKWQQVWLLALERRRKLNDALDRLEELREF ANFDFDIWRKKYMRWMNHKKSRVMDFFRIDK DQDGKITRQEFIDGILSSKFPTSRLMSAVADIFD RDGDGYIDYYEFVAALHPNKDAYKPIDADKIE DEVTRQVAKCKCAKRFQVEQIGDNKYRFFLGNO FGDSQQLRLVRILRSTVMVRVGGGWMLDEFL VKNDPCRAKGRTNMELREKFLADGASQGMMA FRPRGRSRPSSRGASPNRSTSVSSQAAQASPO VPATTPKILHPLTRNYGKPWLTSKMSTPCKAA ECSDFPVPSAEGTPIQGSKLRLPGYLSGKGFHSGE DSGLITTAARVRTQFADSKKTPSRPGSRAGSKA GSRASSRRGSDASDFDISEIQSVCSDEVTPQTHR PTPRAGSRPSTAKPSKIPTQRKSPASKLDKSSKR
2982	A	1	2065	MAAGGAEGGSGPGAAMGDCAEIKSQFRTREGF YKLLPGDGAARRSGPASAQTPVPPQPPQPPGPA SASGPGAAGPASSPPAGPGPGPALPAVRLSLVR LGEPSAGAGEPPATPAGLGSGGDRVCFNLGRE LPGCCRPGSQRWTPITPFLPLYSTINKPI DRIYKGTQPTCEHNCATETISLLVGSAG QVQYLDLIKDKTSLNEERLIDKTKVTYLKWLP ESESFLASHASGHLYLVVSHPCASAPPQYSLL KQAWGFSFYAAKSKAPRNLAKWAVGEGPLNE FAFSPDGRHLACVSQDGLRVFHFDSMLLRGLM KSYFGLLCVCWSPDGRYVVTGGEDDLVTVWS FTEGRVVARGHGHKSWVNAVAFDPYTTRAEEA ATAAGADGERSGEEEEEPAAAGTGSAGGAPLSP LPKAGSITYRFGSAGQDTQFCLWDLTEDVLYPHP PLARTRTLPGTPGTTTPAASSSRGGEPGPGPLPRS LSRSNSLPHPAGGGKAGGPGVAAEPGTPFSIGRF ATLTLQERRDRGAKEHKRYHSLGNISRGSGSG SGSGGEKPSGPVPRSRLDPAKVLGTALCPRIHEV PLLEPLVCKKIAQERLTVLLFLEDCTITACQEGLIC TWARPGKAFTDEETEAQTGEGSWPRSPSKSVVE GISSQPGNSPSGTVV
2983	A	3855	220	RRFRLSAHRAQPCRCRGLMPRGVVFQQLSNLV LQELNANLSNLTSAFEKATAEKIKCQQEADATN RVILLANRLVGGLASENIRWAESVENFRSQGVTL CGDVLLISAFVSYVGYFTKKYRNELMEKFWIPYI HNLKVPITNGLDPLSLLTDDADVATWNNQGLP SDRMSTENATILGNTERWPLIVDAQLGKIKWIKN KYRSELKAIRLGQKSYLDVIEQATSEGDLLIENI GETVDPALDPLLGRNTIKKGKIKIGDKEVGVP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QVPPDPPTHQVLQPTLQARDAGSVHILINFLVTRD GLEDQLLAADVAKERPDLEQLKANLTKSQNEFK IVLKELEDSELLARLSAASGNFLGDTALVENLETT KHTASEIEEKVVEAKITEVKINEARENYRPAER ASLLYFILNDLNKINPVYQFSLKAFNVVFEKAIQR TTPANEVKQRVINLTDEITYSVMYTARGLFERD KLIFLAQVTFQVLSMKKELNPVELDFLLRFPFKA GVVSPVDLFHQHGWGGIKALSEMDEFKNLSDI EGSAKRWKKLVESEAPEKEIFPKWKNTALQK LCMVRCLRPDRMTYAIKNFVEEKMGSKFVEGRS VEFSKSYEESPSTSIFFILSPGVDPLKDVEALGKK LGFTIDNGKLHNVS LGQGQEVVAENALDVAEEK GHWVILQNIHLVARWGLTLDKKLERYSTGRHED YRVFIRAEPAPSPETHIIPQGIENAIKITNEPTGM YANLYKALDLFTQDTLEMCTKEMEFCMLFAL CYFHAVVAERRKFGAQGWNRSYFPNNGDLTISI NVLNYLEANPKVPWDDLRYLFGEIMYGGHITD DWDRRLCRTYLAEYIRTEMLEGDVLLAPGFQIPP NLDYKGYHEYIDENLPPESPYLYGLHPNAEIGFL TVTSEKLFRTVLEMOPKETDSGAGTGVSRREEKV KAVLDDILEKIPETFNMAEIMAKAAEKTYPVVV AFQECERMNLTNEMRRSLKELNLGLKGELTITT DVEDLSTALFYDTPDTPWVARAYPSMMGLAAW YANLLLRIRELEAWTTDFALPTTVWLAGFFNPQS FLTAIMQSMARKNEWPLDKMCLSVEVTKKNRE DMTAPPREGSYVYGLFMEGARWDTQTGVIAEA RLKELTPAMPVIFKAIPVARMETKNIYECVPYKT RIRGPTYVWTFNLKTKEKAAKWILAAVALLQV
2984	A	2	1464	FVLFPGIAMETPGASASSLLPAASRPKRREAGE AGAATSKQVRLEDEEYIEGLQTVIQRDFPDVEK LQAQKEYLEAEERHDLERMFQIAIKFCRAI GKM SREP...VTPAT: STEPVHAGTG... PKG RGLEDGEAGEEBEKEPLPSLDVFLSRY SEDNAS FQIMEVAKERSRARHAWLYQAEBEE...KDN LELPSAEHQAISSQASVETWKYKAKNSLNY -P EGVPDEEQLFKKPRQVVHKNTRFLRDPFSQALS CQLQQAALNAQHKQKVGPDGKELIPQESPRV GGGFVATPSPAPGVNESPMMTWGEVENTPLRV EGSETPYVDRTPGPAFKILEPGRRLGLKMANE AAAKNRKKQEALRRVTENLASLTPKGLSPAMS PALQRLVSRTASKYTDRLRASYTPSPARSTHLK NPGPVGCRPPQSTPGA/PGSATRPLTQDPAISIT DNLLQLPARRKASDFF
2985	A	1890	178	ASTQEAGLLSPPGVGAQRCWNFVACLFPVRACAD MASNDYTQATQSYGAYPTQPGQGYSSQSSQP YGQSSYSGYSQSTDTSGYGQSSYSSYGQSSQNSY GTQSTPQGYGSTGGYGSSQSSQSSYGQSSYPGY GQQPAPSSSTSGSYGSSSSQSSYGQPSGSYSQPS YGGQQSYGQQQSYNPPRGYGGQNNQYNSSSGG GGGGGGGSGYGDQSSMSGSGGGGGGGGGGGGS GGGGGYGNQDQTGAAGSRGYRQIQDRGGRCRG GSGGGGSIGGAAGYNRSGGYEPRGRGGGRGR GGMGGSDRGGFNKFGGPRDQGSRRHDEQDNSD NNTIFVQGLGENVTIESVADYFKQIGIKTNKKTG QPMINLYTDRETGKLKGEATVSFDDPPSAKAAD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=-possible nucleotide deletion, \=possible nucleotide insertion)
				WFDGKEFSGNPIKVSFATRRADFNRRGGNGRGG RGRGGPMGRGGYGGGGSGGGGRGGFSGGGGG GGQQRAGDWKCPNPTCENMNFWRNECNQCK APKPDGPGGGPGGSHMGGNYGDDRRGGRRGGYD RGGYRGRGGDRGGFRGGRRGGGDRGGFGPGKM DSRGEHRQDRRERP
2986	A	1890	178	ASTQEAGLLSPPGVGAQRCWNFVACLPRACAD MASNDYTQQAATQSYGAYPTQPGQGYQQSSQP YQQSYSGYSQSTDTSYGGQSSYSSYGQSSNSY GTQSTPQYGSTGGYSSQSSSYGQSSYPGY GQQPAPSSSTSGSYGSSSSQSSSYGQPQSGSYSQQPS YGGQQQSYGQQQSYNPPRGYGGQNQYNSSSGG GGGGGGGGSYGQDQSSMSGSGGGGGGGGGGG GGGGGYGNQDQTGAAGSRGYRQDRGGRCRG GSGGGGSGGAAGYNRSSGGYEPGRGGRRGGRR GGMGSDRGGFNKFGGPRDQGSRDHSEQDNSD NNTIFVQGLGENVTIESVADYFKQIGIKTNKKTG QPMINLYTDRETGKLKGEATVSFDDPPSAKAID WFDGKEFSGNPIKVSFATRRADFNRRGGNGRGG RGRGGPMGRGGYGGGGSGGGGRGGFSGGGGG GGQQRAGDWKCPNPTCENMNFWRNECNQCK APKPDGPGGGPGGSHMGGNYGDDRRGGRRGGYD RGGYRGRGGDRGGFRGGRRGGGDRGGFGPGKM DSRGEHRQDRRERP
2987	A	1376	898	GGAKAGGAPHPFTLPFRHVGGLSAAPEEVEGML WAGARQHGRNWRKRETSPTGTQGPLPVPVR/VPP GPDGPHAIAPTLWAIPRQCCSPQPGRLNALPPD RCSGPHFGDRAPESCFPGACSVSGACAFKGRPA CPPQEPSLRSSRNRLREGQTFGRMEI
2988	A	1	1011	MGNDSVSYEYGDYSDLSDRPVDCLDGACLAIDP YAPLPYAAIFLNGTGNAMVAVVAGKVAR RRVGATWLLHEAVAEACCLSLPILAVPIARGGH WPYGAVGCRAFLFSELLTMYASVLLAALSADLC FLALGPAWCLRFS/CACGVQVACGAAWTLALL LTVPSAIYRRLHQEHFARLQCVVDYGGSSSTEN AVTAIRFLFGFLGPLVAVASCHSALLCWAARRC RPLGTAIVVGFFVCWAPYHLLGLVLTVAAPNSA LLARALRAEPLIVGLALAHSCLNPMFLYFGRAQ LRRSLPAACHWALRESQQQDESVDSSKSTSHDL VSEMEV
2989	A	27	4074	KSQLCFWVGKAGDILSGDQDKEQKDPYFVETP YGYQLDLDFLKYVDDIQKGNITKRLNIQKRRKPS VPCPEPRTTSGQQGIWTSTESLSSNSDDNKQCP NFLIARSQVTSTPISKPPPLETSLPFLTIPENRQLP PPSPQLPKHNLHVTKTLMETRRRLEQERATMQM TPGEFRRPRLASFGGMGTSSLPSFVSGSNHNP KHQLQNGYQNGDYGSYAPAAPTSSMGSSIRH SPLSSGISTPVTNVSPMHLQHIREQMAIALKRLKE LEEQVRTIPVLQVKISVLQEEKRQLVSQLKNQRA ASQINVCGVRKRSYSAGNASQLEQLSRARRSGG ELYIDYEEEMETVEQSTQRIKEFRQLTADMQA LEQKIQDSSCEASSELRENGECRSVAVGAENMN DIVVYHRGSRCKDAAVGTLVEMRNCGVSVTEA MLGVMTEADKEIELQQQTIESLKEKIYRLEVQLR ETTHDREMTKLKQELQAAGSRKKVDKATMAQP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LVFSKVVEAVVQTRDQMVGSHMDLVDTCVGTS VETNSVGISCQPECKNKVVGPPELPMNWWIVKER VEMHDCAGRSVEMCDKSVSVEVSVCEGTSNTE ESVNDLTLLKTNLNLKEVRSIGCGDCSVDTVCS PKECASRGVNTEAVSQVEAAVMAVPRTADQDT STDLEQVHQFTNTETATLIESCTNTCLSTLTKQTS TQTVETRTVAVGEGRVKDINSSTKTRSIGVGTLL SGHSGFDRPSAVKTKESGVGQININDNYLVGLK MRTIACGPPQLTVGLTASRRSVGVGDDPVGESLE NPQPAPLGMMTGLDHYTERIQKLLAEQQTLLA ENYSELAEAFGEPHSQMGSLNSQLISTLSSINSVM KSASTEELRNPDFQKTSLGKITGSYLGYTCKCGG LQSGSPLSSQTSQPEQEVGTSEGKPISSLDAFPTQ EGTILSPVNLTDQIAAGLYACTNNESTLKSIMKK KDGKDSNGAKKNLQFVGINGGYETTSSDDSSS DESSSESDECDVIBYPLEEEEEEEDEEDTRGMAE GHHAVNIEGLKSARVEDEMQVQCEPEKVEIRE RYELSEKMLSACNLLKNTINDPKALTSKDMRFC LNTLQHEWFRVSSQSAIPAMVGDYIAAFEAISF DVLRYVINLADGNGNTALHYSVSHSNFEIVKLLL DADVCNVDPHQNKAGYTPIMLAALAAVEAKDM RIVEELFGCGDVNAKASQAGQTALMLAVSHGRI DMVKGLLACGADVNIQDDEGSTALMCASEHGH VEIVKLLLAQPGCNGHLEDNDGSTALSIALEAGH KDIAVLLYAHVNFKAQSPGTPRLGRKTSPPGPTH RGSFD
2990	A	69	1687	ERLRPGQRAIRGPVPAAGACASLPPRAGPAQGRH AALGGAEPGSHLHCGVRLQRREPPGGQQRLLPQ RGGSAQTGHQHPGYECQCPGPQPGGTTPALLSL ILEETRGPASANPDKDHSTQPGTMGRKKIQISRI EDORNRQVTFYRKFGLMKKAYFLSVLCDCFIA LLETSATRLFYASTDMDRVLLKYTSYSEFESR ENTDILETLKRRGIGLDGPELEPDEGPPEPGEKFR NAGEGGDPALPRPRLYPAPAMPSPDVVYGAL PPPGCLPSGLGEALPAQSRPSPFRPAAPKAGPPG LGHPFLFSPSHLTSKTPPPLYLPTGRRSDLPGGLA GPRGGLNTRSLSYGLQNPCSTATPGPPLGSFPFL PGGPPVGABAWARRVPQPAAPRRPQSSIKSER LFLRPPGAPATFLRPSPIPCSSPGPWQSLCGLGPP CAGCPWPTAGPGRSPGGTSPERSPGTARARGDP VSLQAFSEKTHVTAPLRGGGLEVGWGTQSSAG GLLSFFFLFVCISTNKNARGVRGPEKK
2991	A	3	1159	IPQPLHCASPKEMSRLRCGDAARTLGRVFGRYF CSPVRPLSSLPDKKKELLQNGPDLQDFVSGDLAD RSTWDEYKGNLKRQKGERLRLPPWLKTEIPMGK NYNKLKNTLRNLNLHTVCEEARCPNIGECWGGG EYATATATIMLMGDTCTRGCRFCSVKTARNPPP LDASEPYNTAKAIAEWGLDYVVLTSVDRDDMP DGAEHIAKTVSYLKERNPKILVECLTPDFRGDL KAIEKVALSGLDVYAHNVETVPELQSKVRDPRA NFDQSLRVLKHAKKVQPDVSKTSIMLGLGENDE QVYATMKALREADVDCLTLGQYMQPTRRHLKV BEYITPEKFKYWEKVGNELGFHYTASGPLVRSS YKAGEFFLKNLVAKRKTDL
2992	A	3	1636	PVPGVPTSPSPCCPQDMQGPWVLLLLGLRLQLSL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GVIPAEENPAFWNRQAAEALDAAKKLQPIQKV AKNLILFLGDGLGVPTVTATRLKGQKNGKLGPE TPLAMDRFPYLALSKTYNVDRQVPDSAATATAY LCGVKANFQTIGLSAAARFNQCNTTRGNEVISV MNRKQAGKSVGVVTTTRVQHASPAGTYAHTV NRNWYSADAMPASARQEGCQDIATQLISNMDID VILGGGRKYMFPMTGPDPEYPADASQNGIRLDG KNLVQEWLAKHQGAWYVWNRTELMQASLDQS VTHLMGLFEPGDTKYEIHRDPTLDPSLMEMTEA ALRLSRNPRGFYLFVEGGRIDHGHHEGVAYQA LTEAVMFDDAIERAGQLTSEEDTLTLVTADHSH VFSFGGYTLRGSSIFGLAPSKAQDSKAYTSILYGN GPGYVFNSGVRPDVNESESGSPDYHQAGIVPLS SETHGGEDVAVFARGPQAHLVHGVQEQSFVAH VMAFAACLEPYTACDLAPPACTTDAHPVAASL PLLAGTLLLLGASAAP
2993	A	3	685	DAWARLLKMNRLFGKAKPKAPPSLTDCIGTVD SRAESIDKKISRLDAELVKYKDQIKKMREGPAKN MVKQKALRVLKQKRMVYEQQRDNLA\NSHSTW TS\HYTIQSLKDTKTVDAMKLGVKEMKKA\YKQ VKIDQIEDLQDQLEDMMEDANEIQEALSRSYGT ELDEDDLEAELDALGDELLADESSYLDEAASA PAIPEGVPTDTKNKDGVLVDEFGLPQIPAS
2994	A	1710	161	RRCELTPFIKTLILPKSWGAPPEDVVMQHVSSSQ SSQRHVQWPGACPGAGEEQPACSQPSLPLTLPS SHQLQQLMVRGGPAGGQNMNVLDLQGVGPGGLQ GSPQVTLAPLPLSPPTSPGFQFSAQPRRFEHGS YIQVTSPLSQVQTSPTQSPSPGQALQNVRA APGPGGLGLCSSSPTGDFVDASVLVRQISLSPSSGG HFVFQDGSGLTQIAQGAQVQLQHPGTPITVRERR PSQFHTOSGGTHHQQSPAAGGAGLOPIASF SHITATLTPQISSEKQLVQQQQVAGGTPPL GFERTFGVLLPGAGGAAGFGMTSPPTSPSRTA VPPGLSSLPLTSVGNMGKVPKKLEELPASPE MAQMRKQCLDYHHQEMQALKEVFKEYLIEL LQHFQGNMMDFLAFKERLYGPLQAYLRQNDLDI EEEEEEVHFEVINDEVKVVARKHGQPGTPVAIAT QLPPRTSAAFPAAQQQLQVLSDGSTVQLPRLSSL GFEDSMC
2995	A	3	924	SAPSGIDASTHAFARCKHPINVRDPSIPTYGLRQS ILLNTRLQDCYVDSPALTNWMARTCAKQINAP APATTSSWEVVRNPLIASSFLVKLVLRRLKKNK CCPPCKFGGKLSKRLKHKDDSVMKATQQARK RNFISKSKQPAGHRRPAGGIRESSKESSKEKLT VQDLEDRYAEHVAATQALPQDSGTAAWKGVRV LLPETQKRQQLSEDTLTIHGLPTEGYQALYHAVV EPMLWNPSGTPKRYSELGKAIKQKLWEALCSQ GAISEGAQRDRFPGRKQPGVHEEPVLKKWPKLK SKK
2996	A	3	1713	GKFGIKPSQRRISGKSTFHSEMEGEDTRDDSLYSI LEELWQDAEQIKRCQEKHNKLLSRTTFLNKKILN TEWDYIEYKDFGKFVHPSPNLILSQKRPKRDSE KSFKHNLDLHIHNKSNAAKNLDKTIGHGQVFTQ NSSYSHHENTHTGVKFCERNQCGKVLSLKHSLS QNVKFIGEKANTCTEFGKIFTQRSHFFAPQKIHT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VEKPHELKSCVNVFTQKPLLSIYLRVHRDEKLYR CTKM/CGKGLHPRNSELIMHEKTHTREKPYKCNE CGKSFFQVSSLLRHQTTHTEKLFECSECGKGF LNSALNIHQIHTGERHHKCSECGKAFTQKSTLR MHQRIHTGERSYICTQCGQAFIQKAHLIAHQRIH TGEKPYECSDCGKSFPKSKQLQMHKRIHTGEKPY ICTECGKAFTNRSNLNTHQKSHTGEKSYICAECG KAFTDRSNFNKHQTIHTGEKPYVCADCGRAFIQK SELITHQRIHTTEKPYKCPDCEKSFSSKKPHLKVHQ RIHTGEKPYICAECGKAFTDRSNFNKHQTIHTGD KPYKSCDCGKGFTQKSVLSMHRNIHT
2997	A	3	1763	AASTRTMGSRHFEGIYDHFVGHFGRFQRVLYFICA FQNISCGIHYLASVFMGVTPHHVCRPPGNVSQVV FHNHSNWSLEDTGALLSSGQKDYVTVQLQNGEI WELSRCSRKNKRENTSSLGYEYTGSKKEFPVCDG YYTDQNTWKSTAVTQWNLVCDRWLAMLQIPL FMFGGPTGIG/VTFGYFSDRLGRRVVLWATSSS MFLFGIAAAFAVDYYTFMAARFFLAMVASGYLV VGFVYVMEFIGMKSRTWASVHLHSFFAVGTLLV ALTGYLVRTWWLYQMILSTVTPFILCCWVLP TPFWLLSEGRYEEAQKIVDIMAKWNRASSCKLS ELSLDLQGPVSNPTEVQKHNLSYLFYNWSITK RTLTVWLIWFTGSLGFYSFSLNSVNLGGNEYLN FLLGVVEIPAYTFVCIAMDKVGRRTVLAYSFLCS ALACGVVMVIPQKHILGVVTAMVKGILPIGAA FGLIYLYTAELYPTTVRSLAVGSGSMVCRLASIL APFSVDLSSIWIFIPQLFVGTALLSGVLTLKLP TLGKRLATTWEEAAKLESENESSKSKLLTTNNS GLEKTEAITPRDSGLGE
2998	A	3	1441	QRPASQLLAPFAAEALPGAPRAAMAQHFSLAAC D/VVGFDFHTTCRYNPEEAPLIVNSFAQ/LVKP KGYDNEELTGEDWDFCCXGLALDLEDGNFL KLANNGTVLRASHGTKMMTPPEVLAEAYGKKEW KHFLSDTGMACRSGKYFYFDNYFDLPGALLCAR VVDYLTKLNNQKTFDFWKDIVAAIQHNYKMS AFKENCGIYFPEIKRDPGRYLHSRPESVKKWLRQ LKNAGKILLITSSHSDYCRLLCAVILGNDFTDLF DIVITNALKPGFFSHLPSQRPFRTLENDEEQEALP SLDKPGWYSQGNVHLVYELLKMTGKPEPKVV YFGDSMHSDFPARHYSNWETVLIIEELRGDEGT RSQRPEESEPLEKKGKYEGPKAKPLNTSSKKWGS FFIDSVLGLENTEDSLVYTWSCKRISTYSTIAPSI BAIAELPLDYKFTRFSSSNSKTAGYYPNPPLVLSS DETISK
2999	A	320	2417	LRRRKMTQPQSLQTTFLLSLLFLVQGAHGRGRH EDFRFCSQRNQTHRSSLHYKPTDLRISIENSEEA LTVHAPFPAAHPASRSFPDPRGLYHFCLYWNHR AGRLHLLYGKRDFLSDKASSLLCFQHQEESLAQ GPPLLATSVTSWWSPQNISLPSAASFTHSPPH TGAHNASVDMCELKRDQLLSQFLKHPQKASRR PSAAPASQQLQSLESKLTSVRFMGDMGSFEEDRI NATVWKLQPTAGLQDLHIHSRQEEEQSEIMEYS VLLPRTLFRQTKGRSGEAKRLLLVDFSSQALFQ DKNSSQVLGEKVLGIVVQNTKVANLTPVVLT QHQLQPKNVTLCVFWVEDPTLSSPGHWSSAGC

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ETVRRRETQTSFCFNHLTYFAVLMVSSVEVDAVH KHYLSLLSYVGCVVSAALCLVTIAAYLCRVPLP CRRKPRDYTIKVHNMNLLAVFLDTSFLLSEPV LTGSEAGCRASAIHLHFSLLTCLSWMGLEGYNLY RLVVEVFGTYVPGYLLKLSAMGWGFPFLVTLV ALVDVDNYGPILA VHRTPGVTYPSMCWIRDSL VSYITNLGLFSLVFLFNMAMLATMVVQILRLRPH TQKWSHVLTLCLSLVLGLPWALIFFSFASGTFQ LVVLYLFSIITSFQGLIFTWYWSMRLQARGGSP LKNSDSARLPISSGSTSSRI
3000	A	66	1003	SRGQLDAGQSSEQHGGNRQPEQSRSSSSSSSP RRSRSAEPAMALSMPLNGLKEEDKEPLIELFK AGSDGESIGNCPFSQRLFMILWLKGVVFSVTTVD LKRKPADLQNLAPGTHPPFTTFNSEVKTVDVNKIEE FLEEVLCPPKYLKLSPKHPESNTAGMDIFAKFSA YIKNSRPEANEALERGLLKTQKLDEYLN SPLPD EIDENSMEDIKFSTRKFLDGNEMTLADCNLLPKL HIVKVVAKKYRNFDIPKEMTGIWRYLTNAYS RD EFTNTCPSDKEVEIAYS DVAKRHLHQVKSRLKE VSFMSPP
3001	A	779	2006	LALTFRSALSTLPGSPMTSSGSPDLQLAWGPSLLP HPPSVWSPALPSCFAGPCPLPLSDTQGWGPN WLAPPSAALCRPDAAVWPDLPSSNILLVTPPPAK *SAVAV*PCPRGAHSLERAARQYTISGSSTSQSGK CSKRDTKCCAVTTSWGCWFQKH WKGEDSGW AFQEGSHLGEGHL
3002	A	909	2799	VEEAWTVWLHWGVRECLLEETNQKEEAASSN WTKARGPFWQEDWVWDMRLKMTTRNFPEREV PCDVEVERFTREVPCSSLGDGWDENQEGHLR QSALTLEKPGTQEAICEYPGFGEHLIASSDLPPSQ RVLATNGFHPADS KRTGL PESVKSFNHFTSLGHQKIMKRGGKSYECNFENI FTLSSSLNENQRNLPGEKQYRCTECGKCFNNS LVLHHRHTGKPYTCNECGKSFSKNYNLIVIQ RIHTGEKPYECSKCGKAFSDGSALTQHQRIHTGE KPYECLECGKTFNRNSSLILHQRHTGKPYRCN ECGKPFTDISHLTVHLRIHTGEKPYECSKCGKAF RDGSYLTQHERHTGKPFCEACGKSFNRN SHL IVHQKHSGEKPYECKEKGKTFIESAYLIRHQRIH TGEKPYGCNQCCQLFRNIAGLIRHQRHTGKPY ECNQCGKA FRDSSCLTKHQRIHTKETPYQCPECG KSFKQNSHLAVHQR LHSREGPSRCPQCGKMFQK SSSLVRHQRAHLGEQPMET*WLGAT*VFQFTLTP VFRRRVLDLTP LWSVEKNPLSYPVN
3003	A	2	1489	SLTEHLSFFQPTAHSLSLLGTMTTCRQFTSSSS MKGSCGIGGGIGGGSSRISSVLAGGSCRAPSTYG GGLSVSSRFSSGGACGLGGGYGGGFSSSSSPGSG FGGGYGGGLGAGFGGGLGAGFGGGGAGG DGLL VGSEKVTMQNLNDRLASYLDKVRAL EEAADL EVKIRDWYQRQRPSEIKDYSFYKTIEDLRNKIIA ATIENAQPILQIDNARLAADDFTKYEH ELALRQ TVEADVNGLRRLVDELTLARTDLEMQIEGLKEE LAYLRKNH*EEMLALRGQTGGEVNVETDAAPG VDLSCILNEMRNQYEQMAEKNRRDAETWFLSKT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EELNKEVASNSELVQSSRSEVTELRRVLQGLEIEL QSLSMKASLENSLEETKGRYCMQLSQIQGLIGS VEBQLAQLRCMEQQSQEYQILLDVKTRLEQEI TYRRLLEGEDAHLSQASGQSYSSREVFTSSSS SSRQTRPILKEQSSSSFSQGGSS
3004	A	2	940	GCAPDTRFFVPEPGGRGAAPWVALVARGGCTFK DKVLVAARRNASAVVLYNEERYGNITLPM SHAG TGNIVVIMISYPKGREILELVQKGIPVTMTIGVGT RHVQEFISGQSVVFAIAFITMMIISLAWLIFYIQ RFLYTGSQIGSQSHRKETKKVIGQLLLHTVKHGE KGIDVDAENCAVCIEFNVKDILPCKHIFHRIC IDPWLLDHRITCPMCKLDVIKALGYWGEPGDVQE MPAPESPGRDPAANLSLALPDDDDGSDSSPPSA SPAESPQCPSFKGDAGENTALLEAGRSDSRHG GPIS
3005	A	184	2552	TMTHQFLLLFLFWVCLPHFCSPEIMFRRTVPVQQ RILSSRVPRSDGKILHRQKRGWMWNQFFLEEY TGSDYQYVGKLHSDQDKGDGLKYLSGDGAGT LFIIDEKTGDIHATRRIDREEKAFYTLRAQAINRR TLRPVEPESEFVIKIHINDNEPTFPEEY TASVPE MSVVGTSVVQVATDADDPSYGNSARVITYSILQ GQPYFSVEPETGIIRTALPNMNRNREYQVVIQ AKDMGGQMGGLSGTTTNNITLTDVNDNPPRFQ NTHLRVLESSPVGTAIGSVKATDADTGKNAEVE YRIIDGDGTD MF DIVTEKDTQEGITVKKPLDYES RRLYTLKVEAENTHVDPRFY YLGPFKDTTIVKISI EDVDEPPVFSRSSYLFEVHEDIEVGTIIGTVMARD PDSISSPIRFS LDRHTDLDRIFNIHSGNGSLYTSKP LDRELSQWHNLT VIAAEINNPKETTRVAVFVRIL DANDNAPQFAVFYDTFVCENARPGQLIQTISA VD KDEPLGGOKTESLA AVPTTVQDNEDNTARIL TRKNGN... LPVVISDNDYPIQSSGTLYI RVCACDSQG MQSCSAEALLPAGLSTGALIAIL LCIILLVIVVLTAA LKQRQKKEPLILSKEDIRDNI SYNDEGGGEEDTQAFDIGTLRNPAAIEKKLRD IIPETLFIPTPTAPDNTDVRDFINERLKEHDLD TAPPYDSLATYAYEGNDSIAESLSLESGTTEGD QNYDYLRWGPFRNKL PQYGGGESDKDS
3006	A	2	541	GRVDKTWWGKSVGIMLTELEKALNSIIDVYHKY SLIKGNFHAVYRDDLKKLETECPQYIRKKGAD VWFKELDINTDGA VNFQEFLILVIKMGVAALNSII DVYHKYSLIKGNFHAVYRDDLKLETECPQYI RKKGADVWFKELDINTDGA VNFQEFLILVIKMG VGSPQKKVASYF
3007	A	1	1253	MYEGIRCLLKALLGFVSLAIGTLYCPRQYRFPFG SLGIEAINVPEPIPD SYRDMATWPTHAPSVEEG GQGRFGNQADHFLGSLAFALLNRS LA VPSWIE YQHHKPPFTNLHVS YQKYFKLEPLQAYHRVISLE DFMEKLAPTHWPPEKRVAYCFEVA AQRSPDKKT CPMKEGNPFGPFWDQFHVSNKSELFTGISFSAS YREQWSQRFSPEHPVLALPGAPAQFPVLEEHRP LQKYMVWSDVMVKTGEAQIHAHLVRPYVGIHL RIGSDWKNACAM LKDG TAGSHFMA SPQCVGYS RSTAAPLTMTMCLPDLKEIQRAVKLWVRSLDAQ SVYVATDSesyVPELQQLFKGKVKVVS LKPEVA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QVDLYILGQADHFIGNCVSSFTAFVKRERDLQGR PSSFFGMDRPPKLRDEF
3008	A	3136	1898	TARGGGSEPGPTMAANYSSSTRREHV KVKTS QPGFLERLSETSGGMFVGLMAFLLSFYLIPTNEG RALKTATSLAEGLSLVVSPDSIHSVAPENEGRLV HIIGALRTSKLLSDPNYGVHLPAVKLRRHVEMY QWVETESREYTEDGQVKKETRYSYNTEWRSEII NSKNFDRIGHKNPRAMAGESFMATAPFVQIGRF FLSSGLIDKVDNFKSLSLSKLEDPHVDIIRRGDF YHSENPKYPEVGDLRVFSYAGLSGDDPDLGPA HVVTVIARQRGDQLVPFSTKSGDTLLLLHHGDFS AEEVFHRELRSNSMKTWGLRAAGWMAFMGL NLMTRILYTLVDWFPVFRDLVNIGLKAFACVAT SLTLLTVAAGWLFYRPLWALLIAGLALVPILVAR TRVPAKKLE
3009	A	93	659	DAAVAMTAQGGGLVANRGRFRKWAIELSGPGGG SRGRSDRGSGQGDLSYPVGYLDKQVPDTSVQET DRILVEKRCWDIALGPLKQIPMNLFMYMAGNTI SIFPTMMVCMMAWRPIQALMAISATFKMLESS QKFLQGLVYLIGNLMGLALAVYKCQSMGLLP THASDWLAFIEPPERMEFSGGGLL
3010	A	2	1041	LIDSAKARYWTQRGTWVYDNNLLLLLKCLWSN VVPECTMASSNTVLMRLVASAYSIAQKAGMIVR RVIAEGDLGIVEKTCATDLQTKADRLAQMSICSS LARKFPKLTIGBEDLPSEEVDQELIEDSQWEEILK QPCPSQYSAIKEEDLVVWVDPLDGTKEYTEGLL DNVTVLIGIAYEGKAIAGVINQPYNYEAGPDAV LGRITWGVGLGAFGFQKEVPAGKHITTTTRSH SNKLVTDCVAAMNPDAVLRVGGAGNKIIQLIEG KASAYVFASPGCKWDTCAPEVILHAVGGKLTID IHGNVLQYHKDVKHMNSAGVLRNVDYAS RVPESIKNALVP
3011	A	291	1452	SPQKTMRSHTITMTITSVSSWPYSSHRMRFITNH SDQPPQNFSA TPNTTCPMDEKLLSTVLTTSYSVI FIVGLVGNIALYVFLGIHRKRNSIQIYLLNVAIAD LLIFCLPFRIMYHINQNKWTLGVILCKVVGTLFY MNMYSIILLGFISLDRIYKINRSIQQRKAITTKQSI YVCCIVWMLALGGFLTMILLTKKGGHNSTMCF HYRDKHNAKGAEIFNFI VVMFWLIFLLILSYIKI GKNLLRISKRRSKFPNSGKYATTARN SFIVLIIFTI CFVPYHAFRTYISSQLNVSSCYWKEIVHKTNEIM LVLSSFNSCLDPVMYFLMSSNIRKIMCQLLFRF QGEPSRSESTSEFKPGYSLHDTSAVKIQSSSKST
3012	A	246	1346	TEPVGYTKAEPIAMRSLGALLLLSACLAVSAG PVPTPPDNIQVQENFISRIYK WYNLAIGSTCPW LKKIMDRMTVSTLV LGE GATEAEISMTSTRWRK GVCEETSGAYEKTDTDGKFLYHKS KWNITMESY VVHTNYDEYAIFLT KKF SRHHGPTTAKLYGRAP QLRETLQDFRVVAQGVGIPEDSIFTMADRGEV PGEQEPEPILIPRRRAVL PQEEEGSGGGQLVTEV TKKEDSCQLGYSAGPCMGMTSRYFYNGTSMAC ETFQYGGCMGNGNNFVTEKECLQTCRTVAACN LPIVRGPCRAFIQLWAFDAVKGKCVLPYGGCQ GNGNKFYSEKECREYCGVPGDGDEELLRFNS
3013	A	67	379	RQMA LLKANKDLISAGLKEFSVLLNQQVFNDPL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VSEEDMVTVVVEDWMNFYINYRQQVTGEPQER DKALQELRQELNTLANPFLAKYRDFLSHELP SH PPPSS
3014	A	1	373	GTWSWTLRAVMSASVSVSVSRVLEEYLSSTPQRL KLLDAYLLYILLTGALQFGYCLFVLTFFHNSLLLF FFFCVGSFHSNVYFLLFTLSFLCFIAFYFFLRFFS LFIWFFHVFFIELSLFYF
3015	A	2	1321	AAAEGTAPSPGRVSPPTPARGEPEVTVEIGETLYC RRPDSTWHS AEVIQSRVNDQEGREEFYVHYVGF NRRLEWVDKNRLALTKTVKDAVQKNSEKYL ELAEQPERKITRNQKRKHDEINHVQKTYAEMDP TTAALEKEHEAITKVKYVDKIHNIGNYEIDA WYFS PPPEDY GKQPKLWLCEYCLKYMKYEKS YRFHLG QCQWRQPPGKEIYRKS NISVYEVDGKDHIYCY NLCLLAKLFLDHKTL YFDEVPFVFYILTEVDRQG AHIVGYFSKEKESPDGNNVACILTPPYQRRGYG KFLIAFSYELSKLESTVGSPEKPLSDLGKLSYRSY WSWVLLLEILRDFRGTL SIKDLSQMTSITQNDIIST LQSLNMVKYWKGGQHVICVTPKLVEEHLKSAQY KKPPITGGWGAAVCRGRWGSVSIWTGRSQGLLI AVT
3016	A	2	1321	AAAEGTAPSPGRVSPPTPARGEPEVTVEIGETLYC RRPDSTWHS AEVIQSRVNDQEGREEFYVHYVGF NRRLEWVDKNRLALTKTVKDAVQKNSEKYL ELAEQPERKITRNQKRKHDEINHVQKTYAEMDP TTAALEKEHEAITKVKYVDKIHNIGNYEIDA WYFS PPPEDY GKQPKLWLCEYCLKYMKYEKS YRFHLG QCQWRQPPGKEIYRKS NISVYEVDGKDHIYCY NLCLLAKLFLDHKTL YFDEVPFVFYILTEVDRQG AHIVGYFSKEKESPDGNNVACILTPPYQRRGYG KFLIAFSYELSKLESTVGSPEKPLSDLGKLSYRSY WSWVLLLEILRDFRGTL SIKDLSQMTSITQNDIIST LQSLNMVKYWKGGQHVICVTPKLVEEHLKSAQY KKPPITGGWGAAVCRGRWGSVSIWTGRSQGLLI AVT
3017	A	38	704	EAHPGGQLGSENGVRMDEDVLTTLKILIGESG VGKSSLLLRFTDDTFDPELAATIGVDFKVKTISVD GNKAKLAIWDTAGQERFRTLTPSYRGAQGVIL VYDVTRRDTFVKLDNWLNELETYCTRNDIVNM LVGNKIDKENREVDREGLKFARKHSMLFIEAS AKTCDGVQCAFEELVEKIQTPGLWESENQNG VKLSHREEGQGGGACGGYCSVL
3018	A	2640	2861	APVLILQMVKLSIVLTPQFLSHDQGLTKELQQH VKS VTCPEYLRKVSECRQMGP GALEQFPGLSC HTSHSG
3019	A	1307	711	PGITMAASLVGKKIVFVTGNAKKLEEVVQILGDK FPCTLVAQKIDLPEYQGEPDEISIQKCQEA VRQV QGPVLVEDTCLCFNALGGLPGPIYKWFLEKLKPE GLHQLLAGFEDKSAYALCTFALSTGDPSQPVRLF RGRTSGRIVAPRGCDQDFGWDPCFQPDGYEQTYA EMPKA EKNAVSHRFRALLELOEYFGSLAA
3020	A	1202	180	VSCLP TSCKMITLNNQDQVPVFNSSHPDEYKIAA LVFYSCIFIGLFVNITALWVFSCTTKKRITVTIYM MNVALVDLIFIMTL PFRMFYAKDEWPFGYFC QILGALT V FYPSIALWLLAFISADRYMAIVQPKY

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				AKELKNTCKAVLACVGVWIMLTITTTPLLLLYK DPDKDSTPATCLKISDIYLKAVNVNLTRLTFFF LIPLFIMIGCYLVIIHNLHGRTSKLKPKVKEKSIRI IITLLVQVLVCFMPFHICFAFLMLGTGENSYNPW GAFTTFLMNLSTCLDVILYYIVSKQFQARVISVM LYRNYLRSMRRKSFRRSGSLRSLSNINSEML
3021	A	27	1897	EEFCTWIAVRVGEMETAPKPGKDVPKDKDLQT KRKKPRRYWEEETVPTTAGASPGPPRNKKNREL RPQRPKNA YILKKSRIKPKQVPKKPREWKNPES QRGLSGAQDPFPGPAPVPVEVVQKFCRIDKSRKL PHSKAKTRSRLEVAEAEETSIKAARSELLAEE PGFLEGEDGEDTAKICQADIVEAVDIASAAKHFD LNLKQFGPYRLNYSRTGRHLAFGGRGHVAALD WVTKKLMCEINVMEAVRDIRFLHSEALLAVAQN RWLHIYDNQGIELHCIRRCDRVTRLEFLPFHLLA TASETGFLT YLDVSVGKIVAALNARAGRLDVM QNPYNAVHILGHSNGTVSLWSPAMKEPLAKILC HRGGVRAVAVDSTGT YMATSGLDHQLKIFDLRG TYQPLSTRTLPHGAGHLAFSQRGLLVAGMGDVV NIWAGQGKASPPSLEQPYLTHRLSGPVHGLQFCP FEDVLGVGHTGGITSM LVPGAGEPNFDGLESNPY RSRKQRQEWVKALLEKVP AELICLDPRALAEV DVISLEQGKKEQIERLGYDPQAKAPFPKPKQKG RSSTASLVKRKRKVMDEEHRDKVRQSLQQQHH KEAKAKPTGARPSALDRFVR
3022	A	1	2249	MTAQDSNTSAHAQRDGP ELPASSWSRWFPLSC LSSPPVSAVEVATEGRDREVAKV GQRFCDDTSGE LRQARDRCCVRMPAPVGRRSPSPRSSMAAVA LRDSAQGMTFEDVAIYFSQEEWELLDESQRFLYC DVMLENFAHVTSLSGYCHGMENEAIASEQSVSIQ VETSKONTPTOKTHLSEIKMCVITLDTLPA*EH QTTSPVQKSYLGSTNRRFCFSADLHQHQKHYN EEEPWKRKVDEATFVTGCRFHV LNYFTCGEAFP APDLLQHEATPSGEEPHSSSSKHIAFFNAKSY KWGEYRKASSHKHTLVQHQSVCSEGGLYECSK CEKAFTCKNTLVQHQQIHTGQKMFECSECEBSFS KKCHLLHKIHTGERPYECSDREKAFIHKSEFIHH QRRHTGGVRHECGEGRKTF SYKSNLIEHQRVHT GERPYECGECGKSFRQSSSLFRHQRVHSGERP YQ CCECGKSFRQIFNLIRHRRVHTGEMPYQCSDCGK SFSCKSELIQHQRHSGERP YECRECGKSFRQFSN LIRHRSIHTGDRPYECSECEKSFSRKFIHQHQRVH TGERPYECSECGKSFRKSDLIQHRRHTGT RPYE GSECGKSFRQ RSGLIQHRR LHTGERPYECSECGK SFSQSASLIQHQRVHTGERPYQCCECGKSFRQIFN LIRHRRVHTGEMPYQCSDCGKSFSCKSELIQHRR HSGERP YECSECGKSFSRKS NLIRHRRVHTEERP
3023	A	3148	634	AAGALRCLAAFPRAEPASRGQSSPARACAASR AERATAAAMAHRCLRLWGRGGCWPRGLQQLL VPGGVGPGEQCLRLTYRFVTTQARASRNSLLTD IIAAYQRFCSRPPKGFGKYFPNGKNGKKASEPKE VMGEKKESKPAATTRSSGGGGGGGGKRGKKD DSHWWSRFQKGDIPWDDKDFRMFFLWTALFWG GVMFYLLKRSGREITWKDFVNNYLSKGVVDRL EVVNKRFRVTVFTPGKTPVDGQYVWFNIGSVDT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				FERNLETLQQLGIEGENRVPVVYIAESDGSFLLS MLPTVLIHAFLLYTIRRGPAAGRTGRGMGGLFSV GETTAKVLKDEIDVKFKDVAGCEEAKLEIMEFV NFLKNPKQYQDLGAKIPKGAILTGPPGTGKTLA KATAGEANVPFITVSGSEFLEMFGVGPARGVRDL FALARKNAPCILFIDEIDA VGRKRGRGNFGGQSE QENTLNQLLVEMDGFNTTTNVVILAGTNRPDILD PALLRPGRFDRQIFIGPPDIKGRASIFKVHLRPLKL DSTLEKDKLARKLASLTPGFSGADVANCNEAA LIAARHLSDSINQKHFEQA IERVIGGLEKKTQVLQ PEEKKTVA YHEAGHAVAGWYLEHADPLLKVSII PRGKGLGYAQYLPKEQYLYTKEQLLDRMCMTL GGRVSEBIFFGRIITGAQDDLKVTQSAYA QIVQ FGMNEKVGQISFDLPRQGD MVLEKPYSEATARI DDEVRLINDAYKRTVALLTEKKADVEKVALLL LEKEVLDKNDMVVELLGP RPFAEKSTYEEFVEGT GSLDEDTSLPEGLKDWNKEREKEKEEPPGEKVA N
3024	A	274	1455	LRACSLPSMSALEKSMHLGRLPSPPLPGSGGSQ SGAKMRMGPRKRDFSPVPWSQYFESMEDVEV ENETGKDTFRVYKSGSEGPVLLLLHGGGHSALS WAVFTA AII SRVQCRIVALDLRSHGETKVKNPED LSAETMAKDVGNVVEAMYGDLPPIMLIGHSMG GAIAVHTASSNLVPSLLGLCMIDVVEGTAMDAL NSMQNFLRGRPKTFKSL ENAIEWSVKSGQIRNLE SARVSMVGQVKQCEGITSPEGSKSIVEGIIIEEEE DEEGSESISKRKEDDMETKKDHPYTWRIELAKT EKYWDGWFRGLSNLFLSCPIPKLLLLAGVDRLD KDLTIGQM QGKFQMQVLPQCGHAVHEDAPDKV AEAVATLIRHRFAEPIGGFQCVFPGC
3025	A	621	306	YHGGGGRAGGSEFSVQGGGQLRPPFTFSKSL SWKGLSSLLFPLYNLQKATLTKELGRGHSPP HLEBPHMLPSGAARWRVLEAPVLVLEPLVLRPA AAPTP
3026	A	1533	454	AKVPQSTREEKRENGLEARSANLMGFNVEEM YEAHAWIQRLSLQNHIIENNHILYLGRKEHDIL SQLQKTSSVSITEIISPRTELEIEGARADLIEVVM NIEDMLCKVQBEMARKKERGLWRSLGQWTIQQ QKTQDEMKENIFLKCPVPPTQELLDQKKQFEKC GLQVLKVEKIDNEVLMAAFQRKKKMMEELHR QPVSHRLFQQVPYQFCNVVCRVGFQRMYSTPCD PKYGAGIYFTKNLKNLAEKAKKISAADKLIYVE AEVLTGFFCQGHPLNIVPPPLSPGAIDGHDSVVD NVSSPETFVIFSGMQAIPQYLWTCTQEYVQSQDY SSGPMRPFQAHPWRGFASGSPVD
3027	A	179	703	PFLGASSNTFRLQVQTQESKAQKEVKMGFIFSK SMNESMKNQKEFMLMNARLQLERQLMQSEMR ERQMAMQIAWSREFLKYFGTFFGLAAISLTAGAI KKKKPAFLVPIVPLSFILTYQYDLGYGTLLERMK GEAEDILETEKSKLQLPRGMITFESIEKARKEQSR FFIDK
3028	A	876	1226	AVGKEPSSSTWVRDREGHIRSRRSKMMLWKLT DNIKYEDCEVSATPARSSVRSQAPSLTLPLLLSL QPAAKRGWDKLSAQRPSLGFARRTRGRSCRER TWMLPSLVSEFLHRD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * =Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
3029	A	3	1731	FREGRFGSSCAVAAPLAGFQGLIECGYLAVDSP SCWTPGGSNPAAPLPQALLPRLPPTVLPFLGPGL SGELEMFTLPQKDFRAPTTCLGPTCMQDLGSSHG EDLEGECSRKLDQKLPRLRGVGDPAISSNTSYL SSRGRMIKWFWDSEEGYRTYHMDYDEDKNP SGHNLGTSENKLCFDLLSWRLSQRDMQRVEPSL LQYADWRGHLFLREEVAKFLSFYCKSPVPLRPE NVVVLNGGASLFSALATVLCAGEAFLIPTPYG AITQHVCLYGNIRLAYVYLDSEVTGLDTRPFQLT VEKLEMALREAHSEGKVKVGLILISPNPLGDVY SPEELQEYL VFAKRHLRHVTVDVYMLSVFEKSV GYRSVLSLERLPDPQRTHVMWATSKDFGMSGRL FGTLYTENQDVATAVASLCRYHGLSGLVQYQM AQLLRDRDWINQVYLPENHARLKAHTYVSEEL RALGIPFLSRGAGFFIWVDLRKYLLKGTFFEEML LWRRFLDNKVLLSFGKA FECKEPGWFRFVSDQ VHRLCLGMQRVQQVLAGKSQVAEDPRPSQSSEP SDQRR
3030	A	1	584	PWLPWSDGRAARSSRKCPRSRFPVQVGKMAVST VFSTSSMLALSRLSLLSPLSVTSFRRFYRGDSP TDSQKDMIEIPLPWQERTDESIETKRARLLYESR KRGMLENCILLSLFAKEHLQHMTEKQLNLYDRLI NEPSNDWDIYYWATEAKPAPEIFENEVMALLRD FAKNKNKEQRLRAPDLEYLFEKPR
3031	A	1177	359	SLWPWILMDDSLMQISLQLLCVYTANFPNGCSSL CWSSCGQHPVQATHRGAVSNSMLCILKLASQM PLENTTVQQMVFMLLSNLALSHDCKGVIQKSNF LQNFLSLALPKGGNKHLSNLTLWLKLLNLSGGE DGQQMILRLDGCDDLTEMISKYKHKSPLPLLI FHNVCFSANKPKILANEKVITVLAACLESENQ AQRICAAALWALIYNQKAKTALKSPSVKRR EAYSLAKKITFPNSEANPLNAYYLLENLVQL NSS
3032	A	2	1242	GISGRPPRPAKRRMGKNPVRPPRALPPVPSQDDIP LSRPKKKKPRTKNTPASASLEGLAQTAGRRPSEG NEPSTKELKEHPEAPVQRRQKKTRLPLELETST QKKSSSSLLRNENGIDAEPABEAVIQKPRRKT KTQPAELQYANELGVEDEIITDEQTTVEQQSVF TAPTGISQPVGKVFVEKSRRFQAADRSELIKTTEN IDVSMDEVKPSWTTTRDVALTVHRAFRMIGLFSHG FLAGCAVWNIVYVLAGDQLSNLSNLLQQYKT LAYPFQSLLYLLLALSTISAFDRIDFAKISVAIRNF LALDPTALASFLYFTALILSLSQMSTDRHL YTP SSVNGSLWEAGIEEQILQPWIVVNLVVALLVGLS WLFLSYRPGMDLSEELMFSSEVEEYPDKEKEIKA SS
3033	A	3	1436	TATSGGIWLRKWRCHWPRPLPQSCVGTEGGLQ VRDTSSRIAKGGVDHTKMSLHGASGGHERSRDR RRSSDRSRDSSHERTESQLTPCIRNVTSPTROHHV EREKDHSSSRPSPRPQKASPNGSISSAGNSSRNS SQSSSDGCKTAGEMVFFVYENAKEGARNIRTSE VTLIVDNTRFVVDPSIFTAQPNTMLGRMFGSGRE HNFTRPNEKGEYEVAEGIGSTVFRAILDYKGTGII RCPDGISIPELREACDYLCISFEYSTIKCRDLSALM HELSNDGARRQFBFYLEEMILPLMVASAQSGERE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				CHIVVLTDDDVVDWDEEYPPQMGEYSQIYSTK LYRFFKYIENRDVAKSVLKERGLKKIRLGIEGYP TYKEKVKKRPGGRPEVTYNYVQRPFIRMSWEKE EGKSRHVDQFCVKSITSITNLAAAADIPQDQLV VMHPTPQVDEL DILPIHPPSGNSDLDPDAQNPM
3034	A	3	1972	SSLAQHRSVAVLGWPAWAAARAPAMQGGN SGVRKREEEGDGAGAVAAPPAIDFPAEGPDPEY DESDVPAEIQVLKEPLQQTFFFAVANQLLLVSL LEHLSHVHEPNPLRSRQVFKLLCQTFIKMGLLSSF TCSDEFSSRLHNNRAITHLMRSKERVVRQDPCE DISRIQKIRSREVALEAQTSTRYLNEFEELAILGKG GYGRVYKVRNKLDGQYYAIKKILKGATKTVCML KVLREVKVLAGLQHPNIVGYHTAWIEHVHVIQ RADRAAIELPSLEVLSDQEEDEQCGVKNDSSS SSIIFAEPTPEKEKRFESD TENQNNKSVKYTTNL VIRESGELESTLELQENGLAGLSASSIVEQQLPLR RNSHLEESFTSTESSEENVNFGQTEAQYHML HIQMQLCELSLWDWIVERNKRGREYVDESACPY VMANVATKIFQELVEGVFYIHNMGIVHRDLKPR NIFLHGPDQQVKIGDFGLACTDILQKNTDWTNR NGKRTPTHTSRVGTCLYASPEQLEGSEYDAKSD MYSLGVVLELFPFGTEMERAEVL TGLRTGQL PESLRKRCPVQAKYIQHLTRRNSSQRPSAIQLLS ELFQNSGNVNLTLQMKHIEQEKEIAELKKQLNLL SQDKGVRDDGKDGGVG
3035	A	110	1172	KLSCPCSHGTRVTAVRGPRLKAGVQWHDLGSLQ PPPSGLKQSSHLSSSSWDFRHAPHTPETYTCPK MIEMEQAEAQLAELDLLASMFPGENELIVNDQL AVAELKDCIEKKTMEGRSSKVYFTINMNLVDSD EKMAMFSLACILPFKYPAVLPEITVRSVLLSRSOQ TQLNTDLTAFLCHGDCILNATVREHA GYRDTSSSPITGSTVQSVDLINLWVHHIY NKCKRKNILEWAKELSLSGFSMPGK*GVVCVEG PQSACEEFWARLRKLNWKRLIRHREDIPFDGTN DETERQRFKFSIFEKVFVNGARGNHMDGQLY QFLNTKGCVDVFQMFLLWV
3036	A	1	2288	FRFAERRAAAAESDVSAKMAGRSMQAARCPD ELSLTNCAVVNEKDFQSGQHIVVRTSPNHYTFT LKTHPSVVPGSIAFSLPQRKWAGLSIGQEIEVSLY TFDKAKQCIGTMTIEIDFLQKKSIDSNPYDTDKM AAEFIQQFNNQAFSVGQQLVFSFNEKLFGLLVKD IEAMDPSILNGEPATGKRQKIEVGLVVGNSQVAF EKAENSSLNLIGKAKTKENRQSIINPDWNFEKMG IGGLDKEFSDFRRAFAASRVFPPEIVEQMCKHVK GILLYGPPGCGKTLLARQIGKMLNAREPKVVNG PEILNKYVGESEANIRKLFADAEERQRLGANS LHIIIFDEIDAICKQRGSMAGSTGVHDTVVNQLLS KIDGVEQLNNILVIGMTNRPDLIDEALLRPGRLV KMEIGLPDEKGRQLQHHTARMRGHQLLSADV DIKELAVETKNFSGAELEGLVRAAQSTAMNRHI KASTKVEVDMEKAESLQVTRGDFLASLENDIKP AFGTNQEDYASYIMNGIUKWGDVTRVLDDGEL LVQQTKNSDRTPLVSVLLEGPPhSGKTALAARIA EESNFFFIKICSPDKMIGFSETAKCQAMKKIFDDA YKSQLSCVVDDIERLLDYVPIGPRFSNLVLQAL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				LVLLKKAPPQGRKLLIGTTSRKDVLQEMEMLNA FSTIIHVPNIATGEQLLEALLGNFKDKERTTIA QQVKGKKVWIGIKLLMLIEMSLQMDPEYRVRK FLALLREEGASPLDFD
3037	A	1	1347	MLDTGSEHLNRILKALPALQSAGSEGQNGSAESL GEGGTRDSRARRKLGGNKEIPTFYPCLVVRSP VTASDLRGTDFAAYHGLSLILEPLGACNRLSVC VPVHSPPGMRVSPRSPSLRTLVIDPAEPAGAQR RFGSKERSGEAGSAVEGLAVA VSMGDGGAERD RGPARRAESGGGGGRCGDRSGAGDLRADGGGH SPTEVAGTSASSPAGSRESGADSDGQPGPEADH CRRILVRDAKGTIREIVLPKGLDLDRPKRTRTFFT AEQLYRLEMEFQRCQYVVGRETERLARQNLSE TQVKVWFQNRRTKQKKDQSRDLEKRASSASEA FATSNILRLLEQGRLLSVPRAPSLALTPSLPGLP ASHRGTSLGDPNRSSPRLNPLSSASAPPLPPLP AVCFSSAPLLDLPAGYELGSSAFEPYSWLERKVG SASSCKKANT
3038	A	924	501	TELLPLCSRSRGPQSGDPLLQLAQARPRLSGE RLETAPSLLLSRMACVISGWALSRGARTWTWAT PTGPVHRAQPAIRSLAEGALTRLKEEKWPGRYI LPNHLTPPFLYKHLGSVPPSHWRSPLISHSVNILA LNWR
3039	A	1263	111	ACGIRHEGALPGLTATPEAMLRFLPDLAFLIL ALGQAVQFQYVFLQFLGLDKAPSPQKFQVPYI LKKIFQDREAAATGVSRLCYVKELGVRGNVL RFLPDQGFFLYPKKISQASSCLOKLLYFNLSAIKE REQLTLAQLGLDLGPNSSYNLGPPELALFLVQE PHVWGQTPKPGKMFVLRSPWPQGA VHFNL DVAKDWDNDNPRKNFGLFLEILVKEDRDSGVNFQ PPTCARLRCSLHASSLLVVTINPDQCIKTPRA APVPKLSCKNLCHRHQLNDLGLWIKWILAP KGFMANYPCHGECFSLTISLNSNYAFMQALMH AVDPEIPQAVCIPTKLSPISMLYQDNDNDNVILRHY EDMVVDECGCG
3040	A	15	849	ASRLPRGPGCGADMRLGLLLVFAGCTFALYL LSTRLPGRRLGSTEEAGGRSLWFPSDLAELEL SEVLREYRKEHQAYVFLFCGAYLYKQGFAPGS SFLNVLGALFGPWLGLLLCCVLTSGATCCYL LSSIFGKQLVVSYPFDKVALLRKVEENRNSLFF FLLFLRLFPMTNWFNLNSAPILNPIVQFFSVLI GLIPYNFICVQTGSILSTLSTLDALFSWDTVFKLL AIAMVALIPGTLIKKFSQKHLQLNETSTANHIHSR KDT
3041	A	1015	175	GLKRRRLCFAKVGDLGCLSLPPSR SARVLEDISI LSCISVDSRIVRTKVPCSVTMSRPRKRLAGTSGSD KGLSGKRTKTENSGEALAKVEDSNPQKTSATKN CLKNLSSHWMKSEPEsrLEKGVDFKFSIEDLKA QPKQITCWDGVRNYQARNFLRAMKLGEAAFFY HSNCKEPGIAGLMKIVKEAYPDHTQFEKNNPHY DPSSKEDNPKWSMVDVQFVRMMKRFIPLAELKS YHQAHKATGGPLKNMVLFRQRLSIQPLTQEEF DFVLSLEEKEPS
3042	A	1015	175	GLKRRRLCFAKVGDLGCLSLPPSR SARVLEDISI LSCISVDSRIVRTKVPCSVTMSRPRKRLAGTSGSD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KGLSGKRTKTENSGEALAKVEDSNPQKTSATKN CLKNLSSHWMKSEPEsrLEKGVdVKFSIEDLKA QPKQTTTCWDGVRNYQARNFLRAMKLGEAAFFY HSNCKEPGIAGLMKIVKEAYPDHTQFEKNNPHY DPSSKEDNPKWSMVDVQFVRMMKRFIPLAELKS YHQAHKATGGPLKNMVLFTQRLSIQPLTQEEF DFVLSLEEKEPS
3043	A	153	1133	VGTAAPGGRDRAPAMGSFQLEDFAAGWIGGA ASVIVGHPLDVTKTRLQAGVGYGNTLSCIRVVY RRESMFGFFKGMSFPLASIAVYNSVVFVFSNTQ RFLSQHRCGEPEASPPRTLSDLLASMVAGVVS GLGGPVDLIKIRLQMOTQPFrdanLGLKRAVAP AEQPAYQGPVHCITTVRNEGLAGLYRGASAML LRDVPGYCLYFIPYVFLSEWITPEACTGPSPCAV WLAGGMAGAISWGTA TPMDVVKSRLQADGVY LNKYKGVLDcISQSYQKEGLKVFRGITVNAVR GPFMSAAMFLGYELSLQAIRGDHAVTSP
3044	A	41	1316	PPLGAGAGIHARSPHPARRRLTAAGVGGRASG LLPTPWRRHHGPSGAAPYPAARLWQGPWRCRR PQMAQRYDELPHYPGIADGPAALAGFPEAVPA APGPYGPHRPPQPLPPGLDSDGLKRDKEIYGH LFPLLALGFEEKELATCSPRDGAGAGLGTGRGGD VCSSDSFNEDNTAFKQVCSErPFSSNPeldNLM IQAIQVLRfHLLLELEKGMpIDLVIEdRDGGCRE DFEDYPAPCPSLPDQNNIWRDHEDSGSVHLGTP GPSSGGLASQSGDNSSDQGVGLDTSVASPSSGGE DEDLDQEPrrNKKRGIffKVATNIMRAWLFOHL SHPYPSEEQKKQLAQDTGLTILQVNNWFINARRR IVQPMIDQSNRTGQGAAFSPegQPIGGYTETEPH VAFRAPASVGMslNSEGEWHYL
3045	A	3	967	VAHTQWHTTCTQLTHRCLEWYDTHA377 METHTHASLSLPSCQECFPSSPSASLGVSHPHPP PSPRWGQTPEGLPAASPCGPGFRCFSSILPTGDS WGMLACLCTVLWHLPAVPALNRTEOPGPGPSIQ KTYDLTRYLEHQRLSLAGTYLNYLGPFFNEPDFN PRLGAETLPRATVDLEVWRSNDKLRLTQNYE AYSHLLCYLRGLNRQAATAELRRSLAHFCTSLQ GLLSIAGVMAALGYPLPQPLPGTEPTWTPGPAH SDFLQKMDDFWLLKELQTLWRSaKDFNRLKK KMQPPAAAVTLHLGAHGF
3046	A	1185	1584	MYAYMYICTHICICAYRGIHIDVLYMCIYIHIWI HTYLCVHIYVYVYICTHICMCIHTYVYVYTYMY VYTYICLCVYICLCVHIYLCVYIHYMYMCTHICMC IHTYVHMCICVYIHYMYTCVYVYTYTCVYMY
3047	A	811	132	SLDLLGPIGILQEGRDPTGQGPQEKQMPASPM NTDAHLdINFKEGLKKERSYTGQFEANVRDEER QCGCGVVPDSLLMKVLSQRLDQQDCIQKGWVL HGVPRDLDAQHLLNRLGYNPNREFFLNVPFDSI MERLTLRRIDPVTGERYHLMYKPPPTMEIQARLL QNPKDABEQVKLKMdLFYRNSADLEQLYGSait LNGDQDPYTVFEYIESGIINPLPKKIP
3048	A	2	1166	RPRRQGLVQEVQTENVTVAEggVAEITCRlHQ YDGSIVVIQNPARQTLFFNGTRALKDERFQLEEFs PRRVIRLSdarLEDEGGYFCQLYTEDTHHQIAT LTVLVAPENPVVEVREQAVEGGEVELSCLVPRSR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				PAATLRWYRDRKELKGVSSSQENGKVWSVAST VRFRVDRKDDGGIICEAQNALPSGHSKQTQYV LDVQYSPTARIHASQAVVREGDTLVLTCAVTGN PRPNQIRWNRGNESLPERAEAVGETLTLPLVSA DNGTYTCEASNKHGHARALYVLVVYGESRLRPT EGGGGAPDPGAVVEAQTSPYAI VGGILALLVFL IICVLVGMVWCSVRQKGSYLTHEASGLDEQGEA REAFNLNGSDGHRKKEEFFI
3049	A	3159	882	VGCTLRVGVMAAAGSRKRRLAELTVDEFLASGF DSESESESENSQAETREAREAAARSPDKPGGSPSA SRRKGRASEHKDQLSRLKDRDPEFYKFLQENDQ SLLNFSDDSDSSEEEGPFHSLPDVLEEASEEEDGA EEGEDGDRVPRGLKGKKNSVPVTVMVERWKQ AAKQRLTPKLFHEVVQAFRAAVATTRGDQESAE ANKFQVTDSAFNA LVTFCIRDLIGCLOKLLFGK VAKDSSRMLQPSSSPLWGKLRVDIKAYLGSAIL VSCLETTVLAALVRHISVLVPCFLTFFPKQCRML LKRMMVVVWSTGEESLRVLAFLVLSRVCRHKKDT FLGPVLKQMYITYVRNCKFTSPGALPFISFMQWT LTELLEPGVAYQHAFLYIRQLAIHLRNAMTTR KKETYQSVYNWQYVHCLFLWCRVLSTAGPSEA LQPLVYPLAQVIIGCIKLIPTARFYPLRMHCIRALT LLSGSSGAFIPVLPFILEMFQQVDFNRKPGRMSSK PINFSVILKLSNVNLQEKAYRDGLVEQLYDLTLE YLHSAHCIGFPELVLPVVLQKLSFLRECKVANY CRQVQQLLGKVQENSAYICSRQRVSFGVSEQQ AVEAWEKLTREEGTPLTLYYSHWRKLRDREIQL EISGKERLEDLNFPEIKRRKMADRKDEDRKQFKD LFDLNSSEEDDTGFSERGILRPLSTRHGVEDDEE DEEEGEEDSSNSEDGDPDAEAGLAPGELQQLAQ GFELEDLCLSEED
3050	A	370	182	RLDRYKSPGSGSSIPAPPSHLLLYLLHIFQSTR GCGCGSRGCGSGCGGCGSSCGGCGSGCGGCGSG RGCGSGCGGCGSSCGGCGSRYVPVCCCKPVC SWVPACSTSCGSGCGSGKGGCGSGCGSGKGGCGS CGCSQSSCCKPCCSSGCGSSCSQSSCCKPCCSS GCGSSCCQSSCCKPYCCQSSCCKPCSCFSGCGSS CCQSSCYKPCCCQSSCCVPVCCQCKI
3051	A	175	4330	NIPRWNFQKGSFGVVLVHFSSEEVDMA SDSPARS LDEIDL SALRDPAGIFELVELVGNNGTYGQVYKGR HVKTGQLAAIKVMDVTGDEEEIEIKQENMLKKY SHHRNIATYYGAFIKKNPPGMDDQLWLVMFCG AGSVTDLIKNTKGYTLKEEWIAYICREILRGLSHL HQHKVIHRDIKGQNVLLTENAEVKLVDFGVSAQ LDRTVGRNRNTFIGTPYWMAPEVIACDENPDATY DFKSDLWSLGITAIEMAEGAPPLCDMHPMRALF LIPRNPAPRLKSKKWSKKFQSFIESCLVKNHSQRP ATEQLMKHPFIRDQPNRQVRIQLKDHDRTKKK RGEKDETEYBYSGSEEEEEENDSGEPSSILNLPGE STLRRDFLRLQLANKERSEALRRQLEQQQREN EEHKRQLLABRQKRIBEQKEQRRRLEEQQRREKE LRKQQEREQRRHYEEQMRREEERRRAEHEQEYI RRQLEEEQRQLEILQQQLLHEQALLLEYKRKQLE EQRQAERLQRQLKQERDYLVS LQHQRQEQR PVE KKPLYHYKEGMSPSEKPAWAKEVEERSRLNRQS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=/possible nucleotide insertion)
				SPAMPHKVANRISDPNLP RSESFSISGVQPARTP PMLRPVDPQIPHLVAVKSQGPALTASQSVHEQPT KGLSGFQEALNVTSHRVEMPRQNSDPTSENPLP TRIEKFDRSSWLRQEEDIPPKVPQRTTSISPALAR KNSPGNGSALGPRLGSQPIRASNPDLRRTEPILES PLQRTSSGSSSSSTPSSQPSQGGSQPGSQAGSSE RTRVRANSKSEGGSPVLPHEPAKVKPEESRDITRPS RPASYKKAIDEDLTALAKELRELRIETNRPMKK VTDYSSSSESESESESESESESETHDGTVAVSDI PRLIPTGAPGSNEQYNVGMVGTGLETSHADSFS GSISRBGTLMIRETSGEKKRSGHSDSNGFAGHNL PDLVQQSHSPAGTPTTEGLGRVSTHSQEMDSGTE YGMGSSTKASFTPFVDPVRYQTSPTDEDEEDEES SAAALFTSELLRQEQAQKLNARKISVVNVNPTNI RPHSDTPEIRKYKKRFNSEILCAALWGVNLLVGT ENGLMLLDRSGQGVYNLNNRRRFQQMDVLEG LNVLVTISGKKNKLRVYYLSWLRNRLHNDPEV EKKQGWITVGDLEGCIHYKVVKYERIKFLVIALK NAVEIYAWAPKPYHKFMAFKSFADLQHKPLLVD LTVEEGQRLKVIFGSHTGFHVIDVDSGNSYDIYP SHIQGNITPHAIVLPKTDGMEMLVCYEDEGVYV NTYGRITKDVVLQWGEMPTSVAYHSNQIMGW GEKAIEIRSVETGHLDGVMHKRAQLKFLCERN DKVFFASVRSGSSQVFFMTLNRNSMMNW
3052	A	1	615	MGQVECGGQKLGNOLEDDSEPAEGKVYSSDEE KLEASAGDPAGSEQEEEGSGGDSDDGFLDSSA GGPGALLGPKPKLKGSGLTGAEAGAPVTAGVTA PGGKSRRRRTAFTSEQLLELEKEFHCKKYLSTL RSQIAHALKLSEVQVKIWFQNRRAKWKRIKAGN VSSRSGEVVRNPKIVVIPVHVNRFAVRSQHQQM FOGARP
3053	A	203	2257	FGVRVPSN...HCMQITSEW DSECLTSLQP LPLPTPPAANEALQTAASLWTVAAVQAIERK VEIHSRLLHLECRGTGAEKKLASCEKTVTELG QLEGKGAVLGTLQYGLLQRRLENLENLLNR NFWILRLPPGIKGDIPKVPVAFDDVSIYFSTPEWE KLEEWQKELYKNIMKGNYESLISMDYAINQPDV LSQIQPEGEHNTEDQAGPEESEIPTDPSEEPGISTS DILSWIKQEEEPQVGAPPESKESDVYKSTYADEE LVIKAEGLARSSLCPEVPVPFSSPPAAAKDAFSDV AFKSQSTSMTPFGRPATDLPEASEGQVTFQTLG SYPLPPPVGEOVFVCHHCGKNLSQDMLLTHQCS HATEHPLPCAQCCKHFTPQADLSSTSQDHASETP PTCPHCARTFTHPSRLTYHLRVHNSTERPFPCPDC PKRFADQARLTSHRRAHASERPFRCACQGRSFS KISLLLHQRGHAQERPFSCPCGIDFNGHSALIRH QMIHTGERPYPCTDCSKSFMKEHLLNHRRLHT GERPFSCPHCGKSFIRKHHLMKHQRIHTGERPYP CSYCGRSFRYKQTLKDHLRSGHNGGCGGSDPS GQPPNPPGLITGLETSGLGVNTEGLETNQWYGE GSGGGVL
3054	A	3	2212	SCGHKSAYGSYTG LQ LFWEDGQELLQHQQLOD LRLCVHLRPQSEKVELSLWTLFVVVGKGEPSAVR EKLKGAGFAAASGPGGRPGAERASTVLNHLT AESRWEPNACNRVSSSPAGVGPLDLPVGPLLYFF

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				APWARASFLCHAFQRPLTGIGLNTVRFTSEFPLH SKDPTAHKLLFTGNYLCKLHPRPRHAPQGSLSDF CHGTEGKDLPSEHNVSEGV AQDRSPEATLCPQ KTCPCDICGLRLKDILHLAEHQTTTHPRQKPFVCE AYVKGSEFSANLPRKQVQQNVHNPIRTEEGQAS PVKTCRDHTSDQLSTCREGGKDFVATAGFLQCE VTPSDGEPHEATEGVVDHFIALRHNCCESGDAF NNKSTLVQHQRHSRERPYECSKCGIFFTYAADL TQHOKVHNRGKPYECCECGKFFSQHSSLVKHRR VHTGESPHVCGDCGKFFSRSSNLIQHKRVTGK PYECSDCGKFFSQRSNLIHHRVHTGRSAHECSE CGKSFNCNSSLIKHWVHTGERPYKNECGKFFS HIALIQHQIVHTGERPHGCGECGKAFIRSSDLMK HQRVHTGERPYECNECGKLFSSQSSSLNSHRLHT GERPYQCSECGKFFNQSSSLNNHRLHTGERPYE CSECGKTFRRQSNLRQHLKVHKPDRPYECSECG KAFNQRP TLIRHQIHIRERSMENVLLPCSQHTPE ISSENRPYQGA VNYKLKL VHPSTHPGEVP
3055	A	268	2954	ARRSSSSQGSAAPTPCQVVEASRDQLVAGPSGK MGNREMEELIPLVNRLQDAFSALGQSCLELPQI AVVGGQSAGKSSVLENFVGRDFLPRGSGIVTRRP LVLQLVTSKAEYAEFLHCKGKKFTDFDEVRLIE AETDRVTGMNKGISSIPINLRVYSPHVLNLTIDL PGITKVPVGDQPPDIEYQIRMIMQFITRENCLILA VTPANTDLANS DALKLAKEVDPQGLRTIGVITKL DLMDEGTDARDVLENKLLPLRRGYVGVVNRSQ KDIDGKKDIKAAMLAERKFFLSHPAYRIADRM GTPHLQKVLNQQLTNHIRTLPNFRNKLQGQLS IEHEVEAYKNFKPEDPTRKTKALLQM VQQFAVD FEKRIEGSGDQVDTLELSGGA KINRIFHERFPFEIV YMEFEKELPREISYALNEGPTGLFTPDMAED AIVKQIVAKKSPSLKSVDLVIELINTVACIK KLANFPRLCBETERIVANHIREREGKTKDQVLLI DIQVSYINTNHEDFIGFANAQQRSSQVHKKTIVG NQVIRKGWLTISNIGIMKGGSKGYWFLTAESLS WYKDDEEKEKKYMLPLDNLKVRDVEKSFMSK HIFALFNTEQRNVYKDYRFLELACDSQEDVDSW KASLLRAGVYPDKSVGNKAENDENGQAENFS MDPQLERQVETIRNLVDSYMSIINKCIRDLIPKTI MHLMINNVKDFINSSELLAQLYSSDQNTLMES AEQAQRDEMLRMYQALKEALGIIGDIGTATVS TPAPPPVDDSWIQHSRRSPPPSPPTTQRRPTLSAPL ARPTSGRGPAPAIPSPGPHSGAPPVPFRPGPLPPFP SSSDSFGAPPQVPSRPTRAPPSVPSRRPPSPTRPTI IRPLESLLD
3056	A	1674	1839	VVRVTCCPPARSTTERTNAYDEEDCVEMVASGG WNDVACHTTMYFMCEFDKKNM
3057	A	1674	1839	VVRVTCCPPARSTTERTNAYDEEDCVEMVASGG WNDVACHTTMYFMCEFDKKNM
3058	A	3363	2525	FLVKLILILCRCLHSLRSVQQLRTSFQDHAVWK PLMKVLQNAPEILVVASSMLCNLLLEFSPSKEPI LESGAVELLCGLTQSENPALRVNGIWALMNMAF QAEQKIKADILRSLSTEQLFRLLSDSLNVLMT LGLLRNLLSTRPHIDKIMSTHGKQIMQAVTLLEG EHNIEVKEQTLCILANIADGTTAKDLIMTNDILQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KIKYYMGHSHVKLQLAAMFCISNLIWNEEEGSQ ERQDKLRDMGIVDILHKLSQSPDSNLCDAKMA LQYYLA
3059	A	679	167	SSWPSLSSQMHPFSFHLHVAAHYGRDSFVRLLE FKAEDVPLSDKGTTPQLAIIRERSSCVKILLDH ANIDIQNGFLLRYAVIKSNHSYCRMFLQRGADTN LGRLEDGQTPLHLSALRDDVLCARMLYNYGAD TNTRNYEGQTPLAVSISISGSSRPCLDFLQEVTS M
3060	A	30	234	PPLQLDMDPNCYCADGDSCTCAGSCKCKECKCT SCKKSCCSCCPAGCAKCAQGCICKGATDKCSCC A
3061	A	428	720	VRRDVRRQATWAMASDLDFSPPEVPEPTFLENL LRYGLFLGAIFQLICVLAIIVPIPKSHEAEAEPSEPR SAEVTRKPKAAVPSVNRKPKKETKKKR
3062	A	1589	276	WKQKYEPLGLDAAGIEEAITAVGSFILKANELLQ VIDSSMKNFKAFFRWLYVAMLRMTEDHVLPELN KMTQKDIITFVAEFLTEHFNEAPDLYNRKGKYFN VERVGQYLKDEDDDLVSPNTEGNQWYDFLQN SSHLKESPLLPYYPRKSLHFVKRRMENIIDQCLQ KPADVIGKSMNQAICIPLYRDRSEDSTRLFKFP FLWNNKTSNLHYLLFTILEDLSYKMCILRRHTDIS QSVSNGLIAIKFGSFTYATTEKVRRSIYSCDLAQF YDDETVTVVLKDTVGREGRDLLVQLPLSLVYN SEDSAEYQFTGTYSTRLDEQCSAIPTRTMHFEKH WRLLESMKAQYVAGNGFRKVSCLSSNLRHVR VFEMDIDDEWELDESSDEEEASNKPKVKEEVL SESEAEENQAGAAALAPEIVIKVEKLDPELDS
3063	A	50	849	DKMPSIFAYQSSEVDWCESNFQYSELVABFYNTF SNIPFFIFGPLMMLMHPYAKQRSRYIYVWVLF MIIGLFSMYFHMILSFLGQLLDEIALWLLSGYS EMMPRCHEPSELGAMTQFIRLVFTIYACTLLSFL RPTVNAYALNSALHLLAVCQEYRKTSNKLRLH LIEVSVVLWAVALSWSIDRLLCSFWQRIHFFYL HSIWHVLISITFPYGAVTMALVDANYEMPGETL KVRYWPRDSWPVGLPIVEERGDDKDC
3064	A	1523	925	AATMADGQMPFSCHYPSRLRRDPFRDSPLSRLL DDGFGMDPFPDDLTA SWPDWALPRLSSAWPGTL RSGMVPRGPTATARFGVPAEGRTPPPFPGEVWK VCVNVHSFKPEELMVKTKDGYVEVSGKHEEKQ QEGGIVSKNFTKKIQLPAEVDPTVTFASLSPEGLL IEAPQVPPYSTFGESSFNNELPQDSQEVCT
3065	A	230	2929	LSTSLTGSHLFLSLGNHSTRENLNAGNFNFPSEGH LVRSTGPGGSFAKHMVAQCVSPKGPLACSRTYF FGATHVPYLGGSKLPPKTEQIRLLSQIYAAVIE AVLAGIACYAKTSSLTKAKEVAEQTLGSGLDSE LIPFKAALRSKMTFHHAVNNQGRIVPLDSEDSLS FVKTACMAVYDIPDLLGGNGCLGSVVFSESFLTS QILVKEKDGTVTITETSSVVLTA AVPRFCSWLVED NEVKLSEKTHQAVRGDESFLGTLYLTGGEGAYLY SSNLQSWPEEGNVHFFSSGLLFSHCRHGSIIISKD HMNSISFYDGDSTSTVAALLIDFKSSLLPHLPVHF HGSSNFLMIALFPKSKIYQAFYSEVPSLWKQDN SGISLKVIQEDGLSVEQKRLHSSAQKLFSAQSQA GEKRSSLKLLSAKLPELDWFLQHFALSSISQEPVM RTHLPVLLQQAINTTHRIESDKVIISIVTGLPGCH

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ASELCAFLVTLHKECGRWMVYRQIMDSSECFHA AHFQRYLSSALEAQNRSAQSA YIRKKTRLLV VLQGYTDVIDVVQALQTHPDSNVKASFTIGAITA CVEPMSCYMEHRFLFPKCLDQCSQGLVSNVFT SHTTEQRHPLLVLQQLIRAAANPAAAFILAENGIV TRNEDIELILSENSFSSPEMLRSRYLMYPGWYEG KLNAGSVYPLMVQICVWFGRPLEKTRFVAKCKA IQSSIKPSPFSGNIYHILGKVKFSDSERTMEVCYNT LANLSIMPVLEGPTPPDSKSVSQDSSGQCEYL VFIGCSLKEDSIKDWLRQSAKQKPQRKALKTRG MLTQQEIRSIHVKRHLEPLPAGYFYNGTQFVNF GDKTDFHPLMDQFMNDYVEEANREIEKYNQELE QQEYHDLFELKP
3066	A	130	588	LAPLRCQPGTRTQPRSHPAANDPSAAMSAAGAR GLRATYHRLDLKVELMLPEKLRPLYNHPAGPRT VFFWAPIMKWGLVCAGLADMARPAEKLSTAQS AVLMATGFIWSRYSLVIIPKNWSLFAVNFFVGAA GASQLFRIWRYNQLKAKAHK
3067	A	2	1016	EFARRRVFIAAREMSLLRSLRVFLVARTGSPYAG SLLRQSPQPRHTFYAGPRLSASASSKELLMKLR KTGYSFVNCKKALETCCGDLKQAEIWLHKEAQ KEGWSKAAKLQGRKTKEGLIGLLQEGNTTVLVE VNCETDFVSRNLKFQLLVQQVALGTMHHCQTL KDQPSAYSKGFLNSELGSLPAGPDREGSLKDQL ALAIGKLGENMILKRAAWVKVPSGFYVGSYVHG AMQSPSLHKLVLGKYGALVICETSEQKTNLEDV GRRLGQHVVGMAPLSVGLDDEPGGEAETKML SQPYLLDPSITLGQYVQPQGVSVVDFVRFECGEG BEAAETE
3068	A	3	1679	NSRVWGPWTEPSAGSLRPMARKQNRNSKELGL VPLTDDTSHAGPPGPGRALLEDHLRSGPGR KRKDWSCSLLVASLAGAAGDSFLYGYNLVSVNA PTPYIKAFYNESWERRHGRPIDDTLLWSVTV SIFAIGGLVGTILIVKMIGKVLGRKHILLANNGFAI SAALLMACSLQAGAFEMLIVGRFIMGIDGGVALS VLPMYLSEISPKEIRGSLGQVTAIFICIGVFTGQL GLPELLGKESTWPYLFQVIVVPAVVQLLSLPLP DSPRYLLEKHNEARAVKAFQTLGKADVSQEV BEVLAESRVQRSIRLVSVLELLRAPYVRWQVVT VIVTMACYQLCGLNATWIFYNSIFGKAGIPPAKIP YVTLSTGGIETLAAVFSGLVIEHLGRRPLLIGGFG LMGLFFGTLTTTLTQDHAPWVPYLSIVGILAIAS FCSGPGGIPFILTGEFFQQSQRPAAFIAGTVNWS NFAVGLLFPFIQKSLDTYCFLVFATICITGAIYLYF VLPETKNRTYAEISQAFSKRNKAYPPEEKIDSAV TDGKINGRP
3069	A	861	300	AAGAVVSAMPKAKGKTRRQKFGYSVNRKRLNR NARRKAAPRIECSHIRHAWDHAKSVRQNLAE LAVDPNRAVPLRKRKVKAMEVDIEERPKEKLRK PYVLNDLBAEASLPEKKGNLTLSDLDYVRYMV ENHGEDYKAMARDEKNYYQDTPKQIRSKINVY KRFYPAEWQDFLDSLQKRKMEVE
3070	A	325	2019	LAEPEVATDSGQQADLPAEGGDPRAEASCSVLH SKPHAMADSRDPASDQMHWKEQRAAQKADV LTTGAGNPVGDKLNVITVGRGPLLVQDVVFTD

243

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isolucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LQITHKDPKTGKLRTPALHPEQKADRYFVLYKP PPKDNIPALVEEYLERATFVANDLDWLLALPHD KFWCQVIFDETLOKCLDSYLRVPRKFDEGVAS APEVVDMQKRLHRSVFLTFLRMSTHESKDHFIS PSAFGEILYNNFLFDIPKILDLCVLFKGKNSPLLQ KMIGNIFTQPSYYSDDLDETLPILQVFSNQLHC GLQGDGANTTPQKLEERGRLTPSDMPLLELKDIV LYLCDTCTTLWAFLDIFPLACQTFQKHDFCYRLA SFYEAAIPEMESAIAKKRREDSKLLGDLWQRLSH SRKKLMEIFHILNQICLLPILESSCDNIQGFIEEFL QIFSSLLQEKRLRDYDALFPVAEDISLLQQASSV LDETRTAYILQAVESA WEGVDRRKATDAKDPSV IEEPNGEPNGVTVTAAEVSQASSHPENSEEEECM GAAAAVGPAMCGVELDSLISQVKDLLPDLGEGFI LACLEYYHYDPEQVINNLEERLAPTLSQLDRNL DREMCPDPTPLLTSRHNVFQNDDEFVFSRDSVDL SRVHKGKSTRKEENTRSLNDKRAVAAQRQRYE QYSVVVEEVPLQPGESLPYHSVYYEYDDTYD GNQVGANDADSDDELISRRPFTIPQVLRTKVPRE GQEEEDDDDEEDDADEEAPKPDHFVQDPAVLREK AEARRMAFLAKKGYRHSSTAVAGSPRGHGQS RETTQERRKKEANKATRANHNRRTMADRKRKSK GMIPS
3074	A	3	251	GEARSPPPAALLDMDPETCPCPSGGSCCTCADSC KCEGCKCTSCCKSCCSCCPAECCKCAKDCVCKG GEAAEAEEAEKSCCQ
3075	A	255	982	SQFSLSQVLVDSAEEGSLAAAAELAAQKREQL RKFREHLMRNEARKLNHQEVVEEDKRLKLPAN WEAKKARLEWELKEEEKKKECAARGEDYEKVK LLEISAEDAERWERKKRKNPDLGFSYAAAQ RQYHRLTKQIKPDMETYERLREKHGEFFPTSNS LLHGTHVPSTEEIDRMVIDLEKQIEKRDYKYSRRR PYNDDADIDYINERNAKFNKKAERFYGKYTAEI KQNLERGTA
3076	A	255	982	SQFSLSQVLVDSAEEGSLAAAAELAAQKREQL RKFREHLMRNEARKLNHQEVVEEDKRLKLPAN WEAKKARLEWELKEEEKKKECAARGEDYEKVK LLEISAEDAERWERKKRKNPDLGFSYAAAQ RQYHRLTKQIKPDMETYERLREKHGEFFPTSNS LLHGTHVPSTEEIDRMVIDLEKQIEKRDYKYSRRR PYNDDADIDYINERNAKFNKKAERFYGKYTAEI KQNLERGTA
3077	A	1	968	FLRPRRACAQLLWHPAAGMASWAKGRSYLAP GLLQGVVAIVTGGATGIGKATVKELLELSNVVI ASRKLRLKSAADELQANLPPTKQARVPIQCNIR NEEEVNNLVKSTLDTFGKINFLVNNGGGQFLSPA EHISSKGWHAVLETNLTGTFYMCKAVYSSWMK KHGGSIVNIIVPTKAGFPLAVHSGAARAGVYNLT KSLAFEWACSGIRINCVAPGVITYSQTAVENYGSW GQSFFEGSFQKIPAKRIGVPEEVSSVVCFLLSAA SPITGQSVDVDGGRSLYTHSYEVPDHDNWPKGA GDLSVVKKMKETFKEKAKL
3078	A	2	3508	FVRESGKAPVTFDDITVYLLQEEWVLLSQQKEL CGSNKLVAPLGPTVANPELFRKFGRGPEPWLGS VQGQRSLLHHHPGKKQMGYMGEMEVQGPRES

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				<p>GQSLPPQKKAYLSHLSTGSGHIEGDWAGRNRKL LKPRSIQKSWFVQFPWLIMNEEQTALFCSACREY PSIRDKRSRLIEGYTGPFKVETLKYHAKSKAHMF CVNALAARDPIWAARFRSIRDPPGDVLASPEPLF TADCFIFYPGGLGGFDSMAELLPSRAELEDPPG DGAIPAMYLDICISDLRQKEITDGIHSSSDINILYN DAVESCIQDPSAEGLSEEVVVFEELPVVFEEDVA VYFTREEWGMLDKRQKELYRDVMRMNYELLAS LGPAAPKPDLSKLERRAAPWIKDPNGPKWGKG RPPGNKKMVAVREADTQASAADSALLPGSPVEA RASCCSSSICEEGDGPRRIKRTYRPSIQRSWFGQ FPWLVIDPKETKLFCACIERPNLHDKSSRLVRG YTGPFKVETLKYHEVSKAHLRCVNTVEIKEDTPH TALVPEISSDLMANMEHFFNAAYSIAYHSRPLND FEKILQLLQSTGTVLGKYRNRCTACTQFIKYSITL KREILEDVRNSPCVSVLLDSSDASEQACVGIYR YFKQMEVKESYITLAPLYSETADGYFETIVSALD ELDIPFRKPGWVVGGLTDGSAMLSRGGLVEKF QEVIPQLLPVHCVAHRLHLAVVDACGSIDL VKK CDRHIRT VFKFYQSSNKRNLNELQEGAAPLEQEIIR LKDLNAVRWVASRRRTLHALLVSWPALARHLQ RVAEAGGQIGHRAKGMKLMRGHFHVKFCHFL LDFLSIYRPLSEVCQKEIVLITEVNATLGRAYVAL ESLRHQAGPKKEEFNASFKDGRHLHGICLDKLEVA EQRFAQDRERTVLTGIEYLQQRFDADRPPQLKN MEVFDTMAWPSGIELASFGNDDILNLARYFECSL PTGYSEEALLEEGLKLTIAQHLPFSSMLCKNALA QHCRFPLLSKLMAVVVCVPISTSCCERGFKAMN RIRTDERTKLSNEVLNMLMMTAVNGVAVTEYD PQPAIQHWYLTSSGRRFSHVYTCAQVPARSPASA RLRKEEMG...EFPRTQKPP...PFAEV...KT ...LPPPERLIYPHTSQEAPG...</p>
3079	A	343	1513	<p>FSPLERLCSLGGWGALQAGE...PSRAGCGRE GATMGCTLSAEERAALERSKAIE...KEDGISAA KDVKLLLLGAGESGKSTIVKQMKIHE...GSGED VKQYKPVVYSNTIQSLAAIVRAMDTLGIEYGDK ERKADAKMVCDDVSRMEDTEPFAELLSAMMR LWGDSGIQECFNRSREYQLNDSAKYYLDSLDRIG AADYQPTAQDILRTRVKTGIVETHFTFKNLHFR LFDVGGQSRERKKWIHCFEDVTAIIFCVALSGYD QVLHEDETTNRMHESLKLFDSSICNNKWFDTSH LFLNKKDIFEEKIKKSPLTICFPYTGPSAFTEAVA YIQAQYESKNKSAHKEIYSHVTCATDTNNIQFVF DAVTDVIIAKNLRGCGLY</p>
3080	A	41	997	<p>EARTARELTDGVTGDLTMADQPKPISPLKNLLA GGFGGVCLVFVGHPLDVKVRLQTQPPSLPGOPP MYSGTFDCFRKTLFREGITGLYRGMAAPIIGVTP MFAVCFGGFGLGKKLQQKHPEDVLSYPQLFAAG MLSGVFTTGIMTPGERIKCLLQIASSGESKYTG LDCAKKLYQEFGRGIYKGTVLTLMRDVPASGM YFMTYEWLKNIFTPEGKRVSELSAPRILVAGGIA GIFNWAVAIPPDVLKSRFQTAPPGKYPNGFRDVL RELIRDEGVTSLYKGFNAVMIRAFANAACFLGF EVAMKFLNWATPNL</p>
3081	A	3	1996	IMADMEDLFGSDADSEAERKDSDSGSDSDSDQE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				NAASGSNASGSESDQDERGDSGQPSNKEFLGDD SEDEGASHHSGSDNHSESDNRSEASERSDHEDN DPSDVDQHSGSEAPNDDDEGHRSDGGSHHSEA EGSEKAHSDDEK WGREDKSDQSDDEKIQNSDDE ERAQGSDEDKLQNSDDDEKMQNTDDEERPQLS DDERQQLSEEEKANSDDERPVASDNDDEKQNSD DEEQQLSDEEKMQNSDDERPQASDEEHRHSDD EBEQDHKSESARGSDSEDEVLRMKRKNAIASDSE ADSDTEVPKDNSGTMDLFGGADDISSGSDGEDK PPTPGQPVDENGLPQDQEEEEPIPETRIEVEIPKV NTDLGNDLYFVKLPNFLSVEPRPFDPOYYEDEFE DEEMLDEEGRTRLKLKVENTIRWRIRDEEGNEI KESNARIVK WSDGMSLSHLGNEVPDVYKAPLQG DHNHLFIRQGTGLQGQAVFKTKLTFRPHSTDSAT HRKMTLSLADRCSTQKIRILPMAGRDPECQRTE MIKKEEERLRASIRRESQRRRMREKQHQRGLSAS YLEPDRYDEEEEGEESISLAAIKNRYKGGIREERA RIYSSDSDEGSEEDKAQRLLKAKKLTSEVRPNL FNSRGLSCTQEPTALNEELTDQAGTN
3082	A	3	921	VEFCLPASADSSSLVAASLAGVRKMATNFLAHE KIWFDFKFKYDDAERRFYEQMNGPVAGASRQEN GASVILRDIARARENIQKSLAGSSGPGASSGTS HGELVVRIASLEVENQSLRGVVQELQQAISKLEA RLNVLEKSSPGHRATAPQTQHVSPMRQVEPPAK KPATPAEDDEDDIDLFGSDNEEEDKEAAQLREE RLRQYAEKKAKKPAL VAKSSILLDVKPWDDDET MAQLEACVRSIQLDGLVWGASKLVPVGYGIRKL QIQCVVEDDKVGTDLLEEEITKFEHVQSVDIAA FNKI
3083	A	3	921	VEFCLPASADSSSLVAASLAGVRKMATNFLAHE KIWFDFKFKYDDAERRFYEQMNGPVAGASRQEN GASVILRDIARARENIQKSLAGSSGPGASSGTS HGELVVRIASLEVENQSLRGVVQELQQAISKLEA RLNVLEKSSPGHRATAPQTQHVSPMRQVEPPAK KPATPAEDDEDDIDLFGSDNEEEDKEAAQLREE RLRQYAEKKAKKPAL VAKSSILLDVKPWDDDET MAQLEACVRSIQLDGLVWGASKLVPVGYGIRKL QIQCVVEDDKVGTDLLEEEITKFEHVQSVDIAA FNKI
3084	A	128	4050	KSIVKIRKRMAAETQTLNFGPEWLRALSSGGSITS PPLSPALPKYKLADYRYGREEMLALFLKDNKIPS DLDDKEFLPILQEEPLPLALVPFTEEEQRNFMS VNSAAVLRLTGRGGGTVVGAPRGRSSSRGRGR GRGECGFYQRSFDEVEGVFGRGGGREGMHRSQS WEERGDRRFEKPGRKDVGRPNFEEGGPTSVGRK HEFIRSESENWRIFREEQNGEDEDGGWRLAGSRR DGERWRPHSPDGRSAGWREHMERRRRFEFDFR DRDDERGYRRVRSRSGSIDDDRDSLPEWCLEDA BEEMGTFDSSGAFLSLKKVQKEPIEEQEMDFRP VDEGECSDSSESHNEEAKPEPKTNKKEGEKTD RVGVEASEETPQTSSSSARPGTPSDHQSQEASQFE RKDEPKTEQTEKAEETRMENSLPAKVPSRGDE MVADVQQPLSQIPSDTASPLLLPPVPNPSPTLRP VETPVVGAPGMGSVSTEPDDEEGLKHLEQQAQEK MVAYLQDSALDDERLASKLQEHRAKGVSIPLMH

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				<p>EAMQKWYYKDPQGEIQGPFNNQEMAEWFQAG YFTMSLLVKRACDESFQPLGDIMKMWGRVPFSP GPAPPPHMGELDQERLTRQQELTALYQMQHLQY QQFLIQQQYAQVLAQQQKAALSSQQQQQLALLL QQFQTLKMRISDQNIIPSVTRSVSPDTGSIWELQ PTASQPTVWEGGSVWDLPLDTTTPGPALEQLQQ LEKAKAAKLEQERREAEMRAKREEEERKRQEEL RRRQKGILRRQEEERKRREEEELARRKQEEALR RQREQEIALRRQREEEERQQEEALRRLEERRRE EEERRKQEELLRKQEEAAKWAREEEEAQRRLE ENRLRMEEEAARLRHEEEERKRKELEVQRQKEL MRQRQQQEEALRRLQQQQQQQLAQMKLPSSS TWGQQSNTTACQSQATLSLAEIQKLEEEERERQLR EEQRRQQRELMKALQQQQQQQQQKLSGWGNV SKPSGTTKSLLEIQEEARQMOKQQQQQQQHQQ PNRARNNTHSNLHTSIGNSVWGSINTGPPNQWA SDLVSSIWSNADTKNSNMGFWDADVKEVGPRN STNKNKNNASLSKSVGVSNRQNKKEVEEELKLLK LFQGVNKAQDGFQWCEQMLHALNTANNLDVP TFVSFLKEVESPYEVHDIYRAYLGDTSBAKEFAK QFLERRAKQKANQQRQQQLPQQQQPPQPPQ QQPQQQDSVWGMNHSTLHSVFQTNQSNQSSN FEAVQSGKKKKKQKMVRADPSLLGFSVNASSER LNMGEIETLDDY</p>
3085	A	128	4050	<p>KSIVKIRKRMMAETQTLNFGPEWLRALSSGGSITS PPLSPALPKYKLADYRYGREEMLALFLKDNKIPS DLLDKEFLPILQEEPLPPLALVPFTEEEQRNFSMS VNSAAVLRLTGRGGGGTVVGA PRGSSSRGRGR GRGECGFYQRSFDEVEGVFGRGGGREGMHRSQS WEERGDRRFEKPGRKDVGRPNFEEGGPTSVGRK HEFIRSESENWRIFREEQNGDEDCWRIACETD DGEKWRPAGLQPRSAWREHMERRRRLFLTR DRDDERGYKRVRS GSGSIDD DRDSLPEWCLDA EEEMGTFDSSGAFLSLKKVQKEPIEEQEMDFRP VDEGEECSDSEGSHNEEAKDPDKNKKEGEKTD RVGVEASEETPQTSSSSARPGTPSDHQSQEASQFB RKDEPKTEQTEKABBE TRMENS LPAKVPSRGDE MVADVQQPLSQIPSDTASPLLLPPVPNPSPTLRP VETPVVGAPGMGSVSTEPDDEGLKHLEQQA EK MVAYLQDSALDDERLASKLQEHRAKGVSIPLMH EAMQKWYYKDPQGEIQGPFNNQEMAEWFQAG YFTMSLLVKRACDESFQPLGDIMKMWGRVPFSP GPAPPPHMGELDQERLTRQQELTALYQMQHLQY QQFLIQQQYAQVLAQQQKAALSSQQQQQLALLL QQFQTLKMRISDQNIIPSVTRSVSPDTGSIWELQ PTASQPTVWEGGSVWDLPLDTTTPGPALEQLQQ LEKAKAAKLEQERREAEMRAKREEEERKRQEEL RRRQKGILRRQEEERKRREEEELARRKQEEALR RQREQEIALRRQREEEERQQEEALRRLEERRRE EEERRKQEELLRKQEEAAKWAREEEEAQRRLE ENRLRMEEEAARLRHEEEERKRKELEVQRQKEL MRQRQQQEEALRRLQQQQQQQLAQMKLPSSS TWGQQSNTTACQSQATLSLAEIQKLEEEERERQLR EEQRRQQRELMKALQQQQQQQQQKLSGWGNV SKPSGTTKSLLEIQEEARQMOKQQQQQQQHQQ</p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				PNRARNNTHSNLHTSIGNSVWGSINTGPPNQWA SDLVSSIWSNADTKNSNMGFWDVAKEVGPRN STNKNKNNASLSKSVGVSNRQNKKBEEEEKLLK LFQGVNKAQDGFQWCEQMLHALNTANNLDVP TFVSFLKEVESPYEVHDYIRAYLGDTSEAKEFAK QFLERRAKQKANQQRQQQLPQQQQQPPQQPP QQPQQQDSVWGMNHSTLHSVFQTNQSNQSN FEAVQSGKKKKKQKMVRADPSLLGFSVNASSER LNMGEIETLDDY
3086	A	675	1334	LHPAATSTAWLHVPPGLSMALSWVLTVLSLLPL LEAQIPLCANLVPVPITNATLDRITGKWFYIASAF RNEEYNKSVQEIQATFFYFTPNKTEDTIFLREYQT RQDQCIYNTTYLNVQRENGTISRYYVGGQEHFAH LLILRDTKTYMLAFDVNDEKNWGLSVYADKPET TKEQLGEFYEALDCLRIPKSDVVYTDWKDKCE PLEKQHEKERKQEEGES
3087	A	1	1575	CTPVARSMATTATCTRTDDYQLFEELGKGAFS VVRRCVKKTSTQEYAAKIINTKKLSARDHQKLE REARICRLKHPNIVRLHDSISEEGFHYLVFDLVT GGELFEDIVAREYYSEADASHCIHQILES VNHIHQ HDIVHRDLKPENLLASKCKGA AVKLADFGLAIE VQGEQQA WFGFAGTPGYLSPEVLRKDPY GKPVD IWACGVILYILLVGYPPFWDEDQHKLYQQIKAG AYDFPSPEWDTVTPEAKNLINQMLTINPAKRITA DQALKHPWVCQRSTVASMMHRQETVECLRKFEN ARRKLGAILTMTL VSRNFSAKSLLNKKS DGG VKPQSNKNKNSLVSPAQEPAPLQTAMEPQTTVVH NATDGIKGSTESCNTTTEDEDLKVRKQEIHKITEQ LIEAINNGDFEAYTKICDPGLTSFEPEALGNLVEG MDFHKFYFENLLSKNSKPIHTTILNPHVHVIGED AACIAYELTYNYDGGGRPITSQSTTRVWTRRD C WLVNHYHUSGAPAAPLQ
3088	A	12	1039	SSVAEFPERVQLSQPNWNFGAGGAWSLDFAE C LKWSAELARLGESIMDGKQGGMDSKPA GPR DEPGIRLLSNPLMGDAVSDWSPMHEAAIHGHQ SLRNLISQGWAVNIITADHVSPLHEACLGHLSC VKILLKHGAQVNGVTADWHTPLFNACVSGSWD CVNLLQHGASVQPESDLASPIHEAARRGHVEC VNSLIAYGNIDHKISHLGTPLYLACENQQRACV KKLLESGADVNQGGKQDSPLHAVARTASEELAC LLMDFGADTQAKNAEGKRPVELVPESPLAQLF LEREGPPSLMQLCRLRIRKCFGIQQHHKITKL VLP EDLKQFLLHL
3089	A	73	432	DMAGLMTIVTSLFLGVCAHHIPTGSVVLSPCC MFFVSKRIPENRVVSYQLSSRSTCLKAGVFTTKK GQQFCGDPKQEWVQRYMKNLDAKQKKASPR RAVAVKGPVQRYPGNQTTTC
3090	A	4627	611	LMEAGGGGGALPAGVETMVLTLGESWPVLVGR RFLSLSAADGSDGSHDSWDVERVAEWPWLSGTI RAVSHTDVTKKDLKVCVEFDGESWRKRRWIEV YSLLRRAFLVEHNLVLAERKSPEISERIVQWPAIT YKPLLDKAGLSITSVRFLGDQQRVFLSKDLLKP IQDVNSLRSLTDNQIVSKEFQALIVKHLDESHLL KGDKNLVGSEVKIYSLDPSTQWFSATVVNGNPA SKTLQVNCEEIPALKIVDPSLIHVEVVHDNLVTC

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GNSARIGAVKRKSENNGTLVSKQAKSCSEASPS MCPVQSVPTTVFKEILLGCTAATPPSKDPRQQST PQAANSPPNLGAKIPQGCHKQSLPEEISSCLNTKS EALRTKPDVCKAGLLSKSSQIGTGDLKILTEPKGS CTQPKTNTDQENRLESVPQALTGLPKECLPTKAS SKAELEIANPPELQKHLEHAPSPSDVSNAPVKA GVNSDSPNNCSGKKVEPSALACRSQNLKESSVK VDNESCCSRSNNKIQNAPSRKSVLTDPAKLKKLQ QSGEAFVQDDSCVNIVAQLPKCRECRDLSLRKD KEQQKDSPVFCRFFHFRRLQFNKHGVL RVEGFLT PNKYDNEAIGLWLPLTKNVVGIDLD TAKYILANI GDHFCQM VISEKEAMSTIEPHRQVAWKRAVKG VREMCDVCDTTIFNLHWVCPRCGFGVCVDCYR MKRKNCQQGAA YKTF SWLCKCVKSQIHEPENLM PTQIIPGKALYDVGDIVHSVRAKWGKANCPCSN RQKLFPSKPASKEDLKQTS LAGEKPTLGAVLQ NPSVLEPAA VGGEAASKPAGSMKPACPASTSPLN WLADLTSGNVNKENKEKOPTMPILKNEIKCLPPL PPLSKSSTVLHTFNSTILTPVSNNNSGFLRNLLNSS TGKTENGLKNTIPKILDDIFASLVQNKTTSDLSKR PQGLTIKPSILGFDTPHYWLC DNRLCLQDPNNK SNWNVFRECWKQGPVMVSGVHHKLNSELWK PESFRKEFGQEVDLVNCR TNEITGATVGDFWD GFEDVPNRLKNEKEPMVLKLDWPPGEDFRDM MPSRFDDL MANIPLPEYTRRDGKLN LASRLPNYF VRPDLGPKMYNAYGLITPEDRKYGTTNLHLDVS DAANVMVYVGIPKGQCEQEEVLKTIQDGDSD LTIKRFIEGKEKPGALWHIYAAKDTEKIREFLKK VSEEQGOENPADHDPHDQSWYLD RSLRKLHQ EYGVQGWAI VQFLGDVVF PAGAPHQVHNLYSC KVAEDFVSPEHV KHCFFV EFRY SQTH DKLQVKNV... VKDAVAMLKASESSFG...
3091	A	97	1838	KRGARRGGWKRKMPSTDLLMKAFEPYLE... YSTKAKNYVNGHCTKYEPWQLIAWSVWTL... VWGYEFVFPESLWSRFKKKCFKLTRKMPIGRK IQDKLNKTKDDISKNMSFLKVDKEYVKALPSQG LSSSAVLEKLKEYSSMDAFWQEGRASGTVYSGE EKLTELLVKAYGDFAWSNPLHPDIFPGLRKIEAEI VRIACSLFNGGPDSCGCVTSGGTESILMACKAYR DLAFEKGIKTPEIVAPQSAHA AFNKAASYFGMKI VRVPLTKMMEVDVRAMRRAISRNTAMLVCSTP QFPHGVIDPVPEAKLAVKYIPLHVDACLGGFL IVFMEKAGYPLEHPDFR VKGVT SISADTHKYGY APKGSSLVLYSDKKYRNYQFFVDTDWQGGIYAS PTIAGSRPGGISAACWAALMHFGENG YVEATKQI IKTARFLKSELENIKGIFVFGNPQLSVIALGSRDFD IYRLSNLMTAKGWNLNQLQFPFSIHFCITLLHAR KRVAIQFLKDIRESVTQIMKNPKAKTTGMGAIYG MAQTTVDRNMGAELSSVFLDSL YSTDVTVQGSQ MNGSPKPH
3092	A	79	2652	LCSQNSPEDWVNFSEKQKRYPWYWTGRKLRSE RAMKIQKKLTGCSRLMLLCLSLLELLLEAGAGNIH YSVPEETDKGSFVGNIADLGLQPQELADGGVRI VSRGRMPLFALNPRSGSLITARRIDREELCAQSM PCLVSFNILVEDKMKLFPVEVEIIDINDNT PQFQL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EELEFKMNEITTPGTRVSLPFGQDLVDVGMNSLQS YQLSSNPHFSLDVQQGADGPQHPEMVLQSPDR EEEA VHHLILTASDGGEPVRSGLRIYIQVVDAN DNPPAFTQAQYHINVPENVPLGTQLLMVNATDP DEGANGEVITYSFHNVDHRVAQIFRLDSYTGEISN KEPLDFEYKMYSMEVQAQDGAGLMAKVKVL KVLVDVNDNAPEVTITSVTTAVPENFPPTIALISV HDQDSGDNGYTTCTFIPGNLPFKLEKLVNYYRL VTERTLDRRELISGYNITTTAIDQGTALSTETHISL LVTDINDNSPVFHQDSYSA YIPENNPRGASIFS AHDLSNENAQITYSLIEDTIQGAPLSAYLSINSD TGVLYALRSFDYEQFRDMQLKVMARDSGDPPLS SNVLSLSLFLDQNDNAPEILYPALPTDGTGVEL APRSAEPGYLVTKVVA VDRDSGQNAWLSYRL KASEPGLFSVGLHTGEVTRARALLDRDALKQSL VVAVQDHGQPPLSATVILTVAVADRIDLADLG SLEPSAKPNDSDLTYLVVAEAAVSCVFLAFVIV LLAHLRLRRWHKSRLQASGGGLASTPGSHFVGV DGVRAFLQTYSHVSLTADSRKSHLIFPQPNYAD TLISQESCEKKGFLSAPQSLLEDKKEPFSQVNFCD ECISYLEKNNS
3093	A	1	3868	PPDNQKLGLLEALLKIGDWQHAQNMIDQMPYY AASHKLIALAICKLIHITIEPLYRSVTSWAVDHAG FLESDPCDSTVGHLLSRVGVPGKAGKSPVNALQ NKRAPKQAESFEDLRRDVFNMF CYLGPLSHDPI LPAKVVRIGKSFMKFQSDGSKQEDKEKTEVILS CLLSITDQVLLPSLSLMDCNACMSEELWGMFKT FPYQHR YRLYGQWKNETYN SHPLLKVKQAQTD RAKYIMKRLTKENVKPSGRQIGKLSHSNPTILFD YVCFEILSQIKYDNLITPVVDSLKYLTSLNYDVL A TILSNCTEALANPEKERAHNDYTTISSW LQSLA SFCGA Y KNTDOLAGLLQYVANQLKAGKSF DL LILKEVVQFMAGIEITEEMTMEQLEAMTGGEQL KAEGGYFGQIRNTKKSSQRLKDALLDHDALPL CLLMAQQRNCFQEGGEKHLKLVGKLYDQCH DTLVQFGGFLASNSTEDYIKRVPSIDVLCNEFHT PHDAAFFLSRPMYAHHISKYDELKKSEKSGKQ QHKVHKYITSCEMVMA PVHEAVVSLHVSKVWD DISPQFYATFWSLTMYDLAVPHTSYEREVNKLK VQMKAIIDNQEMPPNKKKKEKERCTALQDKLL BEEKKQMEHVQRVLQRLKLEKDNWLLAKSTKN ETITKFLQLCIFPRCIFSAIDAVYCARFVELVHQ KTPNFSTLLCYDRVFSDIITYTVASCTENEASRYGR FLCCMLETVTRWHSRATYEKECGNYPGFLTIL RATGFDGGNKADQLDYENFRHV VHKWHYKLT KASVHCLETGEYTHIRNILLVLT KILPWYPKVLNL GQALERRVHKICQEEKEKRPDLALAMGYSGQL KSRKSYM PENEFHHKDPPPRNAVASVQNGPGG GPSSSSIGSASKSDESSTEETDKSRERSQCGVKAV NKASSTPKGNSSNGNSGNSNKA VKENDKEG KEKEKEKKEKTPATTPEARVLGKDGKEKPKBER PNKDEKARETKERTPKSDKEKEKFKKEEKAKDE KFKTTVPNAESKSTQEREREKEPSRERDIAKEMK SKENVKGGEKTPVSGSLKSPVPRSDIPEPEREQKR RKIDTHPSPSHSSTVKDSLIELKESSAKLYNHTPP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				PLSKSKEREMDKKDLKSRERSREREKKDEKDR KERKRDHSNNDREVPPDLTKRRKEENGTMGVSK HKSESPCESPYNEKDKEKNKSKSSGKEKGSDFS KSEKMDKISSGGKESRHDKEKIEKKEKRDSSGG KEEKKHHKSSDKHR
3094	A	2	891	AMLGTRPSRRGAGAVQAEVSERLAMAGPQQQ PPYLHLAELTASQFLEIWKHFDADGNGYIEGKEL ENFFQELEKARKGSGMMSKSDNFGEKMKEFMQ KYDKNSDGKIEMAELAQLPTEENFLLCFRQHVG SSAEFMEAWRKYDTRSGYIEANELKGFLSDLL KKANRPYDEPKLQEYQTILRMFDLNGDGKGLGL SEMSRLLPVQENFLLKFQGMKLTSEEFNAIFTFY DKDRSGYIDEHELDALLKDLYEKNKKEMNIQQL TNYRKSVMSLAEAGKLYRKDLEIVLCSEPPM
3095	A	1685	700	RRPTGRPGALGAPAAAGRVMPLHVKWPFAVPP LTWTLASSVVMGLVGTYSCFWTKYMNHLTVHN REVLIELIEKRGPATPLITVSNHQSCMDPHLWG ILKLRIHWNLKLMRWTPAAADICFTKELHSHFFS LGKCVPVCRGAEFFQAENEGKGVLDTGRHMPG AGKRREKGDGVYQKGMDFILEKLNHGDWVHIF PEGKVNMSSEFLRFKWGIGRLIAECHLNPIILPW HVGMDNDVLPNSPPYFPRFGQKITVLIGKPFSAFP VLERLRAENKSAVEMRKALTDIFIQEEFQHLKTQ AEQLHNHLQAWIEGLACCLLDSPAQSWG
3096	A	6642	4022	FVPGLEPQWEPAPQPSATMSAPSEEEYARLVM EAQPEWLRAEVKRLSHELAETTREKIQAAEYGL AVLEEKHQLKLQFEELEVDYEAIRSEMEQLKEAF GQAHTNHKKVAADGESREESLIQESASKEQYYV RKVLELQTELKQLRNVLTNTQSENERLASVAQE LKEINQNVIEIQRGRLRDDIKEYKFRFARLLQDYS FTEENISLQKQVSVLRONQVEFEGTEIFTL EETBYLNSQLEDARLKQLEBALETIKTER EQKNSLRKELSHYMSINDSFYTSHLHVSLDGLKF SDDAAEPNNDAAELVNGFEHGGGLAKLPLDNKTS TPKKEGLAPPSLVSDLLSELNISEIQKLKQQLM QMEREKAGLLATLQDTQKQLEHTRGSLSEQQEK VTRLTENLSALRRLQASKERQTALDNEKDRDSH EDGDYIEVDINGPEILACKYHVAVAEAGELREQ LKALRSTHBAREAQHAEEKGRYEAEGQALTEKV SLLEKASRQDRELLARLEKELKKVSDVAGETQG SLSVAQDELVTFSSEELANLYHHVCMCNNETPNR VMLDYYREGQGGAGRTSPGGRTSPEARGRRSPI LLPKGLLAPEAGRADGGTGDSSPSGSSLPSPSLD PRREPMNIYNLIAIRDQIKHLQAAVDRTTSLSRQ RIASQELGPAVDKDKEALMBEILKLKSLSTKRE QITTLRTVLKANKQTAVALANLKSKEYENKAM VTETMMKLRLNELKALKEDAATFSSLRAMFATRC DEYTTQLDEMQRQLAAAEDEKKTLSLLRMAIQ QKLALTQRLELELDHEQTRRGRAKAAPKTKPA TPSVSHTCACASDRAEGTGLANQVFCSEKHSIYC D
3097	A	1	879	MVKVVPATRGNLPRSQLTGTHQHCQPREPKITA SERLRRRPRTARLRAHAAPPEPLAVFAPPSDR KELLALPVACDPVIASVMSWVQAASLIQGGPGDK GDVFDEEADESLLAQREWQSNMQRRVKEGYRD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GIDAGKAVTLQQGFNQGYKKGADEVILNYGRLRG TSLALLSWCHLHNNNSTLINKINLLDAVGQCEE YVLKHLKSITPPSHVVDLLDSIEDMDLCHVPAE KKIDEAKDERLCENNAEFNKNCSSKSHSGIDCSYV ECCRTOEHAHSGKPKPHMDFGTDTSQF
3098	A	2	505	GAATLLRSASSAARKAAEAEQVWLHLHRYLSA DRRVLGLREWGRPASERECSLCQRLKRELNMGD VEKGKKIFIMKCSQCHTVEKGGKHKTGPNLHGL FGRKTGQAPGYSYTAANKNKGIIWGEDTLMEYL ENPKKYIPGTMIFVGIIKKKEERADLIAYLKKAT NE
3099	A	144	1386	WAVGQARSFSPSHPRMSSWTWSRRWSPSVALRVT CTSTSSQRWTVLALSKPGSQQVSMHTPAPGPPT AGHTEPPSEPPRRARVAKYRAKFDPRVTAKYDIK ALIGRGSFSR VVRVEHRATROPYAIKMIETKYRE GREVCESELRLVRRVRHANIIQLVEVFETQERVY MVMELATGGELFDRIIAKGSFTERDATRVLQMV LDGVRYLHALGITHRDLKPENLLYHPGTDSKI TDFGLASARKKGDDCLMKTTCGTPEYIAPEVLV RKPYTNSVDMWALGVIAIYLLSGTMPFEDDNR RLYRQILRGKYSYSGEPWPSVSNLAKDFIDRLT VDPGARMTALQALRHPWVVSMAASSSMKNLHR SISQNLKRASSRCQSTKSAQSTRSSRSTRSNKSR RVRERELREL
3100	A	3	1500	ARWNGRWVQVPAWPGPGCGTNASGERQRQLPR AWRPVGRTLGEPIALAWSPPLYLFPIPLPSWAVS QPTPTLGTMFADLDYDIEDDKLGITVPGKVTLQ KDAQNLIGISIGGGAQYCPCLYIVQVFDNTPAAL DGTVAAGDEITGVNGRSIKGKTKVEVAKMIEV KGEVTHYINKLQADPKQGMSLDIVLKKVKHRLV ENMSEETADALGLSAILCNDGLVKRLEELERT ELYKSHLELHNNLLRAFYELSQTHRNGIFQSC AFGDVFSIGVREPQPAASEAFVKFADAHRSIEK FGIRLLKTEFMTDLNTYLNKAIPDTRLTIKKYL DVKFEYLSYCLIKYKEMDDEEYSICALGEPLYRV STGNYEYRLILRCRQEARARFSQMRKDVLEKME LLDQKHVQDIVFQLQRLVSTMSKYYNDCYAVLR DADVFPFIEVDLAHTTLAYGLNQEEFTDGESEEEE EDTAAGEPSRDTRGAAGPLDKGGSWCDS
3101	A	1173	197	QGMDSKQQCVKLNLDGHFMPVLGFGTYAPPEVP RSKALEVTKLAIAGFRHIDSAHLYNNEEQVGLA IRSKIADGSVKREDIFYTSKLWSTFHRPELVRPAL ENSLKKAQLDYVDLYLIHSPMSLKPGEELSPDE NGKVIFDIVDLCTTWEAMEKCKDAGLAKSIGVS NFNRRQLEMILNKPLKYKPV CNQVECHPYFNR SKLLDFCKSKDIVLVAYSALGSQRDKRWDPNS PVLLDPVLCALAKKHKRTPALIALRYQLQRGV VVLAKSYNEQRIQNVQVFELQTAEDMKAIDG LDRNLHYFNDSFASHPNYPYSDEY
3102	A	144	1098	EQRPPTPCGRRPLPLGSAPCRVLRAPRQAPAM SMLPSFGFTQEQVACVCEVLQGGNLERLGRFL WSLPACDHLHKNESVLKAKAVVAFHRGNFREL YKILESHQFSPHNPKLQQLWLKAHYVEAEKLR GRPLGAVGKYRVRQKFPLPRTIWDGEETSYCFK EKS RGVLEWYAHNPYPSPREKRELAETGLTT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TQVSNWFKNRRQRDRAAEAKERENTENNSSSN KQNQLSPLEGGKPLMSSSEEEFSPQSPDQNSVLL LQGNMGHARSSNYSLPGLTASQPSHGLQTHQHQLQD SLLGPLTSSLVDLGS
3103	A	111	1582	LVYSWGCHIMADNDTDRNQTEKLLKRVRELEQ EVQRLKKEQAKNKEDSNIRENSSGAGKTKRAFD FSAHGRRHVALRIAYMGWGYQGAFSSENTNTI EEKLFEALTKTRLVESRQTSNYHRCGRDCKGVS AFGQVISLDLRSQFPGRDSEDFNVKEEANAAAE EIRYTHILNRVLPDIRILAWAPVEPSFSARFSCLE RTYRYFFPRADLDIVTMDYAAQKYVGTHDFRNL CKMDVANGVINFORILSAQVQLVGQSPGEGRW QEPFQLCQFEVTGQAFLYHQVRCMMAILFLIGQ GMEKPEIIDELLNIEKNPQKPQYSMAVEFPLVLY DCKFENVKWTYDQEAQEFNITHLQQLWANHAV KTHMLYSMLQGLDTPVPVPCGIGPKMDGMTWEG NVKPSVIKQTSFVEGVKMRITYKPLMDRPKCQG LESRIQHFVRRGRIEHPHLFHEEETKAKRDCNDT LEEDNTNLETPTKRVCDTEIKSII
3104	A	227	1519	VTLKMNAMLETPELPAVFDGVKLAAVAALVLY IVRCLNLKSPTAPPDLYFQDSGLSRFLKSCPLLT KEYIPPLIWGKSGHIQTALYGKMGRVRSPPHYGH RKFITMSDGATSTFDLFEPLAEHCVGDDITMVICP GIANHSEKQYIRTFVDYAQKNGYRCAVLNHLGA LPNIELTSPRMFTYGCTWEFGAMVNYIKKTYPLT QLVVVGFSLGGNIVCKYLGETQANQEKVLCVS VCQGYALRAQETFMQWDQCRRFYNFLMADN MKKIILSHRQALFGDHVKKPQSLEDTLSRLYTA TSLMQIDDNVMRKFHGYNSLKEYEEESCMBRYL HRIYVPLMLVNAADDPLVHESLLTIPKSLSEKRE NVVGVPLHGGHLCFFEGEVLFPFPLTWMT VEYANAICQWERNKLCQSDTEADLE
3105	A	1	1211	MGLLLMLASAVLGSFLTLLAQFFLYRRQPEPP ADEAARAGEGFRYIKPVPGLLLRELYYGGGRDE BPSGAAPEGGATPTAAPETPAPPTRETCYFLNATI LFLFRELRDTALTRRWVTKKIKVEFEBLLQTKTA GRLLEGLSLRDVFLGETVPFIKTIRLVRPVPSAT GEPDGPEGEALPAACPEELAFBAEVEYNGGFHLA IDVDLVFGKSAYLFVKLSRVVGRRLRVFTRVPFT HWFFSFVEDPLIDFEVRSQFEGRPMPQLTSIIVNQ LKKIIRKHTLPNYKIRFKPFFPYQTLQGFEEDEE HIHIQQWALTEGRLKVTLLECSRLIFGSYDREA NVHCTLELSSSVWEEKQRSSIKTGITSLTAVFMG WHRVSEAFPLWYKLLVDLPFWGLEDGGPLLT VPLRQCPG
3106	A	972	468	MAAAGAGRLRRVASALLRSPRLPARELSAPAR LYHKKVVDHYENPRNVGSLDKTSKNVGTGLVG APACGDVMKLIQVDEKGIKVDARFKTFGCGSA IASSSLATEWVKGKTVEBALTIKNTDIAKELCLPP VKLHCSMLAEDAIAALADYKCLKQEPKKGEAE KK
3107	A	106	1221	TCQDVRSVFSLVANIFGEESTAGAGWHREEDM RKELQLSLSVTLLLVCGLYQFTLKSSCLFCLPSF KSHQGLEALLSHRRGIVFLETSEMEPHLVSCS VESAAKIYPEWPFVFFMKGLTDSTPMPSNSTYPA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				FSFLSAIDNVFLFPLDMKRILLEDTPLSWYNQINA SAERNWLHISSDASRLAIWKYGGIYMDTDVISIR PIPEENFLAAQASRYSSNGIFGLPHHPFLWECME NFVEHYNSAIWGNQGPPELMTRMLRVWCKLEDF QEVSDLRCLNISFLHPQRFYPISYREWRRYYEVW DTEPSFNVSALHLWNHMQEGRAVIRGSNTLV ENLYRKHCPRTYRDLIKGPEGSVTGELGPGNK
3108	A	1612	839	EVALFCFEMAAGMYLEHYLDSIENLPFELQRNFQ LMRDLQRTEDLKAIDKLATEYMSSARLSSEE KLALLKQIQEAYGKCCEFGDDKVQLAMQTYEM VDKHIRRLDTDLARFEADLKEKQIESSDYDSSSS KGKKKGRTQKEKKAARARSKGKNSDEEAPKTA QKKLKLVRTSPEYGMPSVTFGSVHPSDVLDMFV DPNEPTYCLCHQVSYGEMIGCDNPDCSIEWFHFA CVGLTTKPRGKWFCPRCSQERKKK
3109	A	1	2613	MVAVRAAGPREGASQDEAGTVWAPMTGCPCQC RPGPSWLLVDTLEPETAYPVQRPGEQAGNQRL QMKRAQFGPHDWLSLPVPPGPSWLLVDTLEPET AYQFSVLAQNKLGTSAFSEVTVNTLAFPIITPEP LVLVTPPRCLIANRTQQGVLLSWLPPANHSFPIDR YIMEFRVAERWELDDGIPGTEGEFFAKDLSQDT WYEFRLVAVMQDLISEPSNIAGVSSTDIFPQDLT EDGLARPVLGIVATICFLAAAILFSTLAACFVNK QRKRKLKRKKDPPLSITHCRKSLESPLSSGKVSPE SIRTLRAPSESSDDQGQPAKRMLSPTREKELSL YKKT KRAISSKKYSVAKAEAEAEATTPIELISRGF DGRFVMDPAEMEPLSKSRRIEGFPFAEETDMYPE FRQSDNEEDPLVPTSVAAALKSQLTPLSSSQESYL PPPAYSPRFQPRGLEGGLEGRLQATGQARPPA PRPFHHGQYYGYLSSSSPGEVEPPFFYVPEVGSPL SSWITSPPLPTGPGFCTETENGENACISTITLT QITGGRSPEPWGRMPPFQLETPAMMFHQLP PCDVPESLQPKAGLPRGLPPTSLQVPAAYPGILSL EAPKGWAGKSPGRGPVFAFPAKWQDRPMOPL VSQQLRHTSQGMGIPVLPYFEPAEPGAHHGPST FGLDTRWYEPQPRPRPSRQARRAEP SLHQVVLQ PSRLSPLTQSPSSRTGSPELAARARPRPGLLQQA EMSEITLQPPAAVSFSRKSTPSTGSPSSRSRSGSPS YRPAMGFTTLATGYSPPPGPAPAGPGDSLDFVG QTPSPRRTGEELLRPETPPPTLPTLGKLRRDRPAP ATSPPERALSKL
3110	A	88	924	ILGSRTMSLTNTKTGFSVKDILDLPDNTDEEGSV AEGPEEENEGPEPAKRAGPLGQGDALDAVQSLPL KNPFYDSSDNPYTRWLASTEGLQYSLHGLAAGA PPQDSSSKSPEPSADESPDNDKETPGGGGDAGKK RKRRVLFSKAQTYELERRFRQRYLSAPEREHLLA SLIRLTPTQVKIWFQNHRYKMKRARAEEKGMEVT PLPSRRVA VPVLVRDGKPCHALKAQDLAAATF QAGIPFSA YSAQSLQHMQYNAQYSSASTPQYPT AHLVQAQQWTW
3111	A	595	291	PSVASLARRFSGRALWPPSHSVPGNRALCPRLLH GTTLPGGNQRELARQKNMKQSDSVKGKRRDD GLSAAARKQORDSTPRDSEIMQKQKKANEKKEE PK
3112	A	3641	1555	APMLQIHHSFKLIFQNIHKSFKISQRLSQNADST

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RHTNLSNTHYSDLIVWNCCLFFRNWCNEFFLKS CHFAQEREGSGDLCSRAEKTSAACVIFRRFPV APLIPYPLITKEDINAIEMEEDKRDLSREISKFRDT HKKLEEEKGKKEKERQEIEKERRERERERERERE RREREREREREREREKEKERERERERDRDRDRTK ERDRDRDRERDRDRDRERSSDRNKDRSRSREKS RDRERERERERERERERERERERERERERERE REREKDKKRDREDEEDA YERRKLERKLREKEA AYQERLKNWEIRERKKTREYEKEAEREEERRR MAKEAKRLKEFLEDYDDDRDDPKYYRGSALQK RLRDREKEMEADERDRKREKEELEIRQRLAE GHPDPAELQRMEQEAERRRQPQIKQEPESEEEE EEKQEKEEKREBPMEEBEPEQKPKLPKTLRPISS APSVSSASGNATPNTPGDESPCGIIPHENSPDQ QPEEHRPKIGLSLKL GASNSPGQPN SVKRKKLPV DSVFNKFEDSDDDVPRKRKL VPLDYGEDDKNA TKGTVNTTEKRRKHIKSLIEKIPTAKPELFAYPLDW SIVDSILMERRIRPWINKKIEYIGEEATLVDLVC SKVMAHSPQSI LDDVAMVLDEEA EVFVKMWR LLIYETEA KKIGLVK
3113	A	1	669	VCAGIRDPCSTPLAKPAAGGAENLSFGKQPGLET NILKMTTPNKTTPGADPKQLERTGTVREIGSQAV WSLSCKPGFGVDQLRDDNLETYWQSDGSPHL VNIQFRRKTTVKTL CIYADYKSDESYTPSKISVRV GNNFHNLQEIRQELVEPSGWIHVPLTDNHKKPT RTFMQIAVL ANHQNGRDTHMRQIKIYTPVEESSI GKFRCTTIDFMMYRSIR
3114	A	1	1613	MTSKEESRRQQPTAGPAGQGKLPSPSEPQLPTPP TRSLHHFRRLSPSREAAHIAPSSELHLPQSQSA GPPPLGAGTEVELVVPGRDEGSRGALPGSSGVKF VWRKTVPPVSDQVRLSICRLMPLLEMM VQFIRJWRSLLGRTLGTINIMMAQILASH LIKATVIPNRVKMLPYFGIIRNRMMSTHKSKKKI REYYRLNVEEGCSADEVRESFHKLAQYHPDS GSNTADSATFIRIEKAYRKVL SHVIEQTNASQSK GEEEDVEKFKYKTPQHRHYLSFEGIGFGTPTQR EKHYRQFRADRAAEQVMEYQKQKLQSQYFPDS VIVKNIRQSKQKITQAIERLVEDLIQESMAKGDF DNLSGKGKPLKKFSDCSYIDPMTHNLNRILIDNG YQPEWILKQKEISDTIEQLREAIL VSRKKLGNPMT PTEKKQWNHVCEQFQENIRKLNKRINDFNLIPI LTRQKVHFDAQKEIVRAQKIYETLIKTEVTDNRN PNNLDQGEGETPEIKKGFLNMDLVEIY
3115	A	1	2036	FRHRCGCLSYCRSRRGIRRV EPLRRARARVGPRF RPLCRM EIRSNFKNLHKVYQAIEEADFFAIDGE FSGISDGPSVSALTNGFDTP EERYQKLKKHSMDF LLFQFGLCTFKYDYTDSKYITKSFNFYVFPKPFNR SSPDVKFVCQSSSIDFLASQGFD FNKGFRKGIPYL NQEEERQLREQYDEKRSQANGAGALSYVSPNTS KCPVTIPEDQKKFIDQVVEKIEDLLQSEENKNL DL EPCTGFQRKLIYQTL SWKYPKGIHVETLETEKKE RYTIVISKVDEEERKRREQQKHAKQEELND AVG FSRVIHA IANSGKL VIGHNMLLDVMHTVHQFYC PLPADLSEFKEMTT CVFPRLD TKLMAS TQPFKD IINNTSLAELEKRLKETPFNPPKVESAE GFPSYDT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				ASEQLHEAGYDAYTTGLCFISMANYLGSFLSPPKI HVSARSKLIEPFFNKLFLMRVMDIPYLNLEGPDL QPKRDHVLHVTFPKWKTSPLYQLFSAFGNIQIS WIDDTSAFVSLSQPEQVKIAVNTSKYAESYRIQT YAEYMGRKQEEKQIKRKWTEDSWKEADSKRLN PQCIPYTLQNHYYRNNSTFAPSTVGKRNLSPSQE EAGLEDGVSGEISDTELEQTDSCAEPLSEGRKKA KKLKRMMKKELSPAGSISKNSPATLFEVPTDW
3116	A	3	1443	TREAPMALAVAPWGRQWEEARALGRAVRMLQ RLEEQCVDPRLSVSPSLRDLLPRTAQLLREVAH SRRAAGGGGPGGPGSGDFLLIYLANLEAKSRQ VAALLPPRGRRSANDELFRAGSRLRRQLAKLAI FSHMAELHALFPGGKYCGHMYQLTKAPAHTF WRESCGARCVLPWAEFESLLGTCHPVEPGCTAL ALRTTIDLTCSGHVSIFEFDVFTRLFQWPWTLKN WQLLAVNHPGYMAFLTIDEVQERLQACRDKPG SYIFRPSCTRLGQWAIGYVSSDGSILQTPANKPLS QVLLEGQKDGIFYLYPDGKTHNPDLTELQAEPQ QRIHVSEEQLQLYWAMDSTFELCKICAESNKDV KIEPCGHLLCSCCLAAWQHSDSQTCPCFRCCEIKG WEAVSIYQFHGQATAEDSGNSSDQEGRELELGQ VPLSAPPLPPRPDLPPRKPRNAQPKVRLKGNPSPP AALGPQDPAPA
3117	A	296	3547	ERHSSPLLQHILTHALMRNKKHSNNWLAQHW QSSILCFSPVGRITLVRARKFPAIVNCTAIDWFH AWPQEAALVSVSRRIEETKGIEPVHKDSISLFMAH VHTTVNEMSTRYYQNERRHNYTTPKSFLEQISLF KNLLKKKQNEVSEKKERLVNGIQKLKTASQV DLKARLASQEAELQLRNHDAEALITKIGLQTEKV SREKTIADAEERKVTAIQTEVFQKQRECEADLLK AEP...AATAALNTI...SELK... NVLAVMVLLAPRG...WKAAKVFNGK VDDFLQALINYDKEH...CLKVVNEHYLKDPEF NPNLIRTKSFAAAGLCAWVG...UKFYEVYCDVEP KRQALAQANLELAAATEKLE...KKLVVSANYD IEKSEKIRWGQSIKSFEAQEKTLCGDVLLTAAFVS YVGPFTRQYRQELVHCKWVPFLQQKVSIPLTEG LDLISMLTDDATIAAWNNEGLPSDRMSTENAIL THCERWPLVIDPQQQGIKWKNKYGMDLKVTHL GQKGFLNAIETALAFGDVILNLEETIDPVLDP LGRNTIKKKGKIRIGDKECFNKNFRILHTKLAN PHYKPELQAQTILLNFTVTEDEGLEAQLLAEVVSI ERPDLEKLKLVLTQHQNDFKIELKYLEDDLLRL SAAEGSFLDDTKLVERLEATKTTVAEIEHKVIEA KENERKINEARECYRPVAARASLLYFVINDLQKI NPLYQFSLKAFNVLFHRAIEQADKVEDMQGRISI LMESITHAVFLYTSQALFEKDKLTFLSQMAFQIL LRKKEIDPLELDFLLRFTVEHHLSPVDFTLSQSW SAIKAIAMVEEFRGIDRDVEGSAKQWRKWVESE CPEKEKLPQEWKKKSLIQKLILLRAMRPDRMTY ALRNFEVKLGAKYVERTRLDLVKAFEESPATP IFFILSPGVDALKDLEILGKRLGFTIDSGKFHNVSL GQGQETVAEVALEKASKGGHWVILQNVHLVAK WLGTEKLLERFSQGSHRDYRVFMSAESAPTPD EHHPQGLLENSIKITNEPPTGMLANLHAALYNFD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
3118	A	1	226	Q PYSLSSTCLGSPTSPRLEMDPNCSCATGGSCCTCTG SCKCKECKCNSCKKSECGAISRNGLLSQVRGRKP ELGMEE
3119	A	1254	4133	PLATLTMEEQGHSEMEIIPSESHPHIQLLKSREL LVTHIRNTQCLVDNLLKNDYFSAEDAEIVCACPT QPDKVRKILDVQSKGEEVSEFFLYLLQLADAY VDLRPWLEIGFSPSLLTQSKVVNTDPVSRYTQ QLRHHLGRDSKFVLCYAQKEELLLEEIYMDTIME LVGFSNESLGSLSLACLDDHTTGILNEQGETIFIL GDAGVGKSMMLQRLQSLWATGRDLGAVKFFFH FRCRMFSCFKESDRLCLQDLLFKHYCYPERDPPEE VFAFLRLRPHVALFTFDGLDELHSDLDLSRVPDS SCPWEPAHPLVLLANLLSGKLLKGASKLLTART GIEVPRQFLRKKVLLRGFSPSHLRA YARRMFPER ALQDRLLSQLEANPNLCSLCSVPLFCWIFRCFQH FRAAFEGSPQLPDCTMTLTDVFLLVTEVHLNRM QPSSLVQRNTRSPVETLHAGRDTLCSLGQVAHR GMEKSLFVFTQEEVQASGLQERDMQLGFLRALP ELGPGGDQSYEFFHLTLQAFFTAFFLVDDRVG TQELLRFFQEWMPAGAAATTSCYPFLPFQCLQG SGPAREDLFKNKDHQFTNLFLCGLLSKAKQKLL RHLVPAALRRKRKALWAHLFSSLRGYLNLSLR VQVESFNQVQAMPTFIWMLRCTIYETQSQKVGGQ AARGICANYLKLTYCNACSA DCSALSFVLHHP KRLALDLNNDYGVRELQPCFSRLTVLRLS VNQITDGGVKVLSEELTKYKIVTYLGLYNNQITD VGARYVTKILDECKGLTHLKLKGNKITSEGGKY LALAVKNSKSISEVGMWGNQVGDGAKAFEA LRNHPSLTTLASNGISTEGGKSLARALQNTSL EHLWLTQNELNDEVAESLAEMKDTCTLKTLWL IQNTITAKGTAQLADALQNTGITEICLNGVLP EEAKVYEDEKRIICF
3120	A	43	1004	QLWGFAAGSDSRPAMGCDGGTIPKRHELVKGPK KVEVDKDAELVAQWNYCTLSQEILRRPIVACE LGRLYNKDAVIEFLDKSAEKALGKAASHIKS NVTCLKLSDNPAWEGDKGNTKGDKHDDLQAR FICPVVGLMNGRHRFCFLRCCGCVFSERALK KAEVCHTCGAAFQEDDVIVLNGTKEDVDVLKTR MEERRLRKLEKKTKPKAAESVSKPDVSEAP GPSKVKTGKPEEASLDSREKKTNLAPKSTAMNE SSSGKAGKPPCGATKRSIADSESEAYKSFLTTHS SAKRSKEESAHWVTHTSYCF
3121	A	3	1490	HASGPTRPVSWSFHKLKTMKHLLLLLLCVFLVK SQGVNDNEEGFFSARGHRPLDKKREEAPSLRPAP PPISGGGYRARPAAAAATQKKVERKAPDAGGCL HADPD LGVLCPTGCQLQEALLQGERPIRNSVDEL NNNVEAVSQTSSSFQYMYLLKDLWQKRQKQV KDNENVVNEYSSELEKHQLYIDETVNSNIPTNLR VLSILENLRSKIQKLESVSAQMEYCRTPCTVS CNIPVVS GKECEEIRKGGGETSEMYLIQPDSSVKP YRVYCDMNTENGWTVIQNRQDGSVDFGRKW DPYKQGFGNVATNTDGKNYCGLPGEYWLGNK ISQLTRMGPTELLIEMEDWKGDVKVKAHYGGFTV QNEANKYQISVKNYRGTAGNALMDGASQLMGE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				NRTMTIHNGMFFSTYDRDNDGWLTS DPRKQCSK EDGGGWYNNRCHAANPNGRYYWGGQYTWDM AKHGTDDGVVWMNWKGSWYSMKKMSMKIRP FFPQQ
3122	A	3	1490	HASGPTRPVSWSFHKLKTMKHL L L L L L C V F L V K SQGVNDNEEGFFSARGHRPLDKKREEAPSLRPAP PPISGGGYRARPAAAAATQKKVERKAPDAGGCL HADPDLGVLCPTGCQLQEALLQQERPIRNSVDEL NNNVEAVSQTSSSFQYMYLLKDLWQKRQKQV KDNENVVNEYSSELEKHQLYIDETVNSNIPTNLR VLSILENLRSKIQLKLESDVSAQMEYCRTPCTVS CNIPVVSGKECEEIRKGGGETSEMYLIQPDSSVKP YRVYCDMNTENGGWTVIQNRQDGSVDFGRKW DPYKQGFQGNVATNTDGKNYCGLPGEYWLGN DK ISQLTRMGPTTELLIEMEDWKGDKVKAHYGGFTV QNEANKYQISVKNYRGTAGNALMDGASQLMGE NRTMTIHNGMFFSTYDRDNDGWLTS DPRKQCSK EDGGGWYNNRCHAANPNGRYYWGGQYTWDM AKHGTDDGVVWMNWKGSWYSMKKMSMKIRP FFPQQ
3123	A	3	1490	HASGPTRPVSWSFHKLKTMKHL L L L L L C V F L V K SQGVNDNEEGFFSARGHRPLDKKREEAPSLRPAP PPISGGGYRARPAAAAATQKKVERKAPDAGGCL HADPDLGVLCPTGCQLQEALLQQERPIRNSVDEL NNNVEAVSQTSSSFQYMYLLKDLWQKRQKQV KDNENVVNEYSSELEKHQLYIDETVNSNIPTNLR VLSILENLRSKIQLKLESDVSAQMEYCRTPCTVS CNIPVVSGKECEEIRKGGGETSEMYLIQPDSSVKP YRVYCDMNTENGGWTVIQNRQDGSVDFGRKW DPYKQGFQGNVATNTDGKNYCGLPGEYWLGN DK ISQLTRMGPTTELLIEMEDWKGDKVKAHYGGFTV QNEANKYQISVKNYRGTAGNALMDGASQLMGE NRTMTIHNGMFFSTYDRDNDGWLTS DPRKQCSK EDGGGWYNNRCHAANPNGRYYWGGQYTWDM AKHGTDDGVVWMNWKGSWYSMKKMSMKIRP FFPQQ
3124	A	3	544	RVDDFVLLRSRLALRWLSHVRPSRRVPRMPRG SRSTSRMAPPASRAPQMRAAPRPAPVAQPPAA APPSAVGSSAAAPRQPLMAQMATTAAAGVAVG SAVGHTLGHATTGGFSGGSNAEPARPDITYQEPQ GTQPAQQQPCLYEIKQFLECAQNQGDIKLCEGF NEVLKQCRLANGLA
3125	A	3	571	GNSYNHRS LAAYPYMSHSQHSPYLQSYHNSSAA AQTRGDDTDQKTTVIENGHIRFNGKGKKIRKPR TTYSSLQLQALNHRFQQTQYLALPERAELAASLG LTQTQVKIWFQNKRSKFKLLKQGSNPHESDPL QGSAAALSPRSPALPPVWDVSASAKGVSMPPNSY MPGYSHWYSSPHQDTMQRPM
3126	A	43	5377	LSVFFPIPV DGRDRGSNPSLESTSTSTSEGL SAMSGRNELHSRLHPHPQSSLIPMMFSPPELLAS CILRGNF AEAHQVLF T FNLKSSPSSGELMF MERY QEVIELAQVEHKIENQNSDAGSSTIRRTGSGRST LQAIGSAAAAGMVFYSISDVTDKLLNTSGDPIPM LQEDFWISTALVEPTAPLREVLEDLSPPAMAAFD LACSQCQLWKTCKQLLETAERRLNSSLERRGRI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				DHVLLNADGIRGFPVVLQIQSKSLNLYLLMSASQT KSESVEEKGGGPPRCSTITELLQMCWPSLSEDCVA SHTTLSQQLDQVLQSLREALELPEPRTPLSSLVE QAAQKAPEAEHPVQIQTQLLQKNLGKQTPSGS RQMDYLGTFFSYCSLAAVLLQSLSEPDPHVEVK VGNPFVLLQQSSSQLVSHLLFERQVPPERLAAILL AQENLSLSPVQVIVSCCEPLALCSSRQSQQTSSL LTRLGTLAQLHASHCLDDLPLSTPSSPRTTENPTL ERKPYSSPRDSSLPALTSSALAFKSRSKLLATVA CLGASPRKLVSKPSLSWKELRGRREVPLAAEQV ARECERLLEQFPLFEAFLLAAWEPLRGSLLQQGQS LAVNLGASLSTVLLGLHSPIALDVLSEAFES LVARWDSRALQLTEVYGRDVEDLSSIKDAVLSC AVACDKEGWQYLPVKDASLSRSLALQFVDRW PLESCLEILAYCISDTAVQEGKCELQRKLAELQ VYQKILGLQSPVWCDWQTLRSCCVPSTVMN MILEAQEYELCEEWGCLYPREHLISLHOKHLL HLLERRDHDKALQLLRIPDPTMCLEVTEQSLDQ HTSLATSHFLANYLTTHFYGQLTAVRHREIQALY VGSKILLTLPEQHRASYSHLSSNPLFMLEQLLMN MKVDWATVAVQTLQQLLVGQEIGFTMDEVDSL LSRYA EKALDFPYPQREKRSDSVIHLQEIVHQA DPETLPRSPSAEFSPAAPPGISSIHSPSLRERSFPPT QPSQEFVPPATPPARHQWVPDETESICMVCCREH FTMFNRRHHCRCGRVCSSCSTKMMVVEGCRE NPARVCDQCYSYCNKDVPEEPSEKPEALDSSKSE SPPYSFVVRVPKADDEVWILDLKEEENELVRSEF YYEQAPSASLCIAILNLHRDSIACGHQLIEHCRL SKGLTNPEVDAGLLTDIMKQLLFSAKMMFMKAG QSQDLALCDSYISKVDVLNLAAYRHVPSLDQ ILQPAAVTRLNQLLEAEYYQLNTEVSTKTOLDT TGAWHA WGMACLNLTAAAEKFSRCLKPPF DLNQLNHGSRLVQDVVEYLESTVRPFVSLQDDD YFATLRELEATLRTQSLSLAVIPEGKIMNNTYYQ ECLFYLNHYNSTNLAIISFYVRHSCLREALHLLNK ESPPEVFIEGIFQPSYKSGKLHLENLLESIDPTLES WGKYLIAACQHLQKKNYYHILYELQQFMKDQV RAAMTCIRFFSHKAKSYTELGEKLSWLLKAKDH LKIYQLQETSRSSGRKKTFFRKKMTAADVSRHM NTLQLQMEVTRFLHRCESAGTSQITTLPLPTLFG NNHMKMDVACKVMLGGKNVEDGFGIAFRVLQ DFQLDAAMTYCRAARQLVEKEYSEIQQLKCV SESGMAAKSDGDTILLNCLAEAFKRIPPQCCFCSA QELEGLIQAIHNDNDNKVRAYLICCKLRSAYLIAV KQEHSRATALVQQVQQAASGDAVVQDICAQ WLLTSHPRGAHGPSRK
3127	A	467	1259	HLGPPALWIPAASTSTKGFEFGVEDDRPARGPPP PKSEEASWSESGVSSSSGDGPFAGGEVDKRLHQL KTQLATLTSSLATVTQEKSRMEASYLADKKKMK QDLEDASNKAEEERARLEGELKGLQEQAETKA RLITQQHDRAQEQSDHALMLRELQKLLQBERTQ RQDLELRLEETREALAGRAYAAEQMEGFELQTK QLTREVEELKSELQAIRDEKNQDPRLQELQBEA ARLKSHFQAQLQQEMRKVIIIHSFKHQPLT
3128	A	1854	798	ASGSPAPSSSSAMAAACGPGAAGYCLLLGLHLFL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LTAGPALGWNDPDRMLLRDVKALTLHYDRYTT SRRLDPIQLKCVGGTAGCDSYTPKVIQCQNK WDGYDVQWECKTDLDIA YKFGKTVVSCGYES SEDQYVLRGSCGLEYNLDYTELGLOKLKESGKQ HGFASFSDYYYKWSSADSCNMSGLITIVVLLGIA FVVYKFLSDGQYSPPPYSEYPPFSHRYQRFTNS AGPPPPGFKSEFTGPQNTGHGATSGFGSAFTGQQ GYENSGPGFWTGLGTGGILGYLFGSNRAATPFSD SWYYPSYPPSYPGTWNRAYSPLHGGSGSYVCS NSDTKTRTASGYGGTRRR
3129	A	2340	1192	ELARRPKQSSSEKSRNMIRNWL TIFLFPLKLVEK CESSVSLTVPPVVKLENGSSSTNVSLTRPPLNATL VITFEITFRSKNITILELPDEVVPPGVNTSSSQVT SQNVGQLTVYLHGNHNSNQTGPRJRFVIRSSAISII NQVIGWYFVAWSISFYQVIMNWRKRSVIGLSF DFVALNLTGFVAYSVFNIGLLWVPYIKEQFLLKY PNGVNPVNSNDVFFSLHAVVLTLIIVQCCLYERG GQRVSWPAIGFLVLA WLFVFTMIVA AVGVITW LQFLFCFSYIKLAVTLVKYFPQAYMNFYKSTEG WSIGNVLDFGTGGSFSLQMFLQSYNNDQWTLIF GDPTKFGLGVSIVFDVFFIQHFCLYRKRPGYD QLN
3130	A	31	2026	CWWPPLLQLEPEPPPLRPRVAASQGGGMLGKG VVGGGGGTKAPKPSFVS YVRPEEIHNEKEVTEK EVTLHLLPGEQLLCEASTVLKYVQEDSCQHGVY GRLVCTDFKIAFLGDDESALDNDDETQFKNK VIGE NDITLHCVDQIYGVFDEKKKTLFGQLKKYPEKLI HCKDLRVFQFCLRYTKEEEVKRIVSGIHHHTQAP KLLKRLFLFSYATAAQNNVTVDPKNHTVMFDTL KDWCEWELERTKGNMKYKAVSVNEG YKVCERL PA*EVVPTPLPEEPVORFQGHGIFWCWWSCHNGS ALPKEQDEGILQIQKSFLDGIYKTIH YBWKTEDLSSNFLSLQEIQTAYSFKFKQLFLDINS EFWDTDIKWFSLESSSWLDIIRCLKKAIEITEC MEAQNMENVLLEENASDLCLISLVQLMMDPH CRTRIGFQSLIQKEWVMGGHCFLDRCNHLRQND KEEHQRQLSLPLTQSKSSPKRGFFREETDHLIKNL LGKRISKLINSSDELQDNFREFYDSWHSKSTDYH GLLLPHIEGPEIKVWAQRYLRWIPEAQILGGGQV ATLSKLEMMEEVQSLQEKIDERHHSQQAPQAE APCLLRNSARLSSLFPFALLQRHSSKPVLP TSGW KALGDEDDLAKREDEFVDLGDV
3131	A	126	965	QSRSRPRREGVGTGSRVLCILATCGSKMSDIGD WFRSIPAITRYWFAATVA VPLVGKLG LISPAYLF LWPEAFLYRFQIWRPITATFYFVPGPGTGFLYL NLYFLYQYSTRLETGAFDGRPADYLFMLLFNWI CIVITGLAMDMLMPLIMSVLYVWAQLNRDM IVSFWFGTRFKACYLPWVILGFNYHGGSVINELIG NLVGHLYFFLMFRYPMDLGGRNFLSTPQFLYRW LPSRRGGVSGFGVPPASMRRAADQNGGGGRHN WGQGFR LGDQ
3132	A	2	350	FVAGWRALTAPSTSARLRAFGWQAAARLLVFG ARGVGLGSGAPGSLPCYL RMDALALLGGLVNV ARLPERWGPGRFDYWGNSHQIMHLLSVGSILQL HAGVVPDLLWAAHHACPRD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3133	A	1	2921	MTCFKGQKGEQRSHAFEANKDHHKAKVPSPNLYS QLNALQFTVDERSLWLNQFLDLKQSLNQFMA VYKLNDSKSDHEVDVRVDGLMLKFVIPSEVKS ECHQDQPRAISIQSSEMIATNTRHCPNCRHSDLEA LFQDFKDCDFFSKTYTTSFPKSCDNFNLLHPIQRH AHEQDTKMHEIYKGNITPQLNKNLTKTSAATDV WAVYFSQFWIDYEGMKSGKGRPISEFVDSFPLSIW ICQPTRYAESQKEPQTCNQVSLNTSQSESSDLG RLKRRKLLKEYYSTSEPLTNGGQKPSSSDTFFR FSPSSSEADHLLVHVHKKHVMQINHYQYLLLLL LHESLILLSNLRKDVEAVTGSPASQTSICIGILLR SAELALLLHPVDQANTLKSPVSESVSPVDPDYL TENGDFLSSKRKQISRDINRISVTVNHMSDNRS MSVDLSHIPKDPPLFKSASDTNLQKGISFMDYL SDKHLGKISEDESSGLVYKSGSGEIGSETSDKKDS FYTDSSSVLNYREDSNLSFSDSDGNQNILSSTLS KGNETIESIFKAEDLLPEAASLSENLDISKEETPPV RTLKSQSSLSGKPKERCPPNLAPLCVSYKNMKRS SSQMSLDTISLDSMLEEQLES DGS DSHMFLEKG NKKNSTTNYRGTAESVNAGANLQNYGETSPDAI STNSEGAQENHDDLMSVVVKITGVNGEIDIRGE DTEICLVNQVTPDQLGNISLRHYLCNRPVGS DQ KAVHSKSSPEISLRFESGPGAVIHSLLAEKNGFL QCHIEFSTEF LTSSLMNIQHFEDET VATVMMPM KIQVSNTKINLKDDSPRSSTVSLEPAPVTVIDHL VVERSDDGSFHIRDSHMLNTGNDLKENVKSDSV LLTSGKYDLKKQRSVTQATQTS PGVPWPQSAN FPEFSDF TREQLMEENESLKQELAKAKMALAE AHLEKDALLHHKKMTVE
3134	A	9	1579	EEBGLSGGGPRVPCSLWGKQTM DYDFKAKLAA ERERVIN FEYEGCKVGRGT YC YKARRD G KDEKEYALKQIEGT GMSACREALLRELKHPN VIALQKVFLSHSDRKVWLLFDYAEHDLWHIHKFH RASKANKKPMQLPRSMVKSLLYQILDGIHYLHA NWVLHRDLKPANILVMGEGPERGRVKIADMGF ARLFNSPLKPLADLDPVVVTFWYRAPELLGAR HYTKAIDIWAIGCIFAELLTSEPIFHCQEDIKTSN PFHHDQLDRIFSVMGFPADKDWEDIRKMPEYPT LQKDFRRTTYANSSLIK YMEKHKVKPDSKVFL LQKLLTMDPTKRITSEALQDPYFQEDPLPLDV FAGCQIPYKREFLNEDDPEEKGDKNQQQQQNG HQQPTAPPQAAAAPPQAPPQONSTQNGTAGG AGAGVGGTGAGLQHSQDSSLNQVPPNKKPRLGP SGANS GGPVMPSDYQHSSRLNYQSSVQGSSQS QSTLGYSSSSQSSQYHPHQAHRY
3135	A	3	1111	ERKMAEPSPVHCVAAAAPTATVSEKEPFGKILQ LSSRDPPGSLSAKKVRTEEKKAPRRVNGEGGSG GNSRQLQPPAAPSPQSYGSPASWSFAPLSAAPSPS SSRSSFSFAGTAVPSSASASLSQPGPRKLLVPPTL LHAQPHHLLLPAAAAAASANA KSRPKEKREKE RRRHGLGGAREAGGASREENG EVKPLPRDKIKD KIKERDKEKEREKKKHKVMNEIKKENG EVKILL KSGKEKPKTNIEDLQIKKVKKKKKKKHKENEKR KRPKMYSKSIQTICSGLLTDVEDQAAKGILNDNI KDYVGKNLDTKNYDSKIPENSEFPFVSLKEPRVQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				NNLKRLDTLEFKQLIHIEHQPNGGASVIHCLQ
3136	A	1442	682	TAAMSIFTPTNQIRLTNVAVVRMKRAGKRFEIAC YKNKVVGWRSVGVEKDLDEVLTQTHSVFVNVSKG QVAKKEDLISAFGTDDQTEICKQLTKGEVQVSD KERHTQLEBQMFRIATTIVADKCVNPETKRPYTVI LIERAMKDIHYSVKTNKSTKQQAILEVIKQLKEK MKIERAHMRLRFILPVNEGKKLKEKLKPLIKLVIES EDYGGQLEIVCLIDPGCFREIDELIKKETKGKGS EVLNLKDVEEGDEKFE
3137	A	1	3143	MVEGKRHVLHGGQRQERMRAKQKGKPLIKSSDL VRLIHYHHNSSPLHKQSSGSSSPAAAAAPEKPG PKAAEVGDDFLGDFVVGGERVWVNGVKPGVVQY LGETQFAPGQWAGVVLDDPVGKNDGAVGGVR YFECPALQGIFTRPSKLTROPTAEGSGSDAHSVES LTAQNLSLHSGTATPPLTSRVIPLRESVLNSSVKT GNESGNSLSDSGSVKRGKEDLRLGDRVLVGGTK TGVVRYVGETDFAKGEWCGVELDEPLGKNDGA VAGTRYFQCPPKFGLFAPIHKVIRIGFPSTSPAKA KKTKRMAMGVSAETHSPSSSSISSVSVASSVGG RPSRSGLLTETSSRYARKISGTTALQEALKEKQ HIEQLLAERDLERA EVAKATSHICEVEKEIALLK AQHEQYVAEAEKLRARLLVESVRKEKVDLSN QLEERRKVEDLQFRVEEESITKGDLETOTQLEH ARIGELEQSLLLEKAQAERLLRELADNRLTTVAE KSRVLQLEBELTLRRGEIEELQQCLLHSGPPPDH PDAAEILRLRERLLSASKEHQRESGVLRDKYEKA LKAYQAEVDKLRAANEKYAQEVAGLKDKVQQ ATSENMGMLMDNWKSKLDSLADHQSLEDLKA TLNSGPGAQQKEIGELKAVMEGIKMEHQLELGN LQAKHDLETAMHVKEKEALREKLQEAQEELAG LQRHTRAQLEFQASQHRLELQEAQDORRDALE RVHELEKLELQETGSAQAIEFLKEQISLAEKKML DYERLQRAEAQQKQEVESLREKLLVAENRLQAV EALCSSQHTHMESEEDISEETIRTKETVEGLQDKL NKRDKVETALTSQTEMLRAQVSALESCKSGEK KVDALLKEKRRLEAELETVSRKTHDASGQLVLIS QELLRKERSLNELRVLLLEANRHSPGPERDLSRE VHKAEWRIEQKLKDDIRGLREKLTGLDKEKSL SDQRRYSLIDPSSAPELLRLQHQLMSTEDALRDA LDQAQQVEKLMEAMRSCPDKAQTIGNSGSANGI HQQDKAQKQEDKH
3138	A	110	2499	QDRRLRLLELQKTCQPTSTMSGSHTPACGPFSAL TPSIWPQEILAKYTQKEESAEPFYDEFGRV YKEEGDEPGSSLLANSPLMEDAPQRLRWQAHL FTHNHDVGDLTWDKIAVSLPRSEKLRSLVLAGIP HGMRPQLWMRLSGALQKRNSELSYREIVKNSS NDETIAAKQIEKDLLRTMPNACFASMGSGVPR LRRVLRALAWLYPEIGYCQGTGMVAACLLLFLE EEDAFWMMSAIEDLLPASYSFTLLGVQTDQRV LRHLIVQYLPRLDKLLQEHDIELSLTLHWELTAF ASVVDIKLLLRWDLFFYEGSRVLFQTLGMLHL KEELIQSENSASIFNTLSDIPSQMEDAELLGVA MRLAGSLTDVAVETQRRKHLAYLIADQGQLLGA GTLTNLSQVVRRTQRRKSTTTALLFGEDDLEAL KAKNIKQTELVAIDLREAILRVARHFQCTDPKNCS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VVSRQLPGLLPNTALTPPTPLVGLCSLWQELTPD YSMESHQRDHENYVACSRSHRRRAKALLDFERH DDDELGFRKNDIITTVSQDEHCWVGELNGLRG WFPKAFVEVLDERSEKEYSIAGDDSVTEGVTDLV RGTLCPALKALFEHGLKKPSLLGGACHPWLFIEE AAGREVERDFASVYSRLVLCKTFRLEDGKVLTP PEELLYRAVQSVNVTHDAVHAQMDVKLRSLICV GLNEQVLHLWLEVLCSLLPTVEK WYQPW SFLRS PGWVQIKCELRLVCCFAFSLSDWELPAKREAQ QPLKEGV RDMLVKHHLFSWDVDG
3139	A	110	2499	QDRRLRLLELQKTCQPTSTMSGSHTPACGPFSA TPSIWPQEILAKYTQKEESAEQPEFYDEFGRV YKEEGDEPGSSLLANSPLMEDAPQRLRWQAHLB FTHNHDVGDLTWDKIAVSLPRSEKLRSVLGIP HGMRPQLWMRLSGALQKKRNSLSYREIVKNSS NDETIAAKQIEKDILLRTMPSNACFASMGSGVPR LRRVLRALAWLYPEIGYCQGTGMVAACLLLFLE EEDAFWMMSAIIEDLLPASVYSTTLGVQTDQRV LRHLIVQYLPRLDKLLQEHDIELSLITLHWFLTAF ASVVDIKLLLRWDLFFYEGSRVLFQTLGMLHL KEELIQSENSASIFNTLSDIPSQMEDAELLGVA MRLAGSLTDVAVETQRRKHLAYLIADQGQLLGA GTLTNLSQVRRRTQRRKSTITALLFGEDDLEAL KAKNIKQTELVADLREAILRVARHFQCTDPKNCS VVSRQLPGLLPNTALTPPTPLVGLCSLWQELTPD YSMESHQRDHENYVACSRSHRRRAKALLDFERH DDDELGFRKNDIITTVSQDEHCWVGELNGLRG WFPKAFVEVLDERSEKEYSIAGDDSVTEGVTDLV RGTLCPALKALFEHGLKKPSLLGGACHPWLFIEE AAGREVERDFASVYSRLVLCKTFRLEDGKVLTP PEELLYRAVQSVNVTHDAVHAQMDVKLRSLICV GLNEQVLHLWLEVLCSLLPTVEK WYQPW SFLRS PGWVQIKCELRLVCCFAFSLSDWELPAKREAQ QPLKEGV RDMLVKHHLFSWDVDG
3140	A	1	4939	SAALGASLAIPRGLPGVHGRGPCLESGRAMEG AEPRARPERLAEAETRAADGGRLVEVQLSGGAP WGFTLKGGREHGEPLVITKIEEGSKAAAVDKLL AGDEIVGINDIGLSGFRQEAICLVKGSHKTLKL VKRRSELGWRPHSWHATKFSDSHPELAASPFTST SGCPSWSGRHHASSSSHDLSSSWEQTNLQRTLD HFSSLGSVDSLDHPSSRLSVAKSNSSIDHLGSHSK RDSA YGSFSTSSSTPDHTLSKADTSSAENILYTVG LWEAPRQGGGRQAQAAGDPQGSEKLSCFPPRPV GDSGKGPRPEYNAEPKLAAPGRSNFGPVWYVPD KKKAPSSPPPPPPPLRSDSFAATKSHEKAQGPVFS EAAAAQHFTALAAQAQPRGDRRPELTDRPWRS PGSLGKSGGGPGCPQEAHADGSWPPSKDGASSR LQASLSSSDVRFQSPHSGRHPPLYSDHSPLCADS LGQEPGAASFQNDSPQVRGLSSCDQKLGSGWQ GPRPCVQGDQLAAQLWAGCWPSDTALGALES PPPTVGQSPRHHLQPEGPPDARETGRCYPLDKG AEGCSAGAEPPRASRAEKASQRLAASITWADG ESSRICPQETPLLHSLTQEGKRRPESSPEDSATRPP PFDHVGKPTRRSDFATTLRNEIQMHRAKLQK SRSTVALTAAGEAEDGTGRWRAGLGGGTQEGPL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				AGTYKDHLKEAQARVLRATSFKRRDLDPNPGDL YPESLEHRMGDPDTPVPHFWEAGLAQPPSSTSGGP HPPRIGGRRRFTAQKLSYSEPEKMNEVGLTRG YSPHQHPTSEDVTGTFADRWKFFEETSKPVPQR PAQKQALHGIPRDKPERPRTAGRTCEGTEPWSRT TSLGDSLNAHSAAEKAAGTSDLPRLGTFAEYQAS WKEQRKPLEARSSGRCHSADDILDVSLDPQERPO HVHGRSRSSPSTDHYKQEAASVELRRQAGDPGEP REELPSAVRAEAGQSTPRQADAQCREGSPGSQQ HPPSQKAPNPPTFSELSHCRGAPELPREGGRAG TLPRDYRYSEESTPADLGPRASPGSPLHARGQD SWPVSSALLSKRPAPQRPPPPKREPRRYRATDGA PADAPVGVLGPRPFTPSASLDVYVARLSLSHSPS VFSSAQPDTPKATVCERGSQHVSGDASRPLPEA LLPPKQOHLRLQTATMETSRSPPQFAPQKLTDK PLLIQDEEDSTRIERVMDNNTTVKMVPIKIVHSES QPEKESRQSLACPAEPPALPHGLEKDQIKTLSTSE QFYSRFCLYTRQGAEPHRAQPAEPQPLGTQV PPEKDRCTSPGSLYMKAKEKTVEDLKSEELARE IVGKDKSLADILDPSVKIKTTMDLMGIFPKDEH LLEEAQQRRLPKIPSPRSTEERKEEPSVPAAS LATNSTYYSTAPKAELLKMKDLQEQEHEEDS GSDLDHDLVKKQELIESIRKLQVLRARESLLE DVQANTVLGAEEVAIVKGVCKPSEFDKFRMFIG DLKVVNLLLSLGRRLARVENALNLDGASPG DRQSLLEKQRLIQQHEDAKELKENLDRRERIVF DILANYLSEESLADYEHFVKMKSALIEQRELED KIHLGEEQLKCLLDSLQPERGK
3141	A	97	1894	SPRGATMETPPLPPACTKQGHQKPLDSKDDNTE KHCPVTVPNWHMKAFAKVMNELRSQNLCDVT VAEDMEIAHRVVLAACSPYFHAMTTEMSESR AKRVRIKEVDGWTLRMDTVYTANQVTEENV QVLLPAAGLLQLQDVKKTCCEFLSQLHPVNCL GIRAFADMHACTDLLNKANTYAEQHFADVVLSE EFLNLGIEQVCSLISSDKLTISSEKVF EAVIAWV NHDKDVRQEFMARLMEHVRLPLLPREYLVQRV EBEALVKNSSACKNYLIEAMKYHLLPTEQRILMK SVRTRLRTPMNLPKLMVVVGGQAPKAIRSAECY DFKEQRWHQVAELPSRRCRAGMVYLAGLVFAV GGFNGSLRVRTVDSYDPVKDQWTSVANMRDRR STLGAAVLNGLLYAVGGFDGSTGLSSVEAYNIKS NEWFHVAPMNTRRSSVGVGVGGLLYAVGGYD GASRQYLSTVECYNATTNEWTYIAEMSTRRSGA GVGVLNLLLYAVGGHDGFLVRKSVEVYDPTTN AWRQVADMNMCCRANAGVCAVNGLLYVVGGD DGSCNLASVEYYNPPTDKWTVVSSCMSTGRSYA GVTVIDKPL
3142	A	1211	1311	FSNLTTEKVAHAKEENLSMHQMLDQTLLELNN M
3143	A	1809	1041	SEELDREKKLKEDSPRKTPNKESGVPSLPVSLTSI KEEPKEAKHPDSQSMEESKLKNDDRKTPVNWK DSRGTRVAVSSPMSQHQSIIQYLHAYYPQMYD PSHPAYRAVSPVLMHSYPGA YLSPGFHYPVYK MSGREETEKVNTSPSVNTKTTTESKALDLLQKH ANQYRSKSPAPVEKATAEREREAERERDRHSPFG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QRHLHTHHHTHVGMGYPLIPGQYDPFQGLTSAA LVASQQVAAQASASGMFPGQRR
3144	A	78	604	SVSGIVLDLLPYLHFLSNMNLGSAQDPEKREYS SVCVGREDDIKKSERMTAVVHDREVVFYHKGE YHAMDIRCYHSGGPLHLGDIEDFDGRPCIVCPW HKYKITLATGEGLYQSINPKDPSAKPKWCSKGIK QRIHTVTVDNGNITYVTLSENEPKCDSDFYATGDF KVIKSSS
3145	A	2	333	RNSLLPLPLHLDNSTPAKMSCQONQQCQPPPK CPSPKCPPKSPVQCLPPASSGCAPSSGGCGPSSEG GCFLNHRRRHRCRRQRPNSCDRSGSQGGGS GCGHSGGGCC
3146	A	3	1151	VCTALQEFGRSTLLRCLDSGFRPGASRGLVGSW AAMESTLGAIVIAEALQNQLAWLENVWLWITF LGDPKILFLFYFPAAYYASRRVGVIAVLWISLITEW LNLIFKWFLFGDRPFWVWHESGYYSQAPQVHQ FPSSCETGPGSPSGHCMITGAALWPMTALSSQV ATRARSRWVRVMPSLAYCTFLAVGLSRIFILAH FPHQVLAGLITGAVLGWLMTPRVPMERELSFY LTALALMLGTSIYWLFTLGLDLSWSISLAFKW CERPEWIHVDSPFASLSRDSGAALGLGIALHSPC YAQVRRALGNGQKIACLVLAMGLLGPLDWLG HPPQISLFYIFNFKYTLWPCLVLALVPWAVHMF SAQEAPPIHSS
3147	A	1437	594	RSFSLSFSLLSPSEMMALGAAGATRVFVAMVAA ALGGHPLLGVSATLNSVLNSNAIKNLPPPLGGAA GHPGSAVSAAPGILYPGGNKYQITIDNYQPYPCAE DEECGTDEYCASPTRGGDAGVQICLACRKRKR CMRHAMCCPGNYCKNGICVSSDQNHFRGEIETI TESFGNDHSTLDGYSRRITLSSKMYHTKGQEGS VCLRSSDCASCLCTTFHFWSKICKTVYEGQVC TKHRRKGSHGELFQRYCYGEGLSCRIQKDHQ ASNSSRLHTCQR
3148	A	1	1562	MSTLYDIRAHKAQLRFFASSDSNKALEQRRTLH TPKLEHLDRVLYEWFLEKRSRSEGVPSGPMLEK AKDFYEQMQLTEPCVFSGGWLWRFKARHGIK LDASSEKQSADHQAEEQCAFRSLAAEHGLSA EQVYNADETGLFWRCLPNPTPEGGA VPGPKQKG DRLTVLMCANATGSHRLKPLAIGKCSGPRAFKGI QHLPVAYKAQGNAAVVDKEIFSDWFHIFVPSVR EHFRTIGLPEDSKAVLLDSSRAHPQEAELVSSN VFTIFLPASVASLVQPMEQGIRRD FMRNFINPPVP LQPHARYNMNDALFVACAWNAVPSHVFRRA WRKLWPSVAFAGSSSEEELEAECFPVKPHNKSF AHILELVKEGSSCPGQLRQRQAASWGVAGREAE GGRPPAATSPAEBVWSSEKTPKADQDGRGDPGE GEEVAWEQAAVAFDAVLRFAERQPCFSAQEVG QLRALRAVFRSQQQVRRRRGALGAVVKVEALQ EGPGGCGATAQSPLPCSSTAGDN
3149	A	132	4125	VAVMISTAPLYSGVHNWTSSDRIRMCGINEERRA PLSDEESTTGDCQHFGSQEFCVSSSFVKVELTAV GSGSNARGADPDGSATEKLGHKSEDKPDDPQPK MDYAGNVAEAEGLLVPLSSPGDGLKLPASDSAE ASNSRADCSWTPLTQMSKQVDCSPAGVKALDS RQGVGEKNTFILATLGTGVPVEGTLPLVTTNFSF

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LPAPICPPAPSSASVPHSVPDFAFQAPVPPSAPTLLV APVPTPVLAPMPASTPPAAPAPPSVPMPTPTPSSG PPSTPTLIPAFAPTPVPAPTPAPIFTPAPTMPAATP AAIPTSAPIASFSLSRVCFPAQAQAPAMQKVPLSF QPGTVLTPSQPLVYIPPPSCGQPLSVATLPTTLGV SSTLTLPVLPSYLQDRCLPGVLASPELRSYPYAFS VARPLTSDSKLVSEVNRLPCTSPSGSTTTQPAPD GVPGLADTSLVTASAKVLPTPQPLLAPPSGSSAP PHPAKMPSGTEQQTEGTSVTFSPKSPQLEREM ASPPECSEMPDLSSKSNRQKLPLPNQRKTPMP VLTPTVHTSSKALLSTVLSRSQRTTQAAGGNVTSC LGSTSSPFVIFPEIVRNGDPSTWVKNSTALISTPG TYVGVANPVPASLLNKPDLGLNRDPRHLPKQ EPISIIDQGEKGTGATCGKKGSQAGAEGQPSTV KRYTPARIAAGLPQCQTKELSLWKPTGPANIYPR CSVNGKPTSTQVLPVGWSPYHQASLLSIGISSAG QLTPSQGAPIRPTSVVSEFSGVPSLSSEAVHGLP EGQPRPGGSFVPEQDPVTKNKTCRIAAKPYEEQV NPVLLTLSPQTGTLALSVQPSGGDIRMNQGPES ESHLCSDSTPKMEGPQAGCGLKLAGDTKPKNQV LATYMSHELVLATPQNLPKMPPELPLPHDSHPKE LILDVVPSSRRGSSTERPQLGSQVDLGRVKMEKV DGDVVFNLATCFRADGLPVAPQRGQAEVRKA GQARVKQESVGVFACKNKWQDDVTESLPPKK MKCGKEKDSEEQQLQPQAKAVVRSSHRPKCRK LPSPDQESTKKSPRGASDSGKEHNGVRGKHKHR KPTKPESQSPGKRADSHEEGSLEKAKSSFRDFIP VVLSTRTRSQDLKARKQKTSSSQSLEHRLNRN LLLPNKVQGISDSPNGFLPNNLEEPACLENSEKPS GKRCCKTKHMA TVSEEAKGKGRWSQQKTRSPK SPTPVKPTETPSKSRASSEFACTTARQDPE LKELIVNKNAGETLLQQAARATLTYLCLQK DSEDVNHRDNAGYTALHEACSNWTDILNILLE HGA
3150	A	3	2795	SLRMHNLILVRQIKFYQETLQQLIMMLPNVLI IGKNPFSEQGTEEVKLLLLLLLGCVAQCQKKEEF IERIQGLDFDTKAAVAHHIQEVTHNQENVFDLQ WMEVTDMSQEDIEPLLKNMALHLKRLIDERDEH SETIHELSEERDGLHFLPHASSAQSPCGSPGMR TESRQHLVELADAKAKIRRLRQELEBKTEQLLD CKQELEQMBIELKRLQENMNLLSDARSARMYR DELDALREKAVRVDKLESEVSRYKERLHDIEFY KARVEELKEDNQVLLETKTMLEDQLEGTRARSD KLHELEKENLQLKAKLHDMEMERDMDRKKIEE LMEENMTLEMAQKQSMDESLHLGWELEQISRTS ELSEAPQKSLGHEVNELTSSRLKLEMENQSLTK TVEELRTTVDSVEGNASKILKMEKENQRLSKKV EILENEIVQEKQSLQNCQNLKSKDLMKEKAQLEKT IETLRENSERQIKILEQENEHLNQTVSSLRQRSQIS AEARVKDIEKENKILHESIKETSSKLSKIEFEKROI KKELEHYKEKGERAELENELHHLEKENELLQK KITNLKITCEKIEALEQENSELERENRKLKKTLD FKNLTFQLESLEKENSQLEENLELRRNVESLKC ASMKMAQLQLENKELESEKEQLKKGLELLKASF KKTERLEVSYQGLDIENQRLQKTLSENSKKIQQL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				ESELQDLEMENQTLQKNLEELKISSKRLEQLEKE NKSLEQETSQLEKDKKQLEKENKRLRQQAEIKD TTLEENNVKIGNLEKENKTLKEIGIYKESCVRL ELEKENKELVKRATIDIKTLVTLREDLVSEKLKT QQMNNDLEKLTHELEKIGLNKERLLHDEQSTDD SRYKLLSKLESTLKKSLKEEKIAALEARLEES TNYNQQLRQELKTVKKK
3151	A	2	2515	GFWLHLTLGASLPAALGWMDPGTSRGPDPVGV GESQAEPRSEFEVTRREGLSSHNEILLASCGKKFC SRGSRCLSRKTGEPECQCLEACRPSYVPVCGSD GRFYENHCKLHRAACLLGKRITVIHSDCFLKGD TCTMAGYARLKNVLLALQTRLQPLQEGDSRQDP ASQKRLLVESLFRDLADGNGHLSSELAQHVL KKQDLDEDLGCSPGDLLRFDDYNSDSSLTLREF YMAFQVVQLSLAPEDRVSVTTVTVGLSTVLTCA VHGDLRPPIHWKRNGTLNFDLEDINDFGEDDS LYITKVTIHMGNYTCHASGHEQLFQTHVLQVN VPPVIRVYPESQAQEPGVAASLRCHAEGIPMPRIT WLKNGVDVSTQMSKQLSLLANGSELHISVRYE DTGAYTCIAKNEVGVEDISSLFIEDSARKTLANI LWREGLSVGNMFYVFSDDGIIVHPVDCEIQRH LKPTKIFMSYEEICPQREKNATQPCQWVSAVNV RNRYTYVAQPALSRLVVDIAHKVLQSIGVDPL PAKLSYDKSHDQVWVLSWGDVHKSRPSLQVITE ASTGQSQHLIRTPFAGVDDFFIPPTNLIINHIFGFI FNKSDPAVHKVDLETMMPLKTIGLHHHGCVPA MAHTHLGGYFFIQCRQDSPAARQLLVDSVTD SVLGPNGDVTGTPTSPDGRFTVSAADSPWLHV QEITVRGEIQTLYDLQNSGISDLAFQRSFTESNQ YNYAALHTEPDLLELSTGKVGMLKNLKEPPA GPAQFVGGTHRIMRI-SGLECYLLTPARESLE NGRQNTLRCEVSGIKGGITVWV
3152	A	1	2645	GAGWQVSLTGRWSPGREAGAGEVrQDPGSTAA SPSSCDADLSARMARGERRRRVPAEGVRTAER AARGGPGRRDGRGGGPRSTAGGVALAVVLSL ALGMSGRWVLAWYRARRAVTLHSAPAVLPADS SSPAVAPDLFWGTYRPHVYFGMKTRSPKPLLTG LMWAQQGTTPTGTPKLRTCEQGDGVGPYGWFE HDGLSFGRQHIQDGLRLTEFVKRPGGQHGGD WSWRVTVEPQDSGTSALPLVSLFFYVVDGKEV LLPEVGAQKQLKFISGHTSELGDFRFTLLPPTSPG DTAPKYGSYNVFWTSNPGPLLTEMVKSRLNSW FQHRPPGASPERYGLPGSLKWEDRGPSGQGGQ QFLIQQVTLKIPISIEFVFESGSAQAGGNQALPRLA GSLLTQALESHAEGFRERFEKTFQLKEKGLSSGE QVLGQAALSGLLGIGYFYGGQLVLPDIGVEGSE QKVDPALFPPVPLFTAVPSRSFFPRGFLWDEGFH QLVVQRWDPSTLREALGHWLGLLNADGWIGRE QILGDEARARVPPEFLVQRAVHANPPTLLLPVAH MLEVGDPDDLAFRLKALPRLHAWFSWLHQSA GPLPLSYRWRGRDPALPTLLNPKTLPSGLDDYPR ASHPSVTERHDLRCWVALGARVLTALAEHLGE AEVAAELGPLAASLEAAESLDELHWAPELGVFA DFGNHTKAVQLKPRPPQGLVRVVGPRPQQLQYV DALGYVSLFPLLLRLLDPTSSRLGPLLDLADSRH

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LWSPFGLRSLAASSSFYQQRNSEHDPYPYWRGAV WLVNLYALGALHHYGHLEPHQARAALKHGE LRANVVGNVWRQYQATGFLWEQYSDRDGRGM GCRPFHGWTSVLVLLAMAEDY
3153	A	1	4312	MVIKTDELPAAPADSAREHGSQAGGKGRPGAA AVLLADLERDARQGEALPGAAMAGLAPLKPE ASRSSSPGPTGCIRARVAAEAGTRNPGNAGAELE SWLPCCHGHPETPEPRGGQLPTAPELPSVMLLNG DCPESLKKEAAAAEPPRENGLEAGPGDETTGQ EVTVIQDTGFSVKILAPGIEPFSLQVSPQEMVQEIH QVLMDREDTCHRTCFSLHLDGNVLDHFSSELSRV EGLQEGSVLRVVEEPTVREARHVRHVRDLLKS LDPSDAFNGVDCNSLSFLSVFTDGDLDGDSGKRK KGLEMDPIDCTPPEYILPGSRERPLCPLQPQNRD WKFLQCLKVLTMGSWNPPGNRKMHGDLMYLF VITAEDRQVSITASTRGFYLNQSTAYHFNPKPASP RFLSHSLVELLNQISPTFKKNFAVLQKKRVQRHP FERIATPFQVYSWTAPQAEHAMDCVRAEDAYTS RLGYEEHIPGQTRDWNEELQTTRELPRKNLPERL LRERAIFKVHSDFTAAATRGAMAVIDGNVMAIN PSEETKMOMFIWNNIFFSLGFDVRDHYKDFGGD VAAYVAPTNDLNGVRTYNAVDEGLYTLGTVV VDYRGYRVTAQSIIPGILERDQEQSVIYGSIDFGK TVVSHPRYLELLERTSRPLKILRHQVLNDRDEEV ELCSSVECKGIIGNDGRHYLDLLRTFPPDLNFLP VPGEELPEECARAGFPRAHRHKLCLLQELVDA FVEHRYLLFMKLAALQLMQNASQLETPSSLEN GGPSSLESKSEDPPGQEAGSEEEGSSASGLAKVK ELAETIAADDGTDPRSREVIRNACKAVGSSISTAF DIRFNPDIFSPGVRFPESCQDEVRDQKQLLKDAA AFILSCQIPG... QANLNMRYLGKVLELVLRSPA... ELITRSKHFITYLQGVLSGLSA... SYNPVVAHLPADELVSKKRNRKR... TAWAVMTPQELWKNICQEAKNYFED... DQAVETYGLQKITLLREISLKTGIQVLLKEYSFD RHKPAFTEEDVLNIFPVVKHVNPKASDAFHFFQS GQAKVQQGFLKEGCELINEALNLFNNVYGAMH VETCACLRLARLHYIMGDYAEALSNQQKAVL MSERVMGTEHPNTIQEYMHLLALYCFASSQLSTA LSLLYRARYLMLLVFGEDHPEMALLDNNIGLV HGVMBYDLSLRFLLENALAVSTKYHGPKALKVAL SHHLVARVYESKAEFRSALQHEKEGYTIYKTQL GEDHEKTESSEYLKCLTQQAVALQRTMNEIYR NGSSANIPPLKFTAPSMASVLEQLNVINGILFIPLS QKDLENLKAEVARRHQLQEASRNDRRAEPPMA TEPAPAGAPGDLGSQPPAAKDPSPSVQG
3154	A	416	4082	KFKLIKIMLLTLILLPVVSKFSFVLSAPQHWSCP EGTLAGNGNSTCVGPAPFLIFSHGNSIFRIDTEGT NYEQLVVDAGVSVIMDFHYNEKRIYVVDLERQ LLQRVFLNGSRQERVCNIEKNVSGMAINWINEEV IWSNQQEGITVTDMKGNNSHILLSALKYPANVA VDPVERFIFWSSEVAGSLYRADLDGVGVKALLE TSEKITA VSLDVLDKRLFWIQYNREGSNSLICSD YDGGSVHISKHPTQHNLFAMSLFGDRIFYSTWK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				MKTIWIANKHTGKDMVRINLHSSFVPLGELKVV HPLAQPKAEDDTWEPEQKLCKLRKGNCSSTVCG QDLQSHLCMCAEGYALSRDRKYCEGNDWKYCE DVNECAFWNHGCTLGCKNTPGSYYCTCPVGFVL LPDGKRCHQLVSCPRNVSECSHDCVLTSEGPLCF CPEGSVLERDGTKSCGSSPDNGGCSQLCVPLSP VSWECDCFPGYDLQLDEKSCAASGPQPFLLFANS QDIRHMHFDGTDYGTLLSQQMGMVYALDHDVP ENKIYFAHTALKWIERANMDGSQRERLIEEGVD VPEGLAVDWIGRRFYWTDRGKSLIGRSDLNGKR SKIITIENISQPRGIADVHPMAKRLFWTDTGINPRIE SSSLQGLGRLVIASSDLIWPSTGIDTIDFLDKLYWC DAKQSVIEMANLDGSKRRRLTQNDVGHPPFAVA VFEDYVWFSDWAMPSVIRVNKRTGKDRVRLQG SMLKPSSLVVVHPLAKPGADPCLYQNGGCEHIC KKRLGTAWCSCREGFMKASDGKTCALDGHQL LAGGEVDLKNQVTPDLILSKTRVSEDNITESQHM LVAEIMVSDQDDCAPVGCSMYARCISEGEDATC QCLKGAGDGKLCSDIDECMGVPVPCPASSKCI NTEGGYVCRCEGYQGDGIHCLDIDECQLGVHS CGENASCTNTEGGYTCMCAGRLSEGLICPDSTP PPLREDDHHYSVRNSDSECLSHDGYCLHDGV CMYIEALDKYACNCVVGYGICERCQYRDLKWE LRHAGHGQQQKVTVVAVCVVVLVMLLLSLWG AHYYRTQKLLSKNPKNPYEESRDVRSRRPADT EDGMSSCPQWFVVIKEHQDLKNGGQPVAGED GQAADGSMQPTSWRQEPQLCGMGTEQGCWIPV SSDKGSCPQVMERSFHMPSYGTQTLGGVEKPH SLLSANPLWQQRALDPPHQMELTQ
3155	A	533	212	GTSGWYWERLAERRGRLWSREEAMATMENKVI CALVLSMTALGTAFAYTETCTVAPFERCNCG FPGVETCCANKGCCFDLTVRGVPWCFYPNTID VPPEEECEF
3156	A	2	1585	PRVRAADVAAAGAAVVSAGMAKSNGENGPAP AAGESLSGTRESLAQGPDAATTDELSSLGSDSEA NGFAERRIDKFGFTVGSQGAEGALEEVPLEVLRLQ RESKWLDMLNNWDKWMMAKKHKKIRLRCQKGI PPSLRGRAWQYLSGGKVKLQQNPGKFDELDMSP GDPKWLDVIERDLHRQFFHEMFVSRGGHGQQD LFRVLKAYTLRPEEGYCAQAQPIAAVLLMHMP AEQAFWCLVQICEKYLPGYYSEKLEAIQLDGEIL FSLQKVSPVAHKHLSRQKIDPLLYMTEWFMCA FSRTL PWSSVLRVWDMFFCEGVKIIIFRVLVLLK HALGSPEKVKACQGYETIERLRSLSPKIMQEAF LVQEVVELPVTERQIERHLLQLRRWQETRGELO CRSPRLHGAKAILDAEPGPRPALQSPSIRLPLD APLPGSKAKPKPKQAQKEQRKQMKGRGQLEKP PAPNQAMVVAAGDACPPQHVPKDSAPKDSAP QDLAPQVSAHHRSQESLTSQESEDTYL
3157	A	3	601	SSAMGSRSSHAAVIPDGDSIRRETGFSQASLLRLH HRFRALDRNKKGYLSRMDLQQIGALAVNPLGDR IIESFFPDGSQRVDFPGFVRVLAHFRPVEDEDTET QDPKKPEPLNSRRNKLHYAFQLYDLDRDGKISR HEMLQVLRMLVGVQVTEEQLENIADRTVQEAD EDGDGAVSFVEFTKSLEKMDVEHKMSIRILK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3158	A	2	409	ISSCPHTAYEGSMSTLSNFTQTLEDVFRIRIFTYM DNWRQNTTAEQEALQAKVDAENFYVILYLMV MIGMFSFIIVAILVSTVKSRRHSNDPYHQYIVE DWQEKYKSQILNLEESKATIHENIGAAGFKMSP
3159	A	3	416	PWGAAELDMGRRDAQLLAALLVLGLCALAGSE KPSPCQCSRLSPHNRTNCGFPGITSDQCFDNGCCF DSSVTGVPWCFHPLPKQESDQCVMESDRRNCG YPGISPEECASRKCCFSNFIFVPWCFFPKSVEDC HY
3160	A	179	409	KPKTKILKMVYYPELFWVVSQEPFPNKDMEGRL PKGRLPVPKEVNRKKNDETNAASLTPLGSSELRS PRISYLHFF
3161	A	683	1186	LSSTGGLHAAACAAAMSLVPEKFOHILRVLNTN IDGRRKIAFAITAIKGVGRRYAHVVLKADIDL KRAGELTEDEVERVITIMQNPRQYKIPDWFLNRQ KDVKGDKYSQVLANGLDNKLRDLERLKKIRA HRGLRHFVGLRVRGQHTKTTGRRGRTVGVSKK K
3162	A	1	1938	GMPSRSGRAAPGPPPPPPPGQAPRWSRWVP GRLLLLLLPALCCLPGAARAAAAAAGAGNRAA VAVAVARADEAEAPFAGQNWLSYGYLLPYDS RASALHSAKALQSAVSTMQQFYGIPVTGVLDT TIEWMKKPRCGVPDHPHLSRRRRNKRYALTGQK WRQKHITYSIHNYTPKVGELDRKAIRQAFDVW QKVTPLTFEEVPYHEIKSDRKEADIMFFASGFHG DSSPFDGEGGFLAHAYFPGPGIGGDTHFDSDEPW TLGNANHDGNDLFLVAVHELGHALGLEHSSDPS AIMAPFYQYMETHNFKLPQDDLQGIQKIYGPPAE PLEPTRPLPLPVRRIHSPSERKHQRPRPPPLG DRPSTPGTKPNICDGNFNTVALFRGEMFVKDR WFWPLNNRVQEGYPMETQFWKCI PAPT YERAQGRFVPRGDKYWFKEVTVEPGYMSLC BLGSCLPREGIDTALRWEPVGKTYFFKGERYWR YSEERRATDPGYPKPITVWKGIPQAPQGAFISKE GYTTYFYKGRDYWKFDNQKLSVEPGYPRNLRD WMGCNQKEVERRKERRLPQDDVDIMVTINDVP GSVNAVAVVIPCILSLCILVLVYTIFQFKNKTGPQ PVTYYKRPVQEWV
3163	A	1235	2223	SRLSLQFYVSFRRTGLFTCKLIVEIFFRNYMNDL RTNVFVRFPETIACACIYLAARALQPLTRPHW FLLFGTTEEEIQEICIETLRLYTRKKPNYELLEKEV EKRKVALQEAKLKAKGLNPDGTPALSTLGGFSP ASKPSSPREVKAEKSPISINVKTVKKEPEDRQQA SKSPYNGVRKDSKRSRNSRSASRSRSTRSRSS HTPRRHYNRRSRSGTYSSRSRSTRSHSESRR HHNHGSPHLKAKHTRDDLKSSNRHGHKRRKRS RSQSKSRDHSDAAKKHRRHERGHHRDRRERSRSF ERSHKSXHHGGSRSRSGHGRHRR
3164	A	3	3274	DCRLQAAMPTNFTVVPVEAHADGGGDETAERT EAPGTEGPEPERPSPGDGNPRENSPFLNNVEVE QESFFEGKNMALFEEEMDSNPMVSSLLNKLANY TNLSQGVVEHEDEESRRREKAPRMGTFIGVY LPCLQNILGVILFLRLTWIVGVAGVLESFLIVAMC CTCTMLTAISMSAIAITNGVVPAGGSYYMISRLG PEFGGAVGLCFYLGTTFAGAMYILGTIEIFLTYISP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GAAIFQAEAAGGEAAAMLHNMRVYGTCTLVLM ALVVFVGKYNKLALVFLACVVLSILAIYAGVI KSAFDPPDIPVCLLGNRTLSRRSFDACVKAYGIH NNSATSALWGLFCNGSQPSAACDEYFIQNNVTEI QGIPGAASGVFLENLWSTYAHAGAFVEKKGVPS VPVAEESRASTLPYVLTIDIAASFILLVGIYFPSVT GIMAGSNRSGDLKDAQKSIPTGTILAVTTSFIYLS CIVLFGACIEGVVLRDKFGEALQGNLVIGMLAW PSPWVIVIGSFFSTCGAGLQTLTGAPRLLQAIARD GIVPFLQVFGHGKANGEPTWALLTLVICETGILI ASLDSVAPILSMFFLMCYLFVNLAACAVQTLLRTP NWRPRFKFYHWLSFLGMSLCLALMFICSWYYA LSAMLIAGCIYKYIEYRGAKEWGDGIRGLSLNA ARYALLRVEHGPPHTKNWRPQVLVMLNLDAEQ AMKHPRLLSFTSQLKAGKGLTIVGSVLEGTYLD KHMEAQRAEENIRSLMSTETKKGFCQLVSSSLR DGMSHLIQSAGLGGLKHNTVLMASWASWKQED NPFSWKNFVDTVRDTTAAHQALLVAKNVDSPFQ NQERFGGGHIDVWVWHDGGMMLLPFLLRQH KVWRKCRMRIFTVAQVDDNSIQMKKDLQMFY HLRISAEVEVVEMVENDISAFTYERTLMMEQRS QMLKQMQLSKNEQEREAQLIHDRNTASHTAAA ARTQAPPTPKVQMTWTREKLIAEKYRSRDTSL SGFKDLFSMKPDQSNVRRMHTAVKLNQVVLNK SQDAQVLVLLNMPGPPKNRQGDENYMEFLEVLTE GLNRVLLVRGGGREVITIYS
3165	A	3	2681	GRGARGGSGAGALRGCRGYLQKLSGKGPSRGY RSRWVFDARRCYLYYFKSPQDALPLGHLDIAD ACFSYQGPDEAAEPGTEPPAHFQVHSAGAVTVL KAPNRQLMTYWLQELQQRWEYCNSLDMVKW DERTSPTPGDFKGLVARDNTDLYPHPIAEK ARNVLA VETVPGE LVGEQA QAPGHI NSINF YSLKQWGNELKNSMSSFRPGRGHNSRRTVFYT NEEWELLDPTPKDLEESIVQEEKKKLTPEGNKGV TSGGFPPDFGRNPYKGRPLKDIIGSYKNRHSSG DPSSEGTSGSGSVSIRKPASEMQLQVQSQQEELE QLKKDLSSQKELVRLQQTVRSSQYDKYFTSSRL CEGVPKDTLELLHQKDDQILGLTSQLERFSLEKE SLQQEVRTLKSKVGELNEQLGMLMETIQAQDEV IILKSEGEENGPPPTVAPSSPSVVPVARDQLELDR LKDNLQGYKTQNKFLNKEILELSALRRNPERRER DLMARNSSLEAKLCQIESKYLLQEMKTPVCSE DQGPTREVIAQLLEDALQVESQEQAQAFVKPHL VSEYDIYGFRTVPEDDEEEKLVAKVRALDLKTL YL TENQEVSTGVK WENYFASTVNREMMCSPEL KNLIRAGIPHEHRSKVWKWCVDHRHTRKFKDNTE PGHFQTLQKALEKQNPASKQIELDLLRTLNNK HYSCTSEGIQKLRNVLLAFSWRNPDIGYCQGLN RLVAVALLYEQEDAFWCLVTIVEVFMPRDYIT KTLG SQVDQ RVFRDL MSEKLPRLHGHFEQYKV DYTLITFNWFLVVFVDSVVS DILFKIWDSFLYEGP KVIFRFALALFKYKEEILKLQDSMSIFKYLYFT RTILDARSGTDAPTTWRKSGWS
3166	A	10	4070	FPGPTISSNSQLYRASALFETIRHEAQLSTDYKLS LFDLQTSSYQALQRVLVSLGHHDEALAVAERGR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				TRAFADLLVERQTGQQDSDPYSPVTIDQILEMVN GQRGLVLYYSLAAGLYSWLLAPGAGIVKFHEH YLGENTVENSSDFQASSSVTLPTATGSALEQHIAS VREALGVESHYSRACASSETESEAGDIMDQQFEE MNNKLNSVTDPTGFLRMVRRNNLFNRSCQSMTS LFSNTVSPTQDGTSSLPRRQSSFAKPLRALYDLL IAPMEGGLMHSSGPVGRHRQLLVLEGELYLIPF ALLKGSSSNEYLYERFGLLA VPSIRSLSVQSKSHL RKNPPTYSSSTSMAAVIGNPKLPSAVMDRWLWG PMPSAEEEA YMVSELLGCQPLVGSVATKERVMS ALTQAECVHFATHISWKL SALVLTSPMDGNPASS KSSFGHPYTIPESLRVQDDASDGEISDCPPLQEL LLTAADVLDLQLPVKLVVLGSSQESNSKVAADG VIALTRAFLAAGAQCVLVSLWPVPVAAFKMFH AFYSSLLNGLKASAAALGEAMKVVQSSKAFSHP NWAGFMLIGSDVKLNSPSSLIGQALTEILQHPER ARDALRVLLHLVEKSLQRIQNGQRNAMYTSQQS VENKVG GIPGWQALLTAVGFRLDPPTSGLPAAV FFPTSDPGDRLQCCSSTLQSLGLPNPALQALCK LITASETGBQLISRAVKNMVGMHLHQLVQLQAG EKEQDLASAPIQVSISVQLWRLPGCHEFLAALGF VLCEVGQEEVILKTGKQANRRRTVHFALQSLSLF DSTELPKRLSLDSSSSLESLSASAQSVSNALPLGYQ QPPFSPTGADSIASDAISVYSLSSIASSMSFVSKPE GGSEGGGPGGRQDHDRSKNAYLQRSTLPRSQLP PQTRPAGNKDEEEYEGFSIISNEPLATYQENRNTC FSPDHKQPQPGTAGGMRVSVSSKGSISTPNSPVK MTLIPSPNSPFQKVGKLASSDTGESDQSSTETDST VKSQEESNPKLDPOELAOKILEETQSHLIAVERLQ RSGGQVSKSNPEDGVQAPSSTA VFRASETSAFS RPVI SHQKSGSPVTVKPEPPAPSSSLPKVSSVS SPTTSEMSKSGSPSGRPSPGCDSQTSQLDQPL FKLKYPSSPYSAHISKSPRNMSPSSGHQSPAGSAP SPALSYSSAGSARSSPADAPDIDKLKMAAIDEKV QAVHNLKMFWQSTPGHSTGPMKIFRGAPGTMTS KRDLVSLNLSRPNKKEEGVDKLELKELSLQQH DGAPPKAPPNGHWRTETTSLGSLPLPAGPPATAP ARPLRLPSGNGYKFLSPGRFFPSSKC
3167	A	1	762	AARRRQKGKEENMMMDLFETGSYFFYLDGENV TLQPLEVAEGSPLYPGSDGTLSPCQDQMPPEAGS DSSGEEHVLAPPGLOPPHCPGQCLIWACKTCRK SAPTDRRKAATLRERRRLKINEAFEALKRRTVA NPNQRLPKVEILRSAISYIERLQDLLHRLDQQEK MQELGVDPFSYRPKQENLEGADFLRTCSSQWPS VSDHSRGLVITAKEGGASIDSSASSSLRCLSSIVDS ISSEERKLPCVEEVVEK
3168	A	701	246	TSRRVTMKFNPVFTSDRSKNRKRHFNAPSHVRR KIMSSPLSKELRQKYNVRSMPIRKDDDEVQVVRG HYKGQIGKVQVYRKKYVIYIERVQREKANGT TVHVGHPKVVITRLKLDKDRKKILERKAKSRQ VGKEKGKYKEELIEKMQE
3169	A	156	3168	GPGGAISLSVEAKAGADLLVKGKQARMDIYDTQ TLGVVVFGGFMVVSAGIFLVSTFSMKETS YEEA LANQRKEMAKTHHQKVEKKKKKEKTVEKKGKT KKKEEKPNKIPDHPAPNVTVLLREPVRAPAV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				AVAPTPVQPPHIVAPVATVPAMPQEKCLASSPKDK KKKEKKVAKVEPAVSSVNSIQVLTSKAAILETA PKEGRNTDVAQSPEAPKQEPAPAKKSGSKKGP PDADGPLYLPYKTLVSTVGSVMFNEGEAQRLEI LSEKAGIIQD'TWHKATQKGDPAVAILKRQLEBKEK LLATEQEDAAVAKSKLRELNKEMAAEKAKAAA GEAKVKKQLVAREQEITAVQARMQASYREHVK EVQQLQGGKIRTLQEQLENGPN'TQLARLQQENSIL RDALNQATSQVESKQNAELAKLRQELSKVSKEL VEKSEAVRQDEQQRKALEAKAAAFQVQLQ ASHRESEEALQKRLDEVSRCLHTQSSHASLRAD AEKAQEQQQMAELHSLQSSAEVRSKCEELS GLHGQLQEARAENSQLTERIRSIALLAEGQARD AQDVQASQAEADQQQTRLKELESQVSGLEKBAI ELREAVEQQKVKNNDLREKNWKAMEALATAEQ ACKEKLHSLTQAKEESEKQLCLIEAQ'TMEALLAL LPESVLAQQNYTEWLQDLKEKGPTLLKHPPAP AEPSSDLASKLREAEETQSTLQAECDQYRSILAET EGMLRDLQKSVEEEEQVWRKVGAAEEELQKS RVTVKHLEEIVKLGKELESSDQVREHTSHLEAE LEKHMAAASAEQNYAKEVAGLRQLLESQSQL DAAKSEAQKQSDALVRQQLSEMKSHVEDGDI AGAPASSPEAPAEQDPVQLKTQLEWTEAILEDE QTQRQKLTAEFEEAQTSACRLQEELEKLRTAGPL ESSETEEASQLKERLEKEKLTSDLGRAATRLQE LLKTTQEQQLAREKDTVKKLQEQLEKAEDGSSSK EGTSV
3170	A	6730	4027	THASEKYSYGHLP'THSITAHPMV'TIRISDRQLIQ PYIHNSWLLFAALALYSAHLASAEDVDGEKLD PQTRSSATT'LRSQCMQLVGDCLMKAHQKGKGLK ALALEGVLPDGDSSLEDEALPV'TVPTGASFEQLE KKA'VQGAELSEAGNGKRAVHEIRPV'KQRNK ADKGVSLSKDPSCQ'TQISDSPA'DASPTGLPDAB DSEVSSQKPIEEKAVTPSPEQVFAECSQKRILGLL AAMLPPLKSGPTVPLIDLEHVLPLMFQVVISNAG HLNETYHLTLGLLGQLIRLLPAEVDAAVIKVLSA KHNLFAGDSSIVPDGWKT'THLLFSLGAVCLDS RVGLDWACSMAEILRSLNSAPLWRDVIATFTDH CIKQLPFQLKHTNIFTLLVLVGFPQVLCVGT'RCV YMDNANEPHN'VILKHFT'EK'NRAVIVDVKTRKR KTVKDYQLVQKGGGQECGDSRAQLSQYSQHFA FIASHLLQSSMDSHCPEAVEATWVLSLALKGLY KTLKAHGFEEIRATFLQTDLLKLLVKKCSKGTGF SKTWLLRDLEILSIMLYSSKKEINALAEHGDLEL DERGDREEEVERPVSSPGDPEQKKLDPLEGLDEP TRICFLMAHDALNAPLHILRAIYELQMKKTDYFF LEVQKRFDGDEL'TDERIRSLAQ'RWQPSKSLRLE EQSAKAVDTDMILPCLSRPARCDQATAESNPVT QKLISSESELQSQSYAKQRRSKSAALLHKELNCK SKRAVRDYLF'RVNEATAVL'YARHVLASLLAEWP SHVPVSEDILELSPAHMTYILD'FMFMQLEEKHE WEKVVMQTELVLTHQVLP'PLPHRLPPVSASWSEA TCVAVQLPDRCECSKGRVTVSSPKDWASEELRG PERDFQLNQKALSPSSQFPSAEILRHIR
3171	A	557	89	GTRAGPVK'DREAFQRLN'FLYQAAHCVLAQDPEN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QALARFYCYTERTIAKRLVLRDPSVKRTLRCGC SSLLVPGLTCTQRQRRRCRGQRWTVQTCLTCQRS QRFLNDPGHLLWGDRPEAQLGSQADSKPLQPLP NTAHSISDRLPEEKMQTQGSSNQ
3172	A	2	496	FRAGAGRGRRRGEVTSPLSPEPLAFQSLATSRR PEPQTTQTVRSSALPAPPASPMQYAPSPDFKRA LDSSPEANTEDDKTEEDVPMPPKNYLWLTIVSCFC PAPYPINTVALVFSIMSLNSYNDGDYEGARRLGRN AKWVAIASIIIGLLIIGISCAVHFTRNA
3173	A	2	4048	FRSGGCRRRAWTSRWPQRRRSPESCEAPLSAPL WGPQRGLPGREPLRSRSASAIALRTIGHILALLLR LLHLGLSGGGCREDVPPSGRGKKEKMKKHRRRA LALVSCFLCSLVWLPSWRVCKEASSASASSYY SQDDNCALENEDVQFQKKDEREGPINAESLGKS GSNLPISPKEHKLKDDSIQVQNTESKKLSPPVVE TLPTVDLHEESSNAVVDSETVENISSSTSEITPIS KLDEIEKSGTIPIAKPSETEQSETDCDVGEALDAS APIEQPSFVSPPDSL VGQHIENVSSSHGKGKITKSE FESKVSASEQGGGDPKSALNASDNLKNESSDYT KPGDIDPTSVASPKDPEDIPTFDEWKKKVMEVEK EKSQSMHASSNGGSHATKKVQKNRNNYASVEC GAKILANPEAKSTSAILIENMDLYMLNPCSTKI WVIELCEPIQVKQLDIANYELFSSTPKDFLVSISD RYPTNKWIKLGTFFHGRDERNVQSFPLDEQMYAK YVKMFYIKVELLSHFGSEHFCPLSLIRVFGTSM VEEYEEIADSQYHSERQELFDEDYDYPLDYNTGE DKSSKNLLGSATNAILNMVNIAANILGAKTEDLT EGNKSISENATATAAPKMPSTPVPSTPVPSPPEYVT TEVHTHDMPESTPDTPKESPIVQLVQEEEEASPS TVTLGSGGEQDESSPWFESETQIFCSELTICIS SFSEYKRWCSVRVAJYRCEETALSQCLTVV LAQPIILLPAESVDVSQELTELENTNIEREAE TVVLGDLSSSMHQDDLNVHTVDAVELEPSHSQT LSQSLLLDITPEINPLPKIEVCEVEYEAGHIPSPVI PQESSVEIDNETEQKSESFSSEIENETIYETNKVNE LMDNIIKEDVNSMQIFTKLSETIVPPINTATVPDN EDGEAKMNIADTAKQTLISVVDSSSLPEVKEEEQ SPEDALLRGLQRTATDFYAEQNSTDLGYANGN LVHGSNQKESVFMRLNNRIKALEVNMSLSGRYL EELSQRYSRQMEEMQKAFNKTIVKLQNTSRIAE EQDQRQTEAIQLLQAQLTNMTQLVSNLSATVAE LKREVSDRQSYLVISLVLCVVLGLMLCMQRCRN TSQFDGDYISKLPKSNQYPSPKRCFSSYDDMNLK RRTSFPLMRSKSLQLTGKEVDPNDLYIVEPLKFSP EKKKKRCKYKIEKTIETIKPEEPLHPIANGDIKGRK PFTNQDFSNMGEVYHSSYKGPPEGSSETSSQS EESYFCGISACTSLCNGQSQKTKTEKRALKRRRS KVQDQGKLIKTLIQTKSGSLPSLHDIKGNKEITV GTFGVTA VSGHI
3174	A	485	4668	RKCSKEKASKTPSQKIPTTPCCVLQAGPEPRSLAE RMGADGETVVLKNMLIGVNLILLGSMIKPSECQL EVTTERVQRQSVBEEGGIANYNTSSKEQPVVFNH VYNINVPLDNLCSGLEASAEQEVSAEDELTALEY MGQTS DHESQVTFTHRINFPPKACPCASSAQVLQ ELLSRIEMLEREVSVL RDQC NANCCQESAATGQL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				DYIPHCSGHGNFSFESCGCICNEGWF GKNCSEPY CPLGSSRGVCVDGQCICDSEYSGDDCSELRCPT DCSSRGLCVDGECVCEEPYTGEDCRELRCPGDCS GKGRCA NGTCLCEEGYVGEDCGQRQCLNACSG RGQCEEGLCVCEEGYQGPDCSAVAPPEDLRVAG ISDRSIELEWDGPMAVTEYVISYQPTALGGLQLQ QRVPGDWSGVTTITLEPGLTYNISVYA VISNLSL PITAKVATHLSTPQGLQFKTTITETTVEVQWEPFSF SFDGWEISFIPKNNEGGVIAQVPSDVTSTFNTGLK PGEEYIVNVVALKEQARSPTSASVSTVIDGPTQI LVRDVSDTVAFVEWIPRAKVD FILLKYGLVGGE GGRTTFR LQPPLSQYSVQALRPGSRYEVSVA VR GTNESDSATTQFTTEIDAPKNLRVGSRTATSLDL EWDNSEAEVQBYKV VYTTLAGEQYHEVLVPRGI GPTTRATL TDLVPGTEYGVGISA VMNSQQSVPA T MNARTELDSPRDLMTV TASSETSISLIWTKASGPID HYRITFTPSSGIASEVTVPKDRTSYTLTDLEPGAE YIISVTAERGRQQSLESTVDAFTGFRPISHLHFSH VTSSSVNITWSDPSPPADRLILNYSRDEEEEMME VSLDATKRHA VLMGLQPATEYIVNLVA VHGTVT SEPIVGSITTGIDPPK DITISNVTKDSVMVSWSPPV ASFDYYRVS YRPTQVGRLDSSVVPNTVTEFTITR LNPATEYEISLNSVRGREESERICTLVHTAMDNP VDLIATNITPTEALLQWKAPVGEVENYVIVLTHF AVAGETILVDGVSEEFRLVDLLPSTHYTATMYAT NGPLTSGTISTNFSTLLDPPANLTASEVTRQSALIS WQPPRAEIE NYVLT YKSTDGSRKELIVDAEDTWI RLEGLLENTDYTVLLQAAQDTTWSSITSTAFTTG GRVFFHPQDCAQHLMNGDTLSGVYPIFLNGELS QKLQVYCDMTIDGGGWIVFQRRQNGQTDFFRK WADYEVGFGNVEDEFWLCI DNIHRITSQGRYE RVDMLRQQAFAFASYDRFSVEDS KLRI YNGTAGDSL SYHQGRPFSTEDRDNDVAVTNCA MSYKGA W WYKNCHR TNLNGKYGESRHSQGIN VYHWWKGHEFSIPFVEMKMRPYNHRLMAGRKRO SLQF
3175	A	2	623	RLQLPACPALSAAHPLALPSFSSQCHRAEAAAA AATAEGTMASGVTVNDEVIKVFNDMKVRKSST QEEIKKRKKA VLFCLSDDKRQIIVEBAKQILVGDI GDTVEDPYTSFVKLLPLNDCRYALYDATYETKE SKKEDLVFIFWAPESAPLKSMTYASSKDAIKKK FTGIKHEWQVNGLDDIKDRSTLGEKLGGNVVVS LEGKPL
3176	A	99	1567	PRGCWSSCLDAMFRLNLSALAEAVGSRWYH GGSQPIQIRRLMMVAFLGASAVTASTGLLWKR AHAEPPCVDNLKSDIGDKGNKDEGDVCNHEK KTADLAPHPEEKKKKRS GFRDRKVM EYENRIRA YSTPDKIFRYFATLKVISEPGAEVFMTPEDFVRS ITPNEKQPEHLGLDQYIKRFDGKTEKISQEREKF ADEGSIFYTLGECGLISFSDYIFLTTVLSTPQRNFB IAFKMFDLNGDGEVDMEEFEQVQSIRSQTSMG MRHRDRPTTGNTLKSGLCSALTTYFFGADLK GK LTIKNFLEFQRKLQHDVLKLEFERHDPVDGRITE RQFGGMLLAYSGVQSKKLTAMQRQLKKHFKEG KGLTFQEVENFFTFLKNINDVD TALSFYHMAGAS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				LDKVTMQQVARTVAKVELSDHVCDDVVFALFDC DGNGELSNKEFVSIMKQRLMRGLEKPKDMGFTR LMQAMWKCAQETAWDFALPKQ
3177	A	182	648	LGVVGSGAAVGGGRQAARGAALGRRPMAAVLG ALGATRRLLAALRGQSLGLAAMSSGTHRLTAEE RNQAILDLKAAGWSELSEDAIYKEFSFHNFNQA FGFMSRVALQAEKMNHHPWFNPNVYKQITLTS HDCGELTKKDVKLAKFIEKAAASV
3178	A	8	612	ACGCRSFCGSTVMSLLLYALPALGSYAMLSIFF LRRPHLLHTPRAPTFRIRLGAHRGGSGELLENTM EAMENSMAQRSDLELDCQLTRDRVVVVSHDE NLCRQSGLNRDVGSGLDFEDLPLYKEKLEVYFSPG HFAHGSDDRMVRLEDLFQRFPRTPMSVEIKGKN EELIREIAGLVRRYDRNEITIWASEKSSVMKKCK
3179	A	88	1496	QETSKMETLSFPRYNVABIVHIRNKILTGADGKN LTKNDLYPNPKPEVLHMTYMRALQIVYGIRLEHF YMPVNSEVMYPHLMGFLPFSNLVTHLDSFLPI CRVNDFETADILCPKAKRTSRFLSGIINFHFREAC RETYMEFLWQYKSSADKMQLNAAHQEALMK LERLDSVPVEEQEEFKQLSDGIQELQQLNQDFH QKTTVLQEGNSQKKSNISEKTKRLNELKLSVVS KEIQESLTKIVDSPEKLKKNYKEKMKDVTQKLK NARQEVVEKYEYIGDSVDCLPSCQLEVQLYQKK IQDLSDNREKLASILKESLNLEDQIESDESELKKL KTEENSFKRLMIVKKEKLATAQFKINKKHEDVK QYKRTVIEDCNKVQKEKGAVYERVTTINHEIQKI RLGIQQLKDAADREKLKSQEIFLNKLTALEKYHD GIEKAAEDSYAKIDEKTAELKRKMFKMST
3180	A	298	7086	GNMACWPQLRLLWKNLTFRRRQTCQLLEVA WPLFIFLILISVRLSYPPYEQHECHFPNKAMPSAG TLPWVQGI...NTPCFRYPT...VVG... ...ARLFSDARRLLYSQKDT...KVLRTL QQIKKSSSNLKLQDFLVDNETFS...LYHNLSLPK STVDKMLRADVILHKVFLQGYQL...SLCNGSK SEEMIQLDQEVSELGCLPREKLAAAEVLRSN MDILKPIRLTLNSTSPFPKELAEATKTLHSLGT LAQELFSMRSWSDMRQEVMTNVNSSSSSTQI YQAVSRIVCGHPEGGLKIKSLNWYEDNNYKAL FGNGTEEDAETFYDNSTTPYCNDLMKNLESSPL SRIWKALKPLLVGKILYTPDTPATRQVMAEVNK TFQELAVFHDLEGMWEELSPKIWTFMENSQEMD LVRMLLDSRDNDHFWEQQLDGLDWT AQDIVAF LAKHPEDVQSSNGSVYTWREAFNETNQAIRTISR FMECVNLNLEPIATEVWLINKSMELDERKFW AGIVFTGITPGSIELPHHVKYKIRMGIDNVERTNK IKDGYWDPGPRADPFEDMRYVWGGFAYLQDVV EQAIIRVLTGTEKKTGVYMQMPYPCYVDDIFLR VMSRSMPLFMTLAWTYSVAVIKGIYVEKEARLK ETMRIMGLDNSILWFSWFISLIPLLVSAGLLVVI LKLGNLLPYSDPSVVFVLSVFAVVTLQCFLIST LFSRANLAAACGGIYFTLYLPYVLCVAWQDYV GFTLKIFASLLSPVAFGFGCEYFALFEEQGIGVQW DNLFEPSVEEDGFNLTTSVSMMLFDTFLYGVMT WYIEAVFPQYGI PRPWYFPCTKSYWFGESDEK SHPGSNQKRISEICMEEETHLKLGVSIQNLVKVY

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RDGMKVAVDGLALNFYEGQITSFLGHNGAGKTT TMSILTGLFPPTSGTAYILGKDIRSEMSTIRQNLG VCPQHNVLFDMLTVEEHIWFYARLKGLSEKHVK AEMEQMALDVGLPSSKLKSKTSQLSGGMQRKLS VALAFVGGSKVVILDEPTAGVDPYSRRGIWELLL KYRQGRITILSTHIMDEADVLDRIAISHGKLCC VGSSFLKNQLGTGYLTLVKKDVESSLSSCRNS SSTVSYLKKEDSVSQSSDAGLGSDESHTLTID VSAISNLIRKXHVSEARLVEDIGHELTYYLPYEA KEGAFVELFHEIDDRSLDLGISSYGISETTLEIFL KVAEESGVDAETSDGTLPARRNRRAFGDKQSCL RPFTEDDAADPNDSIDIPESRETDLLSGMDGKGS YQVKGWKLTOQQFVALLWKRLLIARRSRKGFF AQIVLPAVFVCIALVFSLIVPPFGKYPSELPQWM YNEQYTFVSNDAPEDTGTLELLNALTDPGFGT RCMEGNPIPDTPCQAGEEWTAPVPQTIMDLFQ NGNWTMQNPSPACQCSSDKIKKMLPVCPPGAGG LPPPQRKQNTADILQDLTGRNISDYLKTYVQIIA KSLKNKIWVNEFRYGGFSLGVSNTQALPPSQEV NDATKQMKKHLKLAKDSSADRFLNSLGRFMTG LDTRNNVKVWFNNKGWHAISSFLNINNAILRA NLQKGENPSHYGITAFNHLNLTQQLSEVAPM TTSVDVLVSICVIFAMSFVPASFVFLIQRVSKA KHLQFISGVKPVITYWLSNFVWDMCNVYVPA TLV IIIFICFQKSYVSSTNLPVLALLLLYGWSITPLM YPASFVFKIPSTAYVVLTSVNLFIGINGSVATFVL ELFTDNKLNNDILKSVFLIFPHFCLGRGLIDMV KNQAMADALERFGENRFVSPLSWDLVGRNLFA MAVEGVVFFLITVLIQYRFFIRPRPVNAKLSPLND EDEDVRRERQRILDGGGQNDILEIKELTKIYRRK RKPAVDRIVGPPGECEGLLGNGACKSSTFKM LTCDETVTRGDALNKNRSILSNIHEVHQNM QDAITELLTGREHVEFFALLRGVPEKEVGKVG WAIRKLGLVKYGEKYAGNYSGGNKRKLSTAMA LIGGFVVFLEPTTGMDPKARRFLWNCALSV KEGRSVVLTSHSMEECEALCTRMAMVNGRFR LGSVQHLKNRFGDGYTIVVRIAGSNPDLKPQDF FGLAFPGSVPEKHRNMLQYQLPSSLSSLARIFI LSQSKKRLHIEDYSVSQTILDQVFVNFADQSD DHLKDLSLHKNQTVVDVAVLTSFLQDEKVKESY V
3181	A	215	1367	PPATSQAALPEALSKGRETPRPATHPARSQDVRP LSCPFDFLRDNVEWSEEQAAAAERKVQENSIQR VCQEKQVDYEINAHKYWNDFYKIHENGFFKDR HWLFTEFPPELAPSQNQNHLKDWFLNKSEVPEC RNEDGPGLIMEEQHKCSSKSLEHKTQTTPVEEN VTQKISDLEICADEFPGSSATYRILEVGCVGNTV FPILQTNNDPGLFVYCCDFSSTAIELVQTNSEYDP SRCFAFVHDLCDDEKSYVPKGSLLIILFVLSAI VPDKMQKAINRLSRLKPGGMVLLRDYGRYDM AQLRFKKGQCLSGNFYVRGDGTRVYFFTQEELD TLFTTAGLEKVQNLVDRRLQVNRGKQLTMYRV WQCKYCKPLLSSTS
3182	A	3	1289	GSETQHLPRDPQHLPWDPPQHQDRRRPELFHAF ARDSAPPPSMVLAAETTSQERLQAIKRRKQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				AEIENKRRQLEDERRQLQHLKSKALRERWLLBG TPSSASEGDEDLRRQMDDDEQKTRLLEDVSRL KGIEVLERGDSAPAAAKENAAAPSPVRAPAPSPA KEERKTEVVMNSQQTTPVGTPKDKRVSNTPLRTV DGSPMMKAAMYSVEITVEKDKVTGETRVLSSST LLPRQPLPLGIKVYEDETKVVHAVDGTAEANGIHP LSSEVDELIIHKADEVTLSEAGSTAGAAETRGA EGAARTTPSRREITGVQAQPGEATSGPPGIQPGQE PPVTMIFMGYQNVEDEAETKKVLGLQDTITAE LVIEDAAEPKEPAPPNGSAAEPTEAASREENQA GPEATTSDDPQDLDMKKHRCKCCSIM
3183	A	333	1931	IAPTGGSHSEIQQLGSGGSSSQRRARERTEPS APRPRWGRSARSPGAHKLPGPPRRRDPGAWARL EAAAHRHRSRSGSMGRRMRGAAATAGLWLLAL GSLALWGGLLPRTLPASRPEDRLPRRPARS GGPAPAPRFPLPPPLAWDARGGSLKTRALLTLA AGADGPPRQSRSEPRWHVSARQPRPEESA GVFWSRGLEEQVPPGFSEAQAAAWLEAARGAR MVALERGGCGRSSNRLARFADGTRACVRYGINP EQIQGEALSYYLARLLGLQRHVPPLALARVEAR GAQWAQVQEELRAAHWTEGSVVSLTRWLPNLT DVVVPAPWRSEDGRLRPLRDAGGELANLSQAE VDLVQWTDLILFDYLTANFDRLVSNLFSLQWDP RVMQRATSNLHRGPGGALVFLDNEAGLVHGYR VAGMWDKYNEPLLQSVCFRERTARRVLELHR GQDAAARLLRLYRRHEPRFELAALADPHAQLL QRRLDFLAKHILHCKAKYGRRSGDLVSPGGKER DLGLGYG
3184	A	1	1004	GSTHASADAWAQWFCTEALVMGAPVWYLVA ALLVGFIILFTRSRGRAASAGQEPHNEELAGAG RVAQPCPLEPEEPAGGPFRFRDI GSEAGOR RAQR...AEADE...EBEAVILAQBE...ET HLSGKIGAKKLRLKEEKQARKAQREAEAREE RKRLESQREAEWKKEERLRLEEBQKEETEPKA REEQAQREHEEYKLKEAFVVEEGVGETMEE QSQFLTEFINYIKQSKVVLEDLASQVGLRTQD TINRIQDLLAEGTITGVIDDRGKFTYITPEBLAAVA NFIQRGRVSI AELAQASNSLIAWGRESPAQAPA
3185	A	2981	7173	CLLAGKFSSTLYETGGCDMSLVNFEPARRASNI CDTDSHVSSSTSVRFYPHDVLSLPQIRLNRLTID TDLLEQQDIDLSPDLAATYGPTEEAQKVKHYY RFWILPQLWIGINFDRLLALFDRNREILENVLA VILAILVAFLGSILLIQGFFRDIWVFQCLVIASCQ YSLKSVQPDSSSPRHGHNRIIAYSRPVYFCICCG LIWLLDYGSRLTATKFKLYGITFTNPLVFISARD LVIVFTLCFPIVFFIGLLPQVNTFVMYLCEQLDIHI FGGNATTSLLAALYSFICSIVAVALLYGLCYGAL KDSWDGQHVPVLSIFCGLLVAVSYHLSRQSSDP SVLFSLVQSKIFPKTEKNPEDPLSEVKDPLPEKL RNSVSERLQSDLVVCIVIGVLYFAIHVSTVFTVLQ PALKYVLYTLVGFVGFVTHYVLPQVRKQLPWH CFSHPLLKTEYNQYEVRNAATMMWFEKLHVW LLFVEKNIIYPLIVLNLSSSAETIASPKKLNTEL ALMITVAGLKLRRSSFSSPTYQYVTVFTVLFKF DYEAFSETMLLDLFFMSILFNKLWELLYKLQFVY

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TYIAPWQITWGSFAHFAQPFVPHSAMLFIQAA VSAFFSTPLNPFLGSAIFITSYVRPVKFWERDYNT KRVDHSNTRLASQLDRNPGTYCQOREVEAITEG VEEDEGFCCCEPGHIPHMLSFNAAFSQRWLAW VIVTKYILEGYSITDNSAASMLQVFDLRKVLTTY YVKGIYYVTSSKLEEWLANETMQEGLRLCAD RNYVDVDPFTFNPINDEDYDHRLAGISRESFCVIY LNWIEYCSSRRAKPVDVDKSSSLVTLCYGLCVL GRRALGTASHHMSSNLESFLYGLHALFKGDFRIS SIRDEWIFADMELLRKVVVPGIRMSIKLHQDHFT SPDEYDDPTVL YEAIVSHEKNLVIAHEGDPAWRS AVLANSPLLALRHVMDDGTNEYKIIMLNRRYL SFRVIKVNKECVRGLWAGQQQELVFLRNRNPER GSIQNAKQALRNMINSSCDQPIGYPIFVSPLTTSY SDSHEQLKDILGGPISLGNIRNFIVSTWHRLRKGC GAGCNSGGNIEDSDTGGGTSCGNNATTANNPH SNVTQGSIGNPGQSGTGLHPPVTSYPPTLGTSHS SHSVQSGLVRQSPARASVASQSSYCYSSRHSSLR MSTTGFPVPCRRSSTSQISLRNLPSSIQSRLSMVNQ MEPSGQSGLACVQHGLPSSSSSSQSIACKHHTL VGFLATEGGQSSATDAQPGNTLSPANNSHSRKA EVIYRVQIVDPSQILEGINLSKRKELQWPDEGIRL KAGRNSWKDWSPQEGMEGHVHRWVPCSRDPG TRSHIDKAVLLVQIDDKYVTVIETGVLELGAEV
3186	A	3	470	SLSAMRFLAATFLLALSTAAQAEVPQFKDCGSV DGVKEVNVSPCPTQPCQLSKGQSYSVNVTFTSN IQSKSSKAVVHGILMGVPVPFPIPEPDGCKSGINC PIQKDKTYSYLNKLPVKSEYPSIKLVVEWQLQDD KNQSLFCWEIPVQIVSHL
3187	A	3	470	SLSAMRFLAATFLLALSTAAQAEVPQFKDCGSV DGVKEVNVSPCPTQPCQLSKGQSYSVNVTFTSN IQSKSSKAVVHGILMGVPVPFPIPEPDGCKSGINC PIQKDKTYSYLNKLPVKSEYPSIKLVVEWQLQDD KNQSLFCWEIPVQIVSHL
3188	A	2	3483	PRVRTKLILLVNDKKRYERVGGGPKRLGRDVEM EEMIEQLQEKVHELEKQNDTLKNRLISAKQQLQT QGYRQTPYNNVQSRINTGRRKANENAGLQECPR KGIKFQDADV AETPHPMFTKYGNSLLEEARGEIR NLENVISQSRGQIEELEHLAEILKTQLRRKENEIE LSLQLREQQATDQRSNIRDNVEMIKLHKQLVE KSNALSAMEGKFIQLQEKQRTLKISHDALMANG DELNMQLKEQRLKCCSLEKQLHSMKFSERRIEEL QDRINDLEKERELLKENYDKLYDSAFSAHEEQ WKLKEQQLKVQIAQLETAALKSDLTDKTEILDRL KTERDQNEKLVQENRELQLQYLEQKQQLDELKK RIKLYNQENDINADELSEALLIKAQKEBQKNGDL SFLVKVDSEINKDLERSMRELQATHAETVQELEK TRNMLIMQHKINKDYQMEVEAVTRKMENLQQD YELKVEQYVHLLDIRAARIHKLEAQLKDIAYGTK QYKFKPEIMPDDSVDEFDETIHLERGENLFEIHIN KVTFSSEVLQASGDKEBPVTFCTYAFYDFELQTP VVRGLHPEYNFTSQYL VHVNDLFLQYIQKNTITL EVHQAYSTEYETIAACQLKFHEILEKSGRIFCTAS LIGTKGDIPNFGTVEYWFRLRVPMDQAIRLYRER AKALGYTTSNFKGPEHMQSLSQQAPKTAQLSSTD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				STDGNLNLHITIRCCNHLQSRASHLQHPHYVVY KFFDFADHDTAIPSSNDPQFDDHMYFPVPMNM DLDRYLKSESLSFYVFDDSDTQENIYIGKVNPLI SLAHDRCISGIFELTDHQKHPAGTHVILKWKFA YLPSPGSITTEDLGNFIRSEEPVVQRLPPASSVST LVLAPRPKPRQRLTPVDKKVSFVDIMPHQSDVSQ EGSVDEVKENTTEKMQQKDDVSLSEGQLAEQS LASSEDETEITEDLEPEVEEDMSASDSDDCIIPGI SKNIKQPSEKIRIBIALSLNDSQVTMDDTIQRLFV ECRFYSLPAETPVSLPKPKSGQWVYYNYSNVY VDKENNAKRDILKAILQKQEMPNRSLRFTVVS DPPEDEQDLECEDIGVAHVLDLADMFOEGRDLIE QNIDVFARADGEGIGKLRVTVEALHALQSVYK QYRDDLEA
3189	A	476	1175	MLKSGWHLRSGMVGTLLTILPHWRRTAHVGTN ILTAVSYLKGLWMECVWHSTGIYQCQIYRSLLA LPQDLQAARALMGISCLLSGIACACAVIGMKCTR CAKGTPAKTTFAILGGTLFILAGLLCMGAVSWTT NDVVQNFYNPLLPSGMKFEIGQALYLGFISSLSL IGGTLCLCLSCQDEAPYRPYQAPPRATTTTANTAP AYQPPAAKYKDNRAVSVTSATHSGYRLNDYV
3190	A	267	1037	DRMAWQGLVLAACLLMFPSTTADCLSRCSLCA VKTQDGPKPINPLICSLQCAALLPSEEWERQSF LSFFTPSTLGLNDKEDLGSKSVGEGPYSELAKLS GSFLKELEKSKFLPSISTKENTLSKSLEEKLRGLS DGFREGAESELMRDAQLNDGAMETGTYLAEE DPKEQVKRYGGFLRKYPKRSSEVAGEGDGDSM GHEDLYKRYGGFLRRIRPKLWDNQKRYGGFLR RQFKVVTRSQEDFPNAYSGELEDA
3191	A	29	574	GTSAGAQTGKALCQLKVPTEKLPSPPTMADEID FTEGAGASSTPACCSALRKNGFVVLKTEKFK IVEMSTSKTGKHGHAKVHLVQDFTGKKYEDIC PSTHNMVDPNKRNDYQLICIQDGYLSLLTETGE VREDLKLPEGELGKEIEGKYNAGEDVQVSMCA MSEYAVAIKPKK
3192	A	105	1661	KVSADGMQSCSSGDSADDPLSRGLRRRGQPRV VVGAGLAGLAAAKALLEQGFTDVTVLEASSHIG GRVQSVKLGHATFELGATWIHSGHGNPIYHLTE ANGLLEETTDGERSVGRISLYSKNGVACYLTNH GRRIPKDVVEEFSPLYNEVYNLTQEFFRHDKPVN AESQNSVGVFTRREEVRNRIRNDPDDPEATKRLKL AMIQQYLKVESCESSSHSMDEVSLSAFGEWTEIP GAHHIIPSGFMRVVELLAEGIPAHVIQLGKPVRCI HWDQASARPRGPEIEPRGEGDHNHDTGEGGQGG EEPRGGRWDEDEQWSVVECEDCELIPADHVIV TVSLGVLKRQYTSFRPGLPTEKVAAIHRLGIGTT DKIFLEFEFPFWGPECNSLQFVWEDEAESHTLTY PPELWYRKICGFDVLYPPERYGHVLSGWICGEEA LVMEKCDDEAVAIECTEMLRQFTGNPNIPKPRRI LRSWGSNPYFRGSYSYTVQVGSSGADVEKLAKP LPYTESSKTATK
3193	A	1	1928	QLGTRRCLRGDKVTNAMQDFLVTNLEPRFIEPQT ANLSVVFKDSNSTTPLIFVLSPGTDPAADLYKFA EEMKFSKKLSAISLGQGQGPRAEAMMRSSIERGK WVFFQNCHLAPSWMPALERLIEHNPDKVHRDF

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RLWLTSLPSNKFVPSILQNGSKMTIEPPRGVRAN LLKSYSSLGEDFLNSCHKVMEFKSLLLSLCLFHG NALERRKFGPLGFNIPYEFTDGDRLICISQLKMFL DEYDDIPYKVLKYTAGENYGGRTDDWDRRCI MNILEDIFYNPDLVSPEHSYSASGIYHQIPPTYDLH GYLSYIKSLPLNDMPEIFGLHDNANITFAQNETFA LLGTIIQLQPKSSSAGSQGREEIVEDVTQNILLKVP EPINLQWVMAKYPVLYEESMNTVLVQEVIRYNR LLQVITQTLQDLLKALKGLVVMSSQLELMAASL YNNTVPELWSAKAYPSLKPLSSWVMDLLQRLDF LQAWIQDGIPAVFWISGFFFFQAFLTGTLQNFAR KFVISIDTISFDFKVMFEAPSELTORPQVGCYIHG LFLEGARWDPEAFQLAESQPKELYTEMAVIWLL PTPNRKAQDQDFYLCPIYKTLTRAGTLSTTGHS NYVIAVEIPHTQQRHWIKRGVALICALDY
3194	A	1	1023	DGWTVPVHAAVDTGNDVSLKLLMYHRIPAHGNS FNEESESSEVFDLDGGEESPEGISKPVVPADLNH ANREGWTAAHIAASKGFKNCLEILCRHGGLEPE RRDKCNRTVHDVATDDCKHLENLNALKIPLRIS VGEIEPSNYGSDDLECENTICALNIRKQTSWDDFS KAVSQALTNHFQAISDGGWWSLEDVTCNNTDS NIGLSARSIRSIITLGNVPWSVGQSFAQSPWDFMR KNKAEHITVLLSGPQEGCLSSVTYASMIPLQMM QNYLRLVEQYHNVIHFHGPESLQDYIVHQLALCL KHRQMGWQDSPVEIVEELEVGCWFFPREQLLRT CSLVA
3195	A	1	1809	MAASAQVSVTFEDVAVTFTQBEWGQLDAAQRT LYQEVMLETCGLMSLGCPLFKPELIYQLDHRQE LWMATKDLSSQSYPGDNTKPKTTEPTFSLHALPE EVLLQEQLTQGASKNSQLGQSKDQDGPSEMQUEV PLITIGIPQPCYIIEKMEETIGI GSDGVCTKI TQKQVSIWDELSDSGPVTALIREEKNSYK CEECKGVETINALLVQHERIHTQVKPYECTECG KTFKSTHLLQHIHTGEKPYKCMCEGKAFNRR SHLTRHQRIHSGEKPYKCECGKAFTHRSTFVLH HRSHTGEKPFVCKEKGKAFDRPGFIRHYIHTGE KPYECIECECGKAFNRRSYLTWHQQIHTGVKPF ECNECGKAFCESADLIQHYIHTGEKPYKCMCEG KAFNRRSHLKQHRIHTGEKPYECSECGKAFTH CSTFVLHKRTHHTGEKPYECKEKGKAFSDRADLIR HFSIHTGEKPYECVECGKAFNRRSHLTRHQIHT GEKPYECIQCGKAFCRSANLIRHSIHTGEKPYEC SECGKAFNRGSSLTHHQRIHTGRNPTIVTDVGRP FMTAQTSVNIQELLGKEFLNITTEENLW
3196	A	1400	264	VGFWERPLRSSRWFRSLRRWEMLARAARGTG ALLRGSLLASGRAPRRASSGLPRNTVVLFPVQQ EAWVVERMGRFHRILEPGLNILIPVLDRIYVQSL KEIVINVPEQSAVTLDNVTQLQIDGVLYLRIMDPY KASYGVEDPEYAVTQLAQTMRSELGKLSLDKV FRERESLNASIVDAINQAADCWGIRCLRYEIKDIH VPPRVKESMQMQVEAERRKRATVLESEGTRESA INVAEGKKQAQILASEAEKAEQINQAAGEASAVL AKAKAKAEAIRILAAALTQHNGDAAASLTVAEQ YVSAFSLAKDSNTILLPSNPGDVTSMAVQAMG VYGALTKAPVPGTPDSLSSGSSRDVQGTDAASLDE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ELDRVKMS
3197	A	66	3632	<p>LWECAAAAAGQRDGGVTLFLKGRVLGRRCAAS LFAREVCVSTSSSRPACFLHRCARARGEQMHQMA SGVGSMKRSRPMWRPGEKKEPQGVVYEDVRD DTEDFKEPLKVVFEGSAYGLQNFNKQKCLKTCD DMDTFFLHYAAAEGQIELMEKITRDSSLEVLHE MDDYGNTPLHCAVEKNQIESVKFLLSRGANPNL RNFNMMAPLHIAVQGMNNEVMKVLLHRTIDV NLEGENGNTAVIIACTTNNSEALQILLNKGAQPC KSNKWGCFPIHQAAFSGSKECMBILRFGEHGY SRQLHINFMNNGKATPLHLAVQNGDLEMIKMCL DNGAQIDPVEKGRCTAIHFAATQGATEIVKLMIS SYSGSVDIVNTTDGCHETMLHRAFLDFHHELAD YLISVGADINKIDSEGRSPLILATASASWNIVNLL LSKGAQVDIKDNFGRNFLHLTVQQPYGLKNLRP EFMQMQQIKELVMDEDNDGCTPLHYACRQGGP GSVNNLLGFNVSIHSSKSKDKKSPLHFAASYGRIN TCQRLQDISDTRLLNEGDHGMTPHLAAKNG HDKVVQLLLKKGALFLSDHNGWTALHHASMG YTQTMKVILDTNLKCTDRLEDGNTALHFAARE GHAKAVALLSHNADIVLNKQASFLHLALHNK RKEVVLTIIRSKRWDECLKIFSHNSPGNKCPIEM IEYLPECMKVLLDFCMLHSTEDKSCRDYYIEYNF KYLQCPLEFTKKTPTQDVIYEPLTALNAMVQNN RIELLNHPVCKEYLLMKWLA YGFRAHMMNLGS YCLGLIPMTILVVNIKPGMAFNSTGINETS DHSEI LDTTNSYLIKTCMILVFLSSIFGYCKEAGQIFQK RNYFMDISNVLEWITYTTGIIFFVLPFVEIPAHQ WQCGAIAVYFYWMNLLYLQRFENCGIFIVMLE VILKTLRSTVVFFIFLLAFGLSFYILLNLQDPFSS PILSHIQTTSMTGGINVRESFLEWTFNFLAIPV LSFAQLVSFTIFVPIVLLTLLIGLAVGDIAEVQKH ASLKRIAMQVELHTSLEKKLPLWFLRKVDQKSTI VYPNKRSGGMLFHIFCFCTGEIRQEI PNADKS LEMEILKQKYRLKDLTFLEKQHELIKLIQKMEII SETEDDDSHCSFQDRFKKEQMEQRNSRWNTVLR AVKAKTHLEP</p>
3198	A	51	2177	<p>KEKSLHHVDQRPPPLWHPGRPGTSQSAAMNASSE GESFAGSVQIPGGTTVLVELTPDIHCICKQQFN NLDAFVAHKQSGCQLTGTSAAAPSTVQFVSEET VPATQTQTTRITITSETQTTTSAPEFVFEHGYQT YLPTESNENQTATVISLPAKSRTKKPTTPPAQKRL NCCYPGCQFKTAYGMKDMERHLKIHTGDKPHK CEVCGKCFSRKDKLKTTHMRCHTGVPYKCKTC DYAAADSSSLNKHRLHSRDERPFKQCICPYASRN SSQLTVHLRSHTGDAPFQCWLCSAKFKJSSDLKR HMRVHSGEKPFCFCNVRCMTKGNLKSHIRIK HSGNNFKCPHCAFLGDSKATLRKHSRVHQSEHR EKCSECSYSSSKAALRIHERIHCTVRPFKNYCS FDSKQPSNLSKHMKKFHHGDMVKTEALERKDTG RQSSRQVAKLDAKKSFFHCDICDASFMREDSLRS HKRQHSEYNESKNSDVTVLQFQIDPSKQPATPLT VHGLQVPLQPSQVPQFSEGRVKIIVGHQVPQANT IVQAAAAAVNIVPPALVAQNPEELPGNSRLQILR QVSLIAPPQSSRCPSSEAGAMTQPAVLLTTHEQTD</p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GATLHQTLIPTASGGPQEGSGNQTFITSSGITCTD FEGNLALIQEGTAEBTVVSDGGQNIATVATTAPPV FSSSSQQELPKQTYSHQGAHPALLCPADSIDP
3199	A	13	2247	QSFHSMEGDPSGLPLLARGASCYSLICPCPRPAD WSILQGTDWISLQADWCIYNPLARHRALTGVFL QSADWCTYNPLARQKSSPSPHSTQEVQLASPLTR RPNKKDSAERNHRPAREGSVAQRQPNPAALEKA EPAARKRNEREGGGSQEPGREHSLEKGYWAPGL GPDPSMCSKQVDPSEGAASHLKHRGGSRAAHLE VRLLRLVGLVAEAGFCYVQVAEQRVVGV LEVAEAAAAPVQHEPTAAVATQSRWFPRGTRPG LCSLPIA VAALLCPGSGPGAQSGLEFVERPPPSPL AVVLARWPLPPPAGRCPRDAPEARVPEKARAEG SERENNYGCGVVGEMTTLVLDNGAYNAKIGY SHENVSVIPNCQFRSKTARLKTFTANQIDEIKDPS GLFYILPFQKGYLVNWDVQRQVWDYLF GKEMY QVDFLDNTNITEPYFNFTSIQESMNEILFEEYQFQ AVLRVNAGALSAHRYFRDNPSLCCIIVDSGYSF THIVPYCRSKKKKEAIRINVGGKLLTNHLKEHSY RQLHVMDETHVINQVKEDVCYVSQDFYRMDI AKLKGEENTVMIDYVLPDFSTIKKGFCCKPREEMV LSGKYKSGEQILRLANERFAVPEILFNPSDIGIE MGIPEAIVYSIQNLPEEMQPHFFKNIVLTGNSLF PGFRDRVYSEVRCLTPTDYDVSVVLPENPITYAW EGGKLISENDDFEDMVVTREDEENGHSVCEEK FDI
3200	A	3	307	AVQRIHEMNIFRLTGDLSHLAAIVILLKKIWKTR SCAGISGKSQLLFALVFTTRYLDLFTSFISLYNTS MKVWYAIHRNVFHLQCTGLWTLNLCQLCFN
3201	A	1	469	IRHEGRGQRGMELVQVLKRGGLQQTGHGGLRG YLRVFRFTNDKVGTLVGEFKYGNYYEDNKK FFGRHRWVVYTTMNGKNFWLGGSMVPPLE WHRWLHSMITDDPPTTKPLTARKFIWTNHKFNV GTPEQYVPYSTTRKKIQEWIPSPSTPYK
3202	A	144	840	NSSQRIMATHALEIAGLFLGGVGMVGTVAVTVM PQWRVSAFIENNIVVFENFWGLWMNCVRQANI RMQCKIYDSSLALSPDLQAARGLMCAASVMSFL AFMMAILGMKCTRCTGDNEKVKAHILLTAGIIFI TGMVVLPVSWVANAIIRDFYNSIVNVAQKREL EALYLGWTTALVLIVGGALFCCVFCCKNEKSSSYR YSIPSHRTTQKSYHTGKKSPSVYSRSQYV
3203	A	2	473	KYRYRRPYPMRKICQVGPAGLAFILNISPVHR VALCHLAGCQEQAAYHTLQILFFLVSAYFFSCP VPEKYFPGSCDIVGHGHQIFHAFLSICITLSQLEAIL LDYQGRQEIFLQRHGPLSVHMACLSFFFLAACSA ATAALLRHKVKARLTKKDS
3204	A	1808	668	PESAPLPAFISSRILPAAWRNWCSYVVTRTISCHV QNGTYLQRLQNCWPWMSCPGSSYRTVVVRPTYK VMYKIVTAREWRCCPGHSRVSCEEVAGSSASLE PMWSGSTMRMALRPTAFSGCLNCSKVSELT LKVLEAKMTMLTVIEQVPVPTPATPEDAPLWGP PPAQGSPGDGGLQDQVGAWGLPGPTGPKGDAG SRGPMGMRGPPGDPLLSNTFTETNNHWPQGPTG PPGPPGPMGPPGPPGPTGVPGSPGHGPPGPTGPK GISGHPGEKGERGLRGEPPGQGSAGQRGEPPGPKG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				DPGEKSHWGEGLHQLREALKILAERVLIETMIG LYPELGSAGAPAGTGTPSLLRGKRGGHATNYRI VAPRSRDERG
3205	A	2810	1652	RTSTQKWQSVFNDSQEHLERFYCNPENDRMRM KYGGQEFWADLNAMNVYETTEFDQLRRLSTPPS SNVNSIYHTVWKFFCRDHFHWREYPESVIRLIEE ANSRGLKEVRFMWNNHYILHNSFFRREIKRRP LFRSCFILLPYLQTLGGVPTQAPPPLAETSSSQICP DGVTSANFYPETWVYMHPSQDFIQVPVSAEDKS YRIYNLFHKTVPFVKYRILQILRVQNQFLWEKY KRKKEYMNRKMFGRDRIINERHLFHGTSQDVVD GICKHNFDPRVCGKHATMFGQGSYFAKKASYSH NFSKSSKGVHFMFLAKVLTGRYTMGSHGMRR PPPVPNGSVTSDLYDSCVDNFFEPQIFVIFNDDQS YPYFVIQYEEVSNTVSI
3206	A	297	4500	CLVDSKLWKGARSVYHQLFMSSLLMDLKYYKKL FAVRFKANYERLQSDYVTDDHDFSVADLSVQ IFTVPSLARMLITEENLMSIIKTFMDHLRHRDAQ GRFQFERYTALQAFKFRVQSLILDLKYVLISKPT EWSDEL RQKFLEGDAFLELLKCMQGM DPITRQ VGQHIEMEPWEAAFTLQMKLTHVISMMQDWC ASDEKVLIBAYKKCLAVLMQCHGGYTDGEQIPIT LSICGHSVETIRYCVSQEKVSHLPVSRLLAGLHV LLSKSEVAYKFPPELLPLSELSPMLIEHPLRCLVL CAQVHAGMWRRNGFSLVNQIYHHNVKCRRE MFDKDVVMLQTGVSMMDPNHFLMIMLSRFELY QIFSTPDYGKRFSSEITHKDVVQQNNTLIEEMLYL IIMLVGERFSPGVGVNATDEIKREIHHQLSIKPM AHSELVKSLEPENKETGMESVIEAVAHFKKPG TGRGMYELKPECAKEFNLYFYHFSRAEQSKAEE KRLKQONREDALEPPVLETCI FASLVNLIQ SDVMLCIMGTLDWAVEINGYAWSESMLQRL HLIGMALQEEKQHEENVTEEHVVTFTTQKISKP GEAPKNPSILAMLETIONAPYLEVHKDMIRWIL KTFNAVKKMRESSPTSVAETEGTIMEESSRDKD KAERKRKAEIARLRREKIMAOQMSEMQRHFDEN KELFQQTLELDASTSAVLDHSPVASDMTLTALGP AQTQVPEQRQFVTCILQEEQEVKVESRAMVLA AFVQRSTVLSKNRSKFIQDPEKYDPLFMHPDLSC GTHTSSCGHIMHAHCWQRYFDSVQAKEQRRQQ RLRLHTSYDVENGFEFLCPLCECLSNTPVPLLLPR NIFNNRLNFSQPNLTQWIRTISQOIKALQFLRKE ESTPNNASTKNSENVDELQLPEGFRPDRPKIPYS ESIKEMLTTFGTATYKVGKLVHPNEEDPRVPIMC WGSCAYTIQSIERILSDEDKPLFGPLPCRLDDCLR SLTRFAAAHWTVASVSVVQGHFCKPFASLVPND SHEELPCILDIDMFHLLVGLVLAFPALQCQDFSGI SLGTGDLHIFHLVTMAHIIQILLTSCTEENGMDQE NPPCEBESAVLALYKTLHQYTGSALKEIPSGWHL WRSVRAGIMPFLKCSALFFHYLNGVSPDPDIQVP GTSHFEHLCSYLSLPNNLICLFQENSEIMNSLIES WCNRNSEVKRYLEGERDAIRYPRESNKLINLPEDY SSLINQASNFSCPKSGGDKSRAPTLCLVCGSLLCS QSYCCQTELEGEDVGACTAHTYSCGSGVGIFLR VRECQVFLAGKTKGCFYSPPYLDDYGETDQGL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				RRGNPLHLCKERFKKIQLWHQHSVTBEIGHAQ EANQTLVGIDWQHL
3207	A	49	963	QLSPSQAPAGAQEVARRVTVGSASHGGRSTMA TTVSTQRGPVYIGELPQDFLRITPTQQQRQVQLD AQAAQQLQYGGAVGTGVRNLNITVVQAKLAKNY GMTRMDPYCRLRLGYAVYETPTAHNGAKNPRW NKVIHCTVPPGVDSFYLEIFDERAFSMDRIA WTHTIPESLRQGVEDKWYSLSGRQGGDDKEGMNL VMSYALLPAAMVMPPQPVVLMPTVYQQGVGY VPITGMPAVCSPGMVPVALPPAAVNAQPRCSEE DLKAIQDMFPNMDQEVIRSVLEAQRGNKDAAIN SLLQMGEPP
3208	A	54	1196	LERTPASADMAWTKYQLFLAGLMLVTGSINTLS AKWADNFMAGCGGSKEHSFQHPFLQAVGMFL GEFSCLAAFYLLRCRAAGQSDSSVDPQPFNPLL FLPPALCDMTGTSLMYVALNMTSASSFQMLRGA VIIFTGLFSVAFGLGRLVLSQWLGLATIAGLVVV GLADLLSKHDSQHKLSEVITGDLILMAQIIVAIQ MVLEEKFVYKHNHPLRAVGTGLFGFVILSLLL VPMYYIPAGSFGSNPRGTLEDALDAFCQVGGQP LIAVALLGNISSIAFFNFAGISVTKELSATTRMV DSLRTVVIWALSLALGWEAFHALQILGFLILLIGT ALYNGLHRPLLGRLSRGRPLABESEQERLLGGTR TPINDAS
3209	A	104	1999	AKVVSLKEFSCFWRREKPVSSLSLQVKAESW DSAVHGCPLSRGTPVDERLFLIVRVTVQLSHPA DMQLVLRKRICVNVHGRQGFQSLKKMSHRSS IPGCGVTFEIVSNIPEDAQGVVEERALARMAANV ENPASADSEAYIEKYLRSLAVENLLTDLRLRQE VAVKEQLTGKGLSRRSISPNVNRLSGSRQDLIP S' SLCENKGRWESQCDVSOTTVSRGIAPAPAL SPQNNHSDPGLSNLAASYINPVKQVVTQMPK: L KSLFPVRDEKRGKRPSPLAHQPVPRMVQSASPI RVTRMEBAQPEMGPDVLVQTMGAPALKICDKP AKVPSPPVIAVTAVTPAPEAQDGPPSPLSEASSG YFSHSVSTATLSDALGPGLDAAAPPGSMPTAPEA EPEAPISHPPPTAVPAEPPGPQQLVSPGRERPD EAPAPGSPFRVRRVRASELRSFSRMLAGDPGCS GAEGNAPAPGAGGQALASDSEEADEVPEWLREG EFVTVGAKHTGVVRYVGPADFQEGTVWGVELD LPSGKNDGSIGGKQYFRCNPGYGLLVPRSRVRR ATGPVRRRSTGLRLGAPEARRSATLSGSATNLAS LTAALAKADRSKHNPNRKSAS
3210	A	324	694	SPFWTEKRRMEKPLFPLVPLHWFQGYTALVVS GGIVGYVKTGSVPSLAAGLLFGSLAGLGAYQLY QDPRNVWGFLAATSVTFVGMGMRSYYYGKF MPVGLIAGASLLMAAKVGVRMLMTSD
3211	A	1078	594	VGMELPAVNLKVILLGHWLLTTWGCIVPSGSYA WANFTILALGVWAVAQRDSIDAISMFLGGLLATI FLDIVHISIFYPRVSLTDTGRFGVGMAILSLLKPL SCCFVYHMYRERGGELLVHTGFLGSSQDRSAYQ TIDSAEAPADPFAVPEGRSQDARGY
3212	A	1	1962	FRCGLAPKGRPRRRADPVASAIMDPAAVLOEK ALKFMMEFRSWCPGWNTMARSRLTATSTSRVQ CSMPRSLWLGCSLADSMPSLRCLYNPGTGALT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				AFQNSSEREDCNNGEPPRKIIPEKNSLRQTYNSCARLCLNQETVCLASTAMKTENCVAKTKLANGTSSMIVPKQRKLSASYEKEKELCVKYFEQWSESDQVEFVEHLISQMCHYQHGHSYLPMLQRDFTTALPARGLDHIAENILSYLDAKSLCAAELVCKEWYRVTSDGMLWKKLIERMVRTDSLWRGLAERRGWGQYLFKNKPPDGNAPPNSFYRALYPKIIQDIETIESNWRGCRHSLQRIHCRSETSKGVYCLQYDDQKIVSGLRDNTIKIWDKNTLECKRILTGHTGSVLCLOQDERVIITGSSDSTVRVWDVNTGEMLNTLIHHCEAVLHLRFNNGMMVTCSKDRSLAVWDMASPTDITLRRVLVGHRAAVNVVDFDDKYIVSASGDRTIKVWNTSTCEFVRTLNHGKRGIACLQYRDRLVVSGSDNTIRLWDIECGACLRVLEGHEELVRCIRFDNKRIVSGAYDGGIKVWDLVAALDPRAPAGTLCRLRLVEHSGRVFRLQFDEFQIVSSSHDDTILIWDFLNDPAAQSEPPRSPSRITYTYISR
3213	A	1	1962	FRCGLAPKGRPRRRADPVASAIMDPAAEVLQEKALKFMMEFRSWCPGWNTMARSRLTATSTSRVQCSMPRSLWLGCSLADSMPSLRCLYNPGTGALTAFQNSSEREDCNNGEPPRKIIPEKNSLRQTYNSCARLCLNQETVCLASTAMKTENCVAKTKLANGTSSMIVPKQRKLSASYEKEKELCVKYFEQWSESDQVEFVEHLISQMCHYQHGHSYLPMLQRDFTTALPARGLDHIAENILSYLDAKSLCAAELVCKEWYRVTSDGMLWKKLIERMVRTDSLWRGLAERRGWGQYLFKNKPPDGNAPPNSFYRALYPKIIQDIETIESNWRGCRHSLQRIHCRSETSKGVYCLQYDDQKIVSGLRDNTIKIWDKNTLECKRILTGHTGSVLCLOQDERVIITGSSDSTVRVWDVNTGEMLNTLIHHCEAVLHLRFNNGMMVTCSKDRSLAVWDMASPTDITLRRVLVGHRAAVNVVDFDDKYIVSASGDRTIKVWNTSTCEFVRTLNHGKRGIACLQYRDRLVVSGSDNTIRLWDIECGACLRVLEGHEELVRCIRFDNKRIVSGAYDGGIKVWDLVAALDPRAPAGTLCRLRLVEHSGRVFRLQFDEFQIVSSSHDDTILIWDFLNDPAAQSEPPRSPSRITYTYISR
3214	A	1	1962	FRCGLAPKGRPRRRADPVASAIMDPAAEVLQEKALKFMMEFRSWCPGWNTMARSRLTATSTSRVQCSMPRSLWLGCSLADSMPSLRCLYNPGTGALTAFQNSSEREDCNNGEPPRKIIPEKNSLRQTYNSCARLCLNQETVCLASTAMKTENCVAKTKLANGTSSMIVPKQRKLSASYEKEKELCVKYFEQWSESDQVEFVEHLISQMCHYQHGHSYLPMLQRDFTTALPARGLDHIAENILSYLDAKSLCAAELVCKEWYRVTSDGMLWKKLIERMVRTDSLWRGLAERRGWGQYLFKNKPPDGNAPPNSFYRALYPKIIQDIETIESNWRGCRHSLQRIHCRSETSKGVYCLQYDDQKIVSGLRDNTIKIWDKNTLECKRILTGHTGSVLCLOQDERVIITGSSDSTVRVWDVNTGEMLNTLIHHCEAVLHLRFNNGMMVTCSKDRSLAVWDMASPTDITLRRVLVGHRAAVNVVDFDDKYIVSASGDRTIKVWNTSTCEFVRTLNHGKRGIACLQYRDRLVVSGSDNTIRLWDIECGACLRVLEGHEELVRCIRFDNKRIVSGAYDGGIKVWDLVAALDPRAPAGTLCRLRLVEHSGRVFRLQFDEFQIVSSSHDDTILIWDFLNDPAAQSEPPRSPSRITYTYISR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				LVEHSGRVFRLQFDEFQIVSSSHDDTLIWDFLND PAAQSEPPRSPSRITYTISR
3215	A	2	1376	EARLVGCQRGGPARPGSYSSGAETAGRAMAAN LSRNGPALQEAYVRVVTEKSPTDVALFTYEGNS NDIRVAGTGEGGLEEMVEELNSGKVMYAFCRV KDPNSGLPKFVLINWTGEGVNDVRKGACASHVS TMASFLKGAHVTTINARAEEDVEPECIMEKVAKA SGANYSFHKESGRFQDVGPPQAPVGSVYQKTNAV SEIKRVGKDSFWAKAEKEEENRRLEKRRRAEEA QRQLEQERRERELREAAARREQRYQEQQGEASPO RTWEQQQEVVSRNRNEQESAVHPREIFKQKERA MSTTSISSPQPGKLRSPFLQKQLTQPTHFGREPA AAISRPRADLPAEPPAPSTPPCLVQAEEEEAVYEPP PEQETFYEQPPLVQQQAGAGSEHIDHHIQGQGLSG QGLCARALYDYQAADDTEISFDPENLITGIEVIDE GWWRGYGPDGHFGMFPANYVELIE
3216	A	936	204	AMASTLEYSPSPLRRLVGPAAGFSRAARADLSW DPMAFFTGLWGPFTCVSRVLSHHCFTTGSLSAI QKMTRVRVVDNSALGNSPYHRAPRCHVYKKN GVGKVGQDQILLAIKGQKKKALIVGHCMGPGRMT PRFDSNNVVLIEDNGNPVGTRIKTPIPTSLRKREG EYSKVLAIQNFV
3217	A	1	1563	MLCALLLLPSLLGATRASPTSGPQECAGKSTVW CQDLQTAARCGAVGYCQGA VWNKPTAKSLPCD VCQDIAAAAGNGLNPDATESDILALVMKTCEWL PSQESSAGCKWMVDAHSSAILSMRGAPDSAPA QVCTALSLCEPLQRHLATLRPLSKEDTFAVAPF MANGPLTFHPRQAPEGALCQDCVRQVSRLQEA VRSNLTLADLNIQECCESLGPGLAVLCKNYLFQFF VPADQALRLLPPQELCRKGGFCEELGAPARLTQ VVAMDGVPSLELGLPRKQSEMOMKAGVTCEVC MNVVQKLDHWMLSNSEL MITHALERVCSVM ASITKECILVDITYSPSLVQLVAKITPEKVCKFIK CGNRRRARAVHDA YAIVPSPWD AENQGSFCNG CKRLLTVSSHNLKSKSTKRDLVAFKGGCSILPLP YMIQCKHFVTQYEPVLIESLKDMMDPVA VCKKV GACHGPRTPLLGTDQCALGPSFWCRSQEAAKLC NAVQHCQKHVWKEMHLHAGEHA
3218	A	1	1563	MLCALLLLPSLLGATRASPTSGPQECAGKSTVW CQDLQTAARCGAVGYCQGA VWNKPTAKSLPCD VCQDIAAAAGNGLNPDATESDILALVMKTCEWL PSQESSAGCKWMVDAHSSAILSMRGAPDSAPA QVCTALSLCEPLQRHLATLRPLSKEDTFAVAPF MANGPLTFHPRQAPEGALCQDCVRQVSRLQEA VRSNLTLADLNIQECCESLGPGLAVLCKNYLFQFF VPADQALRLLPPQELCRKGGFCEELGAPARLTQ VVAMDGVPSLELGLPRKQSEMOMKAGVTCEVC MNVVQKLDHWMLSNSEL MITHALERVCSVM ASITKECILVDITYSPSLVQLVAKITPEKVCKFIRL CGNRRRARAVHDA YAIVPSPWD AENQGSFCNG CKRLLTVSSHNLKSKSTKRDLVAFKGGCSILPLP YMIQCKHFVTQYEPVLIESLKDMMDPVA VCKKV GACHGPRTPLLGTDQCALGPSFWCRSQEAAKLC NAVQHCQKHVWKEMHLHAGEHA
3219	A	1623	572	TSAEGWKGTCTFKDRSKLREHLRSHTQEKVVA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				CPTCGGMFANNTKFLDHRRQTSLDQQHFQCSH CSKRFA TERLLRDHMRNHVNHVKPLCDMTCP PSSLRNHMRFRHSEDRPFKCDCCDYCKNLIDLQ KHLDT HSEEPAYRCDFENCTFSARSLCSIKSHYR KVHEGDSEPRYKCHVCDKCFTRGNNLTVHLRK KHQFKWPSGHPFRFYKEHEDGYMRLQLVRYES VELTQQLLRQPQEGSGLGTSLNESSLQGILETVP GEPGRKEEEEEEGKGSEGTALSASQDNPSVHV NQTNAQGQQEIVYYVLSEAPGEPVPVPEPPSGGI MEKLQGLAEPEIQMV
3220	A	2760	745	SLGIPSGNTRGTGLVLDGDTSYTYHLVCMGPEAS GWGQDEPQTWPTDHRAQQGVQRQGVSYVHA YTGQPSRGLHSENREDEGWQVYRLGARDAHQ GRPTWALRPEDGEDKEMKTYRLDAGDADPRRL CDLERERWAVIQGQAVRKSSTVATLQGTDPDHGD PRTPGPPRSTPLEENVVDREQIDFLAARQQFLSLE QANKGAPHSSPARGTPAGTTPGASQAPKAFNKP HLANGHVVPKPVKGVVREENKVRAVPTWAS VQVVD DPGSLASVESPGTPKETPIEREIRLAQERE ADLREQRGLRQATDHQELVEIPTRPLLTKLSLITA PRRERGRPSLYVQRDIVQETQREEDHRREGLHV GRASTPDWVSEGPQGLRRALSSDSILSPAPDAR AADPAPEVRKVNRIIPDAYQPYLSPGTPQLEFSA FGAFGKPSSLSTAEAKAATSPKATMSPRHLSSE GKPLSTKQEASKPPRGCPQANRGVVRWEYFRLR PLRFRAPDEPQQAQVPHVWGWEVAGAPALRLQ KSQSSDLLERERESVLRREQEVAEERNALFPEV FSPTPDENSQNSRSSQASGITGSYSVSESPFFSPI HLHSNVAWTVEDPVDAPPQQRKKEQWYAGIN PSDGINSEVLEAIRVTRHKNAMEAERWESRIYASE EDD
3221	A	15	478	LVVFFVFFFPFAFKMSKRGRGGSSGAKF VGAVINCADNTGAKNLYISVKGIKGRNLRLPAA GVGDMVMATVKKGKPELRKKVHPAVVIRQRKS YRREDGVFLYFEDNAGVIVNNKGEMKGSAITGP VAKECADLWPRIASNAGSIA
3222	A	207	1321	PLIPLHPANRSPATMAELQEVQITEEKPLLPQGT EAAKTHSVETPYGSVTFVYGTTPKPKRPAILTYH DVGLNYKSCFQPLQFEDMQEIIQNFRVHVHDAP GMEEGAPVFPLGYQYPSLDQLADMIPCVLQYLN FSTIIGVGVGAGAYILARYALNHPDTEGLVLINI DPNAKGWMDWAAHKLTLTSSIPEMILGHLFSQ EELSGNSELIQKYRNIITHAPNLDNIELYWN SYN RRDLNFERGGDITLRCPVMLVVGDAQPHEDAVV ECNSKLDPTQTSFLKMADSGGQPQLTQPGKLTE AFKYFLQGMGYMASSCMTRLSRSRTASLTSAAS VDGNRSRSRTLSSQSESGTLSSGPPGHTMEVSC
3223	A	132	1664	SARRWGAAGAGPHGLHLRAHGPRPSVRTGLPSV GRQAAGAAMGRGWGFLFGLLGAVWLLSSGHGE EQPPETAARCFQCVSGYLDCTCDVETIDRFNN YRLFPRLOKLLSDYFRYYKVNLRPCPFWNDIS QCGRRDCAVKPCQSDEVPDGIKSASYKYSEAN NLIEBCEQAERLGA VDESLEETQKAVLQWTKH DDSSDNFCEADDIQSPEAEYVDLLNPERYTGK GPDAWKIWNVIYEENCFKPQTKRPLNPLASGQG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TSEENTFYSWLEGLCVEKRAFYLRLISGLHASINV HLSARYLLQETWLEKKWGHNITEFQQRFDGILTE GEGPRRLKNLYFLYLIELRALSKVLPFFERPDFQL FTGNKIQDEENKMLLEILHEIKSFPLHFDENSFF AGDKKEAHKLKEDFRLHFRNISRIMDCVGCFC RLWGKLQTQGLGTALKILFSEKLIANMPESGPSY EFHLTRQEIVSLFNAFGRISYKCEIRKTSRNLQ NIH
3224	A	2	803	PGSTISWDRDAAGESGTRAASPSPSGSRTAGRLP SPSYSPLPAPSLFPPPLPAPAASTMSAGGDFGNP LRKFKLVFLGEQSVGKTSLITRFMYDSFDNTYQA TIGIDFLSKTMYLEDRTVRLQLWDTAGQERFRSL IPSYIRDSTVAVVVYDITNLNSFQQTSKWIDDVRT ERGSDVIIIMLVGNKTDLADKRQITIEEGEQRAKE LSVMFIETSAKTGYNVKQLFRRVASALPGMENV QEKSKEGMIDIKLDPQEPPEASEGGCSC
3225	A	3	5054	PEVTKPSLSQPTAASPIGSSPSPVNGGNNAKRVA VPNGQPPSAARYMPREVPPRFRCQQDHKVLKLR GQPPPPSCMLLGGGAGPPPCTAPGANPNNAQVT GALLQSESGTAPDSTLGGAAASNYANSTWGSGA SSNNGTSPNPIHIWDKVIDGSDMEEWPCLASKD TESSSENTTDNNSASNPGSEKSTLPGSTTSNKGK GSQCQSASSGNECNLGVWKS DPKAKSVQSSNST TENNNGLGNWRNVSGQDRIGPGSGFSNPNNSN PSAWPALVQEGTSRKGALETDNSNSSAQVSTVG QTSREQQSKMENAGVNFVVS GREQAQIHNTDGP KNGNTNSLNLSSPNPMENKGMPPGMGLNTRSRS TDAPSQSTGDRKTGSGVSGWGAARGPSGTDTVSG QNSNGNNGNNGKEREDSWKGASVQKSTGSKND SWDNNNRSTGGSWNFGPQDSNDNKWGEGNKM TSGVSGQEWKC...SDFLKGEW...QPNSSST GAF...NKGHPLLENQNAQAP...GRT...TGS EVEGQSTGSHKAGSSDSHNSGRREYRPTHPDC QAVLQTLRLDLDPRVLSNTGWGQ...QDTV WDIEEVPRPEGKSDKGTEGWESAATQTK...SGG WGDAPSQSNQMKSGWGELSASTEWKDPKNTGG WNDYKNNSSNWGGGRPDEKTPSSWNEPNSKD QGWWGGGRQPNQGWSSGKNGWGEEVDQTKNSN WESSASKPVSGWEGGQNEIGTWGNGGNASLA SKGGWEDCKRSPA WNETGRQPN SWNKQHQQQ QPPQPPPPQPEASGSWGGPPPPPPGNVRPSNSS WSSGPQPATPKDEBPSGWEEPPSPQSIRKMDIDD GTSAWGDPNSYNYKNVNLWDKNSQGGPAPREP NLPTPMTSKSASDSKSMQD GWGESDGPVTGARH PSWEEEEEDGGVWNTTGSQGSASSHNSASWGQG GKKQMKCSLKGGNDSWMNPLAKQFSNMGLL SQTEDNPSSKMDLSVGLSDKKFDVDKRAMNLG DFNDIMRKDRSGFRPPNSKDMGTTDSGPYFEKG GSHGLFGNSTAQSRGLHTPVQPLNSSPSLRAQVP PQFISPVQSASMLKQFPNSGLSPGLFNVGPQLSPQ QIAMLSQLPQIPQFLACQLLQQQQQQQLLQN QRKISQAVRQQQEQLARMVSALQQQQQQQQR QPGMKHSPSHVGPKPHLDNMVFNALNVGLPDL QTKGPIPGYGSGFSSGMDYGMVGGKEAGTESR FKQWTSMMEGLPSVATQEANMHKNGAIVAPGK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TRGGSPYNQFDIIPGDTLGGHTGPAGDSWLPKAS PPTNKIGSKSSNASWPPEFQPGVPWKGIQNIDPES DPYVTPGSLGGTATSPIVDTHQLLRDNTTGSN SSLNTSLPSPGAWPYASDNSFTNVHSTSAKFPD YKSTWSPDPIGHNPHTLSNKMWKNHISSRNTTPL PRPPPGLTNPKPSSPWSTAPRSVRGWGTQDSRL ASASTWSDGGSVRPSYWLVLHNLTPQIDGSTLRT ICMQHGPLLTFLHNLQGTALIRYSTKQEAAKAQ TALHMCVLGNTTILAEFATDDEVSRFLAQAPPT PAATPSAPAAGWQSLETGQNSDPVGPALNLF GSTGLGQWSSSAGGSSGADLAGASLWGPPNYSS SLWGVPTVEDPHRMGSPAPLLPGDLLGGGSDSI
3226	A	200	1387	VPWKRQDEQLSLQVETLYLDSPAVIHLLSPTFLP PSSLPPFLQIVDSSSSACTLDSFFFLAPWDSQDC GFKDHQPLTLQALTVELARWTLMLLLSTAMYG AHAPLLALCHVDGRVPPRPSSAVLLTELTKLLC AFSLLVGWQAWPQGPPWRQAAPFALSALLYG ANNLVIYLQRYMDPSTYQVLSNLKIGSTAVLY CLCLRHRLSVROGLALLLMAAGACYAAGGLQ VPGNTLPSPPAAAAASPMLHITPLGLLLLYCLI SGLSSVYTELLMKRQRLPLALQNLFLYTFGVLLN LGLHAGGGSGPGLLEGFSGWAALVLSQALNGL LMSAVMKHGSSITRLFVVSCLVNAVLSAVLL RLQLTAAFLATLLIGLAMRLYYGSR
3227	A	1	679	RSTRARTRRPGLRAVPLPVGGFLGKMKVWVAL LLLAALGSGRAERDCRVSSFRVKENFDKARFSGT WYAMAKKDPEGLFLQDNIVAESVDETQMSA TAKGRVRLNNWDVCADMVGTFTDTPAKFK MKYWGVASFLQKGNDDHWIVDTDYDTYAVQY SCRLLNLDGTCADSYSFVFSRDPNGLPPEAQKIV RQRQFLCAARQYRLIVHNGETGPSEP NLL
3228	A	25	1104	QQESPAACAAARLLEGTDESCGCRGNDEKK LKC VVG DGAVGKTCLLMSYANDAPPEYVPT VFDHYAVTVTVGGKQHLGLYDTAGQEDYNQL RPLSYPTNDVFLICFSVVPASYHNVQEEWVPEL KDCMPHVPLYLIGTQIDLRDDPKTLARLLYMKE KPLTYEHGVKLAKAIGAQCYLECSALTQKGLKA VFDEAILTIFHPKKKKKRCSEGHSCCSII
3229	A	25	722	AISAGRSKMQKLPMEINPEMLNKVLSRLGVAG QWRFVDVLGLEEESLGSVPAPACALLFLPTAQ HENFRKKQIEELKGQEVSPKVYFMKTIGNSCGT IGLIHAVANNQDKLGFEDGSVLKQFLSETEKMSP EDRAKCFEKNEAIQAAHDAVAQEGQCRVDDKV NFHFILFNNVDGHL YELDGRMPFPVNHGASSED LLKDAAKVCREFTEREQGEVRFSAVALCKAA
3230	A	282	1479	GDAATTACAPPDWFLGPRKLAAGPAGGGMLPR RLAAWLAGTRGGGLLALLANQCRFVTGLRVR RAQQIAQLYGRLYSESSRRVLLGRLWRLHGRP GHASALMAALAGVFVWDEERIQEELQRSINEM KRLEEMSNMFQSSGVQHHPPEPKAQTEGNEDE GKEQRWEMVMDKKHFKLWRRPITGTHLYQYRV FGTYTDVTPRQFFNVQLDTEYRKKWDALVIKLE VIERDVVSGSEVLHWVTHFPYPMYSRDYVYVRR YSVDQENNMMLVLSRAVEHPSVPESPEFVRVRS YESQMVRPHKSFDENGFDYLLTYSNPNQTVFPR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				YCVSWMVSSGMPDFLEKLHMATLKAKNMEIKV KDYISAKPLEMSSEAKATSQSSERKNEGSCGP IEYA
3231	A	2117	590	FVPEPPEAGASSPCAPGDPDMSFRKVVVRQSKFRH VFGQPVKNDQCYEDIRVSRVTWDSFCAVNPKE LAVIVEASGGGAFLVLPLSKTGRIDKAYPTVCGH TGPVLDIDWCPHNDEVIASGSEDCTVMVWQIPE NGLTSPLTEPVVLEGHTRVGIIAWHPTARNVL LSAGCDNVVLIWNVGTAEELYRLDSLHPDLIYN VSWNHNGSLFCSACKDKSVRIIDPRRGTLVAERE KAHEGARPMRAIFLADGKVFTTGFSRMSERQLA LWDPENLEEPMALQELDSSNGALLPFYDPDTSV VYVCGKGDSSIRYFEITEPPYIHFLNTFTSKEPQR GMGSMKRGLEVSKCEIARFYKLHERKCEPIVM TVPRKSDLFQDDLYPDTAGPEAALEAEWVSGR DADPILISLREAYVPSKQRDLKISRRNVLSDSRPA MAPGSSHLGAPASTTTAADATPSGSLARAGEAG KLEEVMQELRALRALVKEQGDRCRLEEQLGRM ENGDA
3232	A	3	718	RLREDDRRGLPLSSPLWTEPPLSCCLPATYPADM GTAGAMQLCWVILGFLFRGHNSQPTMTQTSSS QGGLGGLSLTTEPVSSNPGYIPSEANRPSHLSST GTPGAGVPSSGRDGGTSRDITFQTVPPNSTTMSLS MREDATILPSPTSETVLTVAAGVISFIVILVVVVI ILVGVVSLRFKCRKSKESEDPQKPGSSGLSESCST ANGEKDSITLISMKNINMNNGKQSLSAEKVL
3233	A	3	718	RLREDDRRGLPLSSPLWTEPPLSCCLPATYPADM GTAGAMQLCWVILGFLFRGHNSQPTMTQTSSS QGGLGGLSLTTEPVSSNPGYIPSEANRPSHLSST GTPGAGVPSSGRDGGTSRDITFQTVPPNSTTMSLS MREDATILPSPTSETVLTVAAGVISFIVILVVVVI ILVGVVSLRFKCRKSKESEDPQKPGSSGLSESCST ANGEKDSITLISMKNINMNNGKQSLSAEKVL
3234	A	1169	4292	AGDCGRLGVGGSEFPWEGSALGASPLPPICLQSR TWLLRAPAPAELEEEVAAGRGDVWEPFLDSP GEEESLQEASPRLADHGSSSGGGWEVKRSQRLR RGPSSRRPYQDMEYERRGGRDRTGRYGATDR SQDDGGENRSRDHDYRDMDYRSYPREYGSQEG KHDYDDSSSEEQAEDSYEASPGSETQRRRRRRH RHSPTGPPGPRDGDYRDQDYRTEQGEEEEEEED EEEEKASNIVMLRMLPQAATEDDIRGQLQSHG VQAREVRLMRNKSSGQSRGFAFVEFSLQDATR WMEANQHSLNILGQKVS MHYSDPKPKINEDWL CNKCGVQNFKRREKCFKCGVPKSEAEQKLPLGT RLDQQTLP LGGRELSQGLPLPQPYQAQGVLAS QALSQGSEPSSSENANDTILRLNPNHSTMD SILGA LAPYAVLSSSNVRVIKDKQTQLNRGFAFIQLSTIE AAQLLQILQALHPPLTIDGKTINVEFAKGSKRDM ASNEGSRIASAVASTAIAAAQWAIQSASQGEG TWATSEPPVDYSYYQQDEGYGNSQGTESSLYA HGYLKGTGPGITGTGKDGPTGAGPEASLEPGADS VSMQAFSRPQGAAPGIYQSAEASSSQGTAANS QSYTIMSPA VLKSELQSPHTPSSALPPATSPTAQE SYSQYPVPDVSTYQYDETSGYYYDPQTGLYYDP NSQYYNAQSQQYLYWDGERRTYVPALEQ SAD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GHKETGAPSKEGKEKKEKHKTCTAQQIAKDME RWARSLNKQKENFKNSFQPISSLRDDERRESATA DAGYAILEKKGALAERQHTSMDLPKLASDDRP PPRGLVAAYSGESDSEEEQERGGPEREELTDW QKLACLLCRRQFPSKEALIRHQQLSGLHKQNLBI HRRHLSENELEALEKNDMEQMKYRDRAAERR EKYGIPEPPEPKRRKYGGISTASVDFEQPTRDGLG SDNIGSRMLQAMGWKEGSGGLGRKKQGIVTPIEA QTRVRGSGLGARGSSYGVSTESYKETLHKTMV TRFNEAQ
3235	A	3	1217	PSFLNTGLGPTALGVLGGAGAGLMSNPSPQVP EASTSVCRPKSSMASTSRQRERRRRYLSAGR LVRAQALLQRHPGLDVDAGQPPPLHRACARHD APALCLLLRLGADPAHQDRHGDTALHAAARQG PDAYTDFFLPLSRCPSSAMGIKNKDGETPGQILG WGPPWDSAEEDDASKEREWQKLOGELED EWQEVMMGRFEGDASHETQEPESFSAWSDRLARE HAQKCQQQQREAEGSCRPPRAEGSSQSWRQEE EQRLFREARAKEEELRESRARRAQEALGDREP KPTRAGPREEHPRGAGRGSLLWFGDVPWPCCGG GDPEAMAAALVARGPPLEEQALRRYL RVQV RWHPRFLQRFRSQIETWELGRVMGAVTALSQA LNRHAEALK
3236	A	3	1416	GPASGMAEPTSDFTPIGWHASPELTPTLGLSDT APPRDRWMFWAMLPPPPPLTSSLPAAGSKPSSE SQPPMEAQSLPGAPPPFDAQILPGAQPPFDAQSPL DSQPQPSGQPWNFHAITSWYWRQSSDRFPRHQK SLNPAVKNSYYPRKYDAKFTDFSLPPSRKQKKK KRKEPVHFHFCDDTCDRGFKNQEKYDKHMSEHTK CPELDSCFTAHEKIVQFHWNRNMHAPGMKKIKLD TPPELAWREPRKCNPTLANIEYKIKIEKEE RGAVLTTTQYQKNEELMSRHSQMAKIRSPGKNH KWKNDNSRQRAVTGSGSHLCDLKLEGPPEANA DPLGVNLNSDESCKEKPQHSVIPKEVTPALCSL MSSYGSLSGSESEPEETPKTEADVLAENQVLDSS APKSPSQDVKATVRNFSEAKSENRRKKSFEKTNPK REKRLSQLSNVIRTKNTPSISLGNASSSGHST
3237	A	3806	2204	FVGEQEGGCEAGAGRGAQTYPGEAGERWFGRR RRRGRVVSRRKKMSLKSERRGIHVDQSDLLCKKG CGYYGNPAWQGFCSKCWREYHKKARQKQIQED WELAERLQREEEAFASSQSSQAQSLTFSKFEE KKTNEKTRKVTTVKKFFSASSRVGSKKEIQEAKA PSPSINRQTSIETDRVSKEFIEFLKTFHKTGQEIYK QTKLFLEGMHYKRDLSIEEQSECAQDFYHNVAE RMQTRGKVPPERVEKIMDQIEKYIMTRLKYVVF CPETTDDEKKDLAIQKRIRALRWVTPQMLCVPV NEDIPEVSDMVVKAITDIEMDSKRVPDRDLACIT KCSKHIFNAIKITKNEPASADDFLPTLIYIVLKG NPRLQSNIQYITFCNPSRLMTGEDGYFTNLCCA VAFIEKLDASLNLSQEDFDYMSGQTSRQKQEA ESWSPDACLGVKQMYKNLDLLSQLNERQERIMN EAKKLEKDLIDWTDGIAREVQDIVEKYPLEIKPP NQPLAIDSENVENDKLPPPLQPQVYAG
3238	A	1373	449	VLSVCPTGVFRPAPCRMAFMKKYLLPILGLFMA YYYYSANEEFRPEMLQGGKVVITGASKGIGREM

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				AYHLAKMGAHVVTARSKETLQKVVSHCLELG AASAHYIAGTMEDMTFAEQFVAQAGKLMGGLD MLILNHITNTSLNLFHDDIHHVRKSMEVNFLSYV VLTVAALPMLKQSNQSIVVSSLAGKVA YPMVA AYSASKFALDGFSSIRKEYSVSRVNVSITLCVLG LIDTETAMKAVSGIVHMQAAPKEECALEIKGGA LRQEEVYDSSLWTTLLIRNPCRKILEFLYSTSYN MDRFINK
3239	A	213	422	ERTMQLEIKVALNFIIFYLYNKLLW/QPLKKK*EA HWYPDKPLKGS GFHT/GEMVDPVGELAAKRSGL TVED
3240	A	1255	1425	HESYHVNPNLCNPVAPTSGAHSIG*KWPSWLGA VAHSCNPSTLVGRGGRIIRGQELR
3241	A	161	547	PAGIGRSTAKTPGTPGSLEMENLKSGVYPLKEAS GCPGADRNLVVSFYEKGPLTFRDVAIEFSLEEW QCLDTAQDLYRKVMLENYRNLVFLAGIAVSKP DLITCLEQGKEPWNMKRHAMVDQPPGR
3242	A	50	241	PLPARGKSTLPATFCSPSAPELASMSVVPNRSQT GWPRGVTQFGNKYIQQT KPLTLERTINL
3243	A	380	702	FVAYLKLFFSQVCLFASSEMFFTSRKNMSQKLS LLLVLFGLIWGLMLLHYTFQQRHQSSVKLREQI LDLSKRYVKALAEENKNTVDVENGASMAGYK ITVEYF
3244	A	37	1391	VLMDGRMMRSMRLREBESPGPSHTASCLCGSAP CILCSCCPASRNSTVSRLITFFLFLGVLVSHMLSP GVESQLYKLPWVCEEAGIPTVLQGHIDCGSLLG YRAVYRMCFATAAFFFFFTLLMLCVSSSRDPRA AIQNGFWFFKFLILVGLTVGAFYIPDGSFTNIWYF FGVVGSLFILIQLVLLIDFAHSWNQRWLKAE CDSRAWYAGLFFFTLLFYLLSIAAVALMFMYT LPSGCHCKVFISLNLTFVCVSIAAVL PKVQ QPNSSLQASVITLYTMFVTWSA LPEQKCI HLPTQLGNETVVAGPEGYETQWWDAPSIVGLIIF LLCTLFISLRSSDHRQVNSLMQTEECPPMLDATQ QQQVAAACEGRAFDNEQDGVITYSYFFHFCLVL ASLHVMMTLTNWYKPGETRKMISTWTA VVVKI CASWAGLLLYL
3245	A	52	426	SSLGNEDDEILSLAKDITGMFVASHRKMRAHQV LTFLLLFVITSVAENASTSRGCGLDLLPQYVSLC DLDAIWGIVVEAAAGAGALITLLMLILLVRLPF FKEKEKKSPVGLHFLFLLGTLGP
3246	A	3	515	HEVCGSGCCCCHCCAGGPVARQKALPRLRGVMS RFLNVLRSWLVMVSIAMGNLQSRDHTFLYEK LYTGKPNLVNGLQARTFGIWTLLSSVIRCLCAIDI HNKTLYHITLWTFLLALGHFLSELFVYGTAAPTI GVLAPLMVASFSILGMLVGLRYLEVFPVSRQKK RN
3247	A	1	932	ERLCFPCMQSKIYSYMSPNKCSGMRFP LQEENS VTHHEVKCQGKPLAGIYRKREEKR NAGNAVRS A MKSEEQKIKDARKGPLVFPNQS EAAEPPKTP SSCDSTNAAIKQALKKPIKQKQAPRKAQKGT QQNRKLTDFYPVRRSSRSKAELQSEERKRIDELI ESGKEEGMKIDLIDGKGRGVIATKQFSRGDFVVE YHGDLEITDAKKREALYAQDPSTGCMYFYFY LSKTYCVDATRETNRLGRLINHSKCGNCQTKLH

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				DIDGVPHLILIASRDIAAGEELLYDYGDGRSKASIE AHPWLKH
3248	A	3	870	PGSTISCSELKGTQCRATAGSRGRPPMTCWLRG VTATFGRPAEWPGYLSHLGRSAAMD LGPMRK SYRGDREAFETHLTSLDPVKQFAAWFEEAVQC PDIGEANAMCLATCTRDGKPSARMLLLKGFQKD GFRFFTNFESRKGKELDSNPFASLVFYWEPLNRQ VRVEGPVKKLPEEEAECYFHSRPKSSQIGAVVSH QSSVIPDREYLRKKNEELEQLYQDQEVPKPKSW GGYVLYPQVMEFWQGGQTNRLHDRVFRRLPTG DSPLGPMTHRGEEDWLYERLAP
3249	A	43	1210	TRVGRGESGLKMEVKPPPGRFPQPSGRRRRRRG EEGHDPKEPEQLRKLFIGGLSFETDDSLREHFEK WGTLDTCVVMRDPQTKRSRGFGFVTYSCVEEV DAAMCARPHKVDGRVVEPKRAVSREDSVKPGA HLT VKKIFVGGIKEDTEEYNLRDYFEKYGKIETIE VMEDRQSGKKRGFAFVTDDHDTVDKIVVQKY HTINGHNCEVKKALSKQEMQSAGSQRGRGGGS GNFMGRGGNFGGGGGNFGRGGNFGGRGGYGG GGGGSRGSYGGGDDGGYNGFGGDDGGNYGGGPG YSSRGGYGGGPGYGNQGGGYGGGGGYDGYN EGGNFGGGNYGGGNYNDFGNYSQQQSNYGP MKGGSFGRSSGSPYGGGYGSGGGSGGYGSRRF
3250	A	32	1175	VAGRGDMAALRDAEIQKDVQTYYGQVLKRSAD LQTNCGVTTARPVPKHIREALQNVHEEVALRYY GCGLVIPEHLENCWILDGSGSGRDCYVLSQLVG EKGHVTGIDMTKGQVEVAEKYLDYHMEKYGFQ ASNVTFIHGYIEKLGEAGIKNESHDIVVSNVINL VPDKQQLQEAAYRVLKHGGELYFSDVYTSLELP EEIRTHKVLWGECLGGALYWKELAVLAQKIGFC PPRLVTANLITIQNKELEBVGDCRFVSATFRLFK HSKTGPTKRCQVIYNGGITGHEKELMFDANFTFK EGEIVEVDEETAAILKNSRFAQDFLIRPIGEKLPTS GGCSALELKDITDPFKLAESDSMKSRCPVDAAGGCCGTTKSC
3251	A	32	1175	VAGRGDMAALRDAEIQKDVQTYYGQVLKRSAD LQTNCGVTTARPVPKHIREALQNVHEEVALRYY GCGLVIPEHLENCWILDGSGSGRDCYVLSQLVG EKGHVTGIDMTKGQVEVAEKYLDYHMEKYGFQ ASNVTFIHGYIEKLGEAGIKNESHDIVVSNVINL VPDKQQLQEAAYRVLKHGGELYFSDVYTSLELP EEIRTHKVLWGECLGGALYWKELAVLAQKIGFC PPRLVTANLITIQNKELEBVGDCRFVSATFRLFK HSKTGPTKRCQVIYNGGITGHEKELMFDANFTFK EGEIVEVDEETAAILKNSRFAQDFLIRPIGEKLPTS GGCSALELKDITDPFKLAESDSMKSRCPVDAAGGCCGTTKSC
3252	A	1	574	PLGSNTAPALRVMVQAWYMDDAPGDPRQPHRP DGRPVGLEQLRRLGVLYWKLDADKYENDPELE KIRRERNYSWMDIITICKDKLPNYEEKIKMFYEE HLHLDDEIRYILDGSGYFDVRDKEDQWIRIFMEK GDMVTLPAGIYHRFTVDEKNYTKAMRLFVGEPV WTAYNRPADHFEARGQYVKFLAQT
3253	A	2	984	ARAAAHCGICRLVRWWRKRRSVMGIQTSPVLLA SLGVGLVTLLGLAVGSYLVRRSRRPQVTLDPNE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				KYLLRLLDKTTVSHNTRFRFALPTAHHTLGLPV GKHIYLSRIDGSLVIRPYTPVTSDEDDQGYVDLVI KVYLKGVHPKFPEGGKMSQYLDLKVGDVVEF RGPSGLLTYTGKGFHNQPNKKSPPEPRVAKKLG MIAGGTGITPMLQLIRAILKVPEDPTQCFLLFANQ TEKDILREDLEELQARYPNRFLWFTLDHPPKD WAYSKGFVTADMIREHLPAPGDDVLVLLCGPPP MVQLACHPNLDKLGYSQKMRFTY
3254	A	1	968	LQSAGEGVTHVLLLESPARPVAAVTQVQRRRY HRLSDMSMLAERRRKQK WADPQNTA WSNDD SKFGQRMLEKMGWSKGLGAQEQQATDHIKV QVKNNHLGLGATINNEDNWIAHQDDFNQLLAE NTCHGQETTDSSDKKEKKSFSLEEKSKSKNRVH YMKFTKGKDLLSSRSKTDLDICFGKRQSKKTPEG DASPSTPEENETTTTSAFTIQEYFAKRMAALKNK PQVPVPGSDISETQVERKRGKKRKEATGKDVE SYLQPKAKRHTEGKPERAEAQERVAKKKSAPAE EQLRGPCWDQSSKASAQDAGDHVQPA
3255	A	173	439	GSAAMKVKIKCWNGVATWLWVANDENCIGCR MAFNGCCPDCKVPGDDCPLVWGQCSHCFHMC ILKWLHAQQVQQHCPMCRQEWKFKE
3256	A	2	377	TAARRRQKGTAAARRRQKGTLEEVVLPFRSCRVF WIHSGTTMSKVSFKITLTSDPRLPYKVLSPPESTP FTAVLKFAAEFFKVPAATSAITNDGIGINPAQTA GNVFLKHGSELRIIPDRVGS
3257	A	3	1454	GCSAAAAGAGSGPWAAQEQFPFALLSFFIYNPR FGPREGQEENKILFYHPNEVEKNEKIRNVGLCEAI VQFTRTFSPSKPAKSLHTQKNRQFFNEPEENFWM VMVVRNPIIEKQSKDGKPVIEYQEEELDKVYSS VLRQCYSMYKLFNGTFLKAMEDGGVKLLKERL LKIQSFNIGSLNIVKYTAFLYNDQLINQGLEQ DDMRILYKYLTTSLFPRHIEPELAGRDSPIRAEMP GNLQHYGRFLTGPLNLNDPDAKCRFPKIFVNTD DTYEELHLIVYKAMSAAVCFMIDASVHPTLDFC RRLDSIVGPQLTVLASDICEQFNINKRMSGSEKEP QFKFIYFNHMLAEKSTVHMRKTPSVSLTSVHPD LMKILGDINSDFTRVDEDEEIVKAMSDYWVVG KKSDRRELYVILNQKNANLIEVNEEVKKLCATQF NNIFFLD
3258	A	113	1558	APRGCSMPHRKKKPFIEKKKAVSFHLVHRSQRD PLAADESAPQRVLLPTQKIDNEERRAEQRKYGVF FDDDDYDYLQHLKEPSGPSELIPSTFSAHNRREEK EETLVIPSTGIKLPSSVFASEFEEDVGLLNKAAPV SGPRLDFDPDIVAALDDDFDFFDPDNLLEDDFIL QANKATGEEEGMDIQKSENEDDSEWEDVDDEK GDSNDDYDSAGLLSDEDCMSVP GKTHRAIADHL FWSEETKSRFTEYSMTSSVMRRNEQLTLHDERFE KFYEYQYDDDEIGALDNAELEGSIQVDSNRLQEV NDYYKEKAENCVKLNTLEPLEDQDLPNMNELDES EEEMITVVLBEAKEKWDCECISTYSNLYNHPQ LIKYPKPKQIRISSKTGIPLNVLPPKGLTAKQTE RIQMINGSDLPKVSTQPRSKNESKEDKRARKQAI KEERKERRVEKKANKLAFKLEKRRQEKELLNLK KNVEGLKL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
3259	A	3	964	QMEPGNDTQISEFLLLGFSQEPGLQPFLLGFLFSM YLVTVLGNLLIILATISDSHLHTPMYFFLSNLSFA DICVTSTTTPKMLMNIQTQNKVITYIACLMQMYF FILFAGFENFLLSVMAYDRFVAICHPLHYMVIMN PHLCGLLVLASWTMSALYSLLQILMVVRLSFCT ALEIPHFFCELNQVIQLACSDSFLNHMVIYFTVAL LGGGPLTGILYSYSKIISIIHAISSAQGKYKAFSTC ASHLSVVSFLFYGAILGVYLSAATRNSHSSATAS VMYTVVTPMLNPFTYSLRNKDIKRALGIHLLWGT MKGQFFKKCP
3260	A	34	2573	IPFLKSCCCCLDFPFPPLDQVQEECEVERVTE HGTPKPFKFDSDVAFGESQSEDEQFENDLETDP NWQQVLVSREVLGLKPCIEKQREVNELFYTERA HVRTLKVLDDQVFYQVRVSREGILSPSELKIFSNLE DILQLHIGLNEQMKAVRKRNETSVIDQIGEDLLT WFSGPGEELKHAAATFCSNQPFALMKSRQK KDSRFQTFVQDAESNPLCRRLQLKDIPTQMQR TKYPLLLDNIAITYTEWPTEREKVKKAADHCRQIL NYVNQAVKEAENKQRLDYQRRLDTSCLKSEY PNVEELRNLDLTKRKMIEGPLVWKVNRDKTID LYTLLEDILVLLQKQDDRLVLRCHSKILASTAD SKHTFSPVIKLVTVLRQVATDNKALFVISMNDN GAQIYELVAQTVSEKTVWQDLICRMAASVKEQS TKPIPLPQSTPGEGDNDEEDPSKLKEEQHGIVTG LQSPDRDLGLESTLISSKPQSHSLSTSGKSEVRDL FVAERQFAKEQHTDGTGTLKEVGEDYQIAPDSHLP VSEERWALDALRNLGLLKQLLVQQLGLETSVQ EDWQHFPRTASQGPQTDVQNSNIKAYHSG EGHMPFRTGTGDIATCYSPTSTESFAPRDSVGL APQDSQASNILVMDHMIMTPEMPTMEPEGGLDD SGEIPAREANSDENPSEGDGAVTKEEYVNL RISGNYLILDGYDPVQSSSTLQVASSLTLPMT GIPAVESTHQQHPQNTISDGAISPFTPEFLVQQ RWGAMBEYSCEFISQSSCADSQSQIMEYIHKIEA DLEHLKKVEESYTLQRLAGSALTDKHSDEK
3261	A	1	2100	AVEFAEGALTMAPWPELGDAQPNPDKYLEGAA GQOPTAPDKSKETNKTDNTEAPVTKIELLPSYST ATLIDEPTVDDPWNLPQLQDSGIKWSERDTKGK ILCFFQIGRLILLGLFYFFVCSLDILSSAFQLVG GKMAGQFFSNSSIMSNPLGLVIGVLVTVLVQSS STSTSIVSMVSSLLTVRAAPIIMGANIGTSITNT IVALMQVGRSEFRRAFAGATVHDFNWL SVLV LLPVEVATHYLBITQLIVESFHFKNGEDAPDLLK VITKPFKLIVQLDKKVISQIAMNDEKAKNKS LV KIWCKTFINKTQINVTVPSTANCTSPSLCWTDGI QNWTKNVTYKENIAKQHFVNFHLPDLAVGT ILLISLLVLCGLIMTVKILGSVLKGQVATVIKKT INTDFPPFAWLTGYLAILVGAGMTFIVQSSSVFT SALTPLIGIVTIERAYPLTLGSNIGTTTTAILAAL ASPGNALRSSQLALCHFFFNISGILLWYPIPFTRL PIRMAKGLGNISAKYRWFAVFYLIFFFLIPLTVFG LSLAGWRVLVGVGVPVVFHILVLCRLQLQSRCPR VLPKKLQNWNLPLWMRSLKPWDVAVSKFTGC FQMRCCCCRVCCACCLLCGCPKCCRCCKCCE DLEEAQEGQDVPVKAPETFDNITISREAQGEVPA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SDSKTECTAL
3262	A	30	1377	SQQGSQPHRQGPSSLTAPHSLDLPALPPGPRGS QQKLRRVLVPMSVKPSWGPSPSEGVTAVPTSDL GEIHNWTELLDLFNHTLSECHVELSQSTKRVLVLF ALYLAMFVVGLVENLLVICVNWGRSGRAGLMN LYILNMAIADLGIVLSLPVWMLVTLDTYTWLWG SFSCRFTHYFYFVNMYSSIFFLVCLSVDRYVTLTS ASPSWQRYQHRVRRAMCAGIWLVAIPLPEVV HIQLVEGPEPMCLFMAPFETYSTWALAVALSTTI LGFLLPFLITVFNVLTA CLR RQPGQPKSRRHCLL LCAYVAVFVMCWLPHYVTLTLLTLHGTHISLHC HLVHLLYFFYDVIDCF SMLHCVINPILYNFLSPHF RGRLLNAV VHYLPKDQTKAGTCASSSCSTQH SI IITKGDSQPAAPAAHPPEPSLSFQAHHLLPNTSPISP TQPLTPS
3263	A	1	919	QARSPVAAMASQQLCRALVSAQWVAEALRAP RAGQPLQLLDASWYLPKLGRDARREFEERHIPG AAFFDIDQCDRTSPYDHMLPGAHEFAEYAGRL GVGAATHVVIYDASDQGLYSAPRVWWMFRAFG HHAVSLLDGGLRHWRQLPLSSGKSQAPAEF RAQLDPAFIKTYEDIKENLESRRFQVVDSTRATGR FRGTEPEPRDGIPEGHIPGTVNIPFTDFLSQEGLEK SPEIRHLFQEKKVDSLKPLVATCGSGVTACHVA LGAYLCGKPDVPIYDGSWVWEYMRARPEDVISE GRGKTH
3264	A	1	1398	ARRSTPRTAPRASATRSAAGTMREIVHIQAGQCG NQIGAKFWVISDEHGIDPTGSYHGDSDLQLERI NVYYNEAAGNKYVPRAILVDLEPGTMDSVRS GP FGQIFRPDNFVFGQSGAGNNWAKGHYTEGAELV DSVLDVVRKESESCDCLQGFLTHSLGGGTGSG MGTLSKIREEYPLFIMNTFSVMFJFKVSDTVVE ITNATLSVHQLVENTDETYSIDNEALYDIEFRT KLTPTYGDLNHLVSATMSGVTTC LRFPGQLNA DLRKLAVNMVPPRLHFFMPGFAPLT SRGSQQY RALTVPELTQQMFD SKNMMAACDPRHGRYLT AAIFRGRMSMKEVDEQMLNVQNKNSYFVEWIP NNVKTAVCDIPPRGLKMSATFIGNSTAIQELFKRI SEQFTAMFRRKAFLHWYTGE GMDMEFE TEAES NMNDLVSEYQQYQDATADEQGEFEFEFEDEDEA
3265	A	265	862	WWEDARVLGPFHPEEEGHVWMTPEGARAGTG RELEMLDLSLLALGGLVLLRDSVEWEGRSLLKAL VKKSALCGEQVHILGCEVSEEEFREGFDS DINNR LVYHDFRDPLNWSKTEEAFFGGPLGALRAMCK RTDPVPVTIALDSLWLLRLPCTTLCQVLHVS HQDSCPGETPPSLFPLIHLPLPRSVPLFLSTLE
3266	A	2	884	AAGAGADGREPASERASRAEPPAVAMGQNDLM GTAEDFADQFLRVTKQYLPHVARLCLISTFLEDG IRMWFQWSEQRDYIDTTWNCGYLLASSFVFLNL LGQLTGCVLVLSRNFVQYACFLGFHIALQTIAYS ILWDLKFLMRNALGGLLLLLAESRSEGKSMF AGVPTMRESSPKQYMQLGGRVLLVLMFMTLLH FDASFFSIVQNVGTALMILVAIGFKTKLAALTLV VWLFAINVYFNAFWTIPVYKPMHDFLKYDFFQT MSVIGGLLLVVALGPGGVSMDEKKKEW
3267	A	802	1011	ASTFCSAWKRRSTAALWWSGRASRSHPRELGP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LCFVFGTAALSIRSMDVLSLFLEHGKLVFASGLSP RA
3268	A	490	679	EDAWITNPSSLNARSTPSKPLCYTVLKEGQVVG KTTKASNTREKLRPESERRMVKSFGDEV
3269	A	2	796	GSTHASGARPSLKRARSQRGRPLPSRALPSAHKD MTTNAGPLHPYWPQHRLDNFVPNDRPTWHILA GLFSVTGVLVVTWLLSGRAAVVPLGTWRRLSL CWFAVCGFIHLVIEGWFLVLYEDLLGDQAFLSQ LWKEYAKGDSRYILGDNFTVCMETITACLWG PLSLWVVIAFLRQHPLRFILQLVVSVGQIYGDVLYF LTEHRDGFQHGELGHPLYFWFYFVFMNALWL VLPGLVLDAVKHLTHAQSTLDAKATKAKSKKN
3270	A	17	229	GDTGPQILMSYLDVASKLLQMVKKLSQSFC SNFKYLTKYSRKQVSDEIKKSRRTVESNPIFFKKNKKI Q
3271	A	419	553	IQSGLSLCFADLSETPEGRAGVPGCPHSCDGVAS GRPCSPSSAG
3272	A	1211	1450	FQFIQIELLNILQSLIRNQTQSPYNTTAYPAIDSVIT ILPFSFSCFFIITKCFGLSIFPSVIFLHVYFILTLVVF YCC
3273	A	59	1562	QAWSLQVALSPFFFPASPSNSFAAAVPQLLFP ELPLPHVPGQESAKRRSARRFLIMSELTKELMELVW GTKSSPGLSDTIFCRWTQGFVFSESEGSALQEFEG GPCAVIAPVQAFLLKLLFSSEKSSWRDCSQEEQ KELLCHTLCDILESACDHSGSYCLVSWLRGKTT EETASISGSPAESSCQVEHSSALAVBELGFERFHA LIQKRSFRSLPELKDAVLDQYSMWGNKFGVLLF LYSVLLTKGIENIKNEIEDASEPLIDPVYGHGSQS LINLLLTGHAVSNVWDGDRCSGMKLLGIHEQA AVGFLTMEALRYCKVGSYLKISKIPYLDCLASE THITVFFAKDMALVAPSPSEQARRVEGTTT DNGFIPDGLLEDVMKALDLVSDPEYIN DPEGLGILLGPFLLQEFFPDQGSSESPESFT HYNGLKQSNYNEKVMYVEGTAVVMGFEDPMLQ TDTPIKRCLQTKWPYIELLWTTDRSPSLN
3274	A	186	1358	RVVHRFFKSSAFWPAEVKQPRGGPKTGSRK EGA GSRAPQPVVRSFCGSGVGAEGRMEKLRLLGLRYQ EYVTRHPAATAQLETAVRGFSYLLAGRFADSHE LSELVYSASNLLVLLNDGILRKELRKKLPVSLSQ QKLLTWLSVLECVEVFMMGAAGVWGEVGRW LVIALIQLAKAVLRMLLLLWFKAGLQTSPIVPL DRETQAQPPDGDHSPGNHEQSYVGKRSNRVVRT LQNTPSLHSRHWGAPOQREGROQQHHEELSATP TPLGLQETIAEFLYIARPLLHLLSLGLWGQRSWK PWLLAGVVDVTSLSLLSDRKGLTRRERRELRRR TILLYYLLRSPFYDRFSEARILFLLQLLADHVP GVGLVTRPLMDYLPTWQKIYFYSWG
3275	A	575	759	SVYSASSCKCCNYRKTEQIPDCEQPASSM PERPS HESQPTQMMPLSAPSRAEELGQRP G
3276	A	7	258	KAAGHRLLLAAGHPSMPSSDCLLWEGSLE LRPLQHISLLVLVSTTCLFAFPRVPIAFESK SCLYHCHCAFTVRHYMCSSHTG
3277	A	9	2221	KLGVPEEEGGGDDDEDAEAWAMELADV GAAASSQGVDQVLPTPNASSRVIVHVDLDCFYA QVEMISNPELKD KPLGVQKYLVVTCN YEARKLGVK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KLMNVRDAKEKCPQLVLVNGEDLTRYREMSYK VTEELLEFPVVERLGFDFNFVDTLTEMVEKRLQQLQSD ELSAVTVSGHVYNNQSNLLDVLHIRLLVGSQIAAEMRE AMYNQLGLTGCAGVASNKLLAKLVSGVFKPNQQT VLLPESCQHLIHSLNHIKEIPGIGYKTAKCLEALGINS VRDLQTFSPKILEKELGISVQRIOKLSFGEDNSPVIL SGPPQSFSEEDSFKKCSSEVEAKNKIEELLASLLNRLCQ DERKPHTVRLIRYSSEKHYGRESRQCPIPSHVIQKLGT GNYDVMTPMVDILMKLFRNMVNVKMPFHLTLLSVCF CNLKALNTAKKGLIDYYLMPSLSTTSRSGKHSFKMKDTH MEDFPKDKETNRDFLPSGRIESTRTRESPLDTTNF SKEKDINEFPLCSLPEGVDQEVFKQLPVDIQEILSGKS REKFQKGKSVSCPLHASRGVLSFFSKKQMQDIPINPRD HLSKKQVSSVSPCEPGTSGFNSSSSYMSQKDYSYLDN RLKDERISQGPKEPQGFHF TNSNPAVSASFHSFNLQSE QLFSRNHTTDSHKQTVATDSHEGLTENREPDVDEKIT FPSDIDPQVFYELPEAVQKELLAEWKRTGSDFHIGHK
3278	A	1	876	GLRLHVDLVEKPRTGIMAAETRNVAGAEAPPPQ KRYRQRAHSNPMADHTLRYPVKPEMDWSEL YPEFFAPLTQNSHDDPKDKKEKRAQAQVEFAD IGCGYGGLLVELSPLFPDTLILGLEIRVKVSDYVQ DRIRALRAAPAGGFQNIACLRSNAMKHLPNFFY KGQLTKMFFLFPDPHFRTKHKWRISPTLLAEY AYVLRVGGVLYTITDVLLEHDMCTHFEHPLF ERVPLEDSEDVVGHLGTSTEEGKKVLRNGGK NFPAIFRRIQDPVLQAVTSQTSPLPGH
3279	A	82	2929	TRTKRRLGREKAMASPPRGWGCGELLFPFMLLG TLCEPGSGQIRYSMPPELDKGSFVGNIAKDLGLE PQELAERGVRIVSRGRTOIFALNPRSGSLV DRESLCAQSPLCVVNFNLVEMMEIYGVVEII DINDNFRFRDEELKVKNENAAAGTRLVLPFA RDADVGVNSLRSYQLSSNLHFSLDVVSQTDGQK YPELVLEQPLDREKETVHDLLLTALDGGDPVLSG TTHIRVTVLDANDNAPLFTPSEYSVSVENIPVGT RLMLTATDPDEGINGKLTYSFRNEEEKISETFQL DSNLGEISTLQSLDYESRFYLMVVAAQDGGAL VASAKVVVTVQDVNDNAPEVILTSLTSSISEDCL PGTVIALFSVHDGDSGENGEIACSIERNLPFKLEK SVDNYYHLLTTRDLREETS DYNITLTVMDHGT PPLSTESHIPLKVADVNDNPNFPQASYSTSVTEN NPRGVSIQSVTAHDPDSDGNARVTYSLAEDTFQG APLSSYVSINSDTGVLIALRSFDYEQRLDLQLWV TASDSGNPPLSSNVSLSLFVLDQNDNTPEILYPAL PTDGSTGVELAPRSAEPGYLVTKVVAVDKDSGQ NAWLSYRLLKASEPGLFAVGLHTGEVRTARALL DRDALKQSLVVAVEDHGOPLSATFTVTVAAD RIPDILADLGSIKTPIDPEDLDLTLYLVVAVAAVS CVFLAFVIVLLVLRRLRWHSRLLQAEGRSLAG VPASHFVGVGVRFLQTYSHEVSLTADSRKSH LIFPQPNYADTLLSEESCEKSEPLMSDKVDANK EERRVQQAPPNTDWRFSQAQRPGTSGSQNGDDT GTWPNNQFDTEMLQAMILASASEAADGSSTLGG GAGTMGLSARYGPQFTLQHVQLQELGSDYRQN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VYIPGSNATLTNAAGKRDGKAPAGGNGNKKKS GKKEKK
3280	A	149	1288	GTSQMSSHKGSVVAQNGAPASNREADTAELAE LGPLLEEKGKRVIANPPKAEETCTCPVQEEEE VRVLTPLQAHHAMEKMEEFVYKVWEGRWVVI PYDVLDPDWLKDNDYLLHGHRRPMPSPFRACFKSIF RIHTETGNIWTHLLGFVFLFLGILTMLRPNMYF MAPLQEKVVFGMPFLGAVLCLSFVSWLFHTVYCH SEKVSRTFSKLDYSGIALLIMGSFVPWLYYSFYCS PQRLIYLSIVCVLGISAIIVAQWDRFATPKHRQT RAGVFLGLGLSGVVPTMHFTIAEGFVKATTVGQ MGWFFLMAVMYITGAGLYAARIPERFFPGKFDI WFQSHQIFHVLVVAFAFVHFYGVSNLQEFYGL EGGCTDDTLL
3281	A	1	557	RPRRRQPSFSCRVLVLEDPPCFRFTNSMNQEKLA KLQAQVRIGGKGTARRKKKVVHRTATADDDKKL QSSLKKLAVNNIAGIEEVNMIKDDGTVIHFNNPK VQASLSANTFAITGHABAKPITEMPLGILSQLGAD SLTSLRKLAEQFPQVLDKAPKPEDIDEEDDDV PDLVENFDEASKNEAN
3282	A	155	1139	HALGRRGGQSQELSAACGCFALRLRAPGSGRPA LAPGAAAFAGLGGAPRFPFRGSAAGRTMLLKEY RICMPLTVDEYKIGQLYMISKHSHEQSDRGEGVE VVQNEPFEDPHHGNGQFTEKRVYLNKLPKSWAR AVVPKIFYVTEKAWNYYPYTITEYTCSFLPKFSIH IETKYEDNKGSDNTIFDNEAKDVEREVCFIDIACD EIPERYYKESEDPKHFKSEKTGRGQLREGWRDSH QPIMCSYKLVTVKFEVWGLQTRVEQFVHKVVR DILLIGHRQAFWVDEWYDMTMDDVREYKKN MHEQTNKVCNQHSSPVDDIESHAQTST
3283	A	159	547	IKSKLNQQVEVQSEV...EAKG TMGV... DSGRAAV...VGGVVAVGTVLVALS... GIAASSIAAKMMSTAAIANGGGVAAGSLVALQS VGAAAGLSVTSKVIGGFAGTALGAWLGSPPS
3284	A	227	637	TSNSLLRPDRMSVMDLANTCSSFQSDLDFCSDC SVLPLPGAQDVTVCIRCGFNINVRDFEGKVVKTS VVFHQLGTAMPMSVEEGPECQGPVVDRCRCG HEGMAYHTRQMRSADGQTVFYTCTNCKFQEK EDS
3285	A	123	1535	HRLSYDEAFAMANDPLEGFHEVNLASPTSPDLL GVYESGTQEQTTPSVTYRPHPSALSSVPIQANAL DVSELPTQPVYSSPRLNCAEISSISFHVTDPAPCS TSGVTAGLTKLTRKDNYNAREFLQATITEAC DGSDDIFGLSTDLSRLRSPSVLEVREKGYERLKE ELAKAQRELKLDDEECERLSKVRDQLGQELEEL TASLFEEAHKMOVREANIKQATAEKQLKEAQGKI DVLQAEVAALKTLVLSSSPTSPTQEPLPGGKTPF KKGHTRNKSTSSAMSGSHQDLSVIQPIVKDCKEA DLSLYNEFRLWKDEPTMDRTCPLDKIYQEDIFP CLTFSKSELASAVLEAVENNTLSIEPVGLQPIRFV KASAVECGGPKKCALTGQSKSKHRIKLGDSN YYYISFFCRYRITSVCNFFTYIRYIQQGLVKQQDV DQMFWEVMQLRKEMSLAKLGYFKEEL
3286	A	3	589	GPSQSMAGLEGGKPLSGLLNALAQDTFHGYF GITEELLRSQLYPEVPPPEFRPFLAKMRGILKSIA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				ADMDFNQLEAFLTAQTKKQGGITSDQAAVISKFWKSHKTKIRESLMNQSRWNSGLRGLSWRVGDKSQSRHSAQIHPTVAIELELGKYGQSEFLCLEFD EVKVNQILKTLSEVEESISTLISQPN
3287	A	50	390	LGAMAKHHPDLIFCRKQAGVAIGRLCEKCDGKC VICDSYVRPCTLVRICDECNYGSYQGRVCVCGGP GVSDAYYCKECTIQEKDRDGC PKIVNLGSSKTDL FYERKKYGFKKR
3288	A	3	428	RTTFRFRPCESLCGDMKLLTHNLLSSHVRGVGS RGFPLRLQATEVRICPVEFNPNFVARMIPKVEWS AFLEAADNLR LIQVPKGPVVEGYEENEFLRTMH HLLLEVEVIEGTLCPESGRMFPISRGIPNMLSE EETES
3289	A	1	1743	AGCCRDTRFPTPRGPGSLCHNFCRSAACTVTRTI HGSPREDTGTPRSREMMFQDSVAFEDVAVSFTQ EEWALLDPSQKNLYRDMQETFKNLTSVGKTW KVQNIIDEYKNPRRNL SLMREKLCEKESHHC ESFNQIADDMLNRKTLPGITPCESSVCGEVGTGH SSLNTHIRADTGHKSSEYQBYGENPYRNKECKK AFSYLDSFQSHDKACTKEKPYDGEKCTETFISHS CIQRHRVMHSGDGPYKCKFCGKAFYFLNLC LIH ERIHTGVKPYKCKQCGKAFTSTTLPVHERHTTG VNADECKECGNAFSFPSEIRRHKRSHTGEKPYEC KQCGKVFISSSIQYHKMTHTGEKPYECKQCGK AFRCGSHLQKHGRHTHTGEKPYECRQCGKAFTCT SDLQRHEKTHTEDKPYGCKQCGKGFRCASQLQI HERTHSGEKPHECKBCGKVFKYFSSLRIHERHT GEKPYECKQCGKAFTYFSSLHIHERHTGDKPYE CKVCGKAFTCSSIRYHERHTHTGEKPYECKHCGK AFISNYIRYHERHTHTGEKPYQCKQCGKAFTIRASS CRHEHTHTNP
3290	A	2	330	GRPRSSSDNRGLNAGLSSAAVQTRIGNSAAS RRSPAARPPVPAPALPRGRPGTEGSTLSAPAVL VVAVAVVVVVVSAVAVAMANYIHPVPPGSPEVP KLNVTVDQEEHRCREGALSLLQHLRPHWDPOE VTLQLFTDGIITNKLIGCYVGNMTMEDVVLVRIYGN KTELLVDRDEEVKSFRVLQAHGCAPOLYCTFNN GLCYEFIQGEALDPKHVCNPAIFRLIARQLAKIHA IHAHNGWIPKSNLWLKMGKYFSLIPTGFADEDIN KRFLSDIPSSQILQEEMTWMKEILSNLGSPVVLCH NDLLCKNIYNEKQGDVQFIDYEYSYNYLAYDI GNHFNEFAGVSDVDYSLYPDRELQSQWLRAYLE AYKEFKGFGTEVTEKEVEILFIQVNOFALASHFF WGLWALIQA KYSTIEFDLGYAIVRFNQYFKMK PEVTALKVPE
3291	A	102	839	PEAQTS AVLAREKGHLPTMRHEAPMQMASAQD ARYGQKDSSDQNFDMFKLLIIGNSSVGKTSFLF RYADDSFTSAFVSTVGIDFKVKT VFKNEKRIKLQI WDTAGQERYRTITTA YYRGAMGFILMYDITNEE SFNAVQDWSTQIKTYSWDNAQVILVGNKCDME DERVISTERGQHLGEQLGFEFFETSAKDNINVKQ TFERLVDIICDKMSESLETDPAITAAKQNTLKET PPPPQPNAC
3292	A	2	4136	DRPPWNSRVDDFVTNLIHLSSKGHISPAKDTSLQ QRTPAEMSPVLHFYVRPSGHEGAASGHTRRKLQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *-Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				GKLPQLQGVETELCYNVNWTAEALPSAETKKL MWLFGCPLLLDDVARES WLLPGSNDLLLEVGP LNFSTPTSTNIVSVCRA TGLGPVDRVETTRRYRLS FAHPPSAEVEAIALATLHDMTEQHFPPIQSFSP ESMPEPLNGPINILGEGRLALEKANQELGLALDS WDLDFYTKRFQELQRNPSTVEAFDLAQNSEHS RHWFFKQQLHVDGQKL VHS LFESIMSTQESSNP NNVLKFCDNSSAIQKQEVRLRPEDPTRPSRFQQ QQGLRHVVFTAETHNFTGVC PFSGATTGTGGRI RDVQCTGRGAHV VAGTAGYCFGNLHIPGYNLP WEDLSFQYPGNFARPLEVAIEASNGASDYGNKF GEPVLAFARSLGLQLPDGQRREWIKPIMFSGGI GSMEADHISKEAPEPGMEVVKVGGPVYRIGVGG GAASSVQVQGDNTSDLD FGA VQRGDPMEQKM NRVIRACVEAPKGNPICS LHDQ GAGGNGNVLKE LSDPAGAIITYTSRFQLG DPTLNALEIWGA EYQESN ALLRSPNRDFLTHVSARERCPACFVGTTTGDRRI VLVDDRECPVRRNGQGDAPPTPPPTPVDELEW VLGKMPRKEFFLQRKPMLQPLALPPGLSVHQA LERVLR LPAVASKRYL TNKVDRSVGGLVAQQQC VGPLQTP LADVAVVALSHEELIGAATALGEQPV KSLDDPKVAARLAVAEALTNL VFALVTDLRDVK CSGNWMWAAKLPGEGAALADACEAMVAVMA ALGVAVDGGKDSL SMAARVGTETVRAPGSLVIS AYAVCPDITATVTPDLKHPEGRGHLLYVALSPG QHRLGGTALAQCFSQLGEHPDLDLPENLVRAFS ITQGLLKDRLLCSGHDVSDGGLVTCLEMAFAG NCGLQVDVPVPRVDVLSV LFAEEPGLVLEVQEP DLAQVLKRYRDAGLHCL ELGHTGEAGPHAMVR VSVNGAVVLEEPVGELRALWEETS FQLDRLQAE PRCVAE EFRGI RERMGPSYCLPFAKASVPFEP GGPSPRVAILREBGSNDREMA DAFHLAGFEVW DVTMQDLCSGAIGLDIFRGVAFVGGFSYADVLG SAKGWAAAVTFHPRAGAE LRRFRKRPDTFSLGV CNGCQLLALLGWVGGDPNEDAAEMGPDSQPAR PGLLLRHNL SGRYESRWASVRVGPALMLRG MEGAVLPVWSAHGEGYVAFSSPELQAQIEARGL APLHWADDDGNPTEQYPLNPN GSPGGVAGICSC DGRHLAVMPHPERAVRPWQWAWRPPPFDTLTT SPWLQLFINARNWTLEGSC
3293	A	65	642	GVRGFWAGTMASRAGPRAAGTDGSD FQHRERV AMHYQMSVTLKYEIKKLIYVHLVIWLLLVAKMS VHLLRLLSHDQVAMPYQWEYYPYLLSILPSLLGLL SFPRNNISYLVLSMISMGLFSIAPLIYGS MEMFPA AQQLYRHGKAYRFLFGFSAVSIMYLVVLAVQV HAWQLYYSKLLDSWFTSTQEKKKH
3294	A	35	1821	SQRSCPRSPSSPAPPWARCSNPDSRTGGVPVPR WSAGGPALGLMAAPVRLGRKRPLPACPNPLFVR WLTEWRDEATRSRHRTRFVFQKALRSLRRYPLP LRSGKEAKILQHFGDGLCRM LDERLQRHRTSGG DHAPDSPSGENSPAPQGR LAEVQDSSMPVPAQP KAGGSGSYWPARHSGARVILLVLYREHLNPN GH HFLTKEELLQRCAQKSPRVAPGSARPWPALRSL HRNLVLRTHQPARYSLTPEGLELAQKLAESEGLS LLNVGIGPKPEPPGEETA VPGAASAE LASEAGVQQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				QPLELRPGEYRVLLCVDIGETRGGGHRPELLREL QRLHVTHTVRKLVHVGDFVWVAQETNPRDPANP GELVLDHIVERKRLDDLCSSIIDGRFREQKFRLLKR CGLERRVYLVEEHGSVHNLSPSTLLQAVTNTQ VIDGFFVKRTADIKESAAYLALLTRGLQRLYQGH TLRSRPWGTGPNPESGAMTSPNPLCSLLTFSDFN AGAIKKAQSVREVFARQLMQVRGVSGEKAAA LVDRYSTPASLLAAYDACATPKQETLLSTIKCG RLQRNLGPALSRTLSQLYCSYGPLT
3295	A	2	1115	EFHPHTQVSGLLTPQLQEPDVWSPSRGQPVSLHL PGKGAPEVKEMAWWKSWEQEGVTVKSSSHFN PDPDAETLYKAMKGIGTNEQAIDVLTGRSNTQR QQIAKSFKAQFGKDLTETLKSELSGKFERLIVAL MYPPYRYEAKELHDAMKGLGTKEGVHILASRT KNQLREIMKAYEEDYGSSLEEDIQADTSGYLERI LVCLLQGSRDDVSSFVDPALALQDAQDLAAGE KIRGTDEMKTITLCTRSATHLLRVFEEYEKIANK SIEDSIKSETHGSLEEAMLTVVKCTQNLHSYFAE RLYYAMKGAGTRDGTLRNIVSRSEIDLNLKCH FKKMYGKTLSSMIMEDTSGDYKNALLSLVGSDF
3296	A	1	838	GTRGGVPGDNGGVEAGAKPGAAIPLRGDGS GETGPGRVAPGEVRGSPRGHVAGPEGPREVLF FLPSSKPASEVINEYSWKVDFLKGMLQAEKLTSS SEKALANQFLAPGRVPTARERVPATKTVHLQS RARTSEMRSELLGTDSAPEMDVRKRTGVAGS QPVSEKQSAAEGLDLVLRHQNLQEKLAEEMLGL ARSLKTNLAAQSVIKKDNQTLSHSLKMADQNL EKLKTESERLEQHTQKSVNWLLWAMLIVCFIFIS MLFIRIMPKLK
3297	A	46	617	HKQPAGFLGLWLGTTETYTISFPGPETFGGLSHA TGIPCSACROFVGLHSLINVMAMVSAMS VLYLWISACVGLHSLQITFCQHLHRPEGG TCBVIAAHRCCKNRIEERSQTVKCSCLPGKVAG TTRNRPCVDASVIGKWWCEMPCLEGECKTL PDNSGWMCATGKNTTRIHPRT
3298	A	157	748	IQPPDPRNMTLAAYPEKMKELPLVSLFCSCFLAD PLNKSSYKYEADTVDLNWCVISDMEVIELNKCT SGQSFEVILKPPSFDGVPEFNASLPRRRDPSLEEIQ KKLEAAEERRKYQEAELLKHLAEKREHEREVIQ KAIEENNNFIKMAKEKLAQKMESNKENREAHLA AMLERLQEKDKHAEVRKNKELKEEASR
3299	A	5	892	TQLPAPLSGVL SRLQLGSGAPLLTWVQETAGVA GGAPRRRTPTVMWRLLARASAPLLRVPLSDSWA LLPASAGVKTLPLVPSFEDVSIPEKPKLRFIERAPL VPKVRREPKNLSDIRGPSTEATEFTEGNFALG GGYLHWGHFEMMRLTINRSMOPKNNMFAIWRVP APFKPITRKSVGHRMGGGKGGAIDHYVTPVKAGR LVVEMGGRCEFEVVGFLDQVAHKL PFAAKAVS RGTLEKMRKDQEERERNQNPWTFERIATANML GIRKVLSPYDLTHKGKYWGKFYMPKRV
3300	A	2	1847	FVAGGPRGSGSAAETMPEIRVTPLAGQDVGRS CILVSIAGKNVMLDCGMHMGFNDDRRFPDFS TQNGRLTDFLDCVIISHFHLDHCGALPYFSEMVG YDGPYIMTHPTQAICPILLEDYRKIAVDKKGEAN FFTSQMIKDCMKKVAVVHLHQTVQVDDELEIKA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				YYAGHVLGAAMFQIKVGSSESVVYTG DYNMTPD RHLGAAWIDKCRPNLLITESTYATTIRDSKRCRE RDFLKKVHETVERGGKVLIPVFALGRAQELCILL ETFWERMNLKVPIYFSTGLTEKANHYKLFIPWT NQKIRKTFVQRNMFEFKHIKAFDRAFADNPGPM VVFATPGMLHAGQSLQIFRKWAGNEKNMVIMP GYCVQGTVGHKILSGQRKLEMEGRQVLEV KMQ VEYMSFSAHADAKGIMQLVGQAEPESVLLVHGE AKKMEFLKQKIEQELRVNCPMPANGETVTLPTS PSIPVGISLGLLKREMAQGLLPEAKKPRLLHGTLI MKDSNFRLVSSEQALKELGLAEHQLRFTCRVHL HDTRKEQETALRVYSHLKSVLKDHCVQHLPDGS VTVESVLLQAAAPSEDPGTVLLVSWTYQDEEL GSFLTSLKKGLPQAPS
3301	A	2	349	CIRTEPAAAFRRLLGALSGAAALGFASYGAHGAQ FPDAYGKELFDKANKHHFLHSLALLGVPHCRKP LWAGLLLASGTTLFTCSFYQALSGDPSIQTLP AGGTLGLLGLALAL
3302	A	59	1184	LRNCSALGGLFQTIISDMKGSYPVWEDFINKAG KLQSQLRTTVVAAAFLDAFQKVADMATNTRG GTREIGSALTRMCMRHSIEAKLRQFSSALIDCLI NPLQEQMEEWKKVANQLDKDHAKEYKKARQEI KKKSSDTLKLQKKAKKGRGDIQPLDSALQDVN DKYLLLEETEKQAVRKALIEERGRFCTFISMLRP VIEEISMLGETHLQTISEDLKSLTMDPHKLPSSS EQVILDLKGS DYSWSYQTPPSSPSTMSRKSSVC SSLNSVNSSDSRSSGSHSHSPSSHYRYSNLAQQ APVRLSSVSSHDSGFISQDAFQSKSPSPMPPEAPN QRRKEKREPDNPGGPTTASGPPAAAEAAQRPRS M
3303		511	558	AGRGPGPKVSVSWSSCPGSPGQTQRR HSSLPPSQDFVACLSVE VDDRLEWAFNLY DLNKDGCITKEMLDIMKSIYDMMGKYTYPALR EEAPREHVESFFQKMDRNKDG VVTIEEFIESCQK DENIMRSMQLFDNVI
3304	A	40	432	ISEAASGAFQAR*FYQMLEQKTDALGKQSVNRG FTKDKTLSSIFNIEMVKEKTAEIKQIWQYFAA KDTVYAVIPA EKFDLIWNRAQSCPTFLCALPRRE GYEFFVGQWTGTELHFHCTYKYS DPEGKA
3305	A	2	483	LDACSTGPYSRSTHASADAWADAVVVVLKV GMTLFLLYFPQIFNKSNDGFTTTRSYGTVSQIFGS RSPSPNGFTTTRSYGTVCPKDWEFYQARCFLLIHL *SSWNESWDFCKGKGCTLAIVDNSETLKLHDL HDAEKNYIALPYRSSKYMSTCNGTF
3306	A	2	872	TLSSACLIGDAWKELTVAGAVSNQLLVWYPAT ALADNKPVAPDRRISGHVGIIFSMSYLESKGLLA TASEDRSVRIWKGGDLRVPGRVQNIHCFGHS ARVWQVKLENYLISAGEDCVCLVWSHEGEILQ AFRGHQGRGIRAIAAHERQAWVTGGDDSGIRL WHLVGRGYRGLG/DLGSLLQVP**ARYTQGCDS GWLLATAGSD*YRGPVSL*RRQVLGAAARG*T FPVLLPAGGSSWSRGLRIVCYGQWGRSCQGCPH QHSNCCGPDVPSWEGAQLELGPWL
3307	A	2	927	RTSRVEKGLRKAGAAVTMESDEWFSQALPANTS AQKAELIALTQAIRWGKDINVNTDSRYAFATVH

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VRGAICQERRLLTSAEKAIAKKNPPSSKPNRSSSF WGTTCDQVNAKQGPSPGHRLRRNLPGKEKWEI DFTKVKPHQAGYKYLLVLVDTFSGWTEAFATK NETVNMVVKFLLNEIIPRHGLPVAIGSDNGPAFA LSIV*SVSKALNIQWKLHCAYRPQSSGQVERMNC TLKNITLTKLILETGWNWVSLPLALLRVRCTPYW AGFLPFEIMYGRVLPILPKLRDAQLAKISQTNLLQ YLQSP
3308	A	490	1077	NPSLDFNDNEDIPTELSDSSDTHDEGEVQAFYE DLGRQYVNEVFNFSDKLYDLLFTNSPFQRDF MEQRRFSDIIFHPWKKEENGNSRVIPYTITLTNP LEHKTATVRETQTMKASQESECYVIDAEVLTH DVPYHDYFYTINRYLTRVARNKSRLRVSTELRY RKQPWGLVKTFIEKNFWSGLEDFYRHL
3309	A	490	1077	NPSLDFNDNEDIPTELSDSSDTHDEGEVQAFYE DLGRQYVNEVFNFSDKLYDLLFTNSPFQRDF MEQRRFSDIIFHPWKKEENGNSRVIPYTITLTNP LEHKTATVRETQTMKASQESECYVIDAEVLTH DVPYHDYFYTINRYLTRVARNKSRLRVSTELRY RKQPWGLVKTFIEKNFWSGLEDFYRHL
3310	A	2	1198	SPLCHPGLSRER/S*SEAKLRSGRYC*KRQVEAPL *RPLG*TMAASDTERDGLAPEKTSPPDRDKKKEQS EVSVPRAKSHHYSRERSRERKRKSDNEGRKH RSRERSKEGRRHESKDKSSKKHKSEEHNDKEHSS DKGRERLNSSENGEDRHKRKERKSSRGRSHRS RSRERRHRSRERKKSRSRERKKSRSRER KKRSRERSRERKRIRSRERSRHRHRTSRSRTR SRSDRDKKRIEKPFRFSRSLSRTPSPPPFRGRNTA MDAQEALARRLERAKKLQEQREKEMVEKQKQ EIAAAAAATGGSVLNVAAALLASGTQVTPQJAMA AQMAALQ...LAATTGAVPST...AVNP... ...EQKKRKMLWQKKKEG...SAC...MGN
3311	A	177	4	PIQIPPRITPRPSPHLLTPRTGSSPPRAPSPPHPT PGAHDFFPLSAVLSGHTKT
3312	A	3	426	LESPRH*PPCWGPLIWALTVSSVPSPTELSCLKS P/RPACPV/PGLWPSLLSPAPPQSSGPLLGLSPCPG AGQWPSPLSPAPPPSDPLSGLSPCPGAGPRSSPS ASAPCRAVPLSPRRLTWPPHLQVGILIPTGRPWK NL
3313	A	162	2	QLQNLASRGCL*SQLRRLRRENRLNPGGGGCSE IAPCTPAWVTQRDFFRKKK
3314	A	162	2	QLQNLASRGCL*SQLRRLRRENRLNPGGGGCSE IAPCTPAWVTQRDFFRKKK
3315	A	466	1	PRKRESWWGERLP/PRGFPPAAEDAPAPGWKGR KHASRTARAHVFHPIRQSIRSPVGRPGDPRAAH TRSAGTRLQCKASRG*GKGPAPTR*EGGPGSAP APLPASSGCSLFPDSSPWTPPPAPGAAAAQP**T PRCPAALRAGAHIGRVGRPY
3316	A	3	2307	NHLGTLMQNWDSSSRVPFSSGQHSQSFPPSLMS KNSMLQKPTAYVRPMDGQESMEPKLSSEHYSS QSHGNSMTELKPSKAHLTKLKIPSQPLDASASG DVSCVDEILKEMTHSWPPLTAIHTPCKTEPSKFP FPTKESQSNFGTGEQKRYNPSKTSNGHQSKSM LKDDLKLSSESDSDGEQCDKTMPRSTPGSNSEP SHHNEGADNSRDDSSSHSGSESSGSDSESESS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SDSEANEPSQSASPEPEPPPTNKWQLDNWLNKV NPHKVSPASSVDSNIPSSQGYKKEGREQGTGNSY TDTSGPKETSSATPGRVAPKPIQKGSSEGRGRQKS PAQSDSTTQRRTVGKKQPKKAEKAAAEPRGGL KIESETPVDLASSMPSSRHKAATKGSRKPNIKKES KSSPRPTAEKKKYKSTSKSSQKSREHETDTSSSDS DESESLPPSSQTPKYPSNRTVPVKSSVEEEDSFFR QRMFSPMEEKELLSPLSEPDDRYPLIVKIDLNLLT RIPGKPYKETEPKGEKKNVPEKHTREAAQKQASE KVSNGKGRKHKNEEDNRASESKKPKTEDKNSA GHKPSNRESSKQSAAKEKDLLPSPAGVPVPSKDP KTEHGSRKRTISQSSSLKSSSNSNKETSGSSKNSS STSKQKKTGKTSSSSKEVKVKAPSSSSNCPPSAP TLDSSKPRRTKLVFDDRNYSADHYLQEAQKLLKH NADALSDRFKAVYYLDAVVSFIECGNALEKNA QESKSPFFMYSETVDLI
3317	A	496	2	NLLQDEKLVHSYPYDWRQTQETCGYIVPARQWFI NTRDIKTAAKELLKKVKFIPGSALNGMVEMMD RRPYWCISRQVRVWGVPIPVFHHKTKDEYLINSQT TEHIVKLVEQHGSDIWWTLPPQEQLLKEVLSEVG GPDALYVPGQDILDWFDSGTSWSYVLPGBP
3318	A	2	512	AWHEGDSRSDQCHHPYNYGFDYYYGMPFTLVD SCWPDPSRNTLAFESQLWLCVQLVAIALTLTF GKLSGWVSVPWLLIFSMILFIFLLGYA WFSSTSP LYWDCLLMRGHEITEQPMKAEVRAGSIMVKEAIF LFRKGHSKGLFLLFFLPFLQVHKTFPTTDGFHW AP
3319	A	407	1	SSLHRSRPRASPLPVPEAP\SFLPVPAKPSALPPFS LSGAPSSASTFSPHSSPASPTAPSPQSPFSPRPT SPPSLTPTRPPLPADRRGPHELLYQPLHAPLEAAA TGFPSAAACRLRPRP*V*A*Y*A*S*
3320		4037	3432	QMSEAVAEMKMLQYRRDTAGW/KIC*GVSVS WRPSVEFFGNLYRGEGIVYGTLEEVWDCVKPAV GGLRVKWDENVTGFEIJSITDITLCVSRSTSPSAA MKLISPRDFVDLVVKRYEDGTISSNATHVEHPL CPKPGFVRGFNHPGCGCFCEPLGEPKTNLVTF HTDLSGYLPQNVVDSFFPRSMTRFYANLQKAVK
3321	A	37	360	SHSASGAGRPAAPAADLRPAPNGQRPGRPGAR ALWLPGRGRPDEAGRLPGEHLQVPWDPGLTRS PSPRGPCRGAARAGHVGETAPWGCPPPCAWEH KGPGESETP
3322	A	1	420	AIVEDKHSGRSYDITSDLGNVLTSTSIKTVNG*A ESSDSGAESDEEDAQEDLMGAYHSDIDKKMMKI VADHKNLEVIVTNGYDKDGFVHDIQNDIHASSSL NGRSTVHVKPIDENLGQTGKSAVCIHQDINDDH VEDVT
3323	A	8	459	DTLSLNCITLPETLPMTPSF*LSFL*FPGLARAKSIP TKTYSNEVVTLWYRPPDILLGSTDYSTQIDMW*G QVEVWQGPCGKGGGLVTTATQPAALFTVPSLP RGVGCIFYEMATGRPLFPGSTVEEQHLHFIFRILSE EAWALCAVETHR
3324	A	1276	466	PGSTHASARITTY*L*IILSNATEVDNNSKPPPPFP AGAPPASSSSSSSSSPPTVSTAPPLIPPPGFPPPPG APPPSLIPTIESGHSSGYDSRSARAFYGNVAFPH LGSAPSWPSLVDTSKQWDYYARSSSSSSSSSSSS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SSSPRDRDRER*RTREERERERDHSPTSPVFNSEDEE RYRYREYAERGYERHRASREKEERHRERRHREK EETRHKSSRSNSRRRHESEEGDSHRRHKHKKSKR SKEGKEAGSEPAPEQESTEATPAE
3325	A	266	3312	<p>TCLFSASCSSLPSPSSSFALLSTENTQRTYRVNPD GSLRVTFASGMEIGLSSEPHILAGAVNPTLGKCN SLPGEHNANLISVL**GEQGCA*NVFHISFS*AHN RNLLSIDFDHITRTGKIYDDHRKFTLRJLYDQTGR PILWSPVSRYNEVNITYSPSGLVTFIQRGTWNEK MEYDQSFL*SPQL*LSIICYSAFVSFQSVMLLLHS QRRYIFEYDQPDCLLSVTMPMSMVRHSLQTMLSV GYRNTYTPDSSTSFQDYSRDGRLLQTLHLGTG RRVLYKYTKQARLSEVL YDTTQVTLTYEESGD LSDSSTLIA*LLTVFVLVPAGPLIGRQIFRSEEG VNARFDYSYNNFRVTSMQAVINETPLPIDLYRYV DVSGRTEQFGKFSVINYLNQVITTTVMKHTKIF SANGQVIEVQYEILKALAYWMTIQYDENVGRMVI CDIRVGVDANITRYFYEYDADGQLQTVSVNDKT QWRYSYDLNGNINLLSHGKSARLTPLRYDLRDRI TRLGEIQYKMEDEGFLRQRGNDIFEYNSNGLLQ KAYNKASGWTQVQYYDGLGRRVASKSSLGQHL QFFYADLTNPIRVTHLYNHTSSEITSLYDLOGH LIAMELSSGEEYYVACDNTGTPLAVFSSRGQVIK EILYTPYGDYHDTYPDFQVIHFGHGLYDFLTKL VHLGQRDYDVVAGRWTTPNHHIWKQLNLLPKP FNLSTKLIKYGIFHLFLILCLDIRSWLELFGFQL HNVLPGFPPPELENSPSI*QMSNSMLHLLCASLS* TILGIQCELOKQLRNFISSLDQLPMTPRYNDGRCL GGKQPRFAAVPSVFGKGIKFAIKDGIVTADIGVA NEDSRRLAAILNNAHYLENLHFTIEGRDTHYFIK F GSLPFEHNTGNTGREGLENGVNVTVSQMTSV LNGRFGNTLQLQHGALCFNRYGTTVEEEK:MH VLEIARQAVAAQAWTKEQRRLOEGEEGIRAWTE GEKQQLLSTGRVQGYDGYFVLSVEQ</p>
3326	A	290	1041	<p>KACLHLLSSFLTSNLFNPLLPDSLYSVEARSQRA NLGPCRRKRLQILMRLAAGFYSSHKDPSLSAK EKHTDYHNEARGPWPGWVG*RTADGSCGRGPD GAHHPGPKSSSWRASRLPLGLGSHHLDAYVGR DLECGTPAPLQLEIPPQPRGHPAPIPTGQAGPRDS GPGASP*VETRPLTDGRR*PGVRPVGWTPAHPAG TLRPRGAVEPSVSACGKWAPSPTSQCCEGRCD AVPKHRAWRTPLCSQ</p>
3327	A	1	418	<p>CSECGKSFCKKSKFIHQRTHTGEKPYECNQCGK SFCQKGTITVHQRTHTGEKPYECNECGKNFYQK LHLIQHQRTHSGEKPYECYCGKSFCQKTHLTQH QRTHSGERPYVCHDCGKTFQSQSALNDHQKIHT GVKLY</p>
3328	A	1	270	<p>VTRKLPITVDAFTARAFRGSPAADCLENELDED MHQKIAREMNLSETAFIRKLHPTDNFAQRSCFGL IWFPTTDLQILTSSILPSIL</p>
3329	A	45	419	<p>EELSCWQIWQQLANDLTRCQDSMINNSQCHKQOG DFPYQVGTLSIQISEDENYIVNKADGPNNTGNP EFPILRTQDSWRKTFLTESQRLNRDQQISIKNKL CQCKKGVDPIGWISHHDGHRVHKR</p>
3330	A	64	430	FWRNFTGLAPAAAVATTSSSTMRFTSISNSLTST

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				AAIGLSFTTSTTTTATFTTNTTTTTITSGFTVNQNO LLSRGFENLVPTSTVSVVTPVMTYGHLEGLIN EGNLELEIKRRLSSQATQ
3331	A	3	407	TFGCSCSTDCFFQKCCPAEAGVLLAYNKNQKQIP PGTPIYECNSRCQCGPDCPNRIVQKGTQYSLCIFR TSNGRGWGVKTLVKIKRMSFVMEYVGEVITSEE AERRGQFYDNKGITYLFDLDYESDEFTVDAARY
3332	A	25	461	PAADFVLQARPTRADILGIHISKYDEVKAGACFY KMTGLGPGPQALYNGEPFKHEEMNIKELKMAVL QRMMMDASVYLQREVFLGTLNDRTNADFLMDR NNVVPRINTLILRTNQYLNLLSTSVTADAEDFS TFFFLDSQDKSA
3333	A	317	54	AWIFLPLTSCPLWAPGTHKHTLEARSGLGPIK AYPRLGPPTPGEPEAPAQDRTFHCEICNVKVNSK VQLKQHISRRHEIVDPV
3334	A	304	410	AGPSLPSNLQIFQSLPPFMDILLLLFFMIIFAI
3335	A	19	418	VESRNSRVQPRVRLNDRTNADFLMDRNNVVPRI NTLILRTNQYLNLISTSVTADVEDFSTFFFLDSQ DKSAVIAKNMYYLTQDDESIISAATLWIADFDK PSGRKLLFNALKHMITSVHSRVGIIYNPFF
3336	A	1	1003	PSSYSSDELSPGEPLTSPWPAPLGAPEPEHLLNR VLERLAGGATRDSAASDILLDDIVLTHSLFLPTEK FLQELHQYFVRAGGMEGPEGLGRKQACLAMLL HFLDTYQGLLQEEGAGHIKDLYLIMKDESLY QGLREDTLRLHQLVETVELKIPEENQPPSKQVKP LFRHFRIDSCLOTRVAFRGSDIFCRVYMPDHS YVTIRSLSASVQDILGSVTEKLOYSEEPAGREDS LILVAVSSSGEKVLLQPTEDCVFTALGINSHLFAC TRDSYEALVPLPEEIQVSPGDTEIHRVEPEDVANH LTAHFHWELFRVHELEFVDYVFHGE
3337	A	444	43	KILIGLANQFP...CF...LP...VA...DEALY CFEELACRLACNDPGRRLIDQSFLATESSCMTFGD LVNKYCQAAHKLMVAVSEDVLQVADWQRWL FGELPLCYFARVFDVFLVEGYKVLYRVALAXXF
3338	A	1	398	FRGKVRGRSAEMPGSDTALTVDRTYSDPGRHR CKSRVERHDMNTLSLPLNIRRGSDTNLNFDPD GILDFHKVKLTADSLKQKILKVTEQKIEQTSRDG NVAEYLKLVNNADKQAGRIKQVFEKKNQK
3339	A	1	665	AAAASNWGLITNIVNSIVGVSVLTPMPCFKQCGI VLGALLVFCSWMTHQSCMFLVKSASLSKRRTY AGLAFHAYGKAGKMLVETSMIGMLMGTCIAFYV VIGDLGSNFFARLFGFQVGGTFRMFLFAVSLCI VLPLSLQRNMMAISQSFSAAMALLFYTVFMFVIVL SSLKHGLFSGQWLRVSYVRWEGVFRICIPIFGMS FACQSQVLPTDYDSLDEPSV
3340	A	198	367	LLPLQVLQEAFSRCVAVLTRSSKPSDMSVQVCG YISKCYSVAAQFECEKREKITEMP
3341	A	562	277	HSVIKRTPRKYLAELVLDLDFSNKEHLKEKLDEYI KLWNLGVKVRNERREGLIQARSIGAQAQKALGQ VLIYLDACEVAVNWWYAPLVAPISKDR
3342	A	385	2	NLTWWPLFRDVSFYIVDLIMLIIFFLDNVIMWWE SLLLLTAIFYCYVFMKFNQVEKWKQMINRN KVVKVTAPEAQAKPSAARDKDEPTLPAKPRLQR GGSSASLHNSLMRNSIFQNKIHTLDPHV
3343	A	1	385	FRVDNSEBWKDVFISSERSFKLDSLKCGTWYKV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KLAAKNSVSGRRISEIEAKTHGREPSFSKDQHLF THINSTHARLNLQGWNGGCPITAVLEYRPKG T WAWQGLRANSSGEVFLTELREATWY
3344	A	351	147	SPACITSSLSQHIADPRAAPTEVKVRVMNSTAISL QWNRVYSDTVQGQLREYVRKPADSPNYP AH
3345	A	351	147	SPACITSSLSQHIADPRAAPTEVKVRVMNSTAISL QWNRVYSDTVQGQLREYVRKPADSPNYP AH
3346	A	3	1509	AGIRHEAPPTTSNRHRRQIDRGVTHLNISGLKMP RGIADWVAGNVYWDSDGRDVIEVAQMKGENR KTLISGMIDEPHAIIVVDPLRGTMVWSDWGNHPK IETAAMDGTLRETLVQDNIQWPTGLAVDYHNER LYWADAKLSVIGSIRLNGTDPIVAADSKRGLSHP FSIDVFEDYTYGVTYINNRVFKHKFGHSPLVNL T GGLSHASDVVLVYHQHKQPEVTNPCKRCKCEWL CLLSPSGPVCCTCPNGKRLDNGTCVVPVSPPTPPD APRPGTCNLQCFNGGSCFLNARRQPKCRCQPRY TGDKCELDQCWEHCRNGGTCAASPSGMPCTCRCP TGFTGPKCTQQVCAGYCANNSTCTVNQGNQPQ CRCLPGFLGDRQCQYRQCSGYCENFGTCQMAAD GSRQCRCTAYFEGSRCEVNKCSRCLEGACVVNK QSGDVTNCNCTDGRVAPSCCLTCVGHCSNGGSCTM NSKMMPECQCPPHMTGPRCEEHVFSQQPGHIA SILIP
3347	A	974	666	SPEMESHPIQAGVQWHHLSSLQPLPPGPK*FSCF SLPE*LG YRHVPPCLANSVFSVEMGFLHVGQAG LELLTSGDLPALASQSAGITGSHRARPENG FENIF
3348	A	1	1171	LSKITMPVICNEPLSFIQRLTEYM*HTYFIHRPSSL SDPVD RMQCVAFAVSAVASQWERTGKPFNPLL GETYELVRDDLGFRLISEQVSHPPISAFHAEGLN NDFIFHGSIYPKLFKFWGKSVEABPKGTITLELLEH MEAYTWETTCVLENTGWIWIEQYGVVETNI KTGDKOVLNKKPCGLFGKELHKVEGYIQDKSKK KLCALYCKWTECLYSVDPATFDAYKKNDKKNT EEKKNSKQMTSEELDEMPVPDSESVFIIPGSVLL WRIAPRPPNSAQMYNFTSFAMVLNEVDKDMESV IPKTDCLRPDIRAMENGEIDQASEEKKRLEEKQ RAARKNRSKSEEDWKTRWFHQGPNPYNGAQD WYSGSYWDRNYFNLPDIY
3349	A	403	497	NFASSSGKYLRTQKIKCLNNKFTPFPTTEKK*SQS VRPP*SNRIY*ILQS*NISFS*LPN*NFASSSGKYL RTQKIKCLNNKFTPFPTTEKK
3350	A	1	712	GAPAQDCICLPFPFHSSFLES DIRKPARRKIQTINP DFLLLLFMSVPVVSAPPFCPPAEGSRDGRPKASV ARPAAVHEHHSRDCGHLDPVIRSSLGGWQPH*P AQPENRLL*LLPVE*GHQHPTVSPVP*AGSPGGAS GWPGPGQAWRVVP GPHPLCPPASPPSPVQQ**E SVAAGSGLPGCVLCAAGRRPGPLLCVEVGQA LPPGA WVSSSGQRPGLTHPLAYSHGCVPSEG
3351	A	1	428	MAAVVAATALKGRGARNARVLRGILAGATANK ASHNRTRALQSHSSPEGKEEPELSPELEYIPRKR GKNPMKAVGLAWAIGFPCGILLFILTKREVDKDR VKQMKARQNMRLSNTGEYESQRFRASSQSAPSP DVGSGVQT
3352	A	2	841	RTLFRGRRRRREDDRISRPHPSTAESKAPT PKFDLL ASNFPPLPGSSSRMPGELVLENRMSDVVKGVYK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EKDNEELTISCPVPADEQTECTSAQQLNMSTSSP CAEELTALSTTQKEKDLIEDSSVQKDGLNQTTIP VSPSTTKPSRASTASPCNNNNINAATAVALQEP KLSYAEVCQKPPKEPSSVLVQPLRELRSNVVSPT KNEDNGAPENSVEKPHEKPEARASKDYSGFRGN IIPRGAAAGKIREQRRQFSHRAIPQGVTRRNGKEQ YVPPRSPK
3353	A	1054	587	IATPTWTAPLTATPTPAHQYGPAPVPNGAPRLEP PPGKRECRVGQYVVDLTSFEQLALPVLNRADCS SGPGQRVCVIDEIGKMELFSQLFIQAVRQTLSTPG THLGTIPVPKGKPLALVEEIRNRKDVVFNVTK NRNHLDPDIVTCVQSSRK
3354	A	56	1268	GMEPVGCCGECRGSSVDPRSTFVLSNLAEVVER VLTFPLPAKALLRVACVCRWLWRECVRRLRTHRS VTWISAGLAEGHLEGHCLVRVVAEELENVRILP HTVLYMADSETFISLEECRGHKRKRRTSMETA LALEKLFPKQCQVLGIVTPGIVVTPMGSGSNRPQ EIEIGESGFALLFPQIEGIKIPFHFKDPKNLT HQLTEVGLLDNPELRVVLVFGYNCKV GASNYL QQVVSTFSDMNILAGGQVDNLSSLTSEKNPLDI DASGVVGLSFSGHRIQSATVLLNEDVSDEKTAEA AMQRLKAANIPEHNTIGFMFACVGRGFQYRAK GNVEADAFRKFFPSVPLFGFFGNGEIGCDRVTG NFILRKCNVKKDDDLFHSYTTIMALHILGSSK
3355	A	1	707	GTSSGLGGDRLAAGPSPSPFYPQGRGERAYDIY SRLRLRERIVCVMGPIDDSVASLVIAQLFLQSESN KKPIHMYINSPGGVVTAGLAITYDTMQYILNPIC WCVGQAASMGSLLAAGTPGMRHSLPNSRIMIH QPSGGARGQATDIAIQAEEMKLLKQLYNTYAKH TKQSLQVIESAMERDRYMSFMEAQEFGLDKVL VIFPQDGEDEPTLVQKEP...APA...AE...PA... FNYNFCRNLS...SLV*PGMCGLLAKHLS...VG AFLIT/LGVAALCKFAVA*PRKKA YADFYRNY... IKEFEVRKANISQSTK
3356	A	552	338	
3357	A	1	403	ALGSCGGLLTGTLKGTMSGTLWSKGIFAGYKR RIRIQREHTAVLKIEGVYARDETEFYLRMICANV YKANNTVTPVLTDPKTRVMWRKVTQAHGISI MVRAQFRTNLPADAIGHRIRMML*PSRMYTTEPS
3358	A	71	2897	FCSKDKCCLYLPDSINRSKCTAKPGAHSQDRHA VMDSERQVKDITDDIESPKRSIRDSGYIDCWDSER SDSLSPPRHGRDSDSDFSGSRSRQTPSPDVVL RGSSDGRGSDSESDLPKRKLPDVKKDDMSARRT SHGEPKSAVPFNQYLPNKSNTAYVPAPLRKKK AEREEYRKSWSTATSPAGLGKKALQDYGPRTPV SDDAESTSMFDMRCEEEAAVQPHSRARQEQLQ LINNQLREEDDKWQDDLARWKSRRKRSVSQDLIK KEBERKKMEKLLAGEDGTSERRKSIKTYREIVQE KERRERELHEAYKNARSQEEAEGILQYYIERFTIS EAVLERLEMPKILERSHSTBPNLSSFLNDPNPMK YLQQSLPPPKFTATVETTIRASVLDTSMSAGS GSPSKTVTPKAVPMLTPKPYSQPKNSQDVLKTFK VDGKVSVNGETVHREEEKERECPVAPAHSLTK SQMFEGVARVHGSPLELKQDNGSIEINIKKPNSV PQELAAATTEKTEPNSEQEDKNDGGKSRKGNIELAS SEPQHFTTTVTRCSPTVAFVEFPSSQLKNDVSEE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KDQKKPENEMSGKVELVLSQKVVPKPSPEPEAT LTFPFLDKMPEANQLHLPNLNSQVDSSESSEKSPV TTPFKFWAWDPEEBERRRQEKWQQEQERLLQER YQKEQDKLKKEEWEKAQKEVEEEERRRYEEEE* ILEDPVVPFTVSSSSADQLSTSSSMTEGSGTMNKI DLGNCQDEKQDRRWKKSFGQDDSDLLKTRES DRLEEKGSLTEGALAHSGNPVSKGVHEDHQLDT EAGAPHCGTNPQLAQDPSQNQTSNPTHSSDEV KPKTLPLDKSINHQIESPSERRKSISGKKLCSSCGL PLGKGAAMIETLNLVYFHIQCFRCGICKGQLGDA VSGTDVRIIRNGLLNCNDCYMRSRSAGQPTTL
3359	A	3	368	EVTASREGRGACAWECGSSRGPWGLLRGTFAFV RAATP*S*LPKGSRLRHP*/CPPPVHLPKSSCPPR AWAGRATSM*TSSYSSEYQPQT*ALVTLPPrSY YLLTHLLTLTHLHHQLFEP
3360	A	2	392	ARGISLGRDHSAGSGGGTGMAGAWVRKAADYV RSKDFRDYLMSTHFWGPVANWGLPIAAITDMK\I KSPEIISRRMTFAL*CYSLTFVRFAHYVQPNWNL MLGCHTAVDFDQLISSMPCISHGMTASASAL
3361	A	4619	532	LLLGRANSPPYNSVVRTLPATLLLRAGWESF WSCQSRSPWPPRPEVRAPAKGPRGVAGAAGACS AGARLGDAAGGDPASGQAARGCGARAPRGLGR TARARDTAMEDAGAAGPGPEPEPEPEPEPEPEPE PEPEPKPGAGTSEAFSRLWTDVMGILDGSLGNID DLAQQYADYYNTCFSDVCERMEELRKRRVSQD LEVEKPDASPTSLQLRSQIEESLGFCSAVSTPEVE RKNPLHKSNSDSSVGKGDWKKKNKYFWQNF KNQKGIMRQTSKGEDVGYVASEITMSDEERIQ MMMVKEKMITIEBALARLKEYEAQHRQSAALDP ADWPDGSYPTFDGSSNCNSREQSDDTEESVKF KRLHKLNVNSTRVRKKIIRVEEMKKPSTAEF HVTENSPVLDERSALYSGVHIFDGSPEKPP EDDSDSLTTPSSSSLDTWAGRKLVKTFKSGES RGLIKPPKMGTTFFSYPEEKAQKVSRLTEGEM KKGLGSLSHGRTCSFGGFDLTNRSLHVGSNNSDP MGKEGDFVYKEVIKSPTASRISLGKKVKS VKET MRKRMSKKYSSSVSEQDGLDGMGSPPPSPQD PEHLDPKPKLKAGGSVESLSSLSGQSSMSGQTVS TTDSSSTNRESVKSEDGDDEPPYRGPFCGRARV HTDFTSPYDTDLSLKLKGDIIHISKPPMGTWMG LLNNKVGTFNFYVDVLSDEEEKPKRPTRRRRK GRPPQPKSVEDLLDRINLKEHMPITFLNGYEDLD TFKLLEEDLDELNIRDPEHRADLLTAVELLQEY DSNSDQSGSQEKLLVDSQGLSGCSPRDS*CYESS ENLENGKTRKASLLSAKSSTEPSLKAFSRNQLGN YPTLPLMKSGDALKQGQEEGRLGGGLAPDTSKS CDPGPG*LVLNKNRRKPPSPFSCRCAETLAEQP TVDTWPRSHSLDDLQVEPGAQDVPTVTEPPPPQ IVEVPQKTTASSTKAQPLEQDSAVDNALLTQS KRFSEPKLTTKKLEGSIAASGRGLSPFQCLPRNY DAQPPGAKHGLARTPLEGHRKGHEFEGTHPLG TKEGVDAEQRMQPKIPSQPPVPKKSRRERLANG LHPVPMGPGSALPSDAPCLPVKRGSPASPTSPSD CPPALAPRPLSGQALGSPSTRPPPWSELPEPNTS LQEHGVKLGALTRKVSARGVDLETLTENKLA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				HAEGIRSSRREPYS*LRHGRCGAPAEALVQRYAED LDQPERDVAANMDQIRVKQLRKQHRMAIPSGGL TEICRKPVSPGCIS\SVSDWLISIGLPMYAGTLSTA GFSTL\SQVPSLSHTCLQEAG\TEERHIRKLLSAA RLFKLPPGPEAM
3362	A	1	4653	FRGGVGYAHTLHLLPFAGSSVVLARARRTDRWT SGLVEMATLSLTVNSGDPPLGALLAVEHVKDDV SISVEEGKENILHVSENVIFTDVNSILRYLARVAT TAGLYGSNLMHETEIDHWLEFSATKLSSCDSFTS TINELNHCLSLRITYLVGNSLSLADLCVWATLKG NAAWQEQLKQKKAPVHVKRWFGFLEAQQAFQS VGTKWDVSTTKARVAPEKKQDVGKFVELPGAE MGKVTVRFPPEASGYLHIGHAKAALLNQHYQV NFKGKLIMRFDDTNPEKEKEDEFEKVILEDVAML HIKPDQFTYTSDFETIMKYAEKLIQEGKAYVDD TPGEQIKAEREQRIESKHKRKNPIEKNLQMWEEMK KGSQFGHSCCLRAKIDMSSNNGCMRDPTLYRCK IQPHPTGN*YNNVYPTYDFACPIVDSIEGVTHAL RTTEYHDRDEQFYWIEALGIRKPYIWEYSRLNL NNTVLSKRKLTWVFNELVDGWDDPRFPTVRG VLRRGMTVEGLKQFIAAQGSSRSVNMWEDKI WAFNKKVIDPVAPRYVALLKKEVIPVNVPEAQE EMKEVAKHPKNPEVGLKPVWYSPKVFIGADAE TFSEGEMVTFINWGNLNTIKHKNADGKIISLDAK LNLENKDYKTTKVTWLAETTHALPIPVICVITYE HLITKPVLGKDEDFKQYVNKNSKHEELMLGDPC LKDLKKGDIIQLQRRGFFICDQPYEPVSPYSCKEA PCVLIYPDGHTKEMPTSGSKEKTKVEATKNETS APFKERPTPSLNNNCTTSEDSLVLYNRVAVQGD VVRELKAKKAPKEDVDAAVKQLLSLKAEBYKEK TFCQYKPCNPFAETGONISSNSASILFSKSLYDE VAAQGEVVRKLKAEKSPKAKINEAVECLL QYKEKTGKEYIPGPPLSQSSDSSPTRNSEPAGLE TBAKVLFDKVASQGEVVRKLKTEKAPKDQVDI AVQELLQLKAQYKSLIGVEYKPVSATGAEDKDK KKKEKENKSEKQNKPKQNDGQRKDPSKNQGG GLSSSGAGEGQGPKKQTRLGLEAKKEENLADW YSQVITKSEMIEYHDISGCYILRPWAYAIWEAIKD FFDAEIKKLGVENCYFPMFVSQSALEKEKTHVA DFAPEVAWVTRSGKTELAEPPIAIRPTSETVMYPA YAKWVQSHRDLPIKLNQWCNVVRWEFKHPQPF LRTREFLWQEGHSFATMEEAAEEVLQILDLYA QVYEELLAIPVVKGRKTEKEKFAGGDYTTTIEAF ISASGRAIQGGTSHHLGQNFSKMFEIVFEDPKIPG EKQFAYQNSWGLTTRTIGVMTMVHGDNMGLVL PPRVACVQVVIIPCGITNALSEEDKEALIAKNDY RRRLSVNIRVRADLRDNYSPGWKFNHWELKG VPIRLEVGPDMKSCQFVAVRRDTGEKLTVAEN EAETKLQAILEDIQVTLFTRASEDLKTHMVVANT MEDFQKILDSGKIVQIPFCGEIDCEDWIKKTTARD QDLEPGAPSMGAKSLCIPFKPLCELQPGAKCVCG KNPAKYTYTLFGRSY
3363	A	3797	1514	LGGAAPETMPFPVTTQGSQQTQPPQKHYGITSPIS LAAPKETDCVLTQKLVETLKPFGGFLKKEEGTA SRRNFNFQGN*INLVKEWIRRNQ*KAKNLPQSVI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ENVVGGKIFT/FLGSYRL/GEVHTKGADIDGVCVF APRHVDRSDFFTSFYDKLKLQEEVKDLRAVEEA FVPVIKLCFDGIEIDILFARLALQTIPEDLDLRDSS LLKNLDIRCIRSLNGCRVTDEILHLVPNIDNFRLT LRAIKLWAKRHNTYSNIGFLGGVSWAMLVART CQLYPNAIASTLVHKFFLVFSKWEWPNPVLKQP EECNLNLVWDPRVNPSPDRYHLMPIITPAYPQQN STYNVSVSTRMVMVEEFKQGLAITDEILLSKAE WSKLFEAPNFFQKYKHYTVLLASAPTENQRLEW VGLVESKIRILVGSLEKNEFITLAHVNPQSFPAPK ENPDKEEFRTMWVIGLVFKKTENSENLSVDLTY DIQSFTDVTYRQAINSKMFEVDMKIAAMHVKKR QLHQLLPNHVLQKKKKHSTEGVKLTALNDSSLD LSMDSNDMSVSPSPTSATKTSPLNSSGSSQGRNS PAPAVTAASVTNIQATEVSVPVNSSESSGGTSSE SIPQTATQPAISPPPPTVSRVVSSTRLVNPPPRSS GNAATSGNAATKIPTPIVGVKRTSSPHKEESPCK TKTEEDETSEDANCLALS GHDKTEAKEQLDTETS TTQSETIQTAASLLASQKTSSDLS DIPALPANPIP VIKNSIKLRLNR
3364	A	54	3073	SARTMSYDYHQNWGRDGGPRSSGGGYGGGPAG GHGGNRGSGGGGGGGGGGGG/WQGPASRAPER PRNRHVREKTGAEEQ/WKRRGKREL/LVHMD RREEQIVQLLSVQAKNDKESEAQISWFAPEDHG YGTEVSTKNTPCSENKLDIQEKKLINQEKMFRI RNRSYIDRDSEYLLQENEPDGTLDQKLEDLQKK KNDLRYIEMQHFREKLPSYGMQKELVNLIDNHQ VTVISGETGCGKTTQVTQFILDNYIERGKGSACRI VCTQPRRISAJSAERVAERAESCGSGNSTGYQI RLQSRIPRKQGSILYCTTGILQWLQSDPYLSSVS HVLDEIHERNQCSDVLMTVVKELNFT3DLKVY LMSATLNASKFSSTGNCPMIHIFGTFPVVEYLL EDVIEKIRYVPOQKEHRCQFKRGFMQGHVNSQE KEEKEATYKEKWPDYVRELRRRYSASTVDVIEM MEDDKVDLNLIVALIRYIVLEEDGAILVFLPGW DNISTLHDLMSQVMFKSDKFLIPLHSLMPTVN QTQVFKRTPPGVRKIVATNIAETSIDDDVYVID GGKIKETHFDTQNNISTMSAEWVSKANAKQRKG RAGRVPQGSLLFICINGS*EASLLGWTIQLPEIF/R GTPLEELCLQIKVLRLGGI/GLFLSRLMDPPSNEA VLLSIRQLARSLNALDKQEELTPLGVHLARLPVEP HIGKMILFGALFCCLDPVLTIAASLSFKDPFVPLG KEKIADARRKELAKDTRSDHLTVVNAFEGWEEA RRRGFRYEKDYCWEYFLSSNTLQMLHNMKGQF AEHLLGAGFVSSRNPKDPESNINSNEKIIKAVIC AGLYPKVAKIRLNLGKKRKMVKVYTKTDGLVA VHPKSVNVEQTDHFYNWLIYHLKMRTSSIYLYD CTEVSPYCLLFFGGDISIQKDNDQETIAVDEWIVF QSPARIAHLVKRAVVHMDERREEQIVQLLSVQ AKNDKESEAQISWFAPEDHGYDKKYFFKE
3365	A	439	878	ECCNVRLPLRETDLKMKRKPRASSPVVEEQPRA NTKETRKKKSFSQPMASSTKEESQDGRRKGL KGRARKKNAPQKSMALRIEESRPTPSGHSDQL NEEL*QNELQLEQ/PEGT*LEQQSEGTQPEQQSGR MPTISTLSLSE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
3366	A	1	827	FRGYWGVREAFTDASWSGGLGPGKPGMKITRQ KHAKKHLGFFRNNGFVREPYQILLDGTFCQAAL RGRIQLREQLPRLMGETQLCTTRCVLKELETLG KDLYGAKLIAQKCQVRNCPHFKNVSGSECLLS MVEEGNPHHYFVATQDQNLVSVKVKKKPGVPLM FIQNTMVLDPSPKTIKFVKA VESGRLSQCMRK KVSNIKSRNRV**KTLNRGRRKKRKKISGPNPLS CLKKKKKAPDTQSSASEKKRKRKRIRNRSNPKV LSEKQNAEGE
3367	A	40	1467	MLWGCRAKACWGPRLSDLVASHSPQRECISVHV GQAGVQIGNACWELFCLEHGIQADGTFDAQASK INDDDSFTTFFSETGNGKHVPRAVMIDLEPTVVD EVRAGTYRQLFHPEQLITGKEDAANNYARGHYT VGKESIDLVLDRIRKLTACSGLQGLIFHSFGGG TGSFTSLLMERLSLDYGKSKLEFAIYPAPQVS TAVVEPYNSILTTHTTLEHSDCAFMDNEAIYDI CRRNLDIERPTYTNLNRLLISQIVSSITASLRFDGAL NVDLTFEQTNLVYPRIHFPLVTYAPIISAKEYH EQLSVAEITSSCFEPNSQMVKCDPRHGKYMACE MLYRGDVVPKDVNVAIAAIKTKRTIQFVDWCPT GFKVGINYQPPTVVPGGDLAKVQRAVCMLSNNT AIAEAWARLDHKFDLMAKRAVHVWYVGEEM EEGFS*RPGEDLAIALEKDYEEVGTDSFEEENE GEEF
3368	A	3	2597	SLLEETMDESSRLREYTVSLDSDMDDASKCLQE YDSGTGNTREALRPCPRTVSTKAQPGRSASSSSG DKTTSFABEQKIRKLNHTDGESSGSSSQKTTPGSE LNIPHAGAWAQIPEETGLPQGRDITQLLASEMV HLMMLKEKRRAI*AQKKKMEAAFTKQRQKM GRTAFLTVVKKKG DGISPLREEAAGAEDKVVY DRAKEKLAQKTGGRSKGALLES MENFOAET LKSPTTFDPEKQGNLASFSEETLNEGEILEYTKSI EKLNSSLHFLQEQMQRSLQDEMMLMQMREQQS WVISPPQSPQKQIRDFKPSKQAGSSAIAFPSSD SPR\PHPSSTSLNRKSASFVKSQRTPRPNELKI TPLNRTLTPPRSVDLPLRLRRFSPSQVPIQTRSFVC FGDDGEPQLKESKPKBEEVKKELESKGTLEQRG HNPEKEIKPFESTVSEVLSLPTVETVCLTPNEDQ LNQPTPPPKPVFPPTAPKNVNLIEVSLSDLKPE KADVPVEKYDGEDKEQFDDDDQVCCGFFFKD DQKAENDMAMKRAALLEKRLRREKETQLRKQQ LEAEMEKKKEETRRKTEBERQKKEDEARRERFIR QEYMRRKQLKLMEDMDTVIKPRPQVVKQKKQR PKSIHRDHIESPKTPIKGPPVSSLASLNTGDNES VHSGKRTPRSESVGEFLSPSRCSGRNGEKDWEN ASTTSSVASGTEYTGPKLYKEPSAKSNKHIIQNAL AHCCLAGKVNEGQKKKILEMEKSDANNFLILF RDSGCQFRSLYTYCPETEEINKLTGIGPKSITKKM IEGLYKYNSDRKQFSHIPAKTLSASVDAITISHSL WQTKRPVTPKKLLPTKA
3369	A	977	594	RSGSLTQEPGSGQLALACAEGAVEWLYPAGAL RLTLGGPDPRARPGIACLRPVRPFAQAQVFAERA GGALELLLAEGPGPAGGRCVRWGPRERRALFLQ ATPHQDISRRVAARFELREDGRPEIAP
3370	A	345	1383	DLSLECTGFKETNLGVYFLSSKWVLRLYALHIID

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				YSAVLFPC*AMDHLESFIAECDRRTELAKKRLAE TQEEISAEVSAKAQKVHLENEEIGKLLAKAEQLG AEGNVDESQKILMEVEKVRAKKKEAEKTVAEK QEKRNQDRLRRREEREREERLSRRSGSRTDRRR SRSRDRRRRRSRSTSRERRKLSRSRSDRHRHR SRSRSHSRGHRRASRDRSAKYKFSRERASREESW ESGRSERGPPDWRLESSNGKMASRRSEEKEAG/G DLLNRMIVWKHGLLI
3371	A	345	1383	DLSELECTGFKETNLGVYFLSSKWVLRLYALHIID YSAVLFPC*AMDHLESFIAECDRRTELAKKRLAE TQEEISAEVSAKAQKVHLENEEIGKLLAKAEQLG AEGNVDESQKILMEVEKVRAKKKEAEKTVAEK QEKRNQDRLRRREEREREERLSRRSGSRTDRRR SRSRDRRRRRSRSTSRERRKLSRSRSDRHRHR SRSRSHSRGHRRASRDRSAKYKFSRERASREESW ESGRSERGPPDWRLESSNGKMASRRSEEKEAG/G DLLNRMIVWKHGLLI
3372	A	239	3348	PMQNCMCSTLSVLPLGPQPPVPEKRPPEIQHFR MSDDVHSLGKVTSDLAKRRKLTST*GGLSEELGS ARRSGEVTLTKGDPGSLLEEWETVVGDDFSLYYD SYSVDERVSDSKSEVEALTEQLSEEEEEEEEEEE EEEEEEEEEEEEDEESGNQSDRGSSGRRKAKK KWRKDSPWVKPSRKRKRREPPRAKEPRGVNGV GSSGPSEYMEVPLGSLELPSEGTLSPNHAGVSN TSSLETERGFEELPLCSCRMEAPKIDRISERAGHK CMATESVDGELSGCNAAILKRETMRPSSRVALM VLCETHRARMVKHHCCPGCGYFCTAGTFLECHP DFRVAHRFHAKCVSQLNGMVFCPHCGEDASEA QEVTPRGDGVTPPAGTAAPAPPLSQDVPGRAD TSQPSARMRGHGEPRRPPCDPLADTIDSSGSLTL FNGGCLSAVGLPLGPGLEKAL*QESPE LRFEPRL*QKQGLQKQVILMLLDNLN* DQQSKRTPLHAAQKGSVEICHVLLQAGANNA VDKQQRTPLMBAVNNHLEVARVMVQRGGCV YSKEEDGSTCLHHAAKIGNLEMVSLLLSTGQVD VNAQDSGGWTPITWAAEHKHIEVIRMLLTRGAD VTLDNEENICLHWASFTGSAALAEVLLNARCDL HAVNYHGDTPHIAARESYPHDCVLLFLSRGANP ELRNKEGDTAWDLTPERSDVWFALQLNRKLRL GVGNRAIRTEKICRDVARGYENVPIPCVNGVDG EPCPEDYKYISENCETSTMNIDRNITHLQHCTCV DDCSSNCLCGQLSIRCWYDKDGRLLQEFNKIEP PLIFECNQACSCWRNCKNRVVQSGIKVRLQLYR TAKMGWGVRLQTIPQGTICEYVVELISDAEAD VREDDSYLFDLDNKDGEVYCIDARYYGNISRFIN HLCDPNIIPVRVFMHLHQDLRFPRIAFFSSRDITGE ELGFDYGDWFWDIKSKYFTCCQCGSEKCKHSAEAI ALEQSRRLARLDHPPELLPELGSPPVNT
3373	A	587	1584	PDGRLIVSCSEDKTIKIWDTTNKQCVNNFSDSVG FANFVDFNPSGTCLASAGSDQTVKVDVVRVNL LQHYQVHSGGVNCSIFHPSGNYLITASSDGTILKIL DLLKGRLIYTLQGHTGPVFTVSFSKGGELFASGG ADTQVLLWRTNFDELHCKGLTKRNLKRLHFDSP PHLLDIYPRTPHPHEEKVETVEDFFLHLRLIQSL R*SICRSLPLWISFLLLPQQQKPVVGLCQTRV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KRPVDIS*TLP*CHQNVCCQPRKRKQKT*VTSPV KVK/VSIPLA VTDALHIMEQLNVLTQT VSVILEQR LTLTEDKLKDCLNQKLFSAVQQKS
3374	A	398	21	WLYPMALSILDIKMSPSWYFHMAIGIINWNTTAG LSGTL YPKVPQKYLFD SVILLGLMLRKIRQVCQ NVYMKGCSPITL FKIVHYWPGA VAHAYNPSTLG GQVG/WQIT*GOEFETSLDYMVKPHLY
3375	A	3	1051	VPTQQILAFPEQNTTKDWTVTPEHVLPEQSLLT FEEVAMYFSQEEWELLDP TQKALYNDVMQENY ETVISLALFVLPKPKVISCLEQGEWPVQVSPEFK DSAGKSPTGLKLKNDTENHQPVSLSDEIQASAG VISKKAKVKVPQKTAGKENHFD MHRVGKWHQ DFPVKKRKKLSTWKQELLKLM DRHKKDCAREK PFKQCQECGKTFRVSSDLVKHQRIHTEEKPYKCC QCDKRFRWSSDLNKLHTTHQG IKPYKCSWGGKS FSQNTNLHTHQRTHTGEKPF TCHCEGKKFSQNS HLIKHRRTHTGEQPYTCSICRRNFSRRSSLLRHQK LHL*REACPVS HFWKTF
3376	A	137	2329	SFESPAPLPSTCFPQERQDPGPCYVSGAMAGLGP GVGDSEGGPRPLFCRK GALRQKVVEVKSHKFT ARFFKQPTFC SHCTDFIWGIGKQGLQCQVCSFV HRRCHEFVTFECPGAGKGPQTD DPRNKHKFR LH SYSSPTFC DHCGSLLYGLVHQGMKCSCEMN V RRCVRSVPSLCGV DHTERRGR LQLEIRAPTADI HVTVGEARNLIPMDPNGLSDPYVKLKLIPDPRNL TKQKTRTVKATLNPVWNETFVFNLP GDVERRL SVEVWDWDRTSRND FMGAMSFGVSELLKAPVD GWYKLLNQEEGEYYNVPVADADNCSLLQKFEA CNYPLELYERVRMG PSSPIPSPSPTDPKRCFFG ASPGRLHISDFSFLMVLGKSGFKVMLAERRGSD ELYAINKLKEWITQDDVDCTLLEKRYVLLGG RGPGRPHFLTQLHSTFQTPDRLYFVMVVTGG DLMYHIQQLGKFKEPHAAFYAAEIAIGLFFLHNQ BIYRDLKLDNVMLDAEGHIKITDFGMCKENVFP GTTTRTFCGTPDYIAPBIIAYQPYGKSVDWWSFG VLLYEMLAGQPPFDGEDEEELFQAIMEQTVTYP KSLSREAVAICKGFLT KHPGEAPGASGP*WGNLT IRAHGFFPLGFDWERLERLAEIPASF SRPRPCGPQR RGIFDKFFTRAAPALT PPARLVLD SIDQADFQGF TYVNPDFVQPDARSPTSTVHVPVM
3377	A	918	738	SSMLWGFSVFRRSWILNCWLSSSQVGISAACKFS TLTHTHTHTHTRHAPFCGTCLYY
3378	A	1126	456	FSKLIMKTFIIGISGV TNSGKTTLAKNLQKHL PNC SVISQDDFFKPESEIETDKNGFLQYDVLEALNME KMMSAISCWESARHSV VSTDQESAEEIPILIEG FLLFNYKPLDTTWNRSYFLTIPYEECKRRRSTRVY QPPDSPGYFDGHVWPMY LKYRQEMQDITWEVV YLDGTKSEEDLFLQVYEDLIQELAKQKCLQVTA* RRNTINPS/CK*IRKLQGV I
3379	A	1126	456	FSKLIMKTFIIGISGV TNSGKTTLAKNLQKHL PNC SVISQDDFFKPESEIETDKNGFLQYDVLEALNME KMMSAISCWESARHSV VSTDQESAEEIPILIEG FLLFNYKPLDTTWNRSYFLTIPYEECKRRRSTRVY QPPDSPGYFDGHVWPMY LKYRQEMQDITWEVV YLDGTKSEEDLFLQVYEDLIQELAKQKCLQVTA*

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RRNTTNPS/CK*IRKLQGV
3380	A	1443	794	ARRGELAGGGGRASGGSGDGGGGGGGARAP VRAPAAGQPRATKGAPPPGTPPPSPMSSA LDPSEEPVDEVLLQIPPSLLTCGGCQQNIG AIDQYWHEDCLSCDLCGCRLEVGRRLYYKL KLCRRDYLRFGQDGLCASC DKRIRAYEMT KDKVYHLECFKCAACQKHFCVGDYLLINS CEQDIYEWTKINGMI
3381	A	945	474	SLKLRKPPLPTDGVHVFVFSQLDFWGPQ QQGMALQNYDNKL VKCIEELCQKQELCW QEEDEKQRLQNEVRQLTEKLACVNEKLAR NLARKIASCSKFYQTIAETATYLKILESF* VRLS VRKREAGNLTKATAPDQKSSGG RDS
3382	A	1	1458	GIRGKMADRGGVGEAAA VGASPASVPGLN PTLG WRERLRAGLAGTGASLWFVAGLGL LYALRIPLR LCENLAAVTVFLNSLTPKFY VALTGTSSLISGLIFI FEWWYFHKHGT SFIEQVSVSHLQPLMGGTSSIS EPGSP SRNRENETSRLNLSECKVWRNPLNFRGA EYRRYTWTGKEPLTYIDMNL SAQDHQT FTFC DTDFLRPSDTVMQKAWRERNPPA RIKAA YQALE LNE* LCHCICSTG*GRS NNYCRC*KVI* TGTQGR RNNL*AVTA VPAPKSSA*SSTEERYQCTGIY*LKI GNVCKKIRKNKRSSKNNERFDE*ISSYH VEHP* KSL\KSLELQAYPDVQAVLAKY DDISLPKSAAIC YTAALLKTRTVSEKFS PETASTRGLSAAEINAVD AIHRAVEFN PHVPKYLLEMKSLILPPEHILKRGDS EAIA YAFFHLQHWKRIEGALNLLQCTW EGSKYS FPKVTLISLTIH
3383	A	282	2443	RGKGFKEFFLGVCQTFIPCLCAEGIQLO FFCSGSG SSPLLKDLESMTGLFFLCULG TAAIPTNARLLS DHSKPTAETVADNTAIP P...TE...KE... TEDDSHHKAI...SVLKSKEESHEQSAEQ G...S...S... QELGIEGFKRDSGSL*VWNLA EYGTNLK GTL... KEDMSEPQEKKLSNTDFLAPGVSSFTDS NQES ITKREENQEQRNYSHHQLNRSSK HSQGLRDQG NQEQDPNISNGEEEEKEP GEVGTNDNQRKTE \PREHANSKQEE DNTQSDDILEESDQPTQVSKM QEDEF DQGNQE QEDNSNAEMEEENASNVNKH IQ ETEWQSQEGKTGLEAISNHKETEEK TVSEALLME PTDDGNTTPRNHGVDDD GDDGDDGTDGPRH SA\SDDYFHPKPL FWEAERA\HSIA YSPSKLREQ REKV HENENIGTTEPGEHQEAKKAENSSNEE TS SEGNMR\VHAVDSCMSFQCKRGHIC KADQQGKT SLVSCQDPVTCPTKPLDQV CGTDNQTYASSCH LPATKCRLEGTKK GHQLQLDYFGASKSIPTCRD FEVIQF PLRMRDWLKNILMQLYEANSEHAGYL NEK\QRNKVKKIYLADEKRLLAGDHPID LLLRDFK KNYHMYVYPVHWQFSELDQ HPMDRVLTHSELA PLRASLVPMEHCIT RFFEECDPNKDKHITLKEWG HCFGIKE EDIDENLLF
3384	A	3166	928	PSRPHPTHAAMAGPEGFOYRALYFRRER PEDLE LLPGDVLVVSRAALQALGVAEG GERCPQSVGW MPGLNERTRQRGDFGT YVEFLGPVALARPGPR PRGPRPLPAR PRDGAPEPGLTLPDLPEQFSPPDVA PPLL VKLVEAIERTGLDSESHYRPEL PAPRTDWSL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SDVDQWDTAALADGKISFLLALPAPLVTPESAE ARRALREAAGPVGPAPLEPPTLPLHRLTLRFLQ HLGRVASRAPALGPAVRALGATFGPLLRAPPPP SSPPPGGAPDGESESPDFPALLVEKLLQEHLEEQE VAPPALPPKPPKAK\ASTVPGPNNGSPPSLAQDA EWYWGDISREEVNEKL RDPDGTFLVRDASSKI QGEYTLTLRKGGNKLIKVFHRDGHYGFSEPLTF CSVVDLINHYRHESLAQYNAKLDTRLLYPVSKY QQDQIVKEDSVEAVGAQLKVYHQYQDKSREY DQLYEEYTRTSQELQMKRTAIEAFNETIKIFEEQG QTQEKCSKEYLERFRREGN/QTKEMQRILLNSER LKSRIA\EIHSRTKL\EQQLLVPRASDNKRD/IDK PH*TSCLKPDLMLRKRIRDQYLVWLTQKGARQKK INEWLGKINETEDQYALMEDEDDLPHHEERTWY VGKINRTQAEEMLSGKRDGTFLIRESSQRGCYAC SVVVDGDTKHCVIYRTATGFGFAEPYNLYGSLK ELVLHYQHASLVQHNDALTVTLAHPVRAPGPGP PPAAR
3385	A	43	2372	TRDVNSWKELCFNHYNKETTNCYRTTRKWTNY KIIFLGPFRRLRSQGNQVILNLGKERCQLRETGLK LYLPGMDSARHHISHSTSAGPIPSQKEEEMTESQ GTVTFKDVAIDFTQEEWKRLDPAQRKLYRNVML *NYYNLITVGYPTKPDVIFKLEQBEKPWVMEEE VLRHWQGEIHWGVDEHQKNQDRLLRQVEVKFQ KTLTEKGNECQKKFANVFPLNSDFFPSRHNLYE YDLFGKCLEHNFDCNNVVKCLMRKEHCEYNP VKS YGNSSSHFVITPFCNHCGKGFNQTLDLIRH LRIHTGEKPYECSNCRKAFSHKEKLIKHYKIHSRE QSYKCNCEGKAFIKMSNLIRHQRIHTGEKPYACK ECEKSFSQKSNLIDHEKIHTGEKPYECNECGKAFS QKQSLIAHQKVHTGEKPYACNECGKAFPRIASLA LHMRSHTEKPYKCDKCGKAFS QFSMLIIVRIH TGEKPYECNECGKAFS QSSALTVMRSHTGEKP YECKECKRKA FSHKKNFITHQKIHTREKPYECNEC GKAFIQMSNLVRHQRIHTGEKPYICECKGKAFSQ KSNLIAHEKIHSGEKPYECNECGKAFSQKQNFIT HQQVHTGEKPYDCNECGKAFSQIASLTLHLRSHT GEKPYECDKCGKAFSQCSLLNLHMRSHTEKPY VCNECGKAFSQRTFLIVHMRGHTGEKPYECNEC GKAFSQSSSLTIHRGHTGEKPYECKECKRKA FSHK KNFITHQKIHTRE/KPFCNHCGKGFNQTLDLIRH LRIHTGEKPYECSNCRKAFSHKEKLIKHYKIHSRE QSYKCNCEGKAFIKMSNLIRHQRIHTGEKPYACK ECEKSFSQKSNLIDHEKIHTGEKPYECNECGKAFS QKQSLIAHQKVHTGEKPYACNECGKAFPRIASLA LHMRSHTEKPYKCDKCGKAFS QFSMLIIVRIH TGEKPYECNECGKAFS QSSALTVMRSHTGEKP YECKECKRKA FSHKKNFITHQKIHTREKPYECNEC GKAFIQMSNLVRHQRIHTGEKPYICECKGKAFSQ KSNLIAHEKIHSGEKPYECNECGKAFSQKQNFIT HQQVHTGEKPYDCNECGKAFSQIASLTLHLRSHT GEKPYECDKCGKAFSQCSLLNLHMRSHTEKPY VCNECGKAFSQRTFLIVHMRGHTGEKPYECNEC GKAFSQSSSLTIHRGHTGEKPYECKECKRKA FSHK KNFITHQKIHTRENPLSVIIEKASIRLWTSSDI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
3386	A	201	1032	WDDYPQGALRRREAEGHLFLGPPGRVRGQLR GITGPAWYCHSPSHSLLSAFCHLPTPSRCPAMAR PPVPGSVVVPNWHE/RRGQGVPLHSAQEPPAG VWAA*ASAAAA\LSIDTASYKIFVSGKSGVGKT ALVAKLAGLEVPPVHHETTGIQTTVFWPAKLQ ASSRVVMFRFEFWDCGESALKKFDHMLLACME NTDAFLFLFSFTDRASFEDLPGQLARIAGEAPGV VRMVIGSKFDQYMHTDVPERDLTAFRQAWELPL LRVKSVPGRRLG
3387	A	86	96	GSSPDPA SLITMKNQDKKNGAAKQSNPKSSPFGQP EAGPEGAQERPSQAAPAVEAEGPGSSQAPRKPEG AQARTAQSGALRDVSEELSRQLEDILSTYCVDDNN QGGPGEDGAQGEPAEPEDA EKSRITYVARNGEPE PTPVVNGEKEPSKGD/PNTEEIRQSDEVGDRDHRR PQEKKKAKGLGKEITLLMQTLNLTSTPEEKLAAL CKKYAELLEHRNSQKQMKLLQKKQSQLVQEK DHLRGEHSKAVLARSKLESLCRELQRHNRSLKE EGVQRAREEEEEKRKEVTSHFQVTLNDIQLQMEQ HNERN SKLRQENMELAE RLKKLIEQYELREEHID KVFKHKDLQQQLVDAKLQQAQEMLKEAEERHQ REKDFLLKEAVESQRMCELMKQOETHLKQOLA LYTEKFEFQNTLSKSSEVFTTFKQEMEKMTKKI KKLEKETTMYSRWESSNKALLEMAEBEKTVRD KELEGLQVKIQRLKLCRALQT/GAQ*PVRGQRW GSHRTSAVRIFS
3388	A	98	3197	ARPEVPAPPAWLSRRGA AKMGDKKDDKDSPPK NKGKERRDLDDLKKEVAMTEHKMSVEEVC RKY NTDCVQGLTHSKAQEILARDGPNALTPPPTTPEW VKFCRQLFGGSILLWIGAILCFLAYGIQAGTEDD PSCDNL YLGIVLAAVVIITGCFSY YQEA KSSKIME SFKASFTQALVIPFGHLLCVNALV V CCLV EIKGGDRVPADLRISAHGCK/DNSSLTGESEPQT RSPDCTHE/NPLKTRNITFSNNFVEGTARGVVVA TGDRTVMGRIATLASGLEV/KTPIAIEIEHFQIT GVAVFLGVSFILSLILGYTWLEAVIFLIGIIVANV PEGLLATVTVCLTLTAKRMARKNCLVKNLEAVE TLGSTSTICSDKTGILTQNRMTVAHMFWDNQIH EADTTEDQSGTSFDKSSHTWVALF*H/LLGFCNR PVFKGGQDNIPVLKRDVAGDASESALLKCIELSS GSVKLMRERNKKVAEIPFNSTNKYQLSIHETEDP NDNRYLLVMKGAPERILDR CSTILLQGKEQPLDE EMKEAFQNAYLELGG LGERVLGFCHYYLP EEQF PKGFAFDCDDVNFTTDNL CFVGLMSMIGPPRAA VPDAVGKCRSAGIKVIMVTGDHPITAKAIAKGV GIIFEGNETVEDIAARLNIPVSQVNP RDAKACVIH GTDLDFTSEQIDEILQNHTETVFARTSPQQLIIV EGCQRQGAIVAVTGDGVNDSPALKKADIGVAM GIAGSDVSKQAADMILLDDNFASIVTGVEEGR LI FDNLKKSIA YTLTSNIPEITPFLLFIMANIPLPLGTI TILCIDLGTDMVPAISLAYEAAESDIMKRQPRNPR TDKLVNERLISMA YGQIGMIQALGGFFSYFVILA ENGFLPGNLVGIRLNWDDRTVNDLEDSYGQQW TYEQRKVVEFTCHTAFFVSIVVVQWADLIICKTR RNSVFQQGMKNKILIFGLFEETALAAFLSYCPGM DVALRMYPLKPSWWFCAPPYSFLIFVYDEIRKLI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LRNPGGWVEKETYY
3389	A	45	5250	VERLLGCRNSKRTWRMLISKNPWRRLLQGISFG MYSAEELKKLSVKISITNPRYLDLGNPSANGLYD LALGPADSKEVCSTCVQDFSNCSGHLGHIELPLT VYNPLLFDKLYLLLRGSCNLCHMLTCPRAVIHLL LCQLRVLEV GALQAVYELERILNRFLEENPDPSA SEIREELEQYTTETIVQNNLLGSQGAHVKNVCESK SKLIALFWKAHMNAKRCPHCKTGRSVVRKEHNS KLITTFPAMVHRTAGQKDSEPLGIEEAQIGKRGY LTPTSAREHLSALWKNEGFFLNLYLFGMDDDDGM ESRFNPSVFFLDFLVVPPSRYPVSRLLGDMFTN GQTVNLQAVMKDVVLRKLLALMAQEQLPEE VATPTTDEEKDSLAIIDRSFLSTLPGQSLIDKLYNI WIRLQSHVNTIVFDESEMDKLMMDKYPGIRQILEK KEGLFRKHMMGKRVDYAARSVICPDMYINTNEI GIPMVFATKLTYQPVTWPWNVQELRQAVINGPN VHPGASMVINEDGSRTALSAVDMTQREAVAKQ LLTPATGAPKPQGTKIVCRHVKNKDILLNRQPT LHRPSIAHRARILPEEKVLRRLHYANCKAYNADF DGDEMNAHFQSELGRAEAYVLACTDQQYLVP KDQGPLAGLIQDHMVSGASMTTRGCFFTREHYM ELVYRGLTDKVGVRVLLSPSILKPFLWTGKQVV STLLNIIPEDHIPLNLSGKAKITGKAWVKETPRSV PGFNPSMCEQVIIEGELLCGVLDKAHYGSSA YGLVHCCYEIYGGETSGKVLTCARLFTAYLQL YRGFTLGVEDILVKPKADVQRQRIIEESTHCGPQ AVRAALNLPEAASYDEVGRGWQDAHLGKDQRD FNMIDLKFKKEEVNHYSNEINKACMPFGLHRQFPE NTLQLMVQSGAKGSTVNTMQISCLLGQIELEGRS TPLMASGKSLPCFEPYEFTPRAGGFVTGRFLTGIK PPEFFFHCMACREGLVDTAVKTDGLGYLQRCTIK HLEGLVVQYDLTVICSDGSVVQFLYGEDGLDIP KTQFLQPKQFPFLASNYEVIMKSQHLHEVLSRAD PKKALHHFRAIKKWQSKHPNTLLRRGAFLSYSQ KIQEAVKALKLESENRRNGR/RPWDS/G/RMLRMW YELDEESRRKYQKAAACDPSPSVWRPDIYFAS VSETFTKVDDYSQEWAAQTEKSYEKSLSLDR LRTLLQLKWQSRSLCEPGEAVGLLAAQSIGEPST QMTLNTFFHAGRGEMNVTLGIPRLREILMVASA NIKTPMMSVPVLNTKKALKRVKSLKKQLTRVCL GEVLQKIDVQESFCMEEKQNKQFQVYQLRFQFLP HAYYQEQECLRPEDILRFMETRFFKLLMESIKKK NNKASAFRNVNTRRATQRDLNAGELGRSRGE QEGDEEEEGHIVDAEAEEGDADASDAKRKEKQE EEVDYSEEEEEEREGEENDDDEDMQBERNPHREG ARKTQEQDEEVGL/GH*GGPVPSRPPDAAPETHP QPGAPGA/EAMERRVQA/VREIHPFIDDYQYDTEE SLWCQVTVKLPLMKINFDMSSLVVS LAHGAVIY ATKGITRCLLNETTNKNEKELVLNTEGINLPELF KYAEVLDLRLYSNDIHALANTYGIEAALRVIEK EIKDVFAVYGIAVDPRHLSLVADYMCFEQVYKP LNRFGIRSNSSPLQQMTFETSQFLKQATMLGSH DELRSPSACL VVGKVVRGGTGLFELKQPLR ILPPLGPPAQASPSSTMLGEGSQPDWPGGSRDYD LDEIDAYWLELINSELKEMERPELDELTLERVLE
3390	A	2	2080	ILPPLGPPAQASPSSTMLGEGSQPDWPGGSRDYD LDEIDAYWLELINSELKEMERPELDELTLERVLE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				ELETLC HQNMARAIETQEG LGIEYDEDV VCDVC RSPEGEDGNEMVFC DKCNVCVHQACYGILKVPT GSWLCRTCALGVQPKCLLCPKRGGALKPTRSGT KWVHVSCALWIPEVSIGCPEKMEPITKISHIPASR WALSCSLCKECTGTCTCQCSMPSCVTAHVTCFAF DHGLEMRTILADNDEVKFKSFCQEHSDGGPRNE PTSEPTEPSQAGEDLEKVTLRKQRLQQLEEDFYE LVEPAEVAERLDLAEALVDFIYQYWKLKRKANA NQPLLTPKTDEVNDLAQQEQDVL YRRLKLFTHL RQDLERVRNLCYMVTRRERTKHAICKLQEQIFH LQMKLIEQDLCRAGLSTSPIDGTFNSWLAQSV QITAENMAMSEWPLNNGHREDPAPGLLSEELLQ DEETLLSFMRDPSLRPGDPARKARGRTRLPAAK KPPPPPPQDGPGRSTTPDKAPKKTWGQDAGSGK GGQGPPTKPPRRTSSHLPSSPAAGDCPILATPES PPPLAPETPDEAASVAADSDVQVP/PGAASPKPLG RLRPPPPREPR*TRRLPGC/ARPDAGDGDHLSAVA ERPKVSLHFDTETDGYFS/DGEMSNS/DVEAED GGVQRGPREAGAKE/VVRMGVLAS
3391	A	1555	327	NSFLHFLHLKVRTMFLFSPFPVLLLSVVTTASCSKT KACADTQKTCSMITCGIPVTNGTPGRDGRDRPK GEKGEPGLGQVSVAS*ISTSGRCSSKS VLEPATRG LKHLRGEAPLSSGPMHLHSEQPL*NAIASKTKLFV DSLGSHISTQELGVCGCPFRGVSVCLVGELALVQA LH*VAGESFFFGSDHWLIGCAGGEQEWSEILLGK KKRVTATGSSSLCLATGQGLRGLQGP PGKMGPP GNTGTSGIPGPRGQKGD RDNSVAEAKLANLER KL*SLRSELDHTKKL*PFSLGKMSGKKLFVTNGE RMPFSKV KALCAGLQATVAAPKNAEENKAIQDV AKDTAFLG/TDEATEGQFM YLTGGRLTYSNWKK DEFNDHGS/EDCWL L ANGLW/ /TSSFLAICE /A
3392	A	218	1773	GGSRNRQRRSIPVLGYFLKQKKMTKAQESLTLE DVAVDFTWEEWQFLSPAQKDLYKJ/MLENYSN LVSVG YQAGKPDALTKLEQGEPLWTLEDEIHSP AHPEIEKADDHLQQPLQNKILKRTGQRYEHGR TLKSYLGLTNQSRRYNRKEPAEFNGDGAFLHDN HEQMPTEIEFPESRKPISTKSQFLKHQQTHNIEKA HECTDCGKAFLKKSQ L TEHKRIHTGKKPHVCSL CGKAFYKKYRLTEHERAHRGEKPHGCSLCGKAF YKRYRLTEHERAHKGEKPYGCSECGKAFPRKSE LTEHQRIHTGIKPHQCSECGRAFSRKSLLVVHQR THTGEKPHTCSECGKGFQKGNLNIHQRTHTGEK PYGCDGCGKAFS QKSCLVAHQRYHTGKTPFVCPE CGQPCSQKSGLRHQIHSGEKPYKCSDCGKAFL TKTMLIVHHRTHTGERPYGCDECEKAYFYMSCL VKHKRIHSREKRGD/CSEGGKSFHKSQ LKS**TC AGEKPC*YGNCGNGGRAV
3393	A	46	1464	ARSLSGAPSGSSRQDGTSLLR TGAGYSSSQSIETL SLPPGPSHLVGDKSQGG RSCQGQITSAASGKTSK SEPNHVIFKKISRDKSVTIYLG NRDYVDHVASQV QPVDGVVLVDPDLVKGKKVYVTLTCAFRYGQE DIDVIGLTFRRDLYFSRVQVYPPVGAASTPTKLQ ESLLKKLGSNTYPFLLTFFDYLPCSVMLQAPQD SGKSCGVDFEVKAFATDSTDAEEDKIPKKSSVRL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Value, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LIRK VQHAPLEMGPQPRAEAAWQFFMFDKPLH LAVSLNKRDLFPMGSPVPVSVANNTEKPVKKI KASVEQVANVVLYS\SDY\YVKPVAMEEAQEKV PPNSTWTKA\LTLLPWL\NNRERRGIALDGGKIKH EDTNLASSTIIKEGIDRKRSWEILVSYPDQR*SSTV SGFLGRASPSQ*SRPT*RSQFRLMHPQ\EDPAK ESYQDANLVFEEFARP*ILKDAGEA*EGKRDQE
3394	A	211	1591	RPPTMAADQRPKADTLALRQRLISSSCRLFFPEDP VKIVRAQQGYMYDEQGAEYIDCISNVAHVGHCH PLVVQAAHEQNQVLNTNSRYLHDNIVDYAQRLS ETLPEQLCVFYFLNSGSEANDLALRLARHYTGH QDVVVLDAHAYHGHLLSLIDISPYKFRNLDDGQKE WVHVAPLPD TYRGPYREDHPATHVEDGLEKAFS* KRVVQGRNRQICRRQIAAFFAESLPSVGGQIIPPA GYFSQVAEHIRKAGGVFVADEIQVGFRVKGHF WAFQLQGGKDFVPDIVTMGKSIGNGHPVACVAAT QPVARAFEATGVEYFNTFGGSPVSCAVGLAVLN VLEKEQLQDHATSVGSFLMQLLGQKIKHPVIG DVRGVGLFIGVDLIKDEATRTPATEEAAYLVSR KENYVLLSTDGPGRNLIKFKPPMCFSLDNARQV VAKLDAILTDMEEKVRSCE TLRLQP
3395	A	1	1424	FRDGFSLRCGCNAELPGRGGDDAADRAIQRLR TGAAVRYKVMKNWGVIGGIAAALAAGIYVTWG PITERKKRRKGLVPGVLNLGNTCFMNSLLQGLSA CPAFIRWLEEFTSQYSRDQKEPPSHOYLSLTLLHL LKALSCQEVTDDEVLHASCLLDVLRMYRWQISS FEEQDAHELPHVITSSLEDERDRQPRVTHLFDVH SLEHSQK*LPKQITCRTRGSPHPTSNHWKSQHPF HGRLTSNMVCKHCEHQSPVRFDTFDSLSPISAA TWGHPLTLDHCLHHFISSESVRDVCDNCTKIF KGTLSGEEKVEHQRTTFVKQ\XJGK\TQCLCH QRLSWSSHGTP LKRHEHVQNEFI\ADYKYEL LGHKPSQHNPKNKNPGPTLELQDGP GAPT PGL NQPGAPKTQIFMNGACSPSLPTLSAPMPFPLPV VPDYSSSTYLFRLMGSCRPPWETWHSGLTCSFTD GPHL
3396	A	109	107	TQEAGLIFFSPFSLSLSLPLSLFLLSHPHSRTTP NRTPRRTRIPQRPVVMYSPLCLTQDEFHPFIEALL PHVRAFAYTWFNLQARKRKYFKKHEKRMSKEE ERAVKDELLSEKPEVKQK WASRL LAKLRKDIRP EYREDFVLTVTGKKPPCCVLSNPDQKGKMRRID CLRQADKVWRDLVMVILFKGIPLESTDGERLV KSPQCSNPGLCVQPHHIGVSVKELDLYLAYFVH AADSSQSESPSAK*R*H*GPARKWDIWGFQDS FVTSGVFSVT*A*LRVSQTP\AAG\TGNFSLSD LESSSYYSMPGAMRRSLPSTSTSTSKRLKSVED EMDSPGEEFFYTGQGRSPGSGSQSSGWHEVEPG MPSPITLKKSEKSGFSSPSQTSLSLGTAFQTQHR PVITGTQSKFHIA TPSIL\HFPRHSPPFQPGPYFSH PAIRYHPQETLKEFVQLVCPDAGQAGQPNGSS QGKVHNPFLLPTPMLPPPPPPMARPVPLVPD TK PPTTSTEGGAASPTSPTRSPGRTRPQQPFL/SYG PP*PSNALIGGGGGGAGERAGERADLEM
3397	A	1	2002	TGTLTEDGLDVMGVVPLKGAFLPLVPEPRRLP VGPLLRLATCHALSRLQDTPVGDPMDLKMVES

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TGWVLEBEPAAADSAFGTQVLAVMRPPLWEPQLQ AMEEPPVPVSVLHRRFPSSALQRMSVVVAWPGA TQPEAYVKGSPSELVAGLCNPETVPTDFAQMLQS YTAAGYRVVALASKPLPSVPSLEAAQQLTRDTV EGDLSLLGLLVMRNLLKPQTPVIQALRRTRIRA VMVTGDNLQTA VTVARGCGMVAPQEHLLIVHA THPERGQPASLEFLPMESPTAVNGVKDPDQAAAS YTVEPDPRSRLALSGPTFGIIVKHFPKLLPKVLV QGTVFARMAPEQKTEL VCELQKLQYCVGMCGD GANDCGALKAADVGLSLSQAEASVVSPTSSMA SIECVPMVIREGRCSLDTSFSVKYMALYSLTQFI SVLILYTINTNLGDLQFLAIDL VITTTVAVLMSRT GPALVLGRVRPPGALLSVPLSSLLQOMVLVTG VQLGGYFLTLAQPFVPLNRTVAAPDNLPNYEN TVVFSLSFQYLILAAAVSKGAPFR/RPLTNNVPF LLASAL*SSVLVVLVLSPLGLHGPLALRNITDTGF KLLLVLGLVTLNFGGLHAGERARPVPPRLPAPPP AQAGSKKRFRKQLERELAEQWPPLPAGPLR
3398	A	758	1368	FPFRMLTGYLYLMWRRKAFWSGTQRHPLPGLL KRRRRPGRGPWPAPGGQGVGPSAL*KAGSPPAN RPGQGE/PGLISPKPVTEVLPDVQGAPVPVPLPT PPSLPHLQNPPTVQHYLLSFSWKPSQGP*RA* PSPLPPAAMRPDG*PGPASQGPDPQGP/PCPPASLP TSPPGKGFQKTETRKHPPRQGHKPKCTANRPLA SFL
3399	A	906	1091	HHHHHHHHHHHHHLLVAFGKVQ*LQNSPSSSSSS SSGCFWQARFSSYRTLHHHHHHHHHHHHHH
3400	A	1838	325	PFLSVHRSPhGPSKLCDDPQASLVPEPVPGGCQE PEEMSWPPSGEIASPPELPSSPPGLPEVAPDATST GLPDTPAAPETSTNYPVECTEGSAGPQSLPLILE FVKNPCSVKDQTPLECEVENTTENTTENTTENTT PETSPPHPPSSSTPCSAHLTPSSLPSSSSSSSE KFYNFVILHARADEHIALRVSGRSWEALC/PDG ATFCEDFQVPGRGELSCLQDAIDHSFILLLTIN VDCRLSLHQVNQAMMSNLTRQGSQDCVIPFLE VLESSPARLSSDTASLLSGLVRLDEHSQIFARKVA NTFKPHRLQARKAMWRKEQDTRALREQSQHLD GERMQAAALNAAYSAYLQSYLSYQAQMEQLQV AFGSHMSFGTGAPYGARMPFGGQVPLGAPPPFP TWPGCPQPPPLHAWQAGTPPPSPQPAAFPQSLP FPAVPKPFPTASTAPPSEPKGWQPLIHHAQMVT SWG*NKHMWNQRGSQAPEDKTQEAE
3401	A	153	1389	EWGWLGAQQPPEEEAEAEQESPSLCREALAEI KKEISPLFIGMEKCSVGGLLETEQTPALLGNMAM ATSLMDIGDSFGHPACPLVSRSRNSPVEDDDDD DVVFIESIQPPSISAPAIADQRNFIFASSKNEKPQ NYSVIPSSRDLASQGNISETIVIDDEDIETNGG AEKKSSCFIEWGLPGTKNKTNLDFTSSLSRSK VNAGMGNSGITTELTKYIITNVTTLETGISSVNA GQDVNIITYKTSL*NTNLGDVAKGLQSSNFGVNI QTYTPSLTPQTKTGVNLLTLVE*MWQETYFRME NLQLII/CPEDASTKKANVILPVESSKSQEFYSTS CLSPCENNWNLLKGVFNKSRCTICKSLAEVWIFI PKLLFRLTVIILTFKCYVFLFHLHNARVLDV
3402	A	153	1389	EWGWLGAQQPPEEEAEAEQESPSLCREALAEI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				KKEISPLFIGMEKCSVGGLELTEQTPALLGNMAM ATSLMDIGDSFGHPACPLVSRSRNSPVEDDDDD DVVFIESIQPPSISAPAIADQRNFIASSKNEKPQG NYSVIPSSRDLASQKGNISSETVIDDEEDIETNGG AEKKSSCFIEWGLPGTKNKTNDLDFSTSSLSRSK VNAGMGNSGITTELTKYIITNVTLETGISSVNA GQDVNIITYKTSL*NTNLGDVAKGLQSSNFGVNI QTYTPSLTPQTKTGVNLLTLVE*MWQETYFRME NLQLII/CPEDASTKKANVILPVESSKSFQEFYST CLSPCENNWNLLKGVFNKSRCTICKLAEVWIFI PKLLFRLTVIILTFKCYVFLHNLHNAVRLDV
3403	A	609	2765	SRHCTPAERQNETHRAPDFAMSAVLGHQPPFFPA LTLPNGAAALSLPGALAKPIMDQLVGAAETGIP FSSLGPQAHRLPLKTMEPEEEVEDDPKVHLEAKE LWDQFHKRGTEMVITKSGRMFPPFKVRCSGLD KKAKYILLMDIIAADDCRYKFHNSRWVMVAGKA DPMPKRMYYHPDSPATGEQWMSKVVTFHKLKL TNNISDKHGFTILNSMHKYQPRFHIVRANDILKLP YSTFRTYLPETEFIAVTAYQNDKITQLKIDNPNF AKGFRDTGNGRREKRKQLTLQSMRVFDERHKK ENGTSDESSSEQAAFNCF\AQASSPAA\PL*RTSNL KDFSPSRG*RATPEAEQRGSTAPRPATRAKISP HPRRRSPAVTRAAPAVKAHLFAAERPRDSGRDL KASPSRHSPTISSSTRGLGAEERRSPVREGQA PAKVEEARALPGKEAFAPLTVQTDAAAAHQAQ PLPGLGFAPLAGQOFFNGHPLFLHPSQFAMGG AFSSMAAAGMGPLLATVSGASTGVSGLDSTAM ASAAAAQGLSGASAATLPFHQQHVLASQGLA MSPFGSLFPYPYTYMAAAAAA/SSAAASASVHRT P\NLNTMRPRLRYSPYSIPVPVDGSSLLTTALPS GAAALGPEDGKAAALASPASVA\SCCELNS R3SVTLSSSSMSLS*VLCALTEAATSELQSIQRLVS GLEAKPDRSRSPASP
3404	A	1082	1308	LKKFLEVPOSYSLLLSFELQWRA*RPQNAIG*Q FIKTLVFFGIMRSAGDVLSTQVSCALRIMRTAGC SHSSP
3405	A	1553	559	PRPPTQRLSRFAPPCRTAEFFPFRRAVVTRPAPPR ACTVVGRRSSPVTGLAVGAAVAMLTVAARSRPFA PVLSATSRGVAGALTP*MQATVPATPEQPVLDL KRPFLSRESLSGQAVRRPLVASVGLNVPASVCYS HTDIKVPDFSEYRRLVLDSTKSSRESSEARKGFS YLVTVGVTTVGVAYAAKNAVTFVSSMSASADV LALAKIEIKLSDIPEGKNMAFKWRGKPLFVRHRT QKEIEQEAAVELSQLRDPQHDLDRVKKPEWVILI GVCTHLGCVPANAGDFGGYYCPCGHSHYDASG RIRLGAPLNLEVPTYEFTSDDMVIVG
3406	A	83	2671	CLYPDFCRSVTCAMPCFTHRSCREDPGTSESREM DPVAFKDVAVNFTQEEWALLDISQKNLYREVML ETFWNLTSIGKKWKDQNIYEYQNP RRNFRSVT EEKVNEIKEDSHCGETFTFPVDDRLNFQKKKASP EVKSCDSFVCEVGLGNSSSNMNRGDTGHKACE CQEYGPKPWKSQQPKKAFRYHPSLRTQERDHTG KKPYACEKCGKNIIYHSSIQRHMVVHSGDGPYK CKFCGKAFHWLSLYLIHERTHTGEKPYECKQCG KSFSYSATHRIHERTHIGKPYECKQCGKAFHSR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				SCHRHESHMGEKAYQCKECGKAFMCPRYVRR HERTHSRKKLYECKQCGKALSSLTSFQTHIRMH GERPYECKTCGKGFYSKSFQRHEKTHSSEKPY KCKQCGKAFTSRGSRFYHERHTTGEKPYECKQC GKAFRSAPNLQSHGRTHHTGEKPYECKECGKAFIF VNNLQSHERTQTHIRHSGERRYKCKICGKGFYC PKSFQRHEKTHHTGEKLYEC/TATFSSSFSSSSSF*Y HERHTTGEKPYKCEQCGKAFAVSIL*MHGRTH PEEKPYECEQ*RKAFRSAPHL*IRGRTHNGEKPY ACKKCGKPFSAQNLRHRTQTHIMHSVERPYK CKICGRGFYSKSFQTHEKSYTGEKPYECKQCG KAFVSFTSFRYHERHTTGENPYECKQFGKAFRSV KNLRFHKTHTGEKPCBYMKRLTLEGNTMNAS NVAKLSLLPVLFNIMKEFTLGRNPISVSNVRKPLF LPLLNFNIMKGLTWERNPMSVCHVGKPSFLLVPFN IMKGLTLERSPMNISNVGKPSDQPRTFKCMGLT LEKNPMNVSSMGKRSDDLTRFFEYR
3407	A	1426	3	PAAPSGASPRVCGVETARPLGVQRRQSADEGP PGVAGLRHEPPTVWLGSAHRTWVCAHRWFG PAVTRAAQAATMVKLLVAKILCMVGVFVFFMLL GSLLPVKIETDFEKAHRSKKILSLCNTFGGGVFL ATC/LTALLARC*GKSSRRSWSLGHISTDYPLAE TILLGFFMTVFLEQLLTFQENAVLHRPGDLQR RIGRGQRLGV*EPLHGGGRAGPRAVRGAPRPRQP ERAGPLA/PSPVRLLSLAFALSAHSVFEGALGLQ EEGEKVVSFLVGVAVHETLVPVALGISMAGSAM PLRDAAKLAVTVSPMIPLGIGLGLGIEKAQGVPG SVASVLLQGPGRHLSLFTFPGKSWPRSWRKKS DRLLKVLFLVVGYYTVLAGMGLPQVVSGLAIVPA AGSPPGAPGRTQAASPGRASPKSEHCGPGPPVH KGFHGTLLCPRSYTLISIRALLFKELLSLKSLL KK
3408	A	106	451	EARDRLAQSRKEKELNSVASELSARQEESEHSH KHLELRREFKKNVPEEIREMVA/VLKS/QAEVV ALSKRSQEAEEAFLSVYKQ/LIEAPALWELKLKSR PALGDSRVQQGQHDPKTDNQNTOQKAGFKEGW LAEASEREAFGPGFKDPVPVFEEARSLDDRLQPP SFDPSGQPRDLHTSWKRNPELLSPKALKATQAE LLELRRKYDEEAASKADEVGLIMTNLEKANQRA EAAQREVESLREQLASVNSSIRLACCSPQGPSGD KVNFTLCSGPRLEAALASKDREILRLKDVQHLQ SSLQEELEASANQIADLERQLTAKSEAIEKLEEL QAQSDYEEIKTELSILKAMKLASSTCSLPQGMAL PEDSLLIAKEAFFPTQKFLLEKPSLLASPEEDPSED DSIKDSLGTESYPSFQQLPPPPGPEPDLSPSPGQP LLGPSLGPDGTRTFSLSPPPSLASGERLMMPPAAF KGEAGLLVFPFAFYGAKPPTAPATPAPGPEPLG GPEPADGGGGAAGPGAEEELDTAEIAFQVKE QLLKHNIQQRVFGHYVLGLSQGSVSEILARPKA WRKLHG**GKEPFIKMKQFLSDEQNVLALRTIQV RQRGSITPRIRTPETGSDDAIKSIQAKKEIESQK GGEPKTSVAPLSIANGTTPASTSEDAIKSILEQAR REMQAQQQALLEMEVAPRGRSVPPSPPERPSLAT ASQNGAPALVKQEEGSGGPAQAPLPVLSPAAFV QSIIRKVKSEIGDAGYFDHHWASDRGLLSRPYAS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VSPSLSSSSSSGYSGQPNGRAWPRGDEAPVPPED EAAAGAEDPPRTGELKAEGATAEAGARLPYYP AYVPRTLKPTVPPLTPEQYEL YMYREVDLTLELTR QVKEKLAKNGICQRIFGKVLGLSQGSVSDMLSR PKPWSKLTQKGREPFIRMQLWLSOQLGQAVGQQ PGASQASPTPEPRSSPPSPTEPEKSSQEPLSLSL SSKENQQPEGRSSSSLSGKMYSGSQAPGGIQEIV AMSPELDTYSITKRVKEVLTNNLQGLRFGESIL GLTQGSVSDLLSRPKPWHKLSLKGREPFVRMQL WLNDPHNVEKLRDMKKLEKKA YLKRRYGLIST GSDSESPATRSECPSPCLQPDLSLLQIKKPRVVL APEEKEALRKA YQLEPYPSQQTIELLSFQLNLKT NTVINWFHNYRSRMRREMLVEGTQDEPDLDPSG GPGILPPGHSHDPDTPQSPDSETEKPTVKELEL QEGPEENSTPLTTQDKA QVRIKQE QMEEDABEE AGSQPDSGELDKGQGPKEEHPDPPGNDGLPK VAPGPLLPGGSTPDPSLHPQEQSEAGERLHPDP LSFKSASESSRCSLEVSLNSPSAASSPGLMMSVSP VPSSAPISPPPGAPPAKVPSASPTADMAGALHP SAKVNPNLQRRHEKMANLNNIYRLERAANREE ALEWEF
3409	A	162	1710	GPLSPGPYQCRPSLPAQLYPQSLMAAATLRTPTQ GTVTFEDVA VHSWEEWGLLDEAQRCLYRDVM LENLALLTSLDVHHQKQHLGEKHFSINVGRALF VKTCTFHVS GEPSTCREVGKDFLAKLGLFHQQA AHTGEQSNKSDGGAISHRGKTHYNWGEHTKAF SGKHTLVQQRTLTTERCYICSECGKSFSKSYSL NDHWRLHTGEKPYECRECGKSFRQSSSLIQHRR GHTAVRPHECDECGKLF SNKSNLIKHRRVHTGE RPYECSECGKSFNQRSALLQHRGVHTGEKPYEC TECGKSFS... ...NSSLIEHLRVHTGERPYK... LQHRGVPTGERPYECSECGKFF... HTGSRPYECSECGKSFTQNSGLIK... YECE*KKSFHNSSLIKHQRIHSR*KPYECKCG NR*HPGESPVHSECQ/KSFS*RPYLECHTVHKG KTLICRDVQLI
3410	A	167	789	LCMKGISGGVRVAALAARAEREELPVPAMEPOP TAWGSPHPEAVLQLEVAPESSGPCTDTAKDQOS DKLPDLMPPA/EPLGSALELRASLEIDVAERGCE HGPSQQLPRCP*SWAWSEFPWCQRP GCAV*APLP Y*REASFTYQSHSPAASGPFHSAGAGAVYLQAGG V/GEQEKEAVRKSGSSSSCSQRGPMPGMEVCPL LGFWAICP
3411	A	1040	887	ASLSKPAGISTMPWALILLFLLTHSAVSVVQAGL TQPPSVSKDLRAQTATLTCTGNSNNVGHQGVWL QQHQGHPPKLLSYRNNNRPSGISERLSAYKSGNA ASLTTYGLQTEHEAD**CRPRRKLIPKTARLFFFFL IDNEEYLLRVY
3412	A	164	83	RRGIPGSASLSLTMCVRS CFQSPRLQWVWRTAFL KHTQRRHQGSHRWTHLGGSTYRAVIFDMGGVLI PSPGRVAAEWEVQNRIPSGTILKALMEGGENG WMRFMRABITAEGFLREFGRLCSEMLKTSVPVD SFFSLLTSERVAKQFPVMTEAITQIRAKGLQTAVL SNNFYLPNQSFPLDRKQFDVIVESCMEGICKP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				DPRIYKLCLEQLGLQPSESIFLDDLGTNLKEAARL GIHTIKVNDPETA VKELEALLGFTLRVGVPNTRP VKKTMEIPKDSLQKYLKDLLGIQTTGPLELLQFD HGQSNPTYIYIRLANRDLVLRKKPPGTLLPSAHAI EREFRIMKALANAGVPVPNVLDLCESSVIGTTF YVMEYCPGLIYKDPSPGLPESHRRAIYTAMNTV LCKIHSVLDLQAVGLEDY GKQGSTTWV/YSSRRA RGALLFLDWELSYWGDPPFADVGYSCLAHYLP SFPVLRGINDCDLTQLGIPAAEEYFRMYCLQMGL PPTENWNFYMAFSFFRVAAILQGVYKRSLTGQA SSTYAEQTGKLTEFVSNLAWDFAVKEGFRVFK MPFTNPLTRSYHTWARPQSQWCPTGSRSYSSVPE ASPAHTSRGGLVISPELSPPVRELYHRLKHFME QRVYPAEPELQSHQASAAWSPSPLEDLKVQKP W*GGRSGRTSWRLLAGCHT
3413	A	105	1573	PESRHQCFSDRSSHFLTMEME QEKMTMNKELSP DAAAYCCSACHGDETWSYNHPIRGRASRSLSA SPALGSTKEFRRTSLHGPCPVITFGPKACVLQ PQTIMHIQDPASQRLTWNKSPKSVLVKKMRDAS LLQPFKELCTHLMENMIVYVEKKVLEDPALASD ESFGAVKKKFC TFREDYDDISNQIDFICLGGDGT LLYASSLFQGSVPPVMAFHLGSLGFLTPFSFENFQ SQVTQVIEGNAAVVL/RGSRLKVRVVKELRGKK TAVHNGLGEKGSQAAGLMDVKGQAMQYQVL NEVVIDRGPSSYLSNVDVYLDGHLITTVQGD/G* GPQHLSWGP*AFLGRE*RLRLSLSGVIVSTPTGST AYAAAAGASMIHPNVPAMITPICPHSLSFPRIVV PAGVELKIMLSPEARNTAWVSFDGRKRQEI RHG DSISITTS CYPLPSICVRDPVSDWFESLAQCLHWN VRKKQAHFEEEEEEEG
3414	A	20	2602	VIVNKNVN.V.NYV.NQQQR...E...K...ISA ALGLPDLTKPFTF...EREKMAVGVL TQT VGPW PRPVA YLSKQLDGVSKGWPPCLRALAATALLAQ EADKLTLGQNLNIKAPHAVVTLMNTKGHHWLT NARLTKYQSLPCENPHITIEVCNTILNPTTLLPVSE SPGEHNCVEVLDSVYSSRPDLRDQPWASSVDWE LYMDGSSFINSQGERCAGYAVVTILDAVIKAKLW LQGTSAQKABELIALTRAVELSEGQESLEELLGRY FYVSHLPFAKAVAQLCITCRQHNAHQSPQTVSPH IQAYGAAPFEDLQVDFTEMPKCGGNKYLLVLT TYSGWVEAYPTRTEKAYEVTRVLLRDLIPRFLP LRIGSHNGPVFVADLDCVEINVDTGVIWATWIKN EKDPVQLQKGKSGPCTKGQCNPFLVITNPLDP RWKKGERVTLGINGAGLNPRVNILVRGEVYKCS LEPVFQTFYDELNVPITEFPKGKTRNLFLQLAEHV AQSLTVTSCYVCGGTVIADQWPWEARELVPTDP VPDEFPAQKNHPDNFVVLKASIRQYYIARVEKD FTLPVGRLHGG/RSNHTEKNPFSKPKLQTV*AHP ESHRDWTAPTGLYWICGHRAYTKLPASSCVIGTI KPSFFLLSIKTGELLGFPVYASRKSIARN*NNDK WPPERIIQYYGPAT*AQDGSWGYRIPIYMINRIRL QAVLKIITATGRALTILAQOETQMRNATYQNRALA LDYLLAAEGEVC RKFNLTNCCLHIDNQGGVVED IVRDMTKVAHVVPVQVWHGFDPGAMFRKWFPAL GGFKTLIRVIIIGITYLLLPRLLPVLLQMIKSFIAT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LVYQNASAQVYYINHY
3415	A	455	108	NMSWRGRSTYRPRRRSLQPPELIGAMLEPTDEE PKEEKPPTKSRNPTPDQKREDDSG/SAADFKWP EPGKPIFQGAMVRPKTGG/CGCEGGY* CQGEDSP KAEHFKMPEAGEGKSQV
3416	A	1	874	FFFFQRINFIEHSGSVSLLALACDLGWCEDWSCC LVQGGGDLVDVVQTNHGEDEAGGDTDSVDEAR CKESQQAQENLREDLCLSFADKILQIEGSER EHEETRTKQAALDGEPLGGGQLTAVHLHPSKEQ QQEGGERQRGARTHWRGWKEGRRVRLRPPS GKLADQPVRKLGGPPTS/TELPGLQPHAPTHT A/PATPTYSPAPDTPNPPVRWKCLPVEPRTRQLC RERTRKACPPKPRPPLGLPGDPTGPVTHAPPVS PTGASGQERRAEPGAVSYAHASATK
3417	A	243	847	CLKYMYTYIFCPNCVSYKMKTDHFSLRYLHSSC AEDNKSSVDSSGQAAHPSKGGFFPHGTHWGTQC RGHISVLGWQCSCPSTGCRVGLGLAMCQTHAYI HTHTHTHTHTPTDYGAAHTDPLQRWGLGPRKS EAGPLQLSRDQSHPGPLSPGASPRSAGLPGWHP AHQEPRARGRCARDGLSLQTRLTNKYDIQCCQE MRK
3418	A	4073	1000	LDEYEARTLANLDDFEEDNEDDDENRVNQEEK AAKITELINKLNFLDEAEKDLATVNSNPFDDPDA AELNPFDPDSEEPITETASPRKTEDSFYNNSYNP FKEVQTPQYLNPFDEPEAFVTIKDSPQSTKRKNI RPVDMSKYLYADSSKTEBEELDESNPFEYEPKSTP PPNNLVNPVQELETERRVKRKAPAPPVLSPKTGV LNENTVSAGKDLSTSPKPSPIPSVLGRKPNASQS LLVWCKEVTKNYRGVKITNFTTSWRNGLSFCAI LHIF/PDLIDYKSLNPQDIKENNKAYDGFASIGI SRLEPSEMVALEPDEPQTYLQIKATTEQC ELNVVQIEENSSKSTYKVGNTETDTNSSVDQEK YAEISDLKREPELQQHIGAVDFLSQDDSVFVND SGVGESESEHQTPDDHLSNSASPYCRRTKSDTEP QKSQSSGRTSGSDDPGICSNTESTQAQVLLGKK RLKAELELSLYVSDKKKDMSPFICEETDEQ KLQTLDIGSNLEKEKLENSRSLCSDPESPIKKT SLSPSTKLGYSSRDLDLAKKKHASLRQTESDPD ADRTTLNHADHSSKIVQHRLLSRQEELKERARVL LEQARRDAALKAGNKHNTNTATPFCNRQLSDQ QDEERRRQLRERARQLIAEARSQVGMSELPSYGE MAAEKLEKERSKASGDENDNIBIDTNEEIPGFFV GGGDELTNLENDLTPEQNSKLVDLKLKLLV QPQVANSPSSAAQKAVTESSEQDMKSGTEDLRT ERLQKTTERFRNPVVFSDSTVRKTQLQSFSQYI ENRPEMKRQRSIQEDTKKGNEEKAAITETQRKPS EDEVLNKGFKDSISQYVVGELAALENEQKQIDTR AALVEKRLRYLMDTGRNTEEEAMMQEWFML VNKKNALIRRMNQLSLEKEHDLERRYELLNRE LRAMLAIEDWQKTEAQKRREQLLLDELVALVN KRDALVRDLDAQEKQAEEDEHLERTLEQNK KMAKKEEKCVLQ
3419	A	4073	1000	LDEYEARTLANLDDFEEDNEDDDENRVNQEEK AAKITELINKLNFLDEAEKDLATVNSNPFDDPDA AELNPFDPDSEEPITETASPRKTEDSFYNNSYNP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				FKEVQTPQYLNPFDEPEAFVTIKDSPPQSTKRKNI RPVDMISKYLYADSSKTEEEELDESPPFYEPKSTP PPNNLVNPVQELETTERRVKRKAPPPVLSPKTGV LNENTVSAGKDLSTSPKPSPIPSVLGRKPNASQS LLVWCKEVTKNYRGVKITNFTTSWRNGLSFCAI LHHFRPDLIDYKSLNPQDIKENNKKAYDGFASIGI SRLLPESDMVLLAIPDKLTVMTYLYQIRAHFSGQ ELNVVQIEENSSKSTYKVGNYETDTNSSVDQEF YAEISDLKREPELQQPISGAVDFLSQDDSVFVND SGVGESESEHQTPDDHLSPTASPYCRRTKSDTEP QKSQQSSGR TSGSDDPGICSN TDSTQAQVLLGKK RLKAETLELSDLVVSDDKKDMSPPFICEETDEQ KLQTLDIGSNLEKEKLENSRSLECRSDPESPIKKT SLSPTSKLGYSSSRDLDAKKKKHASLRQTESDPD ADRTTLNHADHSSKIVQHRLLSRQEELKERARVL LEQARRDAALKAGNKHNTNTATPFCNRQLSDQ QDEERRRQLRERARQLIAEARSQVGMSELPYGE MAAEKLEKERSKASGDENDNIEIDTNEIPEGFVV GGGDEL TNLENDLTPEQNSKLVDLKLKLELV QPQVANSPSSAAQKAVTESSEQDMKSGTEDLRT ERLQKTTERFRNPVVFSSKDVTRKTQLQSFSQYI ENRPEMKRQRSIQEDTKKGNEEKAITETQRKPS EDEVLNKGFKDSISQYVVGELAALENEQKQIDTR AALVEKRLRYLMDTGRNTEEEAMMQEWFML VNKKNALIRRMNQLSLEKEHDLERRYELLNRE LRAMLAIEDWQKTEAQKRREQLLLDELVALVN KRDALVRDLDAQEKQAEEDDEHLERTLEQNKG KMAKKEEKCVLQ
3420	A	612	1058	ENLGPNYSHRLLHHPFTFYKKIHKHHHEWTAFIG VISLYAHPIEHAVSNMLPVIVGPIVMGSHLSITM WFSLAENITTCYICGVILPFLPSFIFDYTHITFN QCYGVLGVLDHLELITMFKQTKAYERHVLL GFTPLSESIPDSPK
3421	A	23	2005	LLTPCDGRIPGRPSVGAESGSDFFQRRRRRRDPE EPEKTELSERELAVAVASQENDEENEERWVGP LPVEATLAKKRKVFLEFERYLDNLPSASMYERS YMRDVIHVCTKTDFITASHDGHVKFWKKIE EGIEFVKHFRSHLGVIESIASVSEGA LFCSVGDDK AMKVFDVVNFDMINMLKLGYPGQCEWIYCPG DAISSVAASEKSTGKIFTYDGRGDNQPLHIFDKLH TSPLTQIRLNPVYKAVVSSDKSGMIEYWTGPPHE YKFPKNVNWEYKTDITLYEFAKCKAYPTSVCF PDGKKIATIGSDRKVRIFRFTGKLMRVFDESLS MFTELQQMRQQLPDMEFGRRMAVERELEKVDA VRLINIVFDETGHFVLYGTMLGIKVINVETNRCV RILGKQENIRVMQLALFQGIKKHRAATTIEMKA SENPVLQNIQADPTIVCTSFKKNRFYMTKREPE DTKSADSDRDVFNEKPSKEEVMAATQAEQPKRV SDSAIHTSMGDIHTKLPVECPKTVENFCVHSRN GYYNGHTFHRIKGFMIQTGDPGTGMGGESIWG GEFEDEFHSTLRHDPYTLSMANAGSNTNGSQFF ITVVPTPWLDNKHTVFGRVTKGMEVVQRISNVK VNPKTDKPYEDVSIINTVK
3422	A	2486	433	FVLVCAPLTWAGARHRRMAASKKPPRVRVNHQ DFQLRNLRIIEPNEVTHSGDGTGVETDGRMPKVT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				SELLRQLRQAMRNSEYVTEPIQAYIIPSGDAHQSE YIAPCDCRRAFVSGFDGSAGTAITEEHAAMWTD GRYFLQAAKQMDSNWTLMKMGLKDTPTQEDW LVSVLPEGSRVGVDPDIPTDYWKMAKVLRSA GHHLIPVKENLVDKIWTDRPERPCKPLLTLGLDY TGISWKDKVADLRLKMAERNVMWFVVTALDEI AWLFNLRGSDVEHNPVFFSYAIGLETIMLFIDGD RIDAPSVKEHLLLDLGLAEYRIQVHPYKSILSEL KALCADLSPREKVWVSDKASYAVSETIPKDHRC CMPYTPICIAKA\VKNSA\ESEGMRRAHIKDAVAL CELFNWLEKEVPKGGVTEISAADKAEFFRRQQA DFVDLSFPTISSTGPNAGIIHYAPVPETNRTLSD VYLIDSGAQYKDGTTDVTRTMHFGTPTAYEKEC FTYVLKGHIAVSAAVFPTGTKGHLLDSFARSAL WDSGLDYLHGTGHGVGSFLNVHEGPCGISYKTF SDEPLEAGMIVTDEPGYYEDGAFGIRIENVVLV PVKTKYNFNRRGSLTFEPLTLVPIQTKMIDVDSL TDKECDWLNNYHLTCRDVIGKELQKQGRQEAL EWLIRETQPIISKQH
3423	A	5515	934	FKMPENPATDKLQVLQVLDRLKMKLQEKGDTS QNEKLSMFYETLKSPLFNQILTLQOSIKQLKGQL NHIPSDCSANFDFSRKGLLVFTDGSITNGNVHRPS NNSTVSGLPFWTPKLGNEFNSVIQMAQGRQIE YDIERPSTGGLGPSVVALRSQNLGKVDIFVKDV QPGSVADRDQRLKENDQILAINHTPLDQNIHQ AIALLOTTGSLRLIVAREPVHTKSSTSSSLNDTT LPETVCWGHVEEVELINDGSGLGFGIVGGKTS VVRTIVPGGLADRDGRLQTGDHILKIGGTNVQG MTSEQVAQVLRNCGNSVRMLVARDPAGDISVTP PAPAALPVALPTVASKGPGSDSSLFETYNVELVR EDQSLGCRNNSF/GTSHTGE\SCN/VKSIIPGS/ AYHNGHIQ\NDK\AVDGVNIQGFANHDVVEVL RNAGQVVHILVRRKTSSSTSPLEPPSDRGTVVE PLKPPALFLTGA\ETETINVDGEDEEIKERIDTLKN DNIQALEKLEKVPDSENELKSRWENLLGPDYEV MVATLDTQIADDAELQKYSKLLPIHTLRLGVEV DSFDGHHYISSIVSGGPVDTLGLLQPEDELLEN GMQLYGKSRREAVSFLKEVPPFTLVCCRRLFDD EASVDEPRRTETSLPETEVDHNMMDVNTTEDDDG ELALWSPEVKIVELVKDCKGLGFSILDYQDPLDP TRSVIVIRSLVADGVAERSGGLLPGDRLVSVNEY CLDNTSLAEAVEILKA\PPGLVHLGICKPLVEDN BEESCYLHSSSNEDKTEFSGTIHDINSSLILEAPK GFRDEPYFKEELVDEPFLDLGKSFHSQQKEIEQS KEAWEMHEFLT\PRLQEMDEEREMLVDEEYELY QDPSPSMELYPLSHIQEATPVPSVNELHFGTQWL HDNEPSESQEARTGRTVYSQEAQPYGYCPENV KENFVMESLPSVPSTEGNSQQGRFDDLENLSLA KTSLLDLGMPNDVQGPSLLIDLPPVVAQRREQEDL PLYQHQA\TRVISKASAYTGMLSSRYATDTCELPE REEGEGETPNF\SHWGPPRIVEIFREP\NSLGISV GGQTVIKRLKNGEELKGIFIKQVLEDS\PAKGTNA LKTGDKILEVSGVDLQ\NASHSEAVEAIKNAGNP VVFIVQSLSS\TPRVIPNVH\NKANKITGNQ\NQTQ EKKEKRQGTAPPPMKLPPPYKALTDDSDENEEE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				DAFTDQKIRQRYADLPGELHIELEKDKNGLGLS LAGNKDRSRMSIFVVGINPEGPAAADGRMHIGD ELLEINNQILYGRSHQNVASAIKTAPSKVKLVFIR NEDAVNQMAVTPFPVPSSSPSSIEDQSGTEPISSEE VDGSLE\VGKQLPESESFKLA\VSQMKQKQYPTKV SFSSQEIPLAPASSYHSTDADFTGYGGFQAPLSVD PATCPIVPGQEMIEISKRRSGLGLSIVGGKDTPLV NGVDLRNSSHEEAITLRQTPQKVRLVVYRDEA HYRDEENLEIFPVDLQKKA\GRGLGLSIVGKR
3424	A	2223	1162	HASERVVQLPDFVWDQYTHSLGRVEREFKNRKR HTRRVKLVFDKGLPARPKSPLDPKKDGESLSYS MLPLSDGPEGSSSRPQMIRGRLCDDTKPETFNQL WTVVEEQKKLEQLLIKYPPEEVESRRWQKIADELG NRTAKQVASRVQKYFIKLTAGIPVPGRTPNLYI YSKKSSTSRQHPLNKH\LFKP\GTFMTSHEPPVY MDEDDDRSCFHSHMNTAVEDASDDESIPIMYRN LPEYKELLQFKKLKQKLQHMQAESGFVQHVGF KCDNCGIEPIQGV\RW\HCR\DCPP\EMSLDFC\DS CSDCLHETDIHKGDH\QLEPIYRS\ETFLDRDYCV SQGTSYN\LDPNYFPANR
3425	A	2223	1162	HASERVVQLPDFVWDQYTHSLGRVEREFKNRKR HTRRVKLVFDKGLPARPKSPLDPKKDGESLSYS MLPLSDGPEGSSSRPQMIRGRLCDDTKPETFNQL WTVVEEQKKLEQLLIKYPPEEVESRRWQKIADELG NRTAKQVASRVQKYFIKLTAGIPVPGRTPNLYI YSKKSSTSRQHPLNKH\LFKP\GTFMTSHEPPVY MDEDDDRSCFHSHMNTAVEDASDDESIPIMYRN LPEYKELLQFKKLKQKLQHMQAESGFVQHVGF KCDNCGIEPIQGV\RW\HCR\DCPP\EMSLDFC\DS CSDCLHETDIHKGDH\QLEPIYRS\ETFLDRDYCV SQGTSYN\LDPNYFPANR
3426	A	2	1553	LFVVVHDDPRWGTPRYW\GALY\RNQSSPTAPP GLLPLEYFPAAPHCSHSRQW\CSQTHRIHHHPQ MLGPCRQEICGITMAAGTLYTY\NWRAF\KALI AAQYSGAQVRVLSAPPHFHFGQTN\IPEFLRKFP AGKVPAFEGDDGFCVFESNAIAYYSNEELRGST PEAAAQVVQWVSFADSDIVPPASTWVFTLGIM HHNKQATENAKEBVRRLGLLDAYLKTRITLVG ERVTLADITVVCTLLWLYKQVLEPSFRQAFPNTN RWFLTCINQPQFRA\VFGEVKLCEKMAQF\DAKK FAETQPKKDTPRKEKGSREEKQKPOAERKEEKK AAPAPEEEMDECEBQALAAEPKAKDPFAHLPKS TFVLDEFKRKYSNEDT\LSVALPYFWEHFDKDGW SLWYSEYRFPEELTQTFMSCNLITGMFQRLDKLR KNAFASVILFGTNNSSSISGVWVFRGQELAFPLSP DWQVDYESYTWKLDPGSEETQTLVREYFSWE GAFQHVKGAFNQKIFK
3427	A	755	52	TAARRRQKGTAAARRRQKGTAAARRRQKGTAAARR RQKGTAAARRRQKGTAAARRRQKGTAAARRRQKGT AARRRQKGTAAARRRQKGTAAARRRQKGTAAARR QKGLSNLDAAEWLPPKKG\GEKKKGPF\LAINEV VTREYPINILKRIHG\VGFKKRAPRALKEIRKFAM KEMGTPDVRI\TRLNKA\WVAKGIRNVPYRIRVR LSRKRNEDEDS\PNKLYTLVTVVPVTTFKNLQTV NV\DEN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
3428	A	4	1939	LPLSLFSFSEMPPLPLPMDLKGEPPGPKPGPWGP PGPPGFPKPGHKGKPLHGQPGAGPPGFSRMG KAGPPGLPGNVGPPGQGLRGEPIRGDQGLRG PGPPGLPGSGITIPGKPGAQGVPPGPGFQGEPP QGEPGPPGDRGLKGDNGVGPGLPGAPGQGGAP GPPGLPGAGLGKPGLDGLPGAPGDKGESGPPG VPGPRGEPGAVGPKGPPGVDGVGVPGAAGLPGP QGPPGAKGEPGTRGPPGLIGPTGYGMPGLPGPKG DRGPAGVPGLLGDREGEDEGEQGPQGLGG PPGLPGSAGLPGRRPPGPKGEAGPGGPPGVPI RGDQGPSGLAGKPGVPGERGLPGAHGPPGPTGP KGEPGFTGRPGPGVAGALGQKGDGLPGQPL RGSPGIPGLQGPAGIPGQGLPGLKGEPLPGPPG EGRAGEPGTAGPARGPPGVPGSPGITGPPGLPGPP GAPGAFDETGIAGLHLPNGGVEGAVLGKGGKPK FGLGELSAHATPAFTAULTSPLASGMPVKFDR LYNGHSGYNPATGIFTCPVGGVYFYAYHVHVKG TNVWVALYKNNVPATYTYDEYKKGYLDAQSG GAVLQLRPNDQVWVQMPDQANGLYSTEYIHSS FSGFLLCPT
3429	A	212	1075	EGLTGPCERVFPFLGRGPPHGATRAGHRAVRW AGPESLPPLPRSLIMDSPRAGTHQGPLDAETEVG ADRCTSTAYQEQRQVEQVGKQAPLSPGLPAMG GPGPGCEDPAGAGGAGAGGSEPLVTVTVQCAF TVALRARRGADLSSLRALLGQALPHQAQLGQLS YLAPGEDGHVPIPEEESLQRAWQDAAACPRGL QLQCRGAGGRPVLYQVVAQHSYSAQGPEDLGF RQGDITVDVLCEVDQAWLEGHCDGRIGIFPKCFV VPAGPRMSGAPGRLPRSQQGDQP
3430	A	799	1989	INKYINIRKKIKLLSPLPWSHLALC ASATKWV ETFAFAGKIPGYNQPLPT WRSLS TFWLLATKRRKQQLVLI DETKEEEDPPLPTT PTSVNYHFTQCNYKCGFCFHTAKTSFVLPLEEA KRGLLLKVEAGLEKINFSGGEPFLQDRGEYLK LVRFCVELRLPSVSVSNGSLIRERWFQNYGE YLDILAISCDSFDEEVNCPIGRGMGKKNHVENL QKLARRWCRDYRVFPKINSVINPFNVEEDMTEQI KALNPVRWKVFQCLLIEGENCGEDA/LREAERFV IGDEEFERFLERHKEVSCLVPESNQKMKDSYLL DEYMRFLNCRKGRKDPKSKILDVGVEEAIKFSGF DEKMFLKRGGKYTWSKADLKLDW
3431	A	5468	2146	ACGFLPGRCHFSTFKQCQEWLSRLSRATARPAKP EDLFAFAYHAWCLGLTEEDQHTLCPGGEHRC RQEAELARMGFDLQNVWRVSHINSNYKLCPSYP QKLLVPVWITDKELNVASFWSKRIPVVVYRH LRNGAAIARCSQPEISWWGWRNADDEYLVTSIA KACALDPGTRATGGSLSGTGNDTSEACDADFDS SLTACSGVESTAAPQKLLILDARSYTA AVANRAK GGGCECEEYYPNCEVFMGMANIHAIRNSFQYL RAVCSQMPDPSNWSALESTKWQLHLSVMLKA AVLVANTVDREGRPVLVHCSGDGWDRTQPIVALA KILLDPYYRTLEGFQVLVESDWLDFGHKFGDCR GHQENVEDQNEQCPVFLQWLDVSHQLLKQFPCL FEFNEAFLVKLVQHTYSCLYGTFLANNPCEREK RNIYK/RGTCSVWALLRAGNKNFHNFLYTPSSD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				MVLHPVCHVRALHLWTAVYLPASSPCTLGEEN MDLYLSPVAQSQEFSGRSLDRLPKTRSMDDL ACDTSSPLTRTSSDPNLNNHCQEVVRVGLEP NPEGSETSFVDSGVGGPQQTVEVGLPPPLSSQ KDYLSNKPFSKSHKSCSPSYKLLNTAVPREMK SDPEIKVLEETKGPAPDPSAQDELGRITLDGIG EHCPETEAVSALSKVSNKCDGVCNFPESQNSPT GTPQQAQPDMSLGVPSKCVLDHSLSTVCNPPSA ACQTPDPSTDFLNQDPSGSVASISHQEQLSSVP DLTHGEEDIGKRGNNRNGQLLENPRFGKMPLEL VRKPISQSQISEFSFLGSNWDSFQGMVTSFSPGEA TPRRLSYGCCSKRPNSKQMRATGPCFGGQWAQ REGVKSPVCSHSHNGHCTGPGGKNQMWLSSH QVSSTKPVPLNCPSPVPPLYLDDDDGLPFTDVIQ HLRQIEAGYKQEVQLRRQVRELQMRDIRHCC APPAEPPMDYEDDFTCLKESDGSDEDFGSDHSE DCLSEASWEPVDKKETEVRWVPDHMASHCYN CDCEFWLAKRRHHCNCGNVFCAGCCHLKLPIP DQQLYDPVLCNSCYEHIQVSRARELMSQQLKK PIATASS
3432	A	36	1873	MTFFSSVADFIGLDPRIAAWLIDPSDATPSFEDLV EKYCEKSITVKVNSTYGNSSRNIVNQNVRENKLT LYRLTMDLCSKLDYGLWQLFRTLELPLILAV MESHAIQVNKEEMEKTSALLGARLKELEQEAHF VAGERFLITSNNQLREILFGKLKLHLLSQNSLPR TGLQKYPSTVSEALNALRDLHPLPKIILEYRQVH KIKSTFVDGLLACMKKGSISSTWNQTGTVTGRLS AKHPNIQGISKHPIQITTPKNFKGKEDKILTISPR MFVSSKGHTFLAADFSQIELRILTHLSGDPELLKL FQESERDDVFSTLTSQWKDVPVEQVTHADREQT KKVVYA...YAGGFRLAAG...QEA... ...QKYKKEKDFARAAIAQ...Q...VSIMGRRK PLPRIHAHDQQLRAQAERQA...NFVVQGSAADLC KLAMIHVFTAVAASHILTARLVAQHDELLFEVE DPQIPECAALVRRTMESLEQVPLKVSLSAGRSWG HLVPLQEAWALRQAHVALSLPATAWLPLGLP APSPHPCIFRLHFVCSPRQQWEERTGFQQSIVWPS PRSPALYAPGRINPLGLGWPAPWSKCLCKALKK K
3433	A	1481	476	IPPKERAPGIRASCLAITAGARPTSYGRVGCEDV RLSPVSPLAPPDRLASRWEGRSRMKGKKGIVA ASGSETEDEDSMDIPLDLSSAGSGKRRRRGNLP KESVQILRDWLYEHRYNAYPSEQEKALLSQQTH LSTLQVCNWFNARRRLLPDMRLKDGKDPNQFTI SRRGAKISETSSVESVMGIKNFMPALBETPFHSFT AGPNPTLGRPLSAKP/SQSPGSVLARPSVICHITTV TAIERLSLSLSCQSVGCGQNTDIQIATRNLRDS SLMYPEDTCKSGPSTNTQSGLFNTPPPTPPDLNQ DFSGFQLLDVVALKRAAEMELQAKLTA
3434	A	1720	1243	NGPVPPGSGSKTKWAGGSAAEGSPRLSPSGAAQ VPALLRGEPRGGAAAGSFWKPLHQHSCGLRPPP/ PPD/RLSRLPGKTLASCDRENGARRPLLGLSTSFIP IGRRTYASAAEPVGSKAVLVTGCDSGFGFSLAKH LHSGFLVFAGCLMKDKGHDGVKELDSLNSDRL RTVQLNVCSSVEVEKV/VGDCPLEPEGPÆKGMW

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				GLVNNAGISTFGEVEFTSLETYKQVAEVNLWGT VRMTKSFLPLIRRAKGRVVNISSMLGRMANPAR SPYCITKFGVEAFSDCLRYEMYPLGVKVSVEPG NFIAATSLYSPESIQAIKKMWHEELPEVVRKDYG KKYFDEKIAKMETYCSSGSTDTSPVIDAVTHALT ATTPYTRYHPMDYYWWLRMQIMTHLPGAISDM IYIR
3435	A	842	3595	ENQQQMLVAKEQRLHFLKQQERRQQQSISENEK LQKLKERVEAQENKLKKIRAMRGQVDYSKIMN GNLSAEIERFSAMFQEKKEVQTAILRVDQLSQQ LEDLKKGKLNGFQSYNGKLTGPAAVELKRLYQE LQIRNQLNQBQNSKLQQQKELLNKRNMVAMM DKRISELRLRYGKKIQACEKVFLNRVNGTSSPQ SPLSTSGRVAAVGPYIQVPSAGSFPVLGDPIKPS LSIASNAAHGRSKSANDGNWPTLKQNSSSSVKP VQVAGADWKDPSVEGSVKQGTVSSQPVFSA LG PTEKPGIEIGKVPPPIPGVGKQLPPSYGTYPSP TPL PGSTSSLERRRKEGSLPRPSAGLPSRQRPTLLPAT GSTPQPGSSQQIQQRISVPPSPTYPAGPPAF PAGD SKPELPLTVAJRPFADKGSRPQSPRKGPTVNSS SIYSMYLQQATPPKNYQPAAHSA LNKS VKAVYG KPVLPSTSPSPPLPFLHGSLSGTPTQPQPPESTE KEPEQDGPAPADGSTVESLPRPLSPTKLTPIVHS PLRYQSDADLEALRRKLANAPRLKKRSSITEPE GPGGPNIQKLLYQRFNTLAGGMEGTFPYQSPSQ DFMVTLADVDNGNTNANGNLEELPPAQPTAPLP AEPAPSSDANDNELPSPEPEELICPQTHQTAEPA EDNNNNVATVPTTEQIPSPVAEAPSPGEEQVPPA PLPPASHPPATSTNKRTNLKKPNSERTGHGLRVR FNPIALLLDASLEGEFDLVQRIIYEVEDPSKPND GITPLHNAVCAHHHIVKILLETGVNRTAADSD GWTPLHCAASCNSVHLCKQLVESGAAASTISD IETAADKCEEMEEGYIQCSQFLYGVQEKLGVMN KGVAYALWDYEAQNSDELSFHEGDALTILRRKD E
3436	A	3	2604	GSTHASEKMKTGSRALVVTDTGDMSVLNSPRHQ SCIMHVDMDCCFFVSVGIRNRPDLKGKPAVTSN RGTGRAPLRPGANPQLEWQYYQNKILKGKADIP DSSLWENPD SAQANGIDSVLSRAELASCSYE ARQ LGKNGMFFGHAKQLCPNLQAVPYDFHAYKEVA QILYETLASVYTHNIEAVSCDEALVDITEILAE TK LTPDEFANAVRMEIKDQTKCAASVVGISNILLAR MATRKAKPDGQYHLKPPEVDDFIRGQLVTNLPG VGHSMESKLASLGIKTCGDLQYMTMAKLQKEF GPKTGQMLYRFCRGLDDRPRVTEKERKSVSAEI NYGIRFTQPKAEAFLLSLSEEIQRREATGMKG KRLTLKIMVRKPGAPVETAKFGGHGICDNIARTV TLDQATDNAKIIKGAMLMFHTMKNISDMRGV GIHVNLQVPTNLNPSTCPSRPSVQSSHFPSGSYSV RDVFQVQKAKKSTEEHKEVFRAAVDLEISSAR TCTFLPPFAHLPTSPDTNKAESSGKWNLHTPV SVQSRLNLSIEVSPSQLDQSVLEALPDLREQVE QVCAVQQAESHGDKKKEPVNGCNTGILPQPVGT VLLQIPEPQESNSDAGINLIALPAFSQVDPEVFAA LPAELQRELKAA YDQRQRQGENSTHQQSASASV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				PKNPLLHLKAAVKEKKRNKKKKKTIGSPKRIQSPL NNKLLNSPAKTLPGACGSPQKLIDGFLKHGPPA EKPLEELSASTSGVPGLSSLQSDPAGCVRPPAPNL AGAVEFNDVKLTLLREWITTISDPMEEDILQVVKY CTDLIEEKDLEKLDLVIKYMKRLMQQSVESVWN MAFDFILDNVQVVLQQTYGSTLKV
3437	A	32	4038	<p> SLLRLKAQWGSSGAASEPVVLGEEGCGFPSTNE YPDLEERATYPQEDRFLTPGRAQLLWSPWSPL DQEEACASRQLHSLASFSTVTARRNPLHNPWGM ELAASENTDSPSPRLRPGVTLPPGALTMNTKDT TEVAENSHHLKIFLPKKLLECLPRCLLPPERLRW NTNEEIASYLITFEKHDEWLSCAPKTRPQNGSIL YNRKKVKYRKDGYLWKKRKDGKTTREDHMKL KVQGMCECLYGCYVHSSIVPTFHRRCYWLLQNP IVLVHYLNPALCEDCGKGCSPIFCSISSDRREW LKWSREELGQLKPMFHHGWSCGNGTEEFV VEHVQILDTHTPKPAPRTHACLCSGGLSGSLTHK C SSTKHRISPKVEPRALTLTSIPHPHPPEPPPLIAPLP PELPKAHTSPSSSSSSSSGFAEPLIRSPPTSRGG SSRGGTAILLTGLEQRAGGLTPTRHLAPQADPR PMSLAVVVGTEPSAPPAPSPAFDPRFLNSPQR GQTYGGGQGVSPDFPEAAEAHTPCSALEPAAAL EPQAAARGPPQSVAGGRRGNCFFIQDDDSGEEL KGHGAAPPISPPSPPPSPAPLEPSSRVGRGEALF GGPVGASELEPFLSSFPDLMGELISDEAPSIPAPT PQLSPALSTITDFPEWSYPEGGVKVLITGPWTEA AEHYSCVFDHIAVPASLVQPGVLRCPAHEVG LVSLQVAGREGPLSASVLFYRARRFLSLPSTQL DWLSLDDNQFRMSILERLEQMEKRMAEIAAAGQ VPCQGPDAAPPVQDEGQGPGEARVVVLVESMIP PSTTTCSTTAHGSFFRCMSLLHLAAACGYARL IETLQWASVETGSLDLEQVDPLNVDHFSCTPL MWACA:GHLEAAVLLFRWNRQALSIPDSLGRLP LSAHSRGRVRLARCLEELQRQEPSVEPPFALSP PSSSPDTGLSSVESPELSDGTFSVTSAYSSAPDGS PPPAPLPASEMTMEDMAPGQLSSGVPEAPLLLM DYEATNSKGPLSSLPALPPASDDGAAPEDADSPQ AVDVIPVDMISLAKQHEATPERIKREDFVGLPEA GASMRERTGAVGLSETMSWLASYLÆENVDFHPS STPPSELAPFERGRLGLSLTAPSWAEFLSCIPPVGK IGKLIIFALLTLSD\QEQRELYEAARVIQTAFRKYK GRRLEQEQEVAAAVIQRKYKQLTWIALKFA LYKKMTQAAILIQSKFRSYYEQKRFQQRRAAV LIQQHYRSYRRRPGPPHRTSATLPARNKGSFLT KQDQAARKIMRFLRRCRHRMRELKQNQEGLP QPGLAT </p>
3438	A	469	2602	<p> FGRLWGTAFKSWKMKAPIHLILLYATFTQSLK VVTKRGSADGCTDWSIDIKYQVLVGEVRIKC ALFYGYRTNYSLAQSAGLSLMWYKSSGPGDFE EPIAFDGSRMKEEDSIWFRPTLLQDSGLYACVIR NSTYCMKVSISLTVGENDTGLCYNSKMKYFEKA ELSKSKEISCRDIEDFLLPTREPEILWYKECRTKT WRPSIVFKRDTLLIREVREDDIGNYTCELKYGGF VVRRTTTELTVTAPLTDKPPKLLYPMESKLTIQET QLGDSANLTCRAFFGYSGDVSPLIYWMKGKGFIE </p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				DLDENRVWESDINKILKEHLGEQEVSSISLIVDSVEE GDLGNYSYCVENGNGRRHASVLLHKRELMYTV ELAGGLGAILLLLVLVTYKCYKIEIMLFYRNHF GAEELDGDNDKYDAYLSYTKVDPDQWNQETGE EERFALEILPDMLEKHYGYKLFIPDRDLIPTGT EDVARCVDQSKRLIIVMTPNYVVRGWSIFELET RLRNMLVTGEIKVILIECSELRGIMNYQEVEALK HTIKLLTVIKWHGPKCNKLNKSKFWKRLQYEMPF KRIEPI THEQALDVSEQPFGELQTVSAISMAAA STALATAHPDLRSTFHNTYHSQMRQKHYYRSYE YDVPPGTGLPLTSIGNQHTYCNIPMTLNGQRPQT KSSREQNPDEAHTNSAILPLPRETSISSVIW
3439	A	251	2037	GPGNSSILIGGGHLFIRSCNLNLLNLSKENTEHT MAKKVAVIGAGVSGLSSIKCCVDELEPTCFERS DDIGGLWKFTERGSSLSVMIWPLALSLLRHGGFC YSDFFHEDYPNFMNHEKFWDYLQFAEHFDLL KYIQFKTTVCGITKRPDSETGQWDVVTEGKQ NRAVFDAMVCTGHFLNPHLPLEAFPGIHKFKG QILHSQEYKIPEGFQGRVLVIGLGNTGGDIAVEL SRTAAQVLLSTRGTGWVLGRSSDWGYPYNMMV TRCCSFIAQVLPFRFLNWQERKLNKRFNHEDY GLSITKGKKAKFIVNDELPCILCGAITMKTSVIE FTETSAVFEDGTVEENIDVVIFFTGYTFSFPFEEP LKSCTKKIFLYKQVPLNLERATLAIGLIGLKGS ILSGTELQARWVTRVFKGLCKRPASQKLMMEAT EKEQLIKRGVFKDTSKDKFDYIAYMDDIAACIGT KPSIPLFLKDPRLAWEVFGPCTPYQYRLMGPG KWDGARNAILTQWDRTLKPLKTRIVPDSSKAWP SMASHYLKAWGAPVLLASLLICKSSFLKLVDRD KLQDRMSPYLVS LWRG
3440	A	1	3533	IMPGGGRLLR... ENSE...LGESMAGISQNAKTGDLPA...GECV...IASK ALCGLTEAAAQAAYLVGIFDPNSQA...HQGLVDP IQFARANQAIQMACQNLVDPGSSPSQV...RATIV AKHTSALCNACRIASSKTANPVAKRHFVQS...KE VANSTANLVKTIKALDGDSEDNRNKCRIATAPL IEAVENLTAFASNPEFVSIPAQISSEGSQAQEPILV SAKPMLESSSYLIRTARSLAINPKDPPTWSVLAG HSHTVSDSIKSLITSIRDKAPGQRECDYSIDGINRC IRDIEQASLAASVQSLATRDDISVEALQEQLTSVV QEIGHLIDPIATAARGEAAQLGHKGTQLASYFEP LILAAVGVASKILDHQQQMTVLDQTKTLAESAL QMLYAAKEGGGNPKAQHTHDAITEAAQLMKEA VDDIMVTLNEAASEVGLVGGMVDAIAEAMSKL DEGTPPEPKGTFVDYQTTVVKYSKAI AVTAQEM MTKSVTNPEELGGLASQMTSDYGH LAFQGQMA AATAEP EIGFQIRTRVQDLGHGCIFLVQKAGAL QVCPTDSYTKRELIECARAVTEKVS LVSALQAG NKG TQACITAATAVSGHADLDTTIFATAGTLN AENSETFADHRENILKTAKALVEDTKLLVSGAAS TPDKLAQAAQSSAATITQLAEVVKLGAASLGSD DPETQVVLINAIKDVAKALSDLISATKGAASKPV DDPSMYQLKGAAKVMVTNVTSLKTVKA VEDE ATRGTRALEATIECIKQELTVFQSKDVPEKTSSPE ESIRMTKGITMATAKAVAAGNSCRQEDVIATAN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				LSRKAVSDMLTACKQASFHPDVSDEVTRALRF GTECTLGYLDLLEHVLVILQKPTPELKQQLA KRVAGAVTELIQAAEAMKGTWVDPEDPTVIAE TELLGAAASIEAAKKLEQLKPRAKPKQADETL DFBEQILEAAKSIAAATSALVKSASAAQRELVAQ GKVGSIANAADDGQWSQGLISAARMVAAATSS LCEANASVQGHASEEKLISSAKQVAASTAQLL VACKVKADQDSEAMRRLQAAGNAVKRASDNL VRAAQKAAF GKADDDVVVKTKFVGGAQIIAA QEEMLKKEREELEARKKLAQIRQQYKFLPTL REDEG
3441	A	3	1584	NSARGGVGVRGARAMATVQEKAALNLSALHS PAHRPPGFSVAQKPFGATYVWSSIINTLQTQVEV KKRRHRLKRHND CFV GSEAVDVIFSHLIQNKYF GDVDIPRAKVVRVCQALMDYKVFEAVPTKVFG KDKKPTFEDSSCSLYRFTTIPNQDSQLGKENKLY SPARYADALFKSSDIRSASLEDLWENLSLKPANS PHVNISTTLPQVINEVWQEETIGRLLQLVDLPLL DSLLKQGEAVPKIPQPKRQSTMVNSSNYLDRGIL KAYSDSQEDELWLSAIDCLEYLPDQMVVEISRSF PEQPDRTDLVKELLFDAIGRYSSREPLLNLHSD VHNGIAELLVNGKTEIALEATQLLLKLLDFQNR EFRRLLYFMAVAANPSEFKLQKESDNRMVVKRI FSKAIVDNKNLSKGKTDLLVFLMDHQKDVFKI PGTLVHKIVSVKVLMAIQNGRDPNRDAGYTYCQRI DQRDYSNITEKTTIDELLYLLKTLDEDSKLSAKE KKKLLGQFYKCHPDIFIEHFGD
3442	A	160	822	SPASGHCLNGAAVAMFGCLVAGRLVQTAAQQ VAEDKFVFDLPDYESINHVVFMLGTIPPEGMG GSVYPSYPDSNGMPVWQLLGFVINGKPSAIFKIS NKSCEGQVTFGALNIVETPSVAQIGISVLELDC MAQQPVGUNA VSSVDSFTQFTQKMLDNFYNF ASSFAVQ/VPDDTQ/RPSEMFIPANVVLKWYENF QRRSTEPSLENIHWIKINF
3443	A	3	1373	SWHVRRRWLEATMAGGMKVAVSPA VGP GPWG SGVGGGGTVRLLLLSGCLVYGTAETDVNVVML QESQVCEKRASQQFCYTNVLPQWHDWTRIQIR VNSSRLVRVTQVENBEKLKELEQFSIWNFFSSFL KEKLN DTYVNVGLYSTKTCLKVEIEKDTKYSVI VIRRFDPKLFVFLGLMLFFCGDLLSRSQIFYYS TGMTVGIVASLALIIIFILSKFMPKKSPIYVILVGGW SFSLYLIQLVFKNLQEIWRCYWQYLLSYVLTVGF MSFAVCYKYGPLENERSINLLTWLQLMGLCFM YSGIQIPHIALAIIIALCTKNLEHPIQWL YTCRKV CKGAEPVPPRLLTEEEYRIQGEVETRKALEELR EFCNSPDCSAWKTVSRIQSPKRFAD FVEGSSHLT PNEVSVHEQEYGLGSIIAQDEIYEEASSEEDSYS RCPAITQNNFLT
3444	A	566	1718	KGLERTCCAMEESDSEKTTEKENLGPRMDPPLG EPG\GSLGWVLPNTAMKKKVLLMGKSGSGKTS MRSIIFANYIARDTRRLGATILDRHSLQINSSLST YSLVDSVGNTKTFDVEHSHVRFLGNLVLNLWDC GGQDTFMENYFTSQRDNIFRNVEVLIYVFDVESR ELEKDMHYYYQSCLEAILQNSPDAKIFCLVHKMD LVQEDQRDLIFKEREEDLRRLSRPLECSCFRTSIW

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				DETL YKAWSSIVYQLIPNVQQLEMNLRNFAEIE ADEVLLFERATFLVISHYQCKEQRDAHRFEKISNI IKQFKLSCSKLAASFQSMVEVRNSNFAAFIDIFTSN TYVMVVMSDPSIPSAATLINIRNARKHFEKLERV DGPKQCCLLMR
3445	A	566	1718	KGLERTCCAMEESDSEKTTEKENLGPRMDPPLG EPGVGSLGWVLPNTAMKKKVLLMGKSGSGKTS MRSIIFANYIARDTRRLGATILDRIHSLQINSSLST YSLVDSVGNTKTFDVEHSHVRLGNLVLNLWDC GGQDTFMENYFTSQRDNI FRNVEVLIYVFDVESR ELEKDMHY YQSCLEAILQNSPDAKIFCLVHKMD LVQEDQRDLIFKEREEDLRRLSRPLECSCFRTSIW DETL YKAWSSIVYQLIPNVQQLEMNLRNFAEIE ADEVLLFERATFLVISHYQCKEQRDAHRFEKISNI IKQFKLSCSKLAASFQSMVEVRNSNFAAFIDIFTSN TYVMVVMSDPSIPSAATLINIRNARKHFEKLERV DGPKQCCLLMR
3446	A	566	1718	KGLERTCCAMEESDSEKTTEKENLGPRMDPPLG EPGVGSLGWVLPNTAMKKKVLLMGKSGSGKTS MRSIIFANYIARDTRRLGATILDRIHSLQINSSLST YSLVDSVGNTKTFDVEHSHVRLGNLVLNLWDC GGQDTFMENYFTSQRDNI FRNVEVLIYVFDVESR ELEKDMHY YQSCLEAILQNSPDAKIFCLVHKMD LVQEDQRDLIFKEREEDLRRLSRPLECSCFRTSIW DETL YKAWSSIVYQLIPNVQQLEMNLRNFAEIE ADEVLLFERATFLVISHYQCKEQRDAHRFEKISNI IKQFKLSCSKLAASFQSMVEVRNSNFAAFIDIFTSN TYVMVVMSDPSIPSAATLINIRNARKHFEKLERV DGPKQCCLLMR
3447	A	1	2930	VLLGPLWDKLSTADHPVIVTMASKRKSTTPCMIP VKT VVLQDASMFAQPAETL...IQCCLPEASA ASSEAAQNPSST...STLANGHRSTLDGYLYSC YCDFRSHDMTQFVGHMNSEHTDFNKDPTFVCSG CSFLAKTPEGLSLHNATCHSGEASFVWNVAKPD NHVVVEQSIPESTSTPDLAGEPSAEGADGQABIII KTPIMKIMKGKAEAKKIHTLKENVSPQPVGEALP KLSTGEMEVREGDHSFINGAVPVRQASASSAKN PHAANGPLIGTVPVLPAGIAQFLSLQQQPPVHAQ HHVHQPLPTAKALPKVMIPLSSIPTYSAAMDNS FLKNSFHKFPYPTKAELCYLTVVTKYPEBQLKIW FTAQRLKQGISWSPEEIEDARKKMFNTVIQSV PQ PITTVLNTPLVASAGNVQHIIQAALPGHVVGQPE GTGGGLLV TQPLMANGLOATSSPLPLTVTSVPK QPGVAPINTVCSNTTSAVKVVNAAQSLLTACPSI TSQAFLDASIYKNKKSHEQLSALKGSFCRNQFFG QSEVEHLTKVTGLSTREVRKWFSDRRYHCRNLK GSRAMIPGDHRSIIDSVPESFSPSSKVPEVTCIPT TATLATHPSAKRQSWHQTDFPTKYKERAPEQ LRALESSFAQNPLPLDEELDRLRSETKMTRREIDS WFSEERRKKVNAEETKKAENASQEEEEAABDEG GEEDLASELRVSGENGSLEMPSSHILAERKVSPK INLKNLRVTEANGRNEIPGLGACDPEDDES NKLA EQLPGKVSCCKTAQQRHLLRQLFVQTQWPSNQD YDSIMAQTGLPRPEVVRWFGDSRYALKNGQLK WYEDYKRGNFPPGLLVIAPGNRELLQDY YMTHK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				MLYEEDLQNLCDKTMSSQQVKQWFAEKMGEETRAVADTGSEDQGGPGTGELTAVHKGMDTYSEVSENSESWEP RVPEASSEPFDTSSPQAGRQLETD
3448	A	2	1324	FVARAEKGFRRTREAHLLQVAGVGTGLQNGASLSGLASGVMAQRAFPNPYADYNKSLAEGYFDAAGRLTPEFSQRLTNKIRELLQQMERGLKSADPRDGTGYTGWAGI AVL YLHLYDVF GDPA YLQLAHGYV KQSLNCLTKRSITFLCGDAGPLA VAAVLYHKMN NEKQAEDCITRLIHLNKIDPHAPNEMLYGRIGYTY ALLFVNKNFGVEKIPQSHIQICETILTSGENLAR KRNFTAKSPLMYEWYQEYYVGAAHGLAGIYYY LMQPSLQVSQGLHSLVKPSVDYVCQLKFPSGN YPPCIGDNRDLLVHWCHGAPGVITYMLIQAYKVF R/BEREYLCADAYQCADVWQYGLLKKGYGLCYA GSAGNAYAF L TLYNL TQDMKYL YRACKFAEWC LEYGEHGCRTPDTPFSLFEGMAGTIYFLADLLFP TKARVFP AFEL
3449	A	3	2389	SRHVTGAARSPSRAGPSDPPAMGDEDDDESCAV ELRITEANLTGHEEKVSVENFELLKVLGTGAYGK VFLVRKAGGHDAKLYAMKVLKKAALVQRAK TQEHTRTERS VLELVRQAPFLVTLHYAFQTD AKL HLILDYVSGGEMFTHLYQRQYFKEAEVRVYGGE IVLAL EHLHLKLGHIYRDLKLENVLLDSEGHIVLTD FGLSKEFLTEEKERTFSFCGTIEYMAPEIIRSKTGH GKAVDWWSL GILLFELLTGASPFTLEGERNTQAE VSRRLKCSPPFPFRIGPVAQDLLQRLCKDPKKR LGAGPQGAQEVNRNHPFFQGLDWVALAARKIPAP FRPQIRSELDVGNFAEEFTRLEPVVSPPGQPPPG DPRIFQGYSFVAPSILFDHNNAVMTDGLEAPGAG DRPGRAAVARSAMMQDSPFFQYELDLREPALG QGSFVCRRCBQSGQEFVKI SPPE EANTOR EVAALRLCQEPNVTALHEVHHDQLHTYL VLEL LRGGELEHIRKLRHFSESEASQILRSLVSAVSFM HEEAGVVHRDLKFMILYADDTPGAPVKIIDFG/F SPRLRPQSPGVPMQTPSFELQYAAPELLAQGGYD ESCDLWSLGVILYMMLSGQAPFQGASGQGGQS QAAEIMCKIREGRFSLDGEAWQGVSEAKELVR GLLTVDPARKRLKLEGLRGSSWLQDGSARSSPPLR TPDVLESSGPAVRSGLNATFMAFNRGKREGFFLK SVENAPLAKRRKQKLRSATASRRGSPAPANPGR APVASKGAPRRANGPLPPS
3450	A	201	1705	KGTEMNKSRRWQSRRRHGRSHQQNPWFRLRDS EDRSDSRAAQAHD SGHGDDSPSTSSGTAGTSS VPPLPGFYFDPEKKRYFRLLPGHNNCNPLTKESIR QKEMESKRLRLQLBEDRRKKIARMGFNASSMLR KSQGLFLNVTNYCHLAHELRLSCMERKKVQIRS MDPSALASDRFNILADTNSDRLFTVNDVTVGGS KYGIINLQSLKTPTLKVFMHENLYFTNRKVNSV CWASLNHLDSHILLCLMGLAETPGCATLLPASLF VN SHPAGIDRPGM/LCSFRIPGAWSCAWSLNIQA NNCFS TGLSRRVLLTNVV TGHRQSFGTNSDVL A QQFALMAPLLFNGCRSGEIFAIDLRCGNQKGKW KATRLFHDSA VTSVRILQDEQYLMASDMAGKIK L WDLRTTKCVRQYEGHVNEYAYLPLHVHEEEGI LVAVGQDCYTRIWSLHDARLLRTIPSPYPASKAD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3451	A	19	6033	<p>IPSVAFSSRLGGSRGAPGLLMAVGQDLYCYSYS</p> <p>LLSAML SHGAGLALWITLSLLQTGLAEPERCNFT</p> <p>LAESKASSHSVSIQWRILGSPCNFSLIYSSDTLGA</p> <p>ALCPTFRIDNTTYGCNLQDLQAGTIYNFKIISLDE</p> <p>ERTVVLQTDPLPPARFGVSKEKTTSTGLHVWWT</p> <p>PSSGKVTSYEVQLFDENNQKIQGVQIQESTSWNE</p> <p>YTFFNLTAGSKYNIATAVSGGKRFSVYTNGST</p> <p>VPSPVKDIGISTKANSLLISWSHSGSNVERYRLM</p> <p>LMDKGILVHGGVVDKHA TSYAFHGLSPGYLYNL</p> <p>TVMTEAAGLQNYRWKLVRTAPMEVSNLKVTND</p> <p>GSLTSLKV KWRPPGNVDSYNITLSHKGITIKESR</p> <p>VLAPWITETHFKELVPGRLYQVTCASVSLGELS</p> <p>AQKMAVGRTPDKVANLEANNNNGRMRLVVS</p> <p>WSPPAGDWEQYRILLFNDVSVLLNITVGKEETQ</p> <p>YVMDGTGLVPGRQYEVVIVESGNLKNSERCQG</p> <p>RTVPLAVLQLRVKHANETSLSIMWQTPVAEWEK</p> <p>YIISLADRDILLIHKSLSKDAKEFTFDLVPGRKY</p> <p>MATVTSISGDLKNSSSVKGRTVPAQVTDLHVAN</p> <p>QGMTSSLFTNWTQAQGDVEFYQVLLIHENVVIK</p> <p>NESISSETSRYSFHSLKSGSLYSVVVTTVSGGISSR</p> <p>QVVVEGRTVPSSVSGVTNNNSGRNDYLSVSWLL</p> <p>APGDVDNYEVTLSHDGKVVQSLVIAKSVRECSF</p> <p>SSLTPGRLYTVTITTRSGKYENHSFSQERTVPDKV</p> <p>QGVSVSNSARS DYLRVSWVHATGDFDHYEVTIK</p> <p>NKNNFIQTKSIPKSENECVFVQLVPGRLYSVTVT</p> <p>TKSGQYEANEQGNRTIPEPVKDLTLRNRSTEDL</p> <p>HVTWSGANGDVDQYEIQLLFNDMKVFPFHLVN</p> <p>TATEYRFTSLTPGRQYKILVLTISGDVQQSAFIEG</p> <p>FTVPSAVKNIHISPNGATDSLTVNWTGPGGDVDS</p> <p>YTVSAFRHSQKVDSTIPKHVFEHTFHRLEAGEQ</p> <p>YQEI IASVBSGLKNCINNVGPII IASVQV IADR</p> <p>AYSSYSLI/SWQAGVAERYDILLTENGILL</p> <p>NTSEPA TTQHKFEDLTPGKKYKIQLTVSGGLFS</p> <p>KEAQTEGRTVPAAVTDLRITENSTRHLSFRWTAS</p> <p>EGELSWYNIFLYNPDGNLQERAQVDPLVQSFSFQ</p> <p>NLLQGRMYKMVIVTHSGELSNESFIFGRTPASV</p> <p>SHLRGNSNRNTDSLWFNWSPASGDFDFYELLYN</p> <p>PNGTKKENWKDKDLTEWRFQGLVPGRKYVLW</p> <p>VVTHSGDLSNKVTAESRTAPSPPSLMSFADIANT</p> <p>SLAITWKGPDPWTDYNDFELQWLPRDALTVFNP</p> <p>YNNRKSEGRIVYGLRPGRSYQFNVKTVSGDSWK</p> <p>TYSKPIFGSVRTKPKIQNLHCRPQNSTAIACSWI</p> <p>PPDSDFDGYSIECRKMDTQEVEFSRKLEKEKSL</p> <p>NIMMLVPHKRYLVSIKVQSAGMTSEVVEDSTIT</p> <p>MIDRPPPPPPHIRVNEKDV LISKSSINFTVNCWF</p> <p>DTNGAVKYFTVVVREADGSDELKPEQQHPLPSY</p> <p>LEYRHNASIRVYQNTYFASKCAENPNSNSKSFNI</p> <p>KLGAEMESLGGKCDPTQKFCDGPLKPHTAYRI</p> <p>SIRAFITQLFDEDLKEFTKPLYSDTFFSLPITTESEP</p> <p>LFGAIEGVSA GLFLIGMLVAVVALLICRQKVSHG</p> <p>RERPSARLSIRDRPLSVHLNLGQKGNRKTSPIK</p> <p>INQFEGHFMKLQADSNYLLSKEYEELKDVGRNQ</p> <p>SCDIALLPENRGKNRYNNILPYDATRVKLSNVDD</p> <p>DPCSDYINASYIPGNFRREYIVTQGPLPGTKDDF</p> <p>WKMVWEQNVHNMVMTQCVEKGRVKCDHYW</p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				PADQDSL YYGDLILQMLSES VLP EWTIREFKICGE EQ LDAHRLIRHFHYTVWPDHGV PETTQSLIQFVR TVRDYINRSPGAGPTVVHCSAGVGR TGTFIALDR ILQQLDSKDSVDIYGAVHDLRLHRVH MVQTEC QYVYLHQCVRDVLRARKLRSEQENLPFIYENV NPEYHRDPVYSRH
3452	A	63	1073	FFRSSSDNGSPIRQYE/HSTPAHQGPVMGLEGS/ ARNSQLRIVLVGKTGAGK SATGNSILGRKVFHSG TAAKSITKKCEKRSSSWKETELVVVDTPGIFDTE VPNAETSKEIIRCILLTSPGPHALLLVVPLGRYTEE EHKATEKILKMFGERARSMILIFTRKDDLGD TN LHDYLRAPEDIQDLMDIFGDRYCALNNKATGA EQEAQRAQLLGLIQRVVRENKEGCYTNRM YQR AEEI IQKTQAMQELHRV ELEREKARIREEYEEK IRKLEDKVEQEKRKKQMEKKLAEQEAHYAVRQ QRARTEVESKDGILELIMTALQIASFILLRLFAED
3453	A	2674	514	GPITFLKKKAKMKDMPLRIHVLLGLAITTLVQAV DKKVDCPRLCTCEIRPWFTPRSIYMEASTVDCND LGLLTFFARLPANTQILLQTNNAIKIEYSTDFPV NL TGLDLSQNNLSSVTNINGKKMPQLLSVYLEEN KLTELPEKCLSELNLQELYNHNLLSTISPGAFIG LHNLLRLHLNSNRLQMNSK WFDALPNLEILMIG ENPIIRIKDMNFKPLINLRSLVIAGINL TEIPDNAL VGLENLESI SFYDNRLIKVPHVALQKVNLKFLD LNKNPINRIRRGDFSNNMLHLKELGINNMP ELISID SLAVDNL PDLRKIEATNNPRLSYIHPNAFFRLPKL ESLMLNSNALSALYHG TIESLPNLKEISHSNPIRC DCVIRWMNMNKTNIRFM EPDSLFCVDPPEFQGG NVRQVHFRDMMEICLPLIAPESFPSNLNVEAGSY VSFHCRA TAIEPQPEIYWITPSGOKLLPNTLTDKF YVREGTLDNGVTPELNYTCIANLNLNCLL SVMIKVDGSFPQDNNGSLNRI RDIQANSVLVSW KASSKILKSSVKWTAFTKTENSHAAQSARIPSDV KVYNLTHLNPSTEYKICIDETIYQKNRKKCVNVT TKGLHPDQKEYEKNNTTTLMAELGGLLGIGVIC LISCLSPEMNCDGGHSYVRNYLQKPTFALGELYP PLINLWEAGKEKSTSLVKATVIGLPTNMS
3454	A	1844	244	ERYLFATYVAPSATLDIGLQQEKKKEIYMKIOPP FEDLFDTAEEYILLLLLEPWTKMVKSDQIAYKKV ELVEETRQLDSTYFRKLQALHKETFSKKAEDTTC EIGTGILSLSNVSKRTEYWDNVPAEYKHFKFSDL LNNKLEFEHFRRQFLETHSSMDLMCWTDIEQFRR ITYRDRNQKAKSIYIKNKYLNKKYFFGPNSPAS LYQQNQVMHLSGGWGKILHEQLDAPVLVEIQK HVQNRLENVWLPLFLASEQFAARQKIKVQMKDI AEELLQKAEKKIGVWKPVESK WISSCKIIAFRK ALLNPVTSRQFQRFVALKGDLL ENGLLFWQEVQ KYKDLCHSHCDESVIQKKITTIINC FINSSIPPALQI DIPVEQAQKIEHRKELGPYVFREAQM TFLGVMP KFWPQFCEFRKNLT DENIMSVLERRQEYNKQKK KLAVL/QNDEKSGKDGKIQYANTSVP AIKTALLS DSFLGLQPYGRQPTWCYSKYIEALEQERILLKIQE ELEK\SLQACNLSQILRLALQLCL
3455	A	228	3330	APTAQAMMSFGGADALLGAPFAPLHGGGSLHY ALARKGGAGGTRSAAGSSSGFHSWTRTSVSSVS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				ASPSRFRGAGAAASSTDSDLTSLNGPEGCMVAVA TSRSEKEQLQALNDRFAGYIDKVRQLEAHNRSL GEAAALRQQQAGRSAMGELYEREVREMRGAVL RLGAARGQLRLEQEHLLEDIAHVQRQLDDEARQ REEAEEAARALARFAQEAAARVDLQKKAQAL QEECGYLRRHHQEVEGELLGQIQSGAAQAQM QAETRDALKCDVTSALREIRAQLEGHAVQSTLQ SEEWFRVRLDRLSEAAKVNTDAMRSAQEEITEY RRQLQARTTELEALKSTKDSLERQRSELEDHRQA DIASYQEAIIQQLDAELRNTKWEMAAQLREYQDL LNVKMALDIEIAAYRKLLGEECRIGFGPIFSLP EGLPKIPSVSTHIKVKSEEKIKVVEKSEKETVIVVE QTEETQVTEEBVEEDKBAKEEEGKEEEGGEEEE AEGGEEETKSPPAEEAASPEKEAKSPVKEEAKSP AEAKSPEKEEAKSPAENVKSPEKAKSPAKEEAKSP PEAKSPEKDGKQNFQAEVKSPEKAKSPAKEEAK SPAEAKSPEKAKSPVKEEAKSPAEEKSPVKEEAK SPAENVKSPEKAKSPTKEEAKSPEKAKSPEKAKSP EKKEEAKSPEKAKSPVKAEEKSPEKAKSPVKAEEA KSPEKAKSPVKEEAKSPEKAKSPVKEEAKSPEKA KSPVKEEAKTPEKAKSPVKEEAKSPEKAKSPEKA KTLDVKSPEAKTPAKEEARSPADKFPKAKSPVK BEVKSPKAKSPLKEDAKAPEKEIPKKEEVKSPV KEEBKQEVKVKPEPKKAEKAPATPKTEKK DSKKEEAPKKEAPKPKVEEKKEPAVEKPKESKV EAKKEEAEDKKKVPTPEKEAPKVEVKEDAKPK EKTEVAKKEPDDAKAKEPSKPAEKKEAAPEKKD TKEEKAKKPEEKPKTEAKAKEDDKTLSKEPSKP KAELAEKSSSTDQKDSKPPEKATEDKAAKKGK
3456	A	258	1463	YLSFIPGHASKSAPMNGHCFANGPSQKSSIPPLL ITPGELGPHLELDQVVGCGFKYKTVNGV TPIKNSPSLFFCAPLCERGSSEPLPISEALSLDDT DCEVEFLTSSDTDFLLEDSTLSDFKYDVPGRRSF RGCGQINYAYFDTPAVSAADLSYVSDQNGIGVP DPNPPPPQTHRRLLRRSHSGPAGSFNKPAPRISNCCI HRASPNSEDKPEVPPRPVPIPPRPVKPDYRRWSA EVTSSSTYSDEDRPPKVPREPLSPSNSRTPSPKSLP SYLNGVMPPTQSFAPDPKYVSSKALQRQNSEGS ASKVPCILPIENGKKVSSTHYLLPERPPYLDKY BKFFREAKKKNGGAQIQPLPADCGISSATEKPD KTKMDLGGHVKRKHL SYVGTP
3457	A	2	4869	FILSSSSASSEHFHHHSFGNW WPGSFKGHRMS LPFYQRCHQHYDLSYRNKDVRSTVSHYQREKKR SAVYTQGSTAYSSRSSAAHRESEAFRRASASS QQASQHALSSEVSRKAASAYDYGSSHGLTDS LLLDDYSSKLSPKPKRAKHSLLSGEKENLPSDY MVPIFSGRQKHVSGITDTEERIKEAAAYIAQRNL LASEEGITTPKQSTASKQTASKQSTASKQSTASK QSTASRQSTASRQSVVSKQATSALQEEETSEKKS RKVVIRGKAERLSLRKLTLEETETIYHAKLNEDHLL HAFEFIKPRSHTVWEKENVKLHCSIAGWPEPRV TWYKNQVPINVHANPGKYIIESRYGMHTLEINAC DFEDTAQYRASAMNVKGELSA YASVVVKRYKG EFDETRFBAGASTMPLSFGVTPYGYASRFEIHFD DKFDVSGREGETMSLGCRVVITPEIKHFQPEIQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				<p>WYRNGVPLSPSKWVQTLWSGERATLTFSHLNKE DEGLYTIRVRMGEYYEQYSAYVFVRDADAEIEG APAAPLDVKCLEANKDYIIISWKQPAVDGGSPIL GYFIDKCEVGTDSWSQCNDTPVKFARFPVTGLIE GRSYIFRVRAVNKMIGIFPSRVSEPVAAALDPAEK ARLKS/PPLSTLDWTIVIVTEEPSEGIVPGPPTDLS VTEATRSYVVLWSKPPGQRGHEGIMYFVEKCEA GTENWQQRVNTLTPVKSPPRFALFDLAEGKSYCFR VRCNSAGVGEPSEATEVTVVGDKLDIPKAPGKI IPSRNTDTSVVVSWEESKDAKELVGYIEANVA GSGKWEPCNNNPVKTHRFTCHGLVTGQSYIFRV RAVNAAGLSEYSQDSEAEVKAIAAPPSPPCDITC LESFRDSMVLGWKQPKIGGAIEITGYVNYREV IDGVPGKWREANVKA VSEEAYKISNLKENMVY QFQVAAMNMAGLGAPSAVSECFKCEEWTIAVP GPPHSLKCSEVRKDSLVLQWKPPVHSGRTPVTG YFVDLKEAKAKEDQWRGLNEAAIKNVYLKVRG LKEGVSYVFRVRINQAGVGKPSDLGPVVAET RPGTKEVVVNVDGDDGVISLNFECDKMTPKSEFS WSKDYYSTEDSPRLEVESKGNKTKMTFKDLGM DDLGIYSCDVTDTGLASSYLIDEEELKRLLALSH EHKFPTVPVKSELA VEILEKGQVRFWMQAEKLS GNAKVNYIFNEKGIFEGPKYKMHIDRNTGIEMF MEKLQDEDEGTYTFQLQDGKATNHSTVVLVGD VFKKLQKEAEFQRQEWIRKQGPHEVYLSWEVT GECNVLLKCKVANIKKETHIVWYKDEREISVDE KHDFKDGICTLLITEFSKKDAGIYEVILKDDRKG DKSRLKLVDFAFKELMMEVCKKIALSATDLKIQ STAEGIQLYSFVTYYVEDLKVNVSHNGSAIRYSD RVKTGVTGEIWLQINEPTNDKGKYVMELFDG EYHQKTVDLSCLYDEAYAEFQRLKQAAIAEK NRARVLGGLPDVVTIQEGKALNLTENVWGLTP EVSWLKNEKALASDDHCNLKFEAGRATAYFTING VSTADSGKYGLVVKNKYGSETSDFTVSVFPIEBE ARMAALESLESLKGGKKAK</p>
3458	A	3963	827	<p>LSRSSSDNNTNTLGRNVMSTATSPLMGAQSFPNL TTPGTTSTVTMTSSVTSSNVATATTVLVSGQS LSNLTITSLTSTSSSDTGQEAESLYDFLDSCRA STLLAELDDDEDLPEPDEEDDENEDDNQEDQEY EEVMILRRPSLQRRAGSRSDVTHAVTSQLPQVP AGAGSRPIGEQEEEEYETKGGRRRTWDDDYVLK RQFSALVPAFDPRPGRITNVQQTIDLEIPPPGTPHS ELLEBEVECTPSRLALTLKVTGLGTREVELPLTN FRSTIFYVYVQKLLQLSCNGNVKSDKLRRRIWEPTY TIMYREMKDSKKEKENGKMGCSIEHVEQYLG TDELPKNDLITYLQKNADAAFLRHWWKLTGTNKS IRKNRNCSQLIAAYWDLGVEHGTKSGLNQGAIST LQSSDILNLTKQPQAKAGNGQNSCGVEDVLQL LRLITYVASDPYSRISQEDGDEQPQFTFPDPDEFTS/ KKITTKILQQIEPLALASGALPDWCEQLTSKCPF LIPFETRQLYFTCTAFGASRAIVWLQNRREATVE RTRTTSSVRRDDPGEFRVGRLLKHERVKVPRGESL MEWAENVMQIHADRKSVEVEFLGEEGTGLGPT LEFYALVAAEFQRTDLGAWLCDDNFPDDESRHV DLGGGLKPPGYVQRSCGLFTAPFPQDSDELERI</p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TKLFHFLGIFLAKCIQDNRLVDLPISKPFKLMCM GDIKSNMSKLIYESRGDRDLHCTESQSEASTE HDSLSVGSFEEDSKSEFILDPPKPKPPAWFNGILT WEDFELVNPHRARFLKEIKDLAIKRRQILSNKGL SEDEKNTKLQELVLKNPSGSGPPLSIEDLGLNFQF CPSSRIYGFTAVIDLKPSGEDEMITMDNAEEYVDL MFDFCMHTGIQKQMEAFRDGFNKVFPMEKLSF SHEEVQMILCGNQSPSWAAEDIINYTEPKLGYTR DSPGFLRFVRVLCGMSSDERKAFLQFTTGCSTLP PGGLANLHPRLTVVRKVDATDASYPVNTCVHY LKLPEYSSEIMRERLLAATMEKGFHLN
3459	A	88	603	SCGPRGLASLGLGFSGRCDQNGKRSIDGPEAQA EACSGERTYQELLVNQNPQAQPLASRRRLTRKLYK CIKKAVKQKQIRRGVKEVQKFVNKGEKGIMVLA GDTLPYEVYCHLPVMCEDRNLPVYVPSKTDLGA AAGSKRPTCVIMVKPHEEYQEA YDECLEEVQSL PLPL
3460	A	139	1997	QVTNMSDKSELKAELERKKQRLAQIREKKRKE EERKKKETDQKKEAVAPVQESDLEKKRREA LLQSMGLTPESPIVPPMSPSSKSVSTPSEAGSQD SGDGAVGSRRGPIKLGMAKITQVDFPPREIVTYT KETQTPVMAQPKEDDEEDDDVVAPKPIEPPEEK TLKKDEENDSKAPPHELTEEEKQQLHSEEFLSFF DHSTRIVERALSEQINIFFDYSGRDF/ENDKEGEIQ AGAKLSLNRQFFDER\WSKASGWVSCLDWSSQ YPPELLVASYNNNEDAPHEPDGVALVWNMKYK KTTPEYVFHCQSAVMSATFAKFHPNLVVG GTYS GQIVLWDNRSNKRTPVQRTPLSAAAHTHPVYCV NVVGTQNAHNLSISTDGKICSWSLDMLSH PQDS MELVHKQSKAVAVTSMSPVGDVNNFVVGSEE GSVYTACRHGSKAGISEMFEHQGPITGIHCHAA VGAVDFSHLYVTSSFDWTVKLWTTKNKNPLYSF EDNAGYVYDVMWSPTHPALFACVDGMGRDL WNLNNDTEVPTASISVEGNEPALNRVRWTHSGRE IAGVDSEGQIVTYDVGEQIAVPRNDEWARFGRIL AEINANRADAEDEEAATRIPA
3461	A	139	1997	QVTNMSDKSELKAELERKKQRLAQIREKKRKE EERKKKETDQKKEAVAPVQESDLEKKRREA LLQSMGLTPESPIVPPMSPSSKSVSTPSEAGSQD SGDGAVGSRRGPIKLGMAKITQVDFPPREIVTYT KETQTPVMAQPKEDDEEDDDVVAPKPIEPPEEK TLKKDEENDSKAPPHELTEEEKQQLHSEEFLSFF DHSTRIVERALSEQINIFFDYSGRDF/ENDKEGEIQ AGAKLSLNRQFFDER\WSKASGWVSCLDWSSQ YPPELLVASYNNNEDAPHEPDGVALVWNMKYK KTTPEYVFHCQSAVMSATFAKFHPNLVVG GTYS GQIVLWDNRSNKRTPVQRTPLSAAAHTHPVYCV NVVGTQNAHNLSISTDGKICSWSLDMLSH PQDS MELVHKQSKAVAVTSMSPVGDVNNFVVGSEE GSVYTACRHGSKAGISEMFEHQGPITGIHCHAA VGAVDFSHLYVTSSFDWTVKLWTTKNKNPLYSF EDNAGYVYDVMWSPTHPALFACVDGMGRDL WNLNNDTEVPTASISVEGNEPALNRVRWTHSGRE IAGVDSEGQIVTYDVGEQIAVPRNDEWARFGRIL AEINANRADAEDEEAATRIPA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
3462	A	2	2643	TAPEFSRSTHASAHASVARVLRNREIAQLKKEQR RQEFQIRALESQKRQQEMVLRRTQEVSA LRRL AKPMSESVAGRAGLKPPMLDSGAEVSA STTSSE AESGARSVSSIVRQWNRKINHFLGDHPAPT VNGT RPARKKFQKKGASQSFSKAARLKWQSLERRIIDI VMQRM TIVNLEADMERLIKKREELFLLQEALRR KRERLQAESP EEEKGLQELAEIEVLAANIDYIND GITDCQATTIVQLEETKEELDSTDTSVVSSCSLAE ARLLLDNFLKASIDKGLQVAQKEAQIRLLEGRLR QTD MAGSSQNHL LLDALREKAEAHPELQALIYN VQQENGYASTDEEISEFSEGSFSQSFTMKGSTSH DDFKFKSEPKLSAQMKAVSAECLGPPLDISTKNI TKSLASLVEIKEDGVGFSVRDPYRDRVSRTVSL PTRGSTFPQRATETSPLTRRKS YDRGQPIRSTD VGFTPPSSPPTRPNDNRNVSRLTSNQSQGSALD KSDDSDSSLSEVLRGIISPVGGA KGARTAPLQCV SMAEGHTKPI LCLDATDELLFTGSKDRSCKMWN LVTGQEIAALKGHPNNVSIKYCSHSGLVFSVST SYIKVWDIRDSAKCIRTLTSSGQVISGDACAATST RAITSAQGEHQINQIALSPSGTMLYAASGN AVRI WELSRFPVPGKLTGHIGPVMCLTVTQTASQHD L VVTGSKDHYVKMFELGECVTGTIGPTHNFEPHP YDGI ECLAIQGDILFSGSRDNGIKKWDLDQQELIQ QIPNAHKDWVCALAFIPGRPMLLSACRAGVIKV WNVDNFTPIGEIKGHDSPINAICTNAKHIFTASSG CRVKVWNYVPGLTPCLPRRVLAIKGRATTLP
3463	A	198	3146	SGERPEPGNMATCIGEKIEDFKVGNLLGKG SFA GVYRAESIHTGLEVAIKMIDKKAMYKAGMVQR VQNEVKIHCQLKHP SILELYNYFEDSNYVYLVE MCHNGEMNRVYLNKRVKPFSENEARHFMHQITG MLYLHSHCGT HPTLTLNLLLTRT NIKIADEGL AQ LKMPHEKHYYTL CGTPNYISPEIATK TQLE SDVWSLGCMFYTL LIGRPPFD TDTVKNTLNKVV LADYEMPTFLSIEAKDLIHLRRNPADRLSLSSV LDHPFMSRNSSTKSKDLGTVEDSIDSGHATISTAI TASSSTSISGSLFDKRRL LIGQPLPNKMTVFPKNK SS TDFSSSGDGN SFYTQWGNQETSNSGRGRVIQD AEERPHSRYLRRAYSSDRSGTSNSQSQA KTYTM ERCHSAEMLSVSKRSGGGENEERYSP TDNNANIF NFFKEKTSSSSGSFERPDNNQALSNHLCPGKTPFP FADPTPQTETVQQWFGNLQINAH LRKTTBYDSIS PNRDFQGH PDLQKDTSKNAWTDTKVKKNSDAS DNAHSV KQNTMKYMTALH SKPEIIQECVFGS DPLSEQSKTRGMEPPWGYQNR TLR SITSPLVAHR LKPIRQKTKKAVVSILDSBEVCVELVKEYASQ EY VKEVLQISSDGN TITTYYPNGGARGFPLA DRPPSP TDNISRYSF DNLPEKYWRKYQYASRFVQLVRS KSPKITYFTRYAKCILMENSPGADFEVWFYDGV KIHKTEDFIQVIEKTGKSYTLKSESEVNSLK EEEK MYMDHANEGHRICLALESIISEEEKTRSAFFPII IGRKPGSTSSPKALSPPPSVDSNYPTRDRASFNRM VMHSAASPTQAPILNPSMVTNEGLGLTTTASGTD ISSNSLKDCLPKSAQLLKS VFVKNVGWATQQLTS GAVVWVQFNDGSQLV VQAGVSSISYTS PNGQITTR VYGENEKL PDYIKQKLQCLSSILLMFSNPTPNFH

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
3464	A	14	348	AVRTVSGTSLGPRSHSRSPGRCHCFSAVTFSSPRL AASEAPDPMEEWDVPMKKEVESLKYQLAFQR EMASKTIPPELLKWIEDGIPKDPFLNPDLMKNPNFW VIEKGKCTIL
3465	A	5537	405	VRKLDREVRGAWWRGAWARHPRQEAGEHAKR RKGHAETPRGRRKGRAGRSAAAVGELRPARRSL ETSRAAAAMAKDSPSPPLGASPKKPGCSSPAAAV LENQRRELEKLRAELEAERAGWRAERRRFAARE RQLREEAERERRQLADRLRSKWEAQSRRELRLQ QEEMQREREAEIRQLLRWKEAEQRQLQLLHRE RDGVVRQARELQRQLAEELVNRGHCSRPGASEV SAAQCRCLQEVLALRLWQTDGEQAARIRYLQ AALEVERQLFLKYLAHFRGHPALSGSPDPQAVH SLEEPLPQTSSGSCHAPKPACQLGSLDSLAEVG VRSRSLGLVSSACSSSPDGLLSTHASSLDCFAPAC SRSLDSTRSLPKASKSEERPSSPDSTPGSRRLSPP PSPLPPPPPSAHRKLSNPRGEGSESQPCVLTSP PPGLGHHELKLNWLLAKALWVLARRCYTLQEE NKQLRRAGCPYQADEKVKRLKVKRAELTGLAR RLADRARELQETNLRAVSAPIPGESCAGLELCQV FARQRARDLSEQASAPLAKDKQIEELRQECHLLQ ARVASGPCSDLHTGRGGPCTQWLNVRDLDRQL RESQREVLRLQRQLMLQQNGGAWPEAGGQSA TCEEVRRQMLALERELDQRRRECQELGAQAAPA RRRGEEAETQLQAALLKNAWLAENGRLQAKT DWVRKVEAENSEVRGHLGRACQERDASGLIAEQ LLQQAARGQDRQQQLQRPQKALCDLHPSWKEL QALQCRPGHPPEQPWETSQMPESQVKGSRPKF HARAEDYAVSQPNRDIQEKREASLEESPVALGES ASVPQVSETVPASQPSKKTSSQSNSSSEGSMWA TVPSPTLDRDTASEVADLEPDSISAEELRLA APAAPKALTFMAQYNINPFEGPNDHPLEPLFLIA GDYIYIFGDMDEDFYEGELEDGRRGLVTSNFVE QIPDSYIPGCLPAKSPDLGPSQLPAGQDEALEES LLSGKAQGVVDRGLCQMVRVGSKTEVATEILD KTEACQLGLLQSMGKQGLSRPLLGTKGVLRMAP MQLHLQNVATATSANITWVYSSHRHPHVYLLD REHALTPAGVSCYTFQGLCPGTHYRAREVRLP RDLLQVYWGTMSTVTFDILLAGPPYPPLDVLV ERHASPGVLVVS WLPVTIDSAGSSNGVQVTGYA VYADGLKVCEVADATAGSTLLEFSQLQVPLTWQ KVSVRTMSLCGESLDSVPAQIPEDFFMCHRWPET PPFSYTCGDPSTYRVTTFVCPQKLSLAPPSAKASP HNPGSCGEPQAKFLEAFFEPPRRQSPVSNLGE GECPSGAGSQAQELAEA WEGCRKDLLFQKSPQ NHRPPSVSDQTGEKENCYQHMGTSKSPAPGFTHL RTECGPRKEPCQEKALERVLRQKQDAQGFTTP QLGASQQYASDFHNVLKEEQALCLDLWGTER EERREPEPHSRQQALGVKRG CQLHEPSSALCPA PSAKVIKMPRGGPQQLGTGANTPARVFVALSDY NPLVMSANLKAABEELVFQKRQLLRVWGSQDT HDFYLSECNRQVGNIPGRLVAEMEVGTEQTD RR WRSPAQGHLPVAHLEDFQGLTIPQGSSLVLQGN SKRLPLWTPKIMIAALDYDPGDGQMGGQKGRL ALRAGDVVMVY\GPMDDQGFFYGGELGGHRL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				VPANLRKMKSSQGH
3466	A	1	1111	MSKPPDLLRLRLRGAPRQRVCTLFHIGFKFTFFVSI MIYWHVVGEPEKEKGQLYNLP AEIPCPITLTPPTPP SHGPTPGNIFFLETSDRTNPNFLFMCVSESAARTH PESHVLVLMKGLPGGNASLPRHLGISLLSCFPNV QMLPLDLRELFRDITPLADWYAAVQGRWEPYLL PVLSDASRIALMWKFGGIYLDITDFIVLKNLRNLT NVLGTQSRVYVLNGAFLAFERRHEFMALCMRDFV DHYNGWIWGHQGPQLLTRVFKKWCSSRLAESR ACRGVTTLPEAFYPIPWQDWKKYFEDINPEELP RLLSATYAVHVWNKKSQGTRFEATSRALLAQLH ARYCPTTHE/DHENVLVKGPAGHLNLLLMGHW
3467	A	1	2175	MAKVILKQSKQCKNLLTCKVAQVCPVCGCLHC YFWWLSGLESRRPSSPLIDIKPIEFGLSAKKEPIQ PSVLRRTYNPDDYFRKFEPHLYSLDSNSDDVDSL TDEILSKYQLGMLHFSTQYDLLHNHLLTVRVIEA RDLPPISHDGSRQDMAHSNPYVKICLLPDQKNS KQTGVKRKTQKPVFEERYTFEIPFLEAQRRTLLL TVVDFDKFSRHCVIGKVSVP LCEVDLVKGGHW WKAHDSQFSAPGLPADQOFFADLFSGLVLNPQL LGRVWFASQPASLPVGSCLDFPRLDIVLRGEYG NLLEAKQQLRVEGEMLFIPARAANLPVNNKPV LLSLVFAPTWLGLSFYDSRTTSLHPARQIQLPSL QRGEGEAMLSALTLSRSPLEQNIIQLVLSLLHL CGSVVNMPPGNSQPRGDFLYHSICTWVQDNYAQ PLTRESVAQFFNITPNHLSKLF AQHGTMRFIEYVR WVRMAKARMILQKYHLSIHEVAQRCPDSDYF CRVFRQFGMDYVDILQIHRWDYNTPIEETLEAL NDVVKAGKARYIGASSMHASQFAQALELQKH GWAQFVSMQDHYNLIYREEREMIPLCYQEGV AVIPWSPLARGLTRPWCEETARL...EVCKN YKESDENDAQIAERL...SELGATEAQVALAW LLSKPGIAAPIIGTSREEQLDELLNAVDITLKPEQI AELETYPKPHPVVGFK
3468	A	147	3209	ALPLPLPTLYPGMSRRKQRKPQQLISDCEGPSASE NGDASEEDHPQVCAKCCAQFTDPTFLAHQNA STDPPVMVIIGQENPNNSSASSEPRPEGHNNPQ VMDTEHSNPPDSGSSVPTDPTWGPERRGEESSGH FLVAATGTAAGGGGGLILASPKLGATPLPPESTP APPPPPPPPPPGVSGHNLNILELRVLQQRQI HQMOMTEQICRQVLLGLSLGQTVGAPASPEL GTGTASSTKPLLPLFSPIKPVQTSKTLASSSSSSS SSGAETPKQAFFHLYHPLGSQHPFSAGGVGRSHK PTPAPSPALPGSTDQLIASPHLAFPSTTGLLAAQC LGAARGLEATASPGLLKPKNGSGELS YGEVMGP LEKPGGRHKCRFCAKVFSGDSALQIHLRSHTGER PYKCNVCGNRFTTRGNLKVHFHRHREKYPHVQ MNPHPVPEHLDYVITSSGLPYGMSVPPEKAEED ATPGGGVERKPLVASTTALSATESLTLLSTAGT ATAPGLPAFNKFLMKAVEPKNKADENTPPGSE GSAISGVAESSTATRMQLSKLVTSLPSWALLTNH FKSTGSFPLPLCARALGASPSSETSKLQQLVEKID RQGAVAVTSAASGAPTTAPAPSSSASSGPNQCV ICLRVLSCPRALRLHYGQHGGGERPFKCKVCGRF STRGNLRAHFVGHKASPAARAQNSCPICQKKFT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				NAVTLQQHVRMHLGGQIPNGGTALPEGGGAAQ ENGSEQSTVSGAGSFPPQQSQSPSEELSEEEEE EDEEEEDVTDEDSLGRGSESGGEKAISVRGDS EEASGAEEVGTVAATAAGKEMDSNEKTTQQS SLPPPPPPDLDQPQPMEQSSGVLGGKEEGGKP ERSSSPASALTPEGEATSVTLVEELSLQEAMRKEP GESSSRKACEVCGQAFPSQAALAEHHQKTHPKEG PLFVTCVFCRQGFLERATLKKHMLLAHHQVQPPA PHGPQNI AALSLVPGCSPSITSTGLSPFPRKDDPTI P
3469	A	3	5664	NLRPLSFALFLGDPNMANLEESFPRGGTRKIHKP EKAQQSVEQDNLFDISTEEGSTKRKKSQKGPAP TKKLKIEKRESSKSAREKFEILSVESLCEGMRILG CVKEVNELELVISLPNGLQGQFVQVTEICDAYTKK LNEQVTQEQLKDLLHLPFLFSPGMLVRCVVSSL GITDRGKKS VKLSLNPKNVNRVLSAEALKPGML LTGTVSSLEDHGVLVDIGVDGTRAFPLPKAQEY IRQKNKGAKLVKGQYLNLCIVEKVKGNGGVVLS VGHSEVSTAIATEQQSWNLNNLLPGLVVKAQVQ KVTPFGLTLNFLTFFTGVDVDFMHLDPKKAGTYFS NQAVRACILCVHPRTRVHLSLRPIFLQPRPLTR LSCQNLGAVLDDVPVQGFKKAGATFRLKDGVL AYARLSHLSDSKNVFNPEAFKPGNTHKCRIDYS QMDELALLSLRTSIEAQYLRHYHDIEPGAVVKG VLTIKSYGMLVKVGEQMRGLVPPMHLADILMK NPEKKYHIGDEVKCRVLLCDPEAKKLMMLTKKT LIESKLPVITCYADAKPGLQTHGFIRVKDYGCIV KFYNVQGLVPKHELSTEYIPDPERVFTYTGQVV KVVVLNCEPSKERMLLSFKLSSDPEPKKEPAGHS QKKKGKAINIGOLVDVKVLEKTKDGLEVA VI PHN IRALPLTSHLEEVANGPLLEHETQDILHRVL QLEQSEGRVLLCRKPALVSTVGGQDNFSEIH PGMILLIGFVKS IKDYGVFIQLPSGLSGLAPKAIMS DKFVTSTSDHFVEGQTVAAKVTVNDEEKQRMML SLRLSDCGLDLAITSLLLNLQCLEELQVYRSLM SNRDSVLIQTLAEMTPGMFLDLVVQEVLDDGSV VFSGGPVFDLVLKASRYHRAGQEVESGQKKKV ILNVDLLKLEVHVS LHQDLVN RKARKLRKGSE HQAIVQHLEKSFAIASLVETGHAAAFSLTSHLND TFRFDSEKLQVGQGVSLTLKTTEPGVTGLLLAVE GPAAKRTMRPTQKDSETVDEDEEVDPALTVGTI KKHTLSIGDMVTGTVKS IKPTHVVVTLDDGIIGCI HASHILDDVPEGTSPTTKLVGKTVTARVIGGRD MKTFKYLPISHPRFVRTIPELSVRPSELEDGHTAL NTHSVSPMEKIKQYQAGQTVTCFLKKYNVVKK WLEVEIAPDIRGRIPLLLTSLSKVLKHPDKKFRV GQALRATVVGPDSSKTFCLSLTGPHKLEEGEVA MGRVVKVTPNEGLTVSFPFGKIGTVSIFHMSDSY SETPLEDFVPQKVVR CYLSTADNVLTSLRSSRT NPETKSKVEDPEINSIQDIKEGQLLRGVGSIQPH GVFFRLGPSVVGLARYSHVSQHSPPSKKALYNKH LPEGKLLTARVLRNLNHQKNLVELSFLPGDTGKPD VLSASLEGQLTKQEERKTEAEERDQKGEKKNQK RNEKKNQKGQEEVEMPSKEKQOPQKPAQKRG GRECRESGSEQERVSKPKKAGLSEEDSLVDV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				YYREGKEEAETNVLPKEKQTKPAEAPRLQLSSG FAWNVGLDSLTPALPLAESSDSEDEKPHQATI KKSKKERELEKQKAEEKLSRTEBALMDPGRQPE SADDFDRLVLSSPNSSILWLQYMAFHLQATEIEK ARAVAERALKTISFREEQEKLNVVWVALLNLENM YGSQESLTKVFERAVQYNEPLKVFLHLADIYAKS EKFQEAGELYNRMLKRFRQEKA VWKYGAFLLR RSQAAAASHRVLQRALECLPSKEHVDVIAKFAQL EFQLGDAERAKAIFENTLSTYPKRTDVWSVYID MTIKHGSQKDVDRDIFERVIHLSLAPKRMKFFKR YLDYEQHGTEKDVQAVKAKALEYVEAKSSVL ED
3470	A	2334	1226	TAAAPVAPGTMDDATVLRKKGYIVGINLGKGSY AKVKSAYSERLKFNVAVKIIARKKTPTDFVERFL PREMDILATVNHGSIKTYEIFETSDGRIYIMELG VQGDLLFEIKCQGALHEDVARKMFRQLSSAVKY CHDLDIVHRDLKCNLLDKDFNKLSDFGFSKR CLRDSNGRIILSKTFCGSAAYAAPEVLQSIYPQPK VYDIWSLGVILYIMVCGSMPYDDSDIRKMLRIQK EHRVDFPRSKNLTCECKDLIYRMLQ/PDVS/KRLH IDEILSHSWLQPPPK/KATSSASFKEGEGKYRAE KLDTKTGLRPDHRPDHKLGAKTQHRLLVVPEN ENRMEDRLAETSRAKDHHISGAEVGKAST
3471	A	537	148	TERGAPQHPTLPLPSLTPSSVHTGQPKTTPSVILFL PSCEEPQANKATLVCLMNN/FYPGILMVTWKAD GTLITQSVEKTTPSKQSNKYVASSYLSLTPEQW RSRRSYSCQVMQEGSTVEKSVAPAEC
3472	A	1	2272	DKPTRHKTYLSSSWAKMAAAEGPVGDELWQT WLPNHVVFLRLREGLKNQSPTEAEKPASSSLPS PPPQLLTRNVVFGGLGELFLWDGEDSSFLVRLR GP.GGGGEEPAISQYQRLLCINPPLFEIYQVLLSPT QHHVALIGIKGLMVLELPKRWGKNSFEFGGKST VNCSTTPVAERFFTSSTSLTLKHAAWYPSEILDPH VLLTSDNVIRIYSLREPQTPTNVILSEAEESLV LNKGRAYTASLGETAVAFDFGPLAAVPKTLFGQ NGKDEVVAYPLYLYENGETFLTYSLLHSPGN/I WKA VGSIAHASAAEDNYGYDACA VCLPCVPN ILVIATESGMLYHCVVLEGEEDDHTSEKSWDSR IDLIPSLYVFECVELELALKLASGEDDPFDSDFSC PVKLHRDPKCPSRYHCTHEAGVHSVGLTWIHKL HKFLGSDEEDKDSLQELSTEQKCFVEHILCTKPLP CRQPAPIRGFVWVPDILGPTMICITSTYECLIWPLL STVHPASPPLLCTREDVEVAESPLRVLAEPTDSFE KHRSILQRSVANPAFLKASEKDIAAPPPEECQLLS RATQVFREQYILKQDLAKEEIQRRVKLLCDQKK KQLEDLSYCREERKSLREMAERLADKYEEAKEK QEDIMNRMKKLLHSFHSPLPVLSDSERDMKKEL QLIPDQLRHLGNAIKQVTMCKDYQQQKMEKVL SLPKPTILSA YQRKCIQSILKEEGEHIREMVKQIN DIRNHVNF
3473	A	1	2272	DKPTRHKTYLSSSWAKMAAAEGPVGDELWQT WLPNHVVFLRLREGLKNQSPTEAEKPASSSLPS PPPQLLTRNVVFGGLGELFLWDGEDSSFLVRLR GPSGGGEEPALSQYQRLLCINPPLFEIYQVLLSPT QHHVALIGIKGLMVLELPKRWGKNSFEFGGKST

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				VNCSTTPVAERFFTSSTSLTLKHAAWYPSEILDPH VLLTSDNVIRIYSLREPQTPTNVILSEAEESLV LNKGRAYTASLGETAVAFDFGPLAAMPKTLFGQ NGKDEVVAYPLIYL YENGETFLTYISLLHSPGN/I WKA VGSIAHASAAEDNYGYDACAVLCLPCVPN ILVIATESGMLYHCVVLEGEEDDHTSEKSWDSR IDLIPSLYVFECVELELALKLASGEDDPFDSDFSC PVKLHRDPKCPSTRYHCTHEAGVHSVGLTWIHKL HKFLGSDEEDKDSLQELSTEQKCFVEHILCTKPLP CRQPAPIRGFVWVPDILGPTMICITSTYECLIWPLL STVHPASPPLLCTREDVEVAESPLRVLAETPDSFE KHRSILQRSVANPAFLKASEKDIAPPPECLQLLS RATQVFREQYILKQDLAKEIQRKVLLCDQKK KQLEDLSYCREERKSLREMAERLADKYEEAKEK QEDIMNRMKKLLHSFHSFELPVLSDSERDMKKEL QLIPDQLRHLGNAIKQVTMKKDYQQQKMEKVL SLPKPTILSA YQRKCIQSILKEEGEHIREMVKQIN DIRNHVNF
3474	A	4344	2550	DRRREPERHVRVKQRTSVLNMRLRDLKIRFRGH KRDDFLDLAESPNASDTECSDEIPLKVPTSPRDS EELRDPAGPGTLMATGVQDFNRTEFDRLENEIKG HLEIALLEKHFLQEELRKLREETNAEMLRQELDR ERQRRMELEQKVQEVLKARTEEQMAQQPPKGG AQASNGAERRSQGLSSRLQKWFYERFGYVEDF RFQPEENTVETEEPLSARRLTENMRRLKRGAKPV TNFVKNLSALSDWYSVYTSIAIAFTVYMNA VWH GWAIPFLFLAILRLSLNYLIARGWRIQWSIVPEV SEPVEPPKEDLTVSEKQLVLDVAQAQNLFGK MADILEKIKNLFMWVQPEITQKLYVALWAAFLA SCFFPYRLVGLAVGLYAGIKFFLIDFIFKRCPRLR AKYDTPYLPWPLPTDPOI KERRSA TTRRI OT S SRSYVPSAPAGLGKEE WLFHSTKLGNFHEIFN LTENERPLAVCENGWRCCLINRDRKMPD YIRN GVLYVTENYLCFESSKSGSSKRNVIKLVDITDI QKYKVLVLPGSGMGIAVSTPSTQKPLVFGAMV HRDEAFETLSQYIKITSAAASGGDS
3475	A	2	1126	TAARRRQKGAAAAAETHGQAKAKSGWLKPYFY IELMESRKDITNQEELWKMKPRRNLEDDYLHK DTGETSMLKRPVLLHLHQTAAHDEFDCPSELQH TQELFPQWHLPIKIAAIIASLTFLYTLLREVHPLA TSHQQYFYKIPILVINKVLPVMSITLLALVYLPGV IAAIVQLHNGTKYKKFPHWLDKWMLTRKQFGL LSFFFAVLHAIYSLSPMRRSYRYKLLNWAYYQQ VQQNKEDALAEHDVWRMEIYVSLGIVGLAILAL LAVTSIPSVSDSLTWREFHYIQSKLGIVSLLGTHI ALIFAWNKWIDIKQFVWYTPPTFMIAVFLPIVLI FKSILFLCLRKKILKIRHGWEDVTINKTEICSQL
3476	A	143	3191	AKAPPTGESSEPEAKVLHTKRLYRAVVEAVHRL DLILCNKTAYQEVFKPENISLRNKLRELCVKLMP LHPVDYGRKAEELLWRKVYYEVIQLIKTNKKHI HSRSTLECA YRTHLVAGIGFYQHLLLYIQSHYQL ELQCCIDWTHVTDPLIGCKKPV SASGKEMDWAQ MACHRCLVYLGDL SRYQNELAGVDTELLAERFY YQALSVAPQIGMPFNQLGTLAGSKYYNVEAMY CYLRCIQSEVSFEGAYGNLRLYDKAAKMYHQL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, Δ=possible nucleotide deletion, ∇=possible nucleotide insertion)
				KKCETRKLSPGKKRCKDIKRLLVNFMYLQSLLO PKSSSVDSSELTSQCQSVLEDFNLCLFYLPSSPNLS LASEDEEEYESGYAFLPDLLIFQMVIICLMCVHSL ERAGSKQYSAAIAFTLALFSLVNHVNIRLQAELE EGENPVPAFQSDGTDEPESKEPVEKEEPPDPEPP PVTPQVGEGRKSRKFSRLSCLRRRRHPPKVGDDSD DLSEGFESDSSHDSARASEGSDSGSDKSLEGGGT AFDAETDSEMNSQESRSDLEDMEEEGTRSPITL PPRGRSEAPDSLNGPLGPSEASIASNLQAMSTQM FQTKRCFRLAPTFSNLLQPTTNPHTSASHRPCV NGDVDPKSEPASEEGSESESGESSGRSCRNERSIQ EKLQVLMAGELLPAVKVFLDWLRTPNDLIIVCA QSSQSLWNRLSVLLNLLPAAGELQESGLALCPEV QDLLEGCELPDLPSLLLPEDMALRNLPLRAAH RRFNFDTRPLLSTLEESVVRICCSFGHFARLQ GSILQFNPEVGIFVSIAQSEQESLLQQAQAQFRMA QEEARRNRLMRDMAQLRLQLEVSQLEGLQPKK AQSAMSPYLVPDQALCHHLPVIRQLATSGRFTVI IPRTVIDGLDLLKKEHPGARDGIRYLEAEFKKGN RYIRCQKEVGKSFERHKLKRODADAWTLYKILD SCKQLTLAQGAGEEDPSGMVTITGLPLDNPSVL SGPMQAALQAAAHASVDIKNVLDIFYKQWKEIG
3477	A	1	3902	MTEPRERRRGYSVPPRPEVGTQATEWRVEESNFN KIFLKDAELGRSNHLPWDPKPEDASWLPQSCL GGDAVATTGEIHEEKA WKTRALEVGQPAQRDIR RGELWGKEHGADQAIQETLEDLSSLERTLVVSES SPLGGDCQEVTTLVKYQVSEEVPSGTVIGKLSQ ELGREERRRQAGAAQVLQPLQALPIQVDSEGL LSTGRRLDREQLCRQWDPCLVSFDVLATGDLALI HVEIQVLDINDHQPRFPKGEQELEISESASLRTRIP LDRAEETCPITLHETLTHFALELVVCE ETKHLELIVVKELDRETFEELTAIDNGNPP KSGTSLVKVNVLDSDNDNFAFAESSLALEIQEDA APGILLIKLTATDPDQGPNGEVEFFLSKHMPEEV LDTFSIDAKTGQVILRRPLDYENFAEYEDVQAR DLGPNPIPAHCKVLKVLVDVNDNIPSIHVTWASQP SLVSEALPKDSFIALVMADDLDSGNNGLVHCWL SQELGHFRLKRTNGNTYMLLTNATLDREQWPK YTLTLAQDQGLQPLSAKKQLSIQSDINDNAPVF EKSRYEVSTRENNLPSLHLITIKAHDADLGINGK VSYRIQDSPVAHLVAIDSNTGEVTAQRSILNYEEM AGFEFQVIAEDSGQPMCLASSVSVWVSLLDANDN APEVVQPVLSDGKASLSVLVNASTGHLLVPIETP NGLGPAGTDTPLATHSSRPFLTTIVARDADSG ANGEPLYSIRSGNEAHLFILNPHTGQLFVNVTNA SSLIGSEWELEIVVEDQGSPLQTRALLRVMFVTS VDHLRDSARKPGALSMSMLTVICLAVLLGIFGLI LALFMSICRTEKKDNRAYNCREAESTYRQPKR PQKHIQKADIHLVPVLRGQAGEPCEVGQSHKDV DKEAMMEAGWDPCQLQAPFHLTPTLYRTLNRQ NQGAPAESREVLQDTVNLLFNHPRQRNASRENL NLPEQPATGQPRSRPLKVAGSPTGRLAGDQSGE EAPQRPPASSATLRRQRHLNGKVSPEKESGPRQI LRSLVRLSVAFAERNPVEELTVDSPPVQQISQLL SLLHQGQFQPKPNHRGNKYLAKPGGSRSAIPDITD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				GPSARAGGQTDPEQEEGPLDPEEDLSVKQLLEEE LSSLLDPSTGLALDRLSAPDPAWMARLSPLTTN YRDNVISPDAAATEEPRTFTQFGKAEAPELSPTG TRLASTFVSEMSSLLEMLLEQRSSMPVEAASEAL RRLSVCGRTLSLDLATSAAASGMKVQGDPPGGKTG TEGKSRGSSSSSRCL
3478	A	13	1620	TLPPPGNSGCHRLCFPEFELQVTKMEFSGRKWR KLRLAGDQRNASYPHCLQFYLPQPSSENISLIEFEN LAIDRVKLLKSVENLGVS YVKGTEQYQSKLESEL RKLKFSYRENLEDEYEPRRRDHISHFILRLAYCQS EELRRWFIQQEMDLLRFRSILPKDKIQDFLKDSQ LQFEAISDEEKTREQEIVASSPSLSGLKLGFEISY KIPFADALDLFRGRKVYLEDFAYVPLKDIVAAIL NEFRAKLSKALALTARSLPAVQSDERLQPLLNHL SHSYTGQDYSTQGNVGKISLDQIDLLSTKSFPFC MRQLHKALRENHHLRHGGRM QYGLFLKGIGLT LEQALQFWKQEFIKGKMDPKDFDKGYSYNIRHS FGKEGKRTDYTPFSCCLKILSNPPSQGDYHGCFFR HSDPELLKQKLQSYKISPGGISQILDVKGTHYQ VACQKYFEMIHTVDDCGFSLSHPNQYFCESQRI LNGGKDIKKEPIQPETPQPKPSVQKTKDASSALA SLNSSLEMDMEGLEDYFSEDS
3479	A	698	138	RPELELWRLRSRWRPLGVPRRCHRRNWKEPVR AQPLSVTVWAPRCQRP/QPPAPEPSSPNAAVPEAI PTPRAAASAALELPLGPAPVSVAPQAEAEARSTP GPAGSRLGPETFRQRFQRYQDAAGPREAFRQL REL/SPRQWLRPDNRKTKEQIVEMLVQEQLLAILP EAARARRIRRRTDVRTG
3480	A	117	2226	RRGSRSRGPFAEPAAPGGLCSSSEEKTEEGGMAV GLCKAMSQGLVTRDVALDFSQEEWEWLKPSQ KELNRDVMLENFELVVLGLSGLKLNLSLQ GKEFWMVERKMSQGHCADWESWWEIEL TFIDEDEISQEMVMERLASHGLECSSFREAWKY KGEFBLHQGNAERHFMQVTA VKEISTGKRDNEF SN/TWEKHTPEISIFNTTESPTIQVHKFDIYDKLF PQNSVII EYKRLHAEKESLIGNECEEFNQSTYLSK DIGIPPGEKPYESHDFSKLLSFHSLFTQHQTTHFG KLPHGYDECDAFSCYSFFTQPORIHSGEKPYAC NDCGKA FSHDFFLSEHQRT HIGEKPYECKECNKA FRQSAHLAQHQRIHTGEKPFACNECGKA FSRYAF LVEHQRIHTGEKPYECKECNKA FRQSAHLNQHQ RIHTGEKPYECNQCGKA FSRRIALTLHQRIHTGE KPFKCSECGKTFGYRSHLNQHQRIHTGEKPYECI KCGKFFRTDSQLNRHRIHTGERPFECCKGKAF SDALVLIHKKRSHAGEKPYECNCKGKA FSCGSY LNQHQRIHTGEKPYECSECGKA FHQILSLRLHQRI HAGEKPYKCNESQVRVRS ELA VSRGLTTKPADT GPDSTLNAAKVAEPARAGTEAALRPALSVAESA TSLGPLHQGRRFPPEAPAAHPGGTGFTVCAS
3481	A	2	1522	ASRHGMTPGALLMLL GALGPPLAPGVRGSEAE RLREKLFSGYDSSVRPAREVGDRVRVSVGLLAQ LISLNEKDEEMSTKVYLDLEWTDYRLSWDPAEH DGIDSLRITAESVWLPDVVLLNNNDGNFDV ALDI SVVVSSDGSVRWQPPGIYRSSCSIQVTFYFFDWQ NCTMVFSSYSYDSSEVSLQTGLGPDGQGHQEIHI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				HEGTFIENGQWENIHKPSRLIQPPGDPGRGGREGQ RQEVIFYLIIRRKPLFYLVNVIAPCILITLLAIFVY LPPDAGEKMGLSIFALLTLTVFLLLLADKVPETSL SVPIIKYLMFTMVLVTFSVLSVVVLNLHHRSPH THQMPLWVRQIFIHKLPLYLRLLKRPKPERDLMPE PPHCSSPGSGWGRGTDEYFIRKPPSDFLPKPNRF QPELSAPDLRRFIDGPNRAVALLPELREVVSISYI ARQLQEEDHDALKEDWQFVAMVVDRLFLWTF IIFTSVGT/LVIFLDATYHLPDPDFP
3482	A	1273	172	ERWDSGGADAEWYALADWTA/VWLP RSDFYTR LQTGEGHVPALRLPAGMPPDSPRELVPKQAPCSP SDPALPWT/LGHGNQPPAVVPEPQGPMPAGVAA RPGRRFGVYLLYCLNPRYRVR/VYVGFTVNTARR VQQHNGGRKKGGA/GRTSGRGPWEM/VLVHGF PSSVAALRFEWA/WQHPHASRRLAHVGPRLRGET AFAFHLRVLAHMLRAPPWARLPLTLRWVRPDLR QDLC/LPPPHVLLAFGPPPAQVPRPQRRRAGPFD DAEPEPDQGDPGACCSLCAQTIQDEEGPLCCPHP GCLLRAHVICLAEFLQEPPGQLLPLEGQCPCCE KSL/WGDLIWLCQMDTEKEVEDSELEEAHWTD LLET
3483	A	230	3686	WRPWPCIDTSWNLQVAARTLRVSSAQCGLVPT MARVESPVPAARASLTGSCVLGQAMPLRGGAGP SPASHGPTHGSPDPRTCLPGRGAGGMRPHGRGA LGCCGLCSFYTCHGAAGDEIMHQDIVPLCAADIQ DQLKKRFAYLSGGRGQDGPVITFPDYPAFSEIPD KEFQNVMTYLTSLPSLQDAGIGFILVIDRRRDKW TSVKASVLRIAASFANLQVLVLRPTGFFQRTLS DIAFKFNRRDDFKMKVPVIMLSSVPDLHGVIDKSQ LTEDLGGTLDYCHSRWLCQRTAIESFALMVKQT AQMLQSW/LIAETELPNDQ/ELRVLCATTEL EDKAKED/LRALKEGHSVLSLRELQAESEPSV NQDQLDNQATVQRLLAQLNET/AAAFDEFWAKH QQKLEQCLQRHFEQGFREVKA/LPAAASQKIATF TDIGNSLAHVEHLLRDLANFQEKSGVTVRARA LSLTASSFIGNKHYAVDSIRPKCQELRHLCDOFSA EIARRRGLLSKSLELHRRLETSMKWCDEGIYLLA SQPVDKCQSQDGAEALQEIEKFLETGAENKIQE LNAIYKEYESILNQDLMEHVRKVFOKQASMEEV FHRRQASLKKLAARQTRPVQPVAPRPEALAKSP CPSPGIRRGSENSSESSEGGALRRGPYRRAKSEMSES RQGRGSAGEEESLAILRRHVMSEL/DTERAYVE ELLCVLEGYAAEMDNPLMAHLLSTGLHNKKDV LFGNMEEIYHFHNRIFLRELENYTDCPELVGRCF LERMEDFQIYEKYCQNKPRSESLWRQCSDCPFFQ ECQRKLDHKLSDSYLLKPVQRITKYQLLLKEM LKYSRNCEGAEDLQEA/SSILGILKAVNDSMHLI AITGYDGNLGD/LGKLLMQGSFSVWTDHKGHT KVKELARFKPMQRHFLHEKAVLFCKKRENGE GYEKAPSYSYKQSLNMAAVGITENVKGD/DAKKFE IWYNAREEVYIVQAPTPEIKAAWVNEIRKVLTSQ LQACREASQHRALEQSQSLPLPAPTSTSPSRGNSR NIKKLEERKTDPLSLEGYVSSAPLTKPEKGKGW SKTSHSLEAPEDDGGWSSAEBQINSSDAEEDGGL GPKKLVPKTYTVVADHEKGGPDALRVRS/SGDVV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
3484	A	208	6103	ELVQEGDEGLW VTMAQQAADKYLVDKNFINNPLAQADWAAK KLVWVPSDKSGFEPASLKEEVGEEAIVELVENGK KVKVNKDDIQKMNPPKFSKVEDMAELTCLNEAS VLNHLKERYYSGLIYTYSGLFCVVINPYKNLPYS EEIVEMYKGKKRHEMPPIYAITDTAYRSMMDQ REDQSILCTGESGAGKTENTKKVIQYLAYVASSH KSKKDQGELERQLLQANPILEAFGNAKTVKNDN SSRFGKFIRINFVDVNGYTVGANIETYLLEKSRAIRQ AKEERTFHIFYLLSGAGEHLKTDLLLEPYNKYR FLSNGHVTIPGQQDKDMFOETMEAMRIMGPIEEE QMGLLRVISGVLQLGNIVFKKERNTDQASMPDN TAAQKVSHLLGINVTDFTRGILTPRIKVGGRDYVQ KAQTKEQADFAIEALAKATYERMFRWLVLRLNK ALDKTKRQGASFIGILDIAGFEIFDLNSFEQLCINY TNEKLQQLFNHTMFILEQEYQREGIEWNFIDFG LDLQPCIDLIEKPA GPPGILALLDEECWFPKATDK SFVEKVMQEQGTHPKFQPKQLKDKADFCIIHY AGKVDYKADEWLMKNMDPLNDNIATLLHQSSD KRVSELWKDVDRIGLDQVAGMSETALPGAFKT RKGMFRTVGQLYKEQLAKLMATLRNTNPNFVR CIIPNHEKKAGKLDPHLVLDQLRCNGVLEGIRICR QGFNRRVVFQEFQRQYEILTPNSIPKGFMDGKQA CVMIKALELDSNLYRIGQSKVFFRAGVLAHLEE ERDLKITDVIIIGFQACCRGYLARKAPAKRQQQLT AMKVLQRNCAAYLKLNRWQWWRLFTKVKPLL QVSRQEEEMMAKEEELVKVREKQLAAENRLTE METLQSQLMAEKLQLEQLQAETELCAEAEELR ARLTAKKQVELEEICHDLERVEEBEERCQHLQA EKKKMQQNIQLEEEQLEEEESARQKLQLEKVT EETVVEEBEELERQNTKLAKENKLLDRIAEF ENLLEEEKSKSLAKLNKHEAMITDLERLK EEXQRQELEKTRRKLEGDSTDLSQIAELQAQVA ELKMQAKKEEELQAALARVEEEAAQKNMALK KIRELESQISELQEDLKCERVASRNKAQKQKRDG EELEALKTELEDTLSTAAQQLRSKREQEVBNI KKTLEEEAKTHEAQIQEMRQKHSQAVEELAEQL EQTKRVKANLEKAKQTLENERGELANEVQVLLQ GKGDSEHKRKKVEAQLQELQVKFNEGERVTEL ADKVTKLQVELDNVTGLLSQSDSKSSKLTKDFS ALESQLODTQELLQEENRQKLSLSTKLKQVEDE KNSVFREQLEEEEEAAHNLEKQIATLHAQVADM KKKMEDSVGCLETAEEVKRKLQKDLEGLSQRHE EKVAAYDKLEKTKTRLQQLDLDLVDLDHQRQ SACNLEKKQKKFDQLLAEKKTISAKYAEERDRA EAEAREKETKALSLARALEEAMBEQKAELERLNK QFRTEMEDLMSSKDDVGKSVHELEKSKRAIEQQ VEEMKTQLEBEDELQATEDAKLRLEVNQAM KAQFERDLQGRDEQSEEKKKQVVRQVREMEAE LEDERKQSRMAVAARKKLEMDLKDLEAHIDSA NKNRDEAIKQLRKLAQMKDCMRELDDBTRASR EEILAQAKENEKKLKSMEAEMIQLQEELAAER AKRQAQQRDELADEIANSSGKGALALEEKRRRL EARIAQLEEELEEEQGNTELINDRLKKNLQIDQI NTDNLNERSHAQKNENARQQLERQNKELKVKL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				QEMEGTVKSKYKASITALEAKIAQLEEQLDNETK ERQAACKQVRRTEKKLKDVLQVDDERRNAEQ YKDQADKASTRLKQLKROLEEAEAEAAQRANASR RKLQRELEDATETADAMNREVSSLKNKLRRGDL PFVPPRRMARKGAGDGSDEEVDGKADGAEAKP AE
3485	A	2	1782	CSTGVSKAPLTYLMSYGFELGWRKGNRAVACR EDRGGESVGMGQESILSQVHWWAEAPVEKTPGR DSEATIMSLRVHTLPTLLGAVVRPGCRELLCLLM ITVTVGPGASGVCPTACICATDIVSCTNKNLSKVP GNLFRLIKRLDLSYNRIGLLDSEWIPVSFAKLNTL ILRHNNITSISTGSFSTTPNLKCLDLSSNKLKTIVK NAVFOELKVLEVLLLYNNHISYLDPSAFGGLSQL QKLYLSGNFLTQFPMDLVYGRFKLAELMFLDVS YNRIPSMPMHHINLVPGKQLRGIYLGHNPFVCD CSLVSLLVFWYRRHFSSVMDFKNDYTCLWSDS RHSRQVLLQLQDSFMNCSDSIINGSFRLGFIHAQ VGERLMVHCDSTGNANTDFIWWGPDNRLLPEP KEMENFYVFHNGSLVIESPRFEDAGVYSCIAMNK QRLNETVDVTINVSNTVSRSHAHEAFNTAFTT LAACVASIVLVLLYLYLTPCPCCKCKTKRQKNML HQSNHSSILSPGPASDASADERKAGAGKRVVFL EPLKDTAAGQNGKVRFLPSEAVIAEGILKSTRGK SDSDSVNSVFSDFPFVAST
3486	A	357	1173	GDPRETKVFPSSRFARNTVGVSHHQSHLFHTVSR IYVEDKHKILYCEVPKAGCSNWKRLMVLNGLA SSAYNISHNAVHYGKHLKLDSDFLKGIYTRLDT YTKLVLRDPMERLVSAFRDKFDHPNSYYHPVF GKAIKKYRPNACEEALINGSGVKFEFIHYLLDS HRPVGMDIHWEKVSCLCYPCLINYDFVGKFETL EECANFYLOMIGAPK...KFP...FDRH... QVVRQY...DLTRTEKQLIYDFYLDY...FNIT PFL
3487	A	2	3281	CDKSGAVPFSTTRSPRRSPRSAGPSLSSVSPRS LWASSGLSEEHAAPLLPAWPRHPCPPSLTPGSPM AQGAMRFCSEGDCAISPPRCPRRWLPEGVPVQSP PASMYGSTGSLRRVAGPGPRGRELGRVTAPCTP LRGPPSPRVAPSPWAPSSPTGQPPPGAQSSVVIFR FVEKASVRPLNGLPAPGGLSRSWDLGGVSPRPT PALGPGSNRKLRLASTSDPLPARGGSALPGSRN LVHGPPAPPQVGADGLYSSLPNGLGDPPELRLATL FGGPADTGFLNQDGTWSSPREVSSHAQRIARAK WEFFYGLDPPSSGAKPPEQAPPSPPGVGSRQGS GVAVGRAAKYSETDLDTVPLRCYRETDIDEVLA EREEADSAIESQPSSEGPPGTAYPPAPRPGPLPGP HPSLGSNGNEDEDDDEAGGEEDVDDEVFEASEGA RPGSRMPLKSPVPFLPGTSPSADGPDSFSCVF EAI LESHRAKGTSYTSLASLEALASPGTQSPFFTFEL PPQPPAPRPDPAPAPLAPLEPDSGTSSAADGPWT QRGEEEEAEARAKLAPGREPPSPCHSEDSLGLGA APLGSEPPLSQLVSDSELDSTERLALGSTDLIS NGQKADLEAAQRLAKRLYRLDGRKADVARHL GKNNDFSKL VAGEYLKFFVTGMTLDQALRVFL KELALMGETQERERVLAHFSQR YFQCNPEALSSE DGAHTLTALMLLNTDLHGHNIGKRMT CGDFIG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				NLEGLNDGGDFPRELLKALYSSIKNEKLQWAIDE EELRRFLSELADPNPKVIKRISGGSGSGSPFLDLT PEPGAAVYKHGALVRKVHADPDCRKTPRGKRG WKSFHGILKGMILYLQKEEYKPGKALSETELKN AISIHHALATRASNYSKRPHVLYLRTADWRVFL FQAPSLEQMQSWITRINVVAAMFSAPPPAAVSS QKKFSRPLLPSAATRLSQEEQVRTHEAKLKAMA SELREHRAAQLGKKGRGKEAEEQRQKEAYLEFE KSRYSTYAALLRVKLKAGSEELDAVEAALAQAG STEDGLPSSHSSPSLQPKPSSQPRAQRHSSEPRPG AGSGRRKP
3488	A	441	1968	GTETPHCWGRGTAGLRRELDREERDGPATMS FPHFGHPYRGAFQFLASASSSTTCCESTLRVS VASGSTPAPALCCAPYDSRLLSARPELGAALGI YGAPYAAAAAASYPGYLPSPPEPSLYGALNP QYEFKEAAGSFTSSLAQPGAYYPYERTLGQYQY ERYGAVELSGAGRRKNATRETTSTLKAWLNEHR KNPYPTKGEKIMLAITKMTLTQVSTWFANARRR LKKENKMTWAPKNKGGEERKAEGGEEDSLGCL TADTKEVTASQEARGRLSDLEDLEEEEEEEEA EDEEVVATAGDRLTEFRKGAQSLPGCAAAREG RLERRECGLAAPRFSFNDPSGSEADFLSAETGSP RLTMHYPCLEKPRIWSLAHTATASAVEGAPPARP RPRSPECRMIPGQPPASARLSVPRDSACDESSCI PKAFGNPKFALQGLPLNCAPCPRRSEPVVQCQYP SGAEGSGPPAALGVSMQKTPTYRPARQLHTLCH SSLP
3489	A	718	2073	IAAYHKALSYRGHVHANNRGTNNVHFTPPSPS RGILPMNPRNMNHSQVGGIGIPSRNTSMSSSG LGSPNRRSPSIICMPKQPSRQPFVNSMSGFGMN PNAQAFGMNNSLNNIFNGTDGSEN ALADNRNRREGSONPTIFLAGRAPYVGMVTK PANEQSQDFSIHNEDFPALPGSSYKDPTSSNDDSK SNLNTSGKTTSTDGPKFPGDKSSTTQNNNQKK GIQVLPDGRVTNIPQGMVTDQFGMIGLLTFIRAA ETDPGMVHLALGSDLTTLGLNLNSPENLYPKFAS PWASSPCRQDIDFHVPSYLTNIHIRDKLEFFFS W/TAIKLGRYGEDLLFYLYYMNGGDVLLQLLAAV ELFNRDWRVYHKEERVWITRAPGMEPTMKNTY ERGTYFFDCLNWRKVAKEFHLEYDKLEERPHL PSTFNYNPAQQA
3490	A	2	2833	FVAKMATSQYFDFAQGGGPQYSTQAPTLPPTV GASYTGQPTPGMDPAVNPAFPAPAGYGGYQP HSGQDFA YGSRPQEPVPTATTMATYQDSYSYQ SAAARSYEDRPYFQSAALQSGRMTAADSGQPGT QEACGQSPHGSHTAQPQAPIVESGQPASTL SSGYTYPTATGVQPESSASIVTSYPPSYNPTCTA YTAPSYPNYDASVYSAAFPYPPAQPPIPPGPPQ QLPPPPAPAGSGSSPRADSKPPLPSKLPKAGPR QLQLHYCDICKISCAGPQTYREHLGGQKHKKE AAQKTGVQPNGPSRQVQAQLHCDLCAVSTGA DAYAAHIRGSKHQKVFKLHAKLGKPIPTLEPALA TESPPGAEEKPTSPTGPSVCASSRPALAKRPVASK ALCEGPPPEQAAGCRPQWGKPAQPKLEGPGAPT QGGSKAPAGCSDAQVGPPEYVEEVFSDEGRVL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				RFHCKLCECSFNDLNAKDLHVRGRRHRLQYRKK VNPDLPIATEPSSRARKVLEERMQRHLAEERL EQLRRWHAERRRLEEEPPQDVPPHAPPDWAQPL LMGRPESAPASAPLQPGRRPASSDDRHVMCKHATI YPTEQELLAVQRAVSHAERALKLVSDTLAEEDR GRREBEGDKRSSVAPQTRVLKGVMRVGLAKGL LLRGDRNVRLALLCSEKPTHSLRRIAQQLPRQL QMVTEDEYEVSSDPEANIVISSCEEPRMQVTISVT SPLMREDPSTDPGVEEPQADAGDVLSPKKCLESL AALRHARWFQARASGLQPCVIVIRVLRDLRRV PTW GALPAWAMELLVEKAVSSAAGPLGPGDAV RRVLECVATGILLTDGPGLDPCERDQTDALP MTLQEREDVTASAQHALRMLAFRQTHKVLGMD LLPPRHRLGARFRKRQRGPGEGEEGAGEKKRGR RGGEGLV
3491	A	2	1321	FVGDGALSGCRRGRAPRVPSMAGSLPPCVVDCG TGYTKLGYAGNTEPQFIIPSCIAIRESAKVVDQAQ RRVLRGVDDLDFFIGDEAIDKPTYATKWPIRHGII EDWDLMERFMEQVVF KYLRAEPEDHYFLMTEP PLNTPENREYLAEIMFESFNVPGLYIAVQAVLAL AASWTSRQVGERTLTGIVIDSGDGVTHVIPVAEG YVIGSCIHIPIAGRDITYFIQQLREREVGIPPEQS LETAKEIKEKYCYICPDIVKEFAKYDVPDRKWIK QYTGINAINQKKFVIDVGYERFLGPEIFFHPEFAN PDFMESISDVVDEVIQNCIDVRRPLYKNVVLGSG GSTMFRDFGRRLQRLKRVVDARLRLSEELSGG\A RIKPKPVEVQVVTHHMQRYA\WFGGSMMLASTP EFFQVCHTKKDYEYGPSICRHNPFVFGVMS
3492	A	3	2024	PNGVALLHLPAAVIPNTNYMFQDALGGRSRGS RESPAPSRAPASASLWRLVVEAKMAAHAAA AAQAAAAAQAHAADSWYLALLGFAEHFRTS SPPKIRLCVHCLQAVFPFKPPQRIEARTHLLQSGV LYHHTKNSEQARSHLEKAWLISQQIPQFEDVKFE AASLLSELYCQENSVDAAKPLLRKAIQISQQTPY WHCRLLFQLAQLHTLEKDLVSACDLLGVGAEY ARVVGSEYTRALFLLSKGMLLLMERKLQEVHPL LTLCGQIVENWQGNPIQKESLRVFFLVLQVTHYL DAGQVKSVPCKLQQLQCCIQTISTLHDDIELPSNP ADLFHWLPKEHMCVLVYLVTVMHSMQAGYLE KAQKYTDKALMQLEKLKMLDCSPILSSFQVILE HIIMCRLVTGHKATALQEISQVCQLCQQSPRLFS NHAAQLHTLLGLYCVSVNCDNAEAQFTTALR LTNHQELWAFIVTNLASVYIREGNRHQEVVLYS LLERINPDHSFPVSSHCLRAAAFYVRGLFSFFQGR YNEAKRFLRETLKMSNAEDLNRLTACSLVLLGHI FYVLGNHRESNNMVVPAMQLASKIPDMSVQLW SSALLRDLNKAACGNAMDAHEAAQMHQNFSSQL LQDHIEACSLPEHNLTWTDGPPPVQFQAQNGPN TSLASLL
3493	A	3	2024	PNGVALLHLPAAVIPNTNYMFQDALGGRSRGS RESPAPSRAPASASLWRLVVEAKMAAHAAA AAQAAAAAQAHAADSWYLALLGFAEHFRTS SPPKIRLCVHCLQAVFPFKPPQRIEARTHLLQSGV LYHHTKNSEQARSHLEKAWLISQQIPQFEDVKFE AASLLSELYCQENSVDAAKPLLRKAIQISQQTPY

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				WHCRLLFQLAQLHTLEKDLVSACDLLGVGA EY ARVVGSEYTRALFLLSKGMLLMERKLQEVHPL LTLGQIVENWQGNPIQKESLRVFFLVLQVTHYL DAGQVKSVPCLKQLQCCIQTISTLHDEILPSNP ADLFHWLPKEHMCVLVYLVTVMHSMQAGYLE KAQKYTDKALMQLEKLMDCSPILSSFQVILLE HIIMCRLVTGHKATALQEISQVCQLCQSPRLFS NHAAQLHTLLGLYCVSVN CMDNAEAQFTALR LTNHQELWAFIVTNLASVYIREGNNRHEVVLYS LLERINPDHSFPVSSHCLRAAAFYVRGLFSFFQGR YNEAKRFLRETLKMSNAEDLNRLTACSLVLLGHI FYVLGNHRESNNMVVPAMQLASKIPDMSVQLW SSALLRDLNKAACGNAMDAHEAAQMHQNFSSQQL LQDHIEACSLPEHNLTITWTDGPPPVQFQAQNGPN TSLASLL
3494	A	2	1615	VLRGQRGPAGGLAEERRRRGRNEWRIHDVTTAPF PGLVQRRSRLLIVSQVRYFLKNKVSPDLCNEDGL TALHQCCIDNFEEIVKLLLSHGANNVNAKDNE LW TPLHAAATCGHINLVKILVQYGADLLAVNSDGN MPYDLCEDEPTLDVIETCMAYQGITEKINEMRV APEQQMIADIHCMAAGQDLDWIDAQGATLLHI AGANGYLRAAELLLDHGVRVDVKDWGWEPL HAAAFWQGMQMAELLVSHGANLNARTSMDE MPIDLCEEEFEKVLLELKVHKHDVIMKSQLRHK SSLRRTSHRQAS/SVGKVVRRTPQVGTGPNL\YR KEYE/GEEAILWQSAVAEDQRTSTYNGDIRETR TDQENKDPNPRLEKIPVLLSEFPTKIPRGELDMPV ENGLRAPVSA YQYALANGDVWKVHEVPDYSM AYGNPGVADATPPWSSYKEQSPQTLLELKRQRA AAKLLSHPLSTHLGSSMARTGESSEGGKAPLIG RTSPYVSTSTSVY\TVITGDPPLLKFKANIEE 1 EEKV
3495	A	327	1078	APMADT PNGPQGAGAVQFM MTNKLDTAMWL SRLFTVYCSA L FVLPLLGLHEAASFYQ RALLANA LTSALRLHQRL F FQLSRAFLAQALLED SCHYLL YSLIFVNSYPVTMSIFPVLLFSLHAATYTKKVL DARGASNSLPLLR/SVLDKLSANQQNLKFIACNEI FLMPATVFMLFSGQGSLLQPFYTYRFLTRYSSRR NPYCRTL FNLRIVVEHIIMKPACPLFVRRRLCLQS IAFISRLAPTVP
3496	A	3	2867	SSRTREMEKEILRRQIRLLQGLIDDYKTLHG NAP APGTPAASGWQPPTYHSGRAFSARYPRPSRRGYS SHHGPSWRKKYSLVNRPPGPSDPPADHAVRPLH GARGGQPPVPQQHVLERQVQLSQGQNVVIVKVP PSKSGSASASGAQRGSLEEFEDTPWSDQRPREG GEPPRGQLQPSRPTRARGTCSVEDPLLVCQKEPG KPRMVKSVGSGVDSPREPRRTVSESIVAVKASFP SSALPPRTGVALGRKLGSHSVASCAPQLGDRRV DAGHTDQVPVPSGSGVGGPARPASGPRQAREASLV VTCRTNKFRRNNYKWWAASSKSPRVARRALS PR VAAENVCKASAGMANKVEKPQLIADPEPKPRKP ATSSKPGSAPSKYKWKASSPSASSSSSFRWQSEA GSKDHASQLSPVLSRSPSGDVRPALAHSGLKPLSG ETPLSAYKVKTRTKIIRRRGSTSLPGDKKSGTSPA ATAKSHLSLRRRQALRGKSSPVLLKKT PNKGLVQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VTKHRLCRLPPSRAHLPTKEASSLHAVRTAPTSK VIKTRYRIVKKTASPLSAPPFSLPSWRARRLS LSRSLVLNRLRPVASGGGKAQPGSPWWSKGYR CIGGVLYKVSANKLSKTSGQPSDAGSRPLLRTGR LDPAGSCSRSLASRAVQSRSLAIRQARQRREKRK EYCMYYNRFGRGNRGERCPYIHDPEKVAVCTRF VRGTCKKTDGTCPFSSHVSKEKMPVCSYFLKGI CSNSNCPYSHVYVSRKAIEVCSDFLKGYCPLGAK CKKKHTLLCPDFARRGACPRGAQCQLLHRTQKR HSRRAATSPAPGPSDATARSRVASASHGPRKPSAS QRPTRQTPSSAALTAAVAAPPHPGGSASPSSS KASSSSSSSSPPASLDHEVAPSLQEALAAACSN RLCKLPFISLQSSPSPGAQPRVRAPRAPLTKDSG KPLHIKPR
3497	A	1586	141	ATARDLGCARRIDRVVMESTPSRGLNRVHLQCR NLQEFLLGGLSPGVLDRLYGHAPATCLAVFRELPSL AKNWVMRMLFLEQLPQAVALWVKKEFSKA QEESTGLLSGLRIWHTQLLPGGLOGLILNPIFRQN LRIALLGGGKAWSDDTSQLGPDKHARDVPSLDK YAEERWEVVLHFMVGSAAVSQDLAQLLSQA GLMKSTEPGEPPCITSAGFQFLLDTPAQLWYFM LQYLQTAQSRGMDLVEILSFLFQLSFSTLTKDYS VEGMSDSLNLFLQHLREFGLVFQRKRKSRYYP T/RALAINLSSGVSGAGGTVHQPGFVIVETNYRL YAYTESELQIALIALFSEMLYPFANMVVARVTR ESVQQAIASGITAQQIIHFLRTRAHPVMLKQTPVL PPTITDQIRLWELERDRLRFTEGVLYNQFLSQVDF ELLLAHAPKLGVLVFE/NTPAKRLMVVTPAGHS DVKRFWKQKHSS
3498	A	790	190	RDI.GPAALMTASASSFSSSQGVQQPSIYSFSQITR SLFENGVAADKILCNGETAIVRASVCCCR LRGLQYIKVPVTDANDSKLFFDFPIADLIHTVS MRQGRILLNCMAGMSASLCLAYLMKYHSM SLLDAHTWA/TKSRPIK/NNGFWEQLINYEK LFNNNTVRMINSFVGNIPDIYE/DLRMMISM
3499	A	31	1586	TAGFLLAPLEMQRLLTPVKRILQLTRAVQETSLT PARLLPVAHQRFSTASAVPLAKTDTWPKDVGL ALEVYFPAQYVDQTDLEKYNNEAGKYTVGLG QTRMGFCSVQEDINSLCLTVVQRLMERIQLPWD SVGRLEVGTETIIDKSKAVKTVLMELFQDSGNTD IEGIDTTNACYGGTASLFNAANWMESSWDGRY AMVVCGDIAVYPSGNARPTGGAGAVAMLIGPK APLALERGLRGTHMENVDYFKPNLASEYPIVD GKLSIQCYLRALDRCYTSYRKKIQNQWKQAGSD RPFLLDDQLQYMFHTPFCKMVQKSLARLMFND LSASSDTQTSLYKGLEAFGGKLLEDITYTNKDLD KALLKASQDMFDDKKTASLYLSTHNGNMYTSSL YGCLASLLSHSAQELAGSRIGAFSYGSGLAASF FSRVVSQDAAPGSPLADKLVSSTSDLPKRLASRKC VSPEEFTEIMNQREQFYHKVNFSPPGDTNSLFPGT WYLERVDEQHRRKYARRPV
3500	A	185	2692	MLPTVPQSHPGPSALLLQLLLPPTSAPFPNIWS LLAAPGSITHQDLTEEAALNVTQLFLEQPPGRP PLRLEDFLGRITLLADDLFAAYFGPGSSRRFRAAL GEVSRANAAQDFLPTSRNDPDLHFAERLGQGR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ARLVGALRETVVAARALDHTLARQRLGAALHA LQDFYSHSNWVELGEQQPHPLLWPRQELQNLA QVADPTCSDCEELSCPRNWLGFLLTSGYFGTHP PKPPGKCSHGHHFDRSSSQPRGGINKDSTSPGFS PHHMLHLQAAKLALLASIQAFSLRLSRLGDRDFS RLLDITPASSLSFVLDTTGSMGEEINAQIARHL VEQRRGSPMEPVHYVLVPFHDPGFGPVFTTSDPD SFWQQLNEIHALGGGDEPEMCLSAQLALLHTTP LSDIFVFTDASPKDAFLTNQVESLTQERRCRVTL VTEDTSRVQGRARREILSPRLFEPYKAVASGSG EVIFTKDQHIRDVAIVGESMAALVTLPLDPPVV VPGQPLVFSVDGLLQKITVRIHGDISSFWIKNPAG VSQGGEEGGGLGHTRRFQFWMVTMDPPQT GTWEIQVTAEDTPGVRVQAQTSLEDLFLHFGIPME DGPHPLGLYPLTQPVAGLQTQLLVEVTGLGSRA PGDPQPHFSHVILRGVPEGAELGQVPLEPVGPPE RGLLAASLSPILLSTPRPFSLELIGQDAAGRRLHR AAPQPSTVVPVLELSGPGFLAPGSKVPLSLRIA SFGPQDLDLRTFVNPSFSLTSNLSRAHLELNEA WGRLWLEVPDSAAPDSVVMVTVTAGGREANPV PTHAFLRLLSAPAPQDRH
3501	A	1245	5815	RRAHPSHSRLSPYLSVSRDPYFFVTVSRTILTSA PAPPRTAPAPMTGALLQRGCGFLCLSLLLGC WAELGSGLEFPGAEGQWTRFPKWNACCESEMSF QLKTRSARGLVLYFDDEGFCDFLELILTRGGRLQ LSPSIFCAEPATLLADTPVNDGA WHSVIRRQFR NTTLFIDQVEAKWVEVKSKRRDMTVFSGLFVGG LPPELRAAALKLTASVREREPFKGWIRDVRVNS SQVLPVDSGEVKLDDEPPNSGGGSPCEAGEEGE GGVCLNGGVCSVDDQAVCDCSRGTGFRGKDCS QEDNNVEGLAHLMMCDQGK...IATTKGEYF CYDLSQNPIQSSSL...LSFKTLQRNGLMLHTGK ADYVNLALKNGAVSLVINLGSFAFEALVEPVNG KFNDNAWHDVKVTRNLRQHSGIGHAMVTISVD GILTTTGYTQEDYTM LGSDDDFFVVGSPSTADLP GSPVSNFMGCLKEVYKNNDRVLELSRLAKQ GDPKMKIHGVVAFKCNVATLDPITFETPESFSL PKWNAKKTGSISDFRTTEPNGLILFSHGKPRHQ KDAKHPQMIKVDFFAIEMLDGHL YLLDMGSGT IKIKALLKKVNDGEWYHVDFQRDGRSGTISVNT LRTPYTAPGESEILDDEL YLGGLPENKAGLVF PTEVWTALLNYGYVGCIRDLFIDGQSKDIRQMA EVQSTAGVKPSCSKETAKPCLSNPCKNNGMCRD GWNRYVDCSGTG YLGRSCEREATVLSYDGSM FMKIQLPVVMHTEADVSLRFRSQRAYGILMAT TSRDSADTLRLDLAGRVKLTVNLD CIRINCNSS KGPETLFAGYNLNDNEWHTVRVVRGKSLKLT VDDQAMTGQ MAGDHTRLFHNIETGHTERRY LSSVPSNFIGHLQSLTFNGMAYIDLCKNGDIDYC ELNARFGFRNIADPVTFKTKSSYVALATLQAYT SMHLFFQFKTSLDGLILYNSGDGNDFIVVELVK GYLHYVFDLGNANLIKSSNKPLNDNQWHNV MISRDTSNLHTVKIDTKITTQITAGARNLDLKS DL YIGGVAKETYKSLPKLVHAKGFGGCLASVDLN GRLPADLISDGSFSCNGTDSRRGMWKGSTTTCQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EDSCSNQGVCLQQWDGFSCDCSMTSFSGPLCND PGTTYIFSKGGGQITYKWPPNDRPSTRADRLAIGF STVQKEA VLVRVDSSSLGDYLELHHQGGKIGVK FNVGTDIAIEESNAIINDGKYHVVRFRTRSGGNA TLQVDSWPVIERYPAGRQLTIFNSQATIIIGGKEQ GQPFQQLSGLYYNGLKVLNMAAENDANIAIVG NVRLVGEVPSSMTTESTATAMQSEMSTSIMETTT TLATSTARRGKPPTKEPISQTTDDILVASAECPSD DEDIDPCEPSSGGLANPTRAGGREPYPGSAEVIRE SSSTTGMVVGIVAAAALCILLYAMYKYRNRDE GSYHVDESRYISNSAQSNAGVVKQKQPSSAKSS NKNKKNKDKEYVY
3502	A	394	72	KPAHLPTVIIMPKRKPSEGAMSDKVKA/KFELQ RRSAGLFSKPTPPKPKETRPKKDPANQRQKLPKVR KGKADA/SKEGNSPAEERCSMVQTQKVEGWRS SELPVALSF
3503	A	43	3358	SGGRGPVRVRSEQLSPSAEQVSQISQISLGRRLS SLPPPPSRALAPTRAPDTALTIMEVAEVESPLNPS CKIMTFRPSMEEFREFNKYLAYMESKGAHRAGL AKVIPPKEWKPRQCYDDIDNLLIPAPIQQMVTGQ SGLFTQYNIQKKAMTVKEFRQLANSKYCTPRY LDYEDLERKYWKNLTFVAPIYGADINGSIYDEGV DEWNIARLNTVLDVVEEECGISIEGVNTPYLYFG MWKTTFAWHTEDMDLYSINYLFHGEPKSWYAI PEHGKRLERLAQGFFPSSSQGCDALRHKMTLIS PSVLKKYGIPFDKITQEAGEFMITFPYGYHAGFN HGFNCAESTNFA TVRWIDYGVAKLCTCRKDM VKISMDIFVRKFQPDYQLWKQGKDIYTIDHTKP TPASTPEVKA WLQRRRKVRKASRSFQCARSTSK RPKADDEBEVSDEVDGAEPNPDSVTDDLKVS KSEA...K...NT...ASS...M...RMQV... KLSGNSCLSTSVTEDIR...S...YAYRSVPSISSE ADDSIPLSTGYEKPEKSD...FLSWPKSPESCSSVA ESNGVLTEGEESDVESHGNCLEPGEIPAVPSGER NSFKVPSIAEGENKTSKSWRHPLSRPPARSPMTL VKQQAPSDEELPEVLSIEEBVEETESWAKPLIHL WQTKPPNFAAEQEYNATVARMKPHCAICTLLMP YHKPDSSNEENDARWETKLDEVVTSEGKTKPLIP EMCFIYSEENIBYSPPNAFLEEDGTSLLISCAKCC VRVHASCYGPSHEICDGLWCARCKRNAWTAEC CLCNLRGGALKQTKNNKWAHVMCVAVPEVR FTNVPERTQIDVGRIPLQRLKLKCIFCRHRVKRVS GACIQCSYGRCPASFHVTCAHAAGVLMPEDDW PYVVNITCFRHKVNPVNVKSKACEKVISVGQTVIT KHNTRYYSRVMVAVTSQTFYEVMFDDGSFSRD TFPEDIVSRDCLKLGPPAEGEVVQKWPDGKLY GAKYFGSNIAHMYQVEFEDGSQIAMKREDIYTL DEELPKRVKARFVSAGRCHLGTQVNSLSSPHVS QAQQETYLGFWINSKKSQCNIFLSGTY
3504	A	1124	139	RGEEQFDAEFRFACLGFGERLQFESRLRAVHR SRAWTCYLAI RMLMATCCPSPTTTACTGPWQRA PPLRLLVQKREADSSGLAFASNSLQRRKKGLLR PVAPLRTRPPLISLPQDFRQYSSVIDVDLLPETH RRVRLHKHGS DRPLGFYIRDGMSVRVAPQGLER VPGIFISRLVRGGLAESTGLLAVSDEILEVNGIEV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=-possible nucleotide deletion, \=possible nucleotide insertion)
				AGKTLNQVTDMMVANSHNLIVTVKPANQRNN VVRGASGRLTGPPSAGPGAEPDSDDDSSDLVIE NRQPPSSNGLSQGPPCWDLHPGCRHPGTRSSLPS LDDQEQAASSGWGSRIRGDGSGFSL
3505	A	3	2898	SCRSATSQSGCGGGRSWLCSSLKMAAQPPRGIRL SALCPKFLHTNSTSHTWPFSAVAELIDNAYDPDV NAKQIWIDKTVINDHICLTFDTNGNGMTSDKLH KMLSFGFSKVTMNGHVPGLYGNGFKSGSMR LGKDAIVFTKNGESMSVGLLSQTYLAEVIKAEHV VVPVAFNKHRRQMINLAESKASLAAILHSLFSTE QKLLAELDAIGKKGTRIIWNLRSYKNATEFD KDKYDIRIPEDLDEITGKKGYKKQERMDQIAPES DYSLAYCSILYLKPRMQILRGQKVKTQLVSKS LAYIERDVYRPFKLSKTVRITFGFNCRNKDHYGI MMYHRNRLIKAYEKVGCQLRANNMGVGVVGGII ECNFLKPTHNKQDFDYTNEYRLTITALGEKLN YWNEMKVKKNTHEYPLNLPVEDIQKRDPQTWVQ CDACKWRKLPDGMQDQLEKWCNNPDPQFR NCEVPEEPEDLDVHPTYEKTYKKTNKEKFRIRQ PEMIPRINAELLFRPTALSTPSFSSPKESVSKR/RH LSEGTNSYATRLLNNHQVPPQSEPESNSLKRRLS TRSSILNAKNRRLASSQFENS VYKGDDDDDEDVII LEENSTPKPAVDHIDMKSEQSHVEQGGVQVEF VGDSEPCGQTGSTSTSSRCDQNTAATQTEVPS LVVKEETVEDEIDVRNDAVILPSCVEABAKIHE TQETTDKSADDAGCQLQELRNQLLVTEEKENY KRQCHMFTDQIKVLQQRILEMNDKYVKKETCH QSTETDAVFLLESINGKSESPDHMVSYQQALEE IERLKKQCSALQHVKAECSCSNNESKSEMDEM AVQLDDVFRQLDKCSIERDQYKSEVELLEM*KS QIRSCQEFLLKTEVEQL*STHQQTATDVSTSE*LE SVNHMDGESLKLRLRVNVGQ*LA*MI*V*PDLDLQ QVNYDVDVDEILQGVVEQMSEISST
3506	A	2	2120	RPPAAGGRYRAGGRRQAAPSRPPLPSRRRLPQG GRTRRAMDRPAAAAAGCEGGGGPNPAGGR RPPRAAGGATAGSRQPSVETLDSPTGSHVEWCK QLIAATISSQISGSVTSENVSRDYKALRDGNKLA QMEEAPLPFGESIKAIKDV MYICPFMGAVSGTL TVTDFKLYFKNVERDPHFILDVPLGVISRVEKIGA QSHGDNSCGIEIVCKDMRNRLRAYKQEBQSKLG IFENLNKHAFLPSNGQALFAPSYKEKFPINGWKV YDPVSEYKRQGLPNESWKISKINSNYEFCDTYPA IIVVPTSVKDDDLKVAFLAKGRVPVLSWIHPE SQATITRCSQPLVGPNDKRCKEDEKYLQTIMDAN AQSHKLIIFDARQNSVADTNKTKGGGYESESAYP NAELVFLEIHNIHVMRESLRKLKEIVYPSIDEARW LSNVDGTHWLEYIRMLLAGAVRIADKIESGKTSV VVHCSDGWDRTAQLTSLAMLMLDSYRTIKGFE TLVEKEWISFGHRFALRVGHGNDNHADADRSPIF LQFVDCVWQMTRQFPFAFEFNEFLITILDHLYS CLFGTFLCNCEQQRFKEDVYTKTISLWSYINSQL DEFSNPFVNYENHVLYPVASLSHLELWVNYVYV RWNPRMRPQMPIHQNLKELLAVRAELQKRVEG LQREVATRAVSSSSERGSSPSHFATSVHTLV GSSIKIRLTVLCAKNLAKKDFRLLPDPFAKIVVD
3507	A	1	2169	

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GSGQCHSTDTVKNLDPKWNQHYDLYVGKTDISI TISVWNHKKIHKKQAGFLGCVRLLSNAISRLKD TGYQRLDLCKLNPSDDTAVRGQIVVSLQTRDRIG TGGSVVDCRGLLENEGTVYEDSGPGRPLSCFME EPAPYTDSTGAAAGGGNCRFVESPSQDQRLQAQ RLRNPDVRGSLQTPQNRPHGHQSPELPEGYEQRT TVQGQVYFLHTQTGVSTWHDPRIPRDLNSVNC ELGPLPPGWEVRSTVSGRIYFVDHNNRTTQFTDP RLHHIMNHQCQLKEPSQPLPLPSEGSLEDEELPA QRYERDLVQKLKVLRLHSLQPPQAGHCRIEVS REEIFEESYRQIMKMRPKDLKKRLMVKFRGEEG LDYGGVAREWL YLLCHEMLNPYYGLFQYSTDN YMLQINPDSSINPDHLSYFHFVGRIMGLAVFHGH YINGGFTVPFYKQLLGKPIQLSDLESVDPELHKSL VWLENDITPVLDTFCVEHNAFGRILQHELKPN GRNVPVTEENKKEYVRLYVNWRFMRGIEAQFL ALQKGFNELIPQHLLKPFQKELELIGGLDKIDL NDWKSNTLRKHCVADSNTVRWFQAVETDEE RRARLLQFVTGSTRVPLQGFKALQGSTGAAGPR LFTIHLIDANTDNLKAHTCFNRIDIPPYESYEKL YEKLLTAVEETCGFAVE
3508	A	3	6388	ILYNPADLGWNPPVSSWIEKREIQTERANLTILF DKYLPCLDLTRTRFKKIPIEQSMVQMCHLLE CLLTEDIPADCPKEIYEHYVFVAAIWAFGGAMV QDQLVDYRAEFSKWWLTEFKTVKFPSQGTIFYD YIDPETKKFEPWSKLVPOFEFDPEMPLQACLVHT SETIRVCYFMERLMARQRPVMLVGTA GTGKSVL VGAKLASLDPEAYLVKNVPFNYYTTSAMLQAVL EKPLEKKAGRNYGPPGNKKLIYFIDDMNMPEVD AYGTVQPHITIRQHLDYGHVYDRSKLSLKEITNV QYVGMNPTAGSTTNRRLQRHF3VFLSTPGAD ALSST3IILTQHLKLGNFASLQKSIPPLIDLAI3F HQKIATTFLTGIFHYIFNLRFANIFQGILFSSV ECVKSTWDLIRL YLHESNRVYRDKMVEEKDFDL FDKIQTEVLKKTFFDDIEDPVEQTQSPNLYCHFAN GIGEPKYPVQSWELLTQTLVEALENHNEVNTV MDLVLFEDAMRHVCHINRILESPRGNALLVGVG GSGKQSLTRLAFAISSMDVVFQITLRKGYQIQDFK MDLASLCLKAGVKNLNTVFLMTDAQVADERFL VLINDLLASGEIPDLYSDDEVENIISNVRNEVKSQ GLVDNRENCWKFFIDRIRRLKVTLCFSPVGNKL RVRSRKFPAINCTAIHWFHEWPQQALESVSLRF LQNTGIEPTVKQSISKMAFVHTSVNQTSQSYLS NEQRYNYTTPKSFLBFIRLYQSLLRHHRKELKCK TERLENGLLKLHSTSAQVDDLKAKLAAQEVELK QKNEDADKLIQVVGVEDKVSREKAMADEEEQ KVAVIMLEVKKQKQKDCEDLAKAEFALTAAQA ALNTLNKTNLTLSFGSPPLAVSNVSAAVMVL MAPRGRVPKDRSWKAAKVTMAKVDGFLDSLIN FNKENIHENCLKAIRPYLQDPEFNPEFVATKSYA AAGLCWVINIVRFYEVFCDVEPKRQALNKATA DLTAQEKLAIAKIAHLNENLAKLTARFEKA TADKLKCCQEAETAVTISLANRLVGGLASENV RWADAVQNFKQQERTLCGDILLITAFISYLGFFT KKYRQSLLDRTWRPYLSQLKTIPIVTPALDPLRM

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				<p>LMDDADVAAWQNEGLPADRMSSVENATILINCE RWFLMVPDQLQGIKWKINKYGEDLRVTQIGQKG YLQIEQALEAGAVVLIENLEESIDPVLGPIIGRE VIKKGRFIKIGDKECEYNPKFRLILHTKLANPHYQ PELQAQATLINFTVTRDGLDQLLAAVVSMEERP DLEQLKSDLTKQONGFKITLKTLEDSSLRLSSAS GNFLGETVLVENLEITKQTAAEVEKKVQEAQVT EVKINEAREHYRPAARASLLYFIMNDLSKIHFM YQFSLKAFSIVFQKAVERAAPDESLRERVANLID SITFSVYQYTIRGLFECDKLTYLAQLTFQILLMNR EVNAVELDFLLRSPVQGTASPVEFLSHQAWGA VKVLSSMEEFSLNDRDIEGSAKSWKKFVESECPE KEKLPQEWKNTALQRLCMLRAMRPDRMTYAL RDFVEEKLGSKYVVGRAIDFATSFEEGSPATPMF FILSPGVDPLKDVESQGRKLGYTFNNQNFHNVSL GQGQEVVAEALDLAAKKGHWVILQNTLEMCS RETEFKSILFALCYFHAVVAERRKFGPQGWNRSY PFNTGDLTISVNVLYNFLEANAKVPYDDLRYLFG EIMYGGHITDDWDRRLCRTYLGEFIRPEMLEGEL SLAPGFPLPGNMDYNGYHQYIDAELPPESPPLYG LHPNAEIGFLTQTSEKLFRTVLELQPRDSQARDG AGATREEKVKALLEILERVTEFNIPELMAKVE ERTPYIVVAFQECGRMNILTREIQRSLRELEGLK GELTMTSHMENLQNALYFDMVPESWARRAYPS TAGLAAWFPDLLNRIKELEAWTGDFTMPSTVWL TGFFNPQSFLTAIMQSTARKNEWPLDQMALQCD MTKKNREEFRSPREGAYIHGLFMEGACWDTQA GIITEAKLKDLTPMPVMFIKAIPADRQDCGHVY SCPVTKTSQRDPTYVWTFNLKTKENPSKWVLA GVALLQI</p>
1522	A	3	633P	<p>ILYNPA...WPPVSSW...TERA... DKYLPTCLDTRTRFKKL...PEQSMVQMVCILLE CLLTEDIPADCPKEIYEHYF...FAAIWAFGGAMV QDQLVDYRAEFSKWWLTEFKIVKFPSQGTIFY YIDPETKKFEPWSKLVPOFEFDEMPPLQACLVHT SETIRVCYFMERLMARQRPVMLVGTAGTGKSVL VGAKLASLDPEAYLVKNVPFNYYTTSAMLQAVL EKPLEKKAGRNYGPPGNKKLIYFIDDMNMPEVD AYGTVPHTIIRQHLDYGHWDYRSKLSLKEITNV QYVSCMNPTAGSFTINPRLQRHFSVFLSFGAD ALSSIYSIILTQHLLKGNFPASLQKSIPPLIDLALAF HQKIATITFLPTGIKFHYIFNLDRDFANIFQGLFSSV ECVKSTWDLIRLYLHESNRVYRDKMVEKDFDL FDKIQTIVLKKTFDDIEDPVEQTQSPNLYCHFAN GIGEPKYMPVQSWELLTQTLVEALENHNEVNTV MDLVLFEDAMRHVCHINRILESPRGNALLVGVG GSGKQSLTRLAAFISSMDVFQITLRKGYQIQDFK MDLASLCLKAGVKNLNTVFLMTDAQVADERFL VLINDLLASGEIPDLYSDDEVENIISNVRNEVKSQ GLVDNRENCWKFFIDRIRRLKVTLCFSPVGNKL RVRSRKFFAIVNCTAIHWFHEWPQQALESVSLRF LQNTGIEPTVKQSISKFMFVHTSVNQTSQSYLS NEQRYNYTTTPKSFLFIRLYQSLLHRHRKELKCK TERLENGLLKLHSTSAQVDDLKAKLAAQVELK QKNEDADKLIQVVGVEDKVSREKAMADEEEQ</p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				KVAVIMLEVKKQKQKDCEEDLAKAEPALTAQA ALNTLNKTNLTELKSFGSPPLAVSNVSAVMVL MAPRGRVPKDRSWKAAKVTMAKVDGFLDSLIN FNKENIHENCLKAIRPYLQDPEFNPEFVATKSYA AAGLCSWVINIVRFYEVFCDVEPKRQALNKATA DLTAQEKLAIAKAKIAHLNENLAKLTARFEKA TADKLKCCQAEVTAVTISLANRLVGGLASENV RWADAVQNFKQQERTLCGDILLITAFISYLGFFT KKYRQSLDRTWRPYLSQLKTPIVTPALDPLRM LMDDADVAAWQNEGLPADRMSVENATILINCE RWPLMVDPLQGIKWIKNYGEDLRVTQIGQKG YLQIEQALEAGAVVLLENLEESIDPVLGPLLGRE VIKKGRFIKIGDKECEYNPKFRLILHTKLANPHYQ PELQAQATLINFTVTRDGLDQLLAADVSMERP DLEQLKSDLTKQQNGFKITLKTLEDSSLSSAS GNFLGETVLVENLEITKQTAAVEKKVQEAQVT EVKINEAREHYRPAAARASLLYFIMNDLSKHPM YQFSLKAFSIVFQKAVERAAPDESLRERVANLID SITFSVYQYTIRGLFECDKLTYLAQLTFQILLMNR EVNAVELDFLLRSPVQTGTASPVEFLSHQAWGA VKVLSSMEEFNLDRLDIEGSAKSWKKFVESECPE KEKLPQEWKNKTALQRLCMLRAMRPDRMTYAL RDFVEEKLGSKYVVGRLDFATSFEESGPATPMF FILSPGVDPLKDVESQGRKLGYTFNNQNFHNSVL GQGQEVVAEALDLAAKKGHWVILQNTLEMCS RETEFKSILFALCYFHAVVAERRKFGPQGWNRSY PFNTGDLTISVNVLYNFLEANAKVPYDDLRYLFG EIMYGGHITDDWDRRLCRTYLGEFIRPEMLEGEL SLAPGFPLPGNMDYNGYHQYIDAELPPESPYL LHPNAEIGFLTQTSEKLFRTVLELQPRDSQARDG ACATEEEKVKALLERFRVTDDEFTELKVF SEIPTEAFQECGRMNILTREIQRSLRELELGE GEIMTSHMENLQNALYFDMVPESWARRAYPS TAGLAAWFPDLLNRIKELEAWTGDFMPSTVWL TGFFNFQSFLTAIMQSTARKNEWPLDQMALQCD MTKKNREEFRSPREGAYIHGLFMEGACWDTQA GIITEAKLKDLTPMPVPMFIKAIPADVRQDCGHVY SCPVTKTSQRDPTYVWTFNLKTENPSKWVLA GVALLLQI
3510	A	390	3330	AAGSGSRPPAPAARKMADLAECNIKVMCRFRPL NESEVNRRGDKYIAKFQGEDTVVIASKPYAFDRVF QSSTSQEQVYNDCAKKIVKDVLEGYNGTIFAYG QTSSGKTHTMEGKLHDPEGMGIIPRIVQDIFNYTY SMDENLEFHIKVSIFYEYLDKIRDLLDVSKTNLSV HEDKNRVYPYVKGCTERFVCSPEVMDTIDEGKS NRHVAVTNMNEHSSSRSHSIFLINVKQENTQTEQK LSGKLYLVDLAGESEKVSKTGAEGAVLDEAKNIN KLSALGNVISALAEGSTYVPYRDSKMTIRLQDS LGGNCRTTIVICCSPPSSYNESETKSTLLFGQRAKTI KNTVCNVNELTAEQWKKKYEKEKEKNKILRNTI QWLENELNRWRNGETVPIDEQFDKEKANLEAFT VDKDITLTNDKPATAIGVIGNFTDAERRKCEEEIA KLYKQLDDKDEEINQQSQLVEKLKTQMLDQEEL LASTRRDQDNMQAELNRLQAENDASKEEVKEV LQALEELAVNYDQKSQEVEDKTIKEYELLSDELN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QKSATLASIDAELQKLKEMTNHQKKRAAEMMA SLKDLAEIGIAVGNNVDVKQPEGTGMIDEEFTVA RLYISKMKSEVKTVMKRCKQLESTQTESNKKME ENEKELAACQLRISQHEAKIKSLTEYLQNVQKK RQLEESVDALSEELVQLRAQEKVHEMEKEHLNK VQTANEVKQAVEQQIQSHRETHQKQISSLRDEVE AKAKLITDLQDQNKMMLEQERLRVHEKLKA TDQEKSRKLHELTVMQDRREQARQDLKGLEETV AKELQTLHNLKLFVQDLATRVKKSAEIDSDDT GGSAAQKQKISFLENNLEQLTKSAQTSWYRDN DLRCELPKLEKRLRATAERVKALESALKEAKEN ASRDRKRYQQEVDRIKEAVRSKNMARRGHSAQI AKPIRPGQHPAASPTHPSAIRGGGAFVQNSQPVA VRGGGGKKQV
3511	A	1	1757	MASVQASRRQWCYLCDLPKMPWAMVWDFSEA VCRGCNVFEGADRIELLIDAAARQLKRSHVLEGR SPGPPALKHPATKDLAAAAAQGPQLPPQAQPPQ SGTGGGVSGQDRYDRATSSGRLPLPSPALEYTLG SRLANGLGREEAVAEGARRALLGSMPLMPPGL LAAAVSGLGSRGLTAPGLSPARPLFGSDFEKEK QQRNADCLAELENEAMRGRAEEWHGRPKAVREQ LLALSACAPFNVRFKKDHGLVGRVFADFATARP PGYEFELKLFTEYPCGSGNVYAGVLAVARQMFH DALREPGKALASSGFKYLEYERRHSGGEWRQLG ELLTDGVRSFREPAPAEALPQQYPEPAPAAALCGP PPRAPSRNLAPTPTRRRKASPEPEGEAAGKMTTEE QQQRHWVAPGGPYSAETPGVPSPIAALKNVAEA LGHSPKDPGGGGGPVRAGGASPAASSTAQPPTG HRLVARNGEAEVSPTAGAEAVSGGGSGTGATPG APLCCTLCRERLEDTHFVQCPVPPEHKFCFPCSR AFIKAQGPAGEVYCPSCNCPJVGSPVPLT GEIATILAGTGVKKERD
3512	A	3	1994	NTNSSSVTNSAAGVEDLNIVQVTVPDNEKERES IEKIKQLREQVNDLFSRKFGAIGVDFPVKVPYK KITFNPGCVVIDGMPPGVVFKAPGYLEISSMRRL EAAEFIKFTVIRPLPGLSNGEYSTVGKRKIDQE GRVFOEKWERAYFFVEVQNICTCLICKRSMVSK EYNLRRHYQTNHSHYDQYMERMRDEKLHELK KGLRKYLLGLSDTECEQKQVFANPSPTQKSPVQ PVEDLAGNLWEKLRKIRSFVAYSIAIDEITDINN TTQLAIFIRGVDFNDVSEELDTVPMTGTGKSGN EISFRVEKSLKNFCINWSKLVSVASTGTPPMVDA NNGLVTKLKSRAVTFCKGAELKSICCIHPESLCA QKLKMDHVMMDVVVKSNNWICSRLNHSEFTTL LYBLDSQYGSLLYYTEIKWLSRGLVLKRFFESLE EIDSFMSRGKPLPQLSSIDWIRDLAFLVDMTMH LNALNISLQGHISQIVTQMYDLIRAFKAKLCLWET HLTRNNLAHFPTLKLVSRLNESDGLNYPKIAELK TEFQKRLSDFKLYESELTFSSPFSTKIDSVHEELQ MEVIDLQCNTVLKTKYDKVGIFEYFYKYLWGSYP KYKHHCakilSMFGSTYICEQLFSIMKLSKTKYC SQLKDSQWDSVLHIAT
3513	A	1836	513	FKSLLSVKWFCFSILVLIFLGTRCYWEMTQSRPSP DPHGRGRWEGGRSRPKGGEEGRRRTRVPGLVTAS GPGNPLPDRLGEMAGGRHRRVVGTLHLLLVAA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LPWASRGVSPSASA WPEEKNYHQPAILNSSALRQ IAEGTSISEMWQNDLQPLLIERYPGSPGSYAARQ HIMQRIQRLQADWVLEIDTFLSQTPYGYRSFSNII STLNPTAKRHLVLACHYDSKYFSHWNNRVFVG ATDSAVPCAMMLELARALDKKLLSLKTVSDSKP DLSLQLIFFDGEEAFLHWSPQDSLYGSRHLAAKM ASTPHPPGARGTSQHLHGMDLLVLLDLIGAPNPTF PNFFPNSARWFERLQAIEHELHELGLLKDHSLG RYFQNYSYGGVIQDDHIPFLRRGVPVHLIPSPFP EVWHTMDDNEENLDESTIDNLNKILQVFVLEYL HL
3514	A	1836	513	FKSLLSVKWFCSILVLIFLGTRCYWEMTQSRPSP DPHRGRWEGGRSRPKGGEGRRRTRVPGLVTAS GPGNPLPDLRGEMAGGRHRRVVGTLHLLLVAA LPWASRGVSPSASA WPEEKNYHQPAILNSSALRQ IAEGTSISEMWQNDLQPLLIERYPGSPGSYAARQ HIMQRIQRLQADWVLEIDTFLSQTPYGYRSFSNII STLNPTAKRHLVLACHYDSKYFSHWNNRVFVG ATDSAVPCAMMLELARALDKKLLSLKTVSDSKP DLSLQLIFFDGEEAFLHWSPQDSLYGSRHLAAKM ASTPHPPGARGTSQHLHGMDLLVLLDLIGAPNPTF PNFFPNSARWFERLQAIEHELHELGLLKDHSLG RYFQNYSYGGVIQDDHIPFLRRGVPVHLIPSPFP EVWHTMDDNEENLDESTIDNLNKILQVFVLEYL HL
3515	A	114	754	LCRDLTMTSSKRTKTKTKRPPQRATSNVFAF DQSQIQEFKEAFNMIDQNRDGFIDKEDLHDMLAS LGKNPTDEYLDAMMNEAPGPINFTMFLTMFGEK LNGTDPEDVIRNAFACFDEEATGTIQEDYLRELL TTVMGDRFVDEVEDELYREAPNDKKGCIFNYIE ETGGPIEKDKKITECPSPNVP ATE VFLEIFLLHGP
3516	A	1	5169	MAAAPSALLLLPPFPVLSTYRLQSRSRPSAPETDD SRVGGIMRGEKNYYFRGAAGDHGSCPTTTSPLA SALLMPSEAVSSWSSESGGGLSGGDEEDTRLLQL LRTARDPSEAFQALQAALPRRGRLGFPRRKEAL YRALGRVLVEGGSDEKRLCLQLSDVLRGQGEA GQLEEAFLALLPQLVVSLREENPALRKDALQIL HICLKRSPGEVLRTLIQQGLESTDARLRASTALLL PILLTTEDLLLGLDLTEVIISLARKLGDQETEESE TAFSALQQIGERLGQDRFQSYISRLPSALRRHYN RRLESQFGSQVPYYLELEASGFPEPLPCAVTSL NSNLKFGIIPQELHSRLLDQEDYKNRTQAVEELK QVLGKFNPSTPHSSLVGFISLLYNLLDDSNFKVV HGTLEVHLVIRLGEVQQFLGPVIAASVKVLA DNKLVIKQBYMKIFLKLMEVGPQVLCLEH LKHKHSRVREEVNICICSLLTYPSEDFDLPKLSF DLAPALVDSKRRVRQAALFAFLASSMGSGKT SILFKAVDTVELQDNGDGMNAVQARLARKTLP RLTEQGFVEYAVLMPSSAGGRSNHLAHGADTD WLLAGNRTQSAHCHCGDHVRDSMHIYGSYSPTI CTRRVLSAGKGKNKLPWENEQPGIMGENQTSTS KDIEQFSTYDFIPSAKLKLSQGMPVNDDLCSRK RVSRNLFQNSRDFNPDCLPLCAAGTTGTHQTNLS GKCAQLGFSQICGKTGSGVSDQLQFLGTTSSHQEK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				VYASLNFGSKTQQTFGSQTECTSSNGQNPSPGAY ILPSYPVSSPRTSPKHTSPLIISPKKSQDNSVNFNS WPLKSFEGLSKPKSHRRSLSAQKSSDPTGRNHHG VENSQEKPPVQLTPALVRSPPSSRRGLNGTKPVPI PRGISLLPDKADLSTVGHKKKEPDDIWKCEKDS LPIDLSELNFKDKDLQDEEMHSSRLSLRNSAAKK RAKLSGSTSDLESPDSAMKLDLTMDSPSLSSSPNI NSYSESGVYSQESLTSSLSTTPQGKRIMSDIFPTFG SKPCPTRLSSAKKKISHIAEQSPSAGSSSNPQOISS FDFTTTKALSEDSVVVVVGKGVFGSLSSAPATCSQ SVISSVENGDTFSIKQSIIEPPSGIYGRSVQQNISSYL DVENEKDAKVSISKSTYNKMRQKRKEEKELFHN KDCEKKEKNSWERMRHTGTEKMASESETPTGAI SQYKERMPSVTHSPEIMDLSELRPFSKPEIALTEA LRLLADEDWEKKIEGLNFIRCLAAHFHSEILNTKL HETNFAVVQEVKNLRSVSRAAVVCLSDLFTYL KKSMDOQLDTTVKVLLHKGESNTFIREDDVKA LRAMVNNVTPARAVVSLINGGQRYVGRKMLFF MMCHPNFEKMLEKYVPKDLPIYKDSVRNLQKQ GLGEIPLDTPSAKGRRSHTGSGVGNTRSSSVSRDA FNSAERA VTEVREVTRKSVPRNSLESAEYLKLIT GLLNAKDFRDRINGIKQLLSDTENNQDLVVGNTV KIFDAFKSRLHDSNSKVNLALETMHKMPIRLRD HLSPIINMLIPAIVDNNLNSKNPGIYAAATNVVQA LSQHVNDNYLLLQPFCTKAQFLNGKAKQDMTEKL ADIVTELYQRKPHATEQKVLVVLWHLLGNMTN SGSLPGAGGNIRTATAKLSKALFAQMGNLLNQ AASQPPHIKKSLEELDMTILNEL
3517	A	1449	252	QDLKPVLDRYLAITYLKMVFFTCNACGESVKKI QVEKHVSVCRNCEC'LCIDCGKDFWGDDYKNH VYCISEDQKYGGKQVWITLIGD/AYTQV IQKISELTPRPNVSPK/VRELLEQISAFDTPQK AKFQNWMMKNSLKVHNEILDQVWNIFSRASNSE PVNKEQDQRPLHPVANPHAEISTKVPASKVKA VEQQGEVKKNKRRERKEERQKKRKREKKELKE NHQENSRNQPKKRRKKQEQADLEAGGEEVPEA NGSAGKRSKKKKQKQKDSASEEERVVGAGKRKR RHSKVETDSKKKKMKLPEHPEGGEPEDEAPAK GKFNWKGTIKAILKQAPDNEITIKLRKKVLAQY YTVTDEHHRSEEELLVIFNKKISKNPFTKLLKDK VKLVK
3518	A	3	635	APDSNARNDHFDACSLRVQAGLSSAGPALGNSG LAALMASPSKAVIVPGNGGGDVTHGWYGVWK KELEKIPGFQCLAKNMPDPITARESIWLPFMETEL HCDEKTHIGHSSGAIAAMRYAETHRVYAIVLVSA YTSDLGDENERASGYFTRPWQWEKIKANCPTYV QFGSTDDPFLPWKEQEVADSWKPNCTNSLTV ATFRTQSFMN
3519	A	81	2277	VRETRREMAMAMSDSGASRLRRQLESGGFEARL YVKQLSQSDGDRDLQHRQRIQALAEETAQNL KRNVYQNYRQFIETAREISYLESEMYQLSHLLTE QKSSLESIPLTLLPAAAAAGAAAASGGGEGVGGGA GGRDHLRGQAGFFSTPGGASRDGSGPGGEGKQR TLTTLLEKVEGCRHLLTPGGQYLVYNGDLVEYD ADHMAQLQRVHGFLMNDCLLVATWLPQRRGM

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				YRYNALYSLDGLAVVNVKDNPPMKDMFKLLMF PENRIFQAENAKIKREWLEVEDTKRALSEKRRR EQEEAAAPRGPPQVTSKATNPFEDDEEEPAVPE VEEEKVDLSMEWIQELPEDLDVCIAQRDFEGAV DLLDKLNHYLEDKPSPPVKELRAKVEERVRL TEVLVFELSPDRSLRGGPKATRRVSQLIRLGQC TKACELFLRNRAAAVHTAIRQLRIEGATLLYHK LCHVFFTSLLETAREFEIDFAGTDSGCYSAFVVW ARSAMGMFVDAFSKQVFDSESLSTAAECVKVA KEHCQQLGDIGLDLTFIHALLVKDIQALHSYK EIIIEATKHRNSEEMWRRMNLMTPEALGKLKEE MKSCGVSNFEQYTGDDCWVNLSTVVAFTKQT MGFLEEALKLYFPELHMLVLESLVEIILVAVQHV DYSLRCEQDPEKKAQIRNASFLYETVLPVVEK RFEEGVGKPAKQLQDLRNASRLIRVNPESTTSVV
3520	A	1706	540	FVAHLAWPWRADGDMEDGVLNEGFLVKRGHIV HNWKARWFILRQNTLVYKLEGGRRVTPPKGRI LLDGCITTCPCLEYENRPLLIKLTQTSTEYFLEA CSRBE/RRDAWAFETGAJHAGQARGKVQQLHS LRNSFKLPPHISLHRIVDKMHDSNTGIRSSPNMEQ GSTYKKTFLGSSLVDWLISNSFTASRLAVALAS MLMEENFLRPVGVSRMGAIKSGDLAEQFLDDST ALYTFAESYKKKISPKEEISLSTVELSGTVVKQGY LAKQGHKRKNWKVRRFVLRKDPALHYDYPSK EENRPVGGFSLRGSLSVALEDNGVPTGVKGNVQ GNLFKVITKDDTHYIQAISKAERAEWIGSL KSLNLMNKDPEGTPDSLPSLPR
3521	A	3	3063	HASVSLSLGCPRPCADTPGPQPMQMDLRVGRPP VEPPPEPTLLALQRPQRLHHHLFLAGLQQRSVE PMRVKMLPACGATLSI.VPSLPAFSIPRHQSST PCFHLGCTPCGQSLMDITAFETQTATQEQLRQJ LHKDKSRKSAVASVVKQLAEVILKKQQAAL RTVHPNSPCLYRTLEPLETEGATRSMLSSFLPPV PSLPSDPPEHFFLKTVSEPNLKLRYKPKKSLERR KNPLLRKESAPPSLRRPAETLGDSSPSSSTPAS GCSSPNDSEHGPNPILGSEALLGQRLRLQETSVAP FALPTVSLPAILGLPAPARADSDRRTHPTLGP GPILGSPHTPLFLPHGLEPEAGGTLP SRLQPILLD PSGSHAPLLTVPLGLPLFHFAQSLMTTERLSGSG LHWPLSRTRSEPLPPSATAPPPPGPMQPRLEQLKT HVQVIKRSAPSEKPRLRQIPSAEDLETGGGGP QVDDGLEHRELGHGQPEARGPAPLQHPQVLL WEQQLRAGRLPRGSTGDTVLLPLAQGGHRPLSR AQSSPAAPASLSAPEPASQARVLSSETPARTLP TTGLIYDSVMLKHQCSCGDNRRHEHAGRIQSIW SRLQERGLRSQCECLRGRKASLEELQSVHSEH LLYGTNPLSRLKLDNGKLAGLLAQRMFVMLPCG GVGVDDTDITWNLHSSNAARWAAGSVTDLAFK VASRELKNGFAVVRPPGHHADHSTAMGFCFFNS VAIACRQLQQQSKASKILVDWDVHHGNGTQQT FYQDPSVLYISLHRHDDGNFFPGSGAVDEVGAGS GEGFNVNVAWAGGLDPPMGDPEYLAARIVVM PIAREFSPDLVLSAGFDAEGHPAPLGGYHVSA KCFGYMTQQLMNLAGGAVVLALEGGHDLTAIC DASEACVAALLGNRVDPLSEEGWKQKPNLNAIR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				SLEAVIRVHSKYWGCMQRLASCPDSWVPRVPG ADKEEVEAVTALASLSVGILAEDRPSEQLVEEEE PMNL
3522	A	9	602	KMAALGEPVRLERDICALLELEKLQRSGEVPPQ KLQALQRLVQSEFCNAVREYEHVYETVDISSP EVRANATAKATVAFAASEGHSHPRVVELPKTE EGLGFNIMGGKEQNSPIYISRIIP/GGIADRHGGLK RGDQLLSVNGVSVVEGEHHEKAVELLKAAQGV KLVVRYTPKVL EEMESRFEKMRSAKRRQQT
3523	A	645	1465	IMAETSLLEAGASAASTAAALENLQVEASCSVCL EYLKEPVIIECGHNFCKACITRWEDLERDFPCP VCRKTSRYRSLRPNRQLGSMVEIAKQLRPSSGRS GMRASAPQHHEALSIFCYEDQEA VCLICAISHTH RAHTVVPDLDATQYKEKLQKCLEALNQKLQEI TRCKSSEKKPGELKRLVESRRQQLREFEELHRR LDEEQVLLSRLEEEEDILQRLRENAHLGDKR RDLAHLAAEVEGKCLQSGFEMLKVRPLPHSPS G
3524	A	3	698	PMVRHEAGEALGAIGDPEVLEILKQYSSDPVIEV AETCQLAVRRLEWLQHHGGEPAAGPYLSVDPAP PAER/DVGRLREALDESRLPFERYRAMFALRN AGGEEAALALAEGLHCGSALFRHEVGYVLGQLQ HEAAVPQLAAALARCTENPMVRHECAEALGAIA RPACLAALQAHADDPERVVRESCKVALDMYEH ETGRAFYADGLEQLRGAPSLGPNPHPELPEDS
3525	A	1452	694	EGLQRPEYLVASAAFGQLAWGGEGRGRAGCS SSGFRDAEPLLLSCPGRNEPLKKEKRLKWKSDYP MTDQGLRSKRDEFWDTAPAFEGRKEIWDALKA AAYAAEANDHELAQAILDGASITLPHGTLCECY DELGNRYQLPIYCLSPVNLLEHTEEFSELEPEP PPSVRREFPKVLEISGKDVLSASLHGLK KQLHAQE/GTPKPSWQRVSGKLLTERTRLQET KIQKDFVIQVIINQPPPPQD
3526	A	123	5441	PGNEGLGLAADHNEDLGHLSADAPWPAVTMAP RKRSHHGLGFLCCFGGSDIPEINLRDNHPLQFME FSSPIPNAEELNIRFAELVDELDLTDKNREAMFAL PPEKKWQIYCSKKKEQEDPNKLATSWPDYYIDRI NSMAAMQSLYAFDEEBETEMRNQVVEDLKTALR TQPMRFVTRFIELEGLTCLLNFLRSMDHATCESRI HTSLIGCHALMNSQGRAHVLAQPEAISTIAQSL RTENSKTKVAVLEILGAVCLVPGGHKKVLQAML HYQVYAAERTRFQTLNELDRLSGRYRDEVNKL TAIMSFINAVLNAGAGEDNLEFRLHLRYEFLMLG IQPVIDKLQHENAILDKHLDFEMVRNEDDLEL ARRFDMVHIDTKSASQMFELIHKKLKYTEAYPC LLSVLHHCLQMPYKRNGGYFQQWQLLDRLQOI VLQDERGVDPDLAPLENFNVKNIVNMLINENEV KQWRDQAEKFRKEHMELVSRLEKEREKETL EKEEMMRTLNLNKMMDKLARESQELRQARGQVA ELVAQSELSTGPVSSPPPPGGPLTLSSMTNDL PPPPPLPFACPPPPPPPLPPGGPPTPPGAPPCFG MGLPLQDPYPSSDVPLRKKRVPQPSHPLKSNW VKLNEERVPGTVWNEIDDMQVFRILDLED FEKM FSAYQRHQELITNPSQQKELGSTEDIYLASRKVK ELSVIDGRRANQNCILLSKLKL SNEIRQAILKMD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				EQEDLAKDMLLEQLLKFIPEKSDIDLLEEHKHEIER MARADRFLEYMSRIDHYQQRQLQALFFKKKFQER LAEAKPKVEAILLASRELVRSKRLRQMLEVILAI GNFMNKGQRGGAYGFRVASLNKIADTKSSIDRN ISLLHYLIMILEKHFPDILNMPSELQHLPEAAKVN LAELEKEVGNLRRGLRAVEVELEYQRRQVREPS DKFVPVMSDFITVSSFSFSELEDQLNEARDKFAK ALMHFGEHDSKMQPDEFFGIFDTFLQAFSEARQD LEAMRRRKEEEEERRARMEAMLKEQRERERWQR QRKVLAAAGSSLEEGGEFDDLVSALRSGEVDFDKD LCKLKRSRKRSGSQALEVTRERAINRLNY
3527	A	1445	714	LLGTRMLAGQLEARDPKEGTHPEDPCPGAGAV MEKTAVAAEVLTEDCNTGEMPPLQQQIIRLHQE LGRQKSLWADVHGKLRSHIDALREQNMELREKL RALQLQRWKARKKSAASPHAGQESHTLALEPAF GKISPLSADEETIPKYAGHKNQSGHSSWGQRSSS NNSAPKPMMLKIERISSWKTPQENRDKNLSRR RQDRRA TPTGRPTPCAERRGVSEDGKVASDTCV TLHWPLGKFRFR
3528	A	484	1777	RISKIQVYYSTGYSSRKMNP TLGLAIFLAVLLTVK GLLKPSFSPRNYKALSEVQGWKQRMAAKELAR QNMDLGFKLLKKLAFYNPGRNIFLSPLSISTAFS MLCLGAQDSTLDEIKQGFNFRKMPEKDLHEGFH YIIHETQKTQDLKLSIGNTLFIDQRLQQRKFLE DAKNFYSAETILT NFQNLMAQKQINDFI/ESKTH GKINNLIENIDPGTVMLLANYIFFRARWKHEFDP NVTKEEDFFLEKNSSVKVPMFMFRSGIYQVGYDD KLSC TLEIPYQKNITAFILPDEGKLKHLEKGLQV DTFSRWKTLLSRRVVDVSVPRLHMTGTDFLKK LSYIGVSKIFEEHGD LTKIAPHRS LKVG EAVNKA EIKZMEERGTEGAAGTGAQTLPMEITLVVVTDKP YMLIYGLNIPSVLFLGRIVNPIGK
3529	A	1	5684	VSSVCHENPTEVFEDGENPPSSRSSES GFTEFIQY QADRTDIDRELSEGQGA AAPIGSTSSETETAST VGSEETIIQTSPVVTQGTATRSRKAQKTAMQCC LEYVQQLTRLNLNLYIIQNNSFSQSLATEHQDGLG REQGETSKWDRNSQGDVKEKNISKQKTSKEYLS AFLAACQLFLECSSFVYIAEGNHTSELRSEKLET DCEHVQPPQWLQ TLMNACSQASDFSQSV AISL VMDLVGLTQSVAMVTGENINSVEPAQPLSPNQG RVAVVIRPPLTQGNLRYIAEKTEFFKHVALTLWD QLGDGTPQHHQKSVELFYQLHNLVPSSICEDVI SQQ LTHKDKKIRMEAHAKFAVLWHLTRDLHINK SSSFVRSFDRSLFIMLDSLNSLDGSTSSVGQAWL NQVLQRHDIARVLEPLLLLLLHPKTQRVSVQRV QAERYWNKSPCYPGEESDKHFMQNFACSNVSQ VQLITSKGNGEKPLTMDIEINFSLTVNPLSDRLSL LSTSSSETIPMVVSDFDLPDQQIEILQSSDSGCSQSS AGDNLSEYVDPETVNAQEDSQMPKESSPDDDVQ QVFDLICKVVSGLVESASVTSQLEIEAMPKPK SDIDPDEETIKIEDDSIQQSONALLSNESSQFLSVS AEGGHECVANGISRNSSSPCISGTTHTLHDSSVAS IETKSQRSHSSIQSFKEKLEK VSEKETIVKESG KQPGAKPKVKLARKKDDDKKKSSNEKLKQTSV FFSDGLDLENWYSCGEGDISEIESDMGSPGSRKSP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				NFNIHPLYQHVLVLYLQLYDSSRTLYAFSAIKAILK TNPIAFVNAISTTSVNNA YTPQLSLLQNLARHRI SVMGKDFYSHIPVDSNHNFRSSMYIELISLCLYY MRSHYPYTHVKVTAQDLIGNRNMQMMSIEILTLL FTELAKVIESSAKGFPSFISDMLSCKKVQKVILHC LLSSIFSAQKWHSEKMAGKNLVAVEEGFSEDSLI NFSEDEFDNGSTLQSLLKVLQRLIVLEHRVMT IPEENETGFDVVSADLEHISPHQPMSTLQYLHAQ SITCQGMFLCAVIRALHQHCACKMHPQWIGLIT STLPYMGKVLQRVVSVTLQLCRNLDNLIQQYK YETGLSDSRPLWMASHPPDMILTLEGITAIHYC LLDPTTQYHQLLVSVQDKHLFEARSGILSILHMI MSSVTLLWSILHQADSSEKMTIAASASLTINLG ATKNLRRQILELLGPISMNHGVHMAAIAFVWN ERRQNKTTTRTKVIPAASEEQLLVELVRSISVM RAETVIQTVKEVLKQPPAIAKDKKHLSEVCM QFFYAYIQRIQVVPNLVDSWASLLILLKDSIQLSLP APGQFLILGVLNEFIMKNPSLENKKDQDRLQDVT HKIVDAIGAIAAGSSLEQTTWLRRNLEVKPSPKIM VDGTLNLESDVEDMLSPAMETANITPSVYSVHAL TLLSEVLAHLLDMVFYSDEKERVIPLLVNIMHYV VPYLRNHSAHNAPSyracVQLSSLSGYQYTRR AWKKEAFDLFMDPSFFQMDASCVNHWRAIMDN LMTHDKTTFRDLMTRVAVAQSSSLNLFANRDVE LEQRAMLLKRLAFAIFSEIDQYQKYLPIQERLV ESLRLPQVPTLHSQVFLFRVLLLRMSPOHLTSL WPTMITELVQVFLMEQELTADEDISRTSGPSVA GLETTYTGNGFSTSYNSQRWNLVLSACKFLD LALALPSENLPQFQMYRWAFFPEASDDSGLEVRR QGIHQREFKPYVVRALAKLLRKRAKKNPEEDNSG RTLGWEPCHLLLTCTVRSMEQLLP NSKVTSRCGGHSGSPILVFPNKDMKLENHKP CSSKARQKIEEMVEKDFLEGMIKT
3530	A	1	5684	VSSVSHENPTEVFEDGENPPSSRSSESGETFIQY QADRTDDIDRELSEGQGAAPIGSTSETETAST VGSEETIIQTPSVVTQGTATRSRKAQKTAMQCC LEYVQQFLTRLINLYIIQNNFSQSLATEHQQDLG REQGETSKWDRNSQGDVKEKNISKQKTSKEYLS AFLAACQLFLECSSFPVYIAEGNHTSELRSEKLET DCEHVQPPQWLQTLMNACQASDFSVQSV AISL VMDLVGLTQSVAMVTGENINSVEPAQPLSPNQG RVAVVIRPPLTQGNLRYIAEKTEFFKHVALTLWD QLGDGTPQHHQKSVELFYQLHNLVPSSSICEDVI SSQLTHKDKKIRMEAHAKFAVLWHLTRDLHINK SSSFVRSFDRSLFIMLDSLNSLDGSTSSVGQAWL NQVLQRHDIARVLEPLLLLLLHPKTQRVSVQRV QAERYWNKSPCYPGESDKHFMQNFACSNVSQ VQLITSKNGEKPLTMDEIENFSLTVNPLSDRLSL LSTSETIPMVVSDFDLPDQQIEILQSSDSGCSQSS AGDNLSYEVDPETVNAQEDSQMPKESSPDDDVQ QVFDLICKVVSGLVESASVTSQLEIAMPKPK SDIDPEETIKIEDDSIQSQNALLSNESSQFLSVS AEGGHECVANGISRNSSSPCISGTTHTLHDSSVAS IETKSRQRSHSSIQFSFKEKLSEKVSEKETIVKESG KQPGAKPKVKLARKKDDDKKSSNEKLKQTSV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				FFSDGLDLENWYSCGEGDISEIESDMGSPGSRKSP NFNHPLYQHVLVLYLYDSSRTLYAFSAIKAILK TNPIAFVNAISTTSVNNAAYTPQLSLLQNLARHRI SVMGKDFYSHIPVDSNHNFRSSMYEILISLCLYY MRSHYPPTHVKVTAQDLIGNRNMQMMSIEILTLL FTELAKVIESSAKGFPSFISDMLSKCKVQKVILHC LLSSIFSAQKWHSEKMAGKNLVAVEEGFSEDLSI NFSEDEFDNGSTLQSLLKVLQRLIVLEHRVMT IPEENETGDFVVSLEHISHPQMTSLQYLHAQ SITCQGMFLCAVIRALHQHCAKMHQPWIGLIT STLPYMGKVLQRVVSVTLQLCRNLNLIQYQK YETGLSDSRPLWMAIIPPDMLTLLEGITAIHYC LLDPTTQYHQLLVSDQKHLFEARSGILSIHMI MSSVTLLWSILHQADSSEKMTIAASASLTITNLG ATKNLRQQLLELLGPISMNHGVHFMAAIAFVWN ERRQNKTTTRTKVIPAASEEQLLVELVRSISVM RAETVIQTVKEVLKQPPAIAKDKKHLSEVCM QFFYAYIQRIQVPNLVDSWASLLILLKDSIQLSLP APGQFLILGVLNEFIMKNPSLENKKDQRDLDQVT HKIVDAIGAIAGSSLEQTTWLRRNLEVKPSPKIM VDGTNLESDVEDMLSPAMETANITPSVYSVHAL TLLSEVLAHLLDMVFYSDEKERVIPLLVNIMHYV VPYLRNHSAHNAPSIRACVQLLSLSGYQYTRR AWKKEAFDLFMDPSFFQMDASCVNHWRAIMDN LMTHDKTTFRDLMTRVAVAQSSSLNLFANRDVE LEQRAMLLKRLAFAIFSSEIDYQKYLQDIQERLV ESLRLPQVPTLHSQVFLFRVLLLRMSPOHLTSL WPTMITELVQVFLMEQELTADEDISRTSGPSVA GLETTYTGNGFSTSYNSQRWLNLYLSACKFLD IALALPSENLPQFQMYRWAFIPASDDSGLEVRR RTLGWEPGHLLNLTNMEQLLPFFNVLSQVF NSKVTSRCGHSGSPLYSNAFPNKMCKLENHKP CSSKARQKIEEMVEKLEFLGMIKT
3531	A	553	2470	LISPSPALSSQDPALSLKENLEDISGWGLPEARSK ESVSFKDVAVDFTQEEWGQLDSPQRALYRDVM LENYQNLLALGPLHHPDVISHLERGEFPWSMQ REVPRGPCPEWELKAVPSQQQGICKEEPAQEPIM BRPLGGAQAWGRQAGALQRSQAAPGRVTRCHG LGRPAVEEFPLRCPLFAQQRVPEGGPLDTRKNV QATEGRITKAPARLCAGENASTPSEPEKFPQVRRQ RGAGAGEGEFVCGECGKAFRQSSSLTLHRRWHS REKAYKCDECGKAFTWSTNLEHRRHTGEKPF CGECGKAFCSSSLNVHQRIHTGERPYKCSACEK AFSCSSLLSMHLRVHTGEKPYRCGECGKAQNR THLTRHHRIHTGEKPYQCGSCGKAFTCHSSLT EKIHS GDKPFKCS DCEKAFNSRLTLHQRTHTG EKPFKCADCGKGFSC HAYLLVHRRHSGEKPFK NECGKAFSSHA YLIVHRRHTGEKPFDCSQCWKA FSCHSSLIVHQRIHTGEKPYKCECGRAFSQNHCL IKHQKHSGEKSFKEKCGEMFNWSSHLTEHQRL HSEGKPLAIQFNKHLSTYYVPGSLLGAGDAGLR DVDPIDALDVAKLLCVVPPRAGRNFSLGSKPRN
3532	A	3931	317	HRELQDSPAEPAGSMPLRHWGMARGSKPVG GAQPMAMGGLKVLLHWAGPGGGEPWVTFSES

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * =Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				SLTAEVCIHIAHKVGITPPCFNLFALFDAQAQV WLPPNHILEIPRDASLMLYFRRHFRYSRNWHGM NPREPAVYRCGPPGTEASSDQTAQGMQLLDPAS FEYLFEGQKHEFVNDVASLWELSTEEIHHFKNE SLGMAFLHLCHLALRHGIPLEEVAKKTSFKDCIP RSFRRHIRQHSALTRLRLRNVRFRFLRDFQPGRLS QQMVMVKYLATLERLAPRFGTERVPVCHLRLLA QAEGEPCYIRD SGVAPTDPGPESAAGPPTHEVLV TGTGGIQWWPVEEEVNKEEGSSGSSGRNPQASL FGKKAKAHKAFGQPADRPREPLGAYFCDFRDT HVGLKEHCVSIHRQDNKCLELSLPSRAAALS FVS LVDGYFRLTADSSHYLCHEVAPRLVMSIRDGIH GPLLEPFVQAKLRPEDGLYLHJHWTSHPYRLILTV AQRSQAPDGMQSLRLRKFPQEQDGAFFVLEGWG RSFPSVRELGAALQGCLLRAGDDCFSLRRCCLPQ PGETSNLIIMRGARASPTLNLSQLSFHRVDQKEI TQLSHLGQGTTRTNVYEGRLRVEGSGDPPEGKMD DEDPLVPGRDRGQELRVVLKVLDP SHHDIALAF YETASLMSQVSHTHLAFVHGV CVRGPENIMVTE YVEHGPLDVWLRRRERGHVPMWKMVVAQQLA SALSYLENKNLVHGNVCGRNILLARLGLAEGTSP FIKLSDPGVGLGALSREERVERIPWLAPECLPGG ANSLSTAMDKWGFATLLEICFDGEAPLQSRSPS EKEHFYQRQHRLPEPSCPQLATLTSQCLTYEPTQ RPSFRITLRLTRLQPHNLADVLTVNPDSPASDPT VFHKRYLKKIRDLGEGHFGKVS LYCYDPTNDGT GEMVAVKALKADCGPQHRSGWKQEIDILRLTYH EHIKYGKCCEDQGEKSLQLVMEYVPLGSLRDYL PRHSIGLAQLLLFAQQICEGMAYLHAQHYYHRLD AARNVLLDNDRLVKIGDFGLAKAVPEGHEYRV REDGDSPEVWYAFVTLKPVKYYA... LYELL...DSSQSP...TKFLELIGIA... ELLERGERLPRPDKCPCVYHLMKNC...TEASF RPTFENLIPILKTVEHYQGGQAPS VFSVC
3533	A	182	3465	FRWLDFFRGINSQFEFGRKKENMTSPAKF... EIIAEYDTQVKEIRAQLTEQMKCLDQQCELRVQL LQDLQDFFRKKAIEIMDYSRNLEKLAERFLAKT RSTKDQQFKKDQNVLSPVNCWNLLLNQVKRES RDHTTSLDIYLNNIIPRFVQVSEDSGRLFKKSKEV GQQLQDDLMKVLNELYSVMKTYHMYNADSISA QSKLKEAEKQEEKQIGKSVKQEDRQTPRSPDSTA NVRIEEKHVRRSSVKKIEKMKERQAKYTENKL KAIKARNEYLLALEATNASVFYKYTHDLSDLIDQ CCDLGYHASLNRALRTLFLSAELNLEQSKHEGLD AIENAVENLDATSDKQRLMEMYNNVFCPPMKFB FQPHMGDMASQLCAQQPVSSELLQRCLQLQSRL STLKIENEEVKKTMEATLQTIQDIVTVEDFDVSD CFQYSNSMESVKSTVSETFMSKPSIAKRRANQQE TEQFYFTKMKEYLEGRNLITKLQAKHDLLQKTL GESQRTDCSLARRSSTVRKQDSSQAIPLVVESCIR FISRHLQHEGIFRVSGSQVEVNDIKNAFERGEDP LAGDQNDHDMDSIAGVLKLYFRGLEHPLFPKDF HDLMACVTMDNLQERALHIRKVLLVLPKTTLLI MRYLFAFLNHL SQFSEENMMDPYNLAICFGPSL MSVPEGHDQVSCQAHVNELIKTHIQHENIFFSPRE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				LEGPVYSRGGSMEDYCDSPHGETTSVEDSTQDV TAEHHTSDDECEPIEAIKFDYVGRtarelsfkk GASLLLYQRASDDWWEGRHNGIDGLIPHQYITV QDTEdgvversspkseievispeepkvtaragas CPSGGHVADIYLANINKQRKRPESGSIRKtfrsds HGLSSSLTDSSSPGVGASCRPSSQPIMSQSLPKEG PDKCSISGHGSLNSISRHSSSLKNRLDSPQIRKTAT AGRSKSFdNHRPMDPEVIAQDIEATMNSALNELR ELERQSSVKHTPDVVLDTLEPLKtSPVVAPTSEPS SPLHTQLLKDPEPAFQRSASTAGDIACAFRPVKS VKMAAPVKPPATRPKPTVFPKTNATSPGVNSST SPQSTDKSCTV
3534	A	1	2640	FRFVCPASRRPAAGLRDAASSAPRGMASEGRE PESEGIKLSADVKPFVPRFAGLNVAWLESSEACV FPSSAATYYPFVQEPVTEQKIYTEDMAFGASTFP PQYLSSEITLHPYAYSPTLDSTQNVYVPGSQY LYNQPSCYRGFQTVKHRNENTCPLPQEMKALFK KKTYDEKKTYDQKFDSEADGTISSEIKSARGS HHLIYAENSLKSDGYHKRTDRKSRIIAKNVST KPEFEFTTLDPELQGAENNMSEIQKQPKWGPVH SVSTDISLLREVVKPAAVLSKGEIVKNNPNESV TANAATNSPCTRELSTWTPMGYVVRQTLSTELS AAPKNVTSMINLKTIASSADPKNVSISSSEALSSD PSYNKEKHIIHPTQKSKASQGSdleQNEASRKNK KKKEKSTSKYEVLTVQEPPIEDAEFPNLAVAS ERRDRIETPKFQSKQPDNFKNNVKKSQLPVQL DLGGMLTALEKKQHSQHAQSSKPVVSVGAV PVLSEKASGERGRRMSQMKTPHNPLDSSAPLM KKGKQREIPKAKKPTSLKKIILKERQERKQRLQE NAVSPAFTSDDTQDGEsggDDQFPEQAELSGPEG MDELSTTSVEDKSEEPGTTELQRDTEASHL * FN HITTPATTEPRFRDYCSQMLSKEVDACVTDLLKE LVRFQORMYQKDPVKAkTKRRLVLGLREVVKH LKLKLLKCVIISPNCCKIQSKGGLDDTLHTIDYA CEQNIPFVTAENRKALGRSLNKAVPVSVVGIFS DGAQDQFHKMVELTVAARQAYKTMLENVQOE LVGEPASLRHLPAYPHRAPAALQKMAPQP/VKEK EAPHYIEIWKHLEAYSGCTLELESLEASTSQM MNLNL
3535	A	1747	983	LFQFQVCRSVLSPRAAGCTWSLAPRSRGAAGSPR RYRGPQPPAPPALPNSRSPVASGREMVVLSV PAEVTVILLDIEGTTTPIAFVKDILFPYIEENVKEY LQTHWEEBECQDVSLLRKQVFAADVPAVRKW REAGMKVYTYSSGSVEAQKLLFGHSTEGDILELV DGHFDTKIGHKVESESyrKIADSIGCSTNNILFLT DVTREASAAEEADVHVAVVVRPGNAGLTDDEK TYYSLITSFSELYLPSSST
3536	A	3	1302	GRPPTAPHTGRPPTANRGDPRLDLKRGCARLLTS IESRGRPAASAGLRRDRCALRRWPLRRAPLARAT RRRAGSPRRCAPRPACPGQWSRARHQPGGLCL LLLLLCQFMEDRSAQAGNCWLRQAKNGRCQVL YKTELSKEECCSTGRLSTSWTEEDVNDNTLfkW MIFNGGAPNCIPCKETCENVDCGPGKKCRMNKK NKPRCVCAPDCSNITWKGpvcGLDGKTYRNECA LLKARCKEQPELEVQYQGRCKKTCRDVFCPGSS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TCVVVDQTNNA YCVTCNRCPEPASSEQYL CGND GVTYSASACHLRKATCLLGRSIGLAYEGKCIKAK SCBDIQCTGGKKCLWDFKVGGRGCSL CDELC PD SKSDEPVCASDNATYASECAMKEAACSSGV LLE VKHSGSCNSISEDTEEEEEDEDQDYSFPISILEW
3537	A	285	2123	IGLFLQVAPLSVMAKSCPSVCRCDAGFIYCND RF LTSIPTGIPEDATTLYLQNNQINNAGIPSDLKNLL KVERIYLYHNSLDEFPTNLPKYVKELHLQENNR TITYDSLKIPYLEELHLDNSVSAVSIEGAFRD SNYLRLLFLSRNHLSTIPWGLPRTIEELRLDDNRIS TISSPSLQGLTSLKRLVLDGNLLNNHGLGDKVFF NLVNLTELSLVRNSLTAAPVNLPGTNLRKLYLQ DNHINRVPPNAFSYLRQLYRLDMSNNNLSNLPQ GIFDDL NITQLILRNPWYCGCKMKWVRDWL QSLPVKVNVRGLMCQAPEKVRGMAIKDLNAELF CDKDSGIVSTIQITTAIPNTVYPAQGWPA PVTK QPDIKNPKLTKDHQTTGSPSRKTTTTVKS VSDTI HISWKLALPMTALRLSWLKLGHSPAFGSITETIVT GERSEYLVTALEPDSPYKVCMPMETSNLYLFD ETPVCIE TETAPLRMYNPTTTLNREQEKEPYKNP NLPLAAIIGGAVALVTIALLALVCWYVHRNGSLF SRNCAYSKGRRRKDDYAEAGTKKDNSILEIRETS FQMLPISNEPISKEEFVIHTIFFPNGMNLYKNNH
3538	A	877	6184	WNVKPSLLVVQLFKFSDKEEHEQNDSISGKTGET GVEEMIA TRKVEQDSKETVKLSHEDDHILEDAGS SDISSDA ACTNPNKTENSLVGLPSCVDEVTECNL ELKDTMG IADKTENTLERNKIEPLGYCEDAESNR QLESTEFNKS NLEVVD TSTFGPESNILENAICDVP DQNSKQLNAIESTKIESHETANLQDDRNSQSSSV SYLESKSVKSKHTKPVHSKQNM TTDAPKKIVAA KYEVHSKTKKNSVKTDTVETQNTVHRPV KVANKQIDKEPIQSCNSGVKSVENQALVLEKK TLQDQTLVQIFKPLTHSLSDKSHAETGCLKEPHH PAQTGHVSHSSQKQCHKPQQQAPAMKNSHVK EELEHPGVEHFKEEDKLKLKKPEKNLQPRQRSS KSFSLDEPPLFPDNIATIRREGSDHSSSFESKYMW TPSKQCGFCKKPHGNRFMVGCGRCD DWFHGDC VGLSLSQAQQMG EEDKEYVCVKCCA EEDKKTEI LDPDTLENQATVEFHSGDKTMECEKLGLSKHTT NDRTKYIDDTVKHKVKILKRESGEGRNSSDCRD NEIKKWQLAPLRKMGQPVLP RRSSEEKSEKIPKE STTVTCTGEKASKPGTHEKQEMKKKKVEKGVL NVHPAASASKPSADQIRQSVRHSLKDILMKRLTD SNLKVPEEKA AKVATKIEKELFSFRD TDAKYKN KYRSLMFNLKDPKNNILFKKVLKGEVTPDHLIR MSPEELASKELAAWRRREN RHTIEMIEKEQREVE RRPITKITHKGEIEIESDAPMKEQEAAMEIQEPAA NKSLEKPEGSEKARKEEVDSMSKDTTSQHRQHLF DLNCKICIGRMAPPVDDLSPKKVKVVVG VARKH SDNEAESIADALSSTSNLASEFEEBEKQESPKSTF SPAPRPEMPGTVEVESTFLARLNFTWKGFINMPS VAKFVTKAYPVSGSPEYLTEDLPDSIQVGGRI SPQ TVWDYVEKIKASGTKEICVVRFTPVTEEDQISYT LLFAYFSSRKRYGVAANNMKQVKDMYLIPLGAT DKIPHPLVPFDGPGLELHRPNLLGLIIRQKLKRQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				HSACASTSHIAETPESAPPIALPPDKKSKIEVSTEE APBEENDFFNSFTTVLHKQRNKPQONLQEDLPTA VEPLMEVTKQEPPKPLRFLPGVLIGWENQPTTLE LANKPLPVDDILQSLGTTGQVYDQAQSVMEQ NTVKEIPFLNEQTSKIEKTDNVEVTDGENKEIK VKVDNISESTDKSAEITSVVGSSSISAGSLTSLSL RGKPPDVSTEAFLTNLSIQSKQETVESKEKTLKR QLQEDQENNLQDNQTSNSSPCRSNVGKGNIDGN VSCSENLVANTARSPQFINLKRDPQAAGRSQPV TTSESKDGDSCRNGEKHMLPGLSHNKEHLTEQIN VEEKLCSAEKNSCVQQSDNLKVAQNSPSVENIQT SQAEQAKPLQEDILMQNIETVHPFRGSAVATSH FEVGNTCPSEFPKSKITFTSRSTSPRTSTNFSMPMRP QQPNIQLKSSPPGFPFPFPNFPQSMFGFPFHL PPPLPPPFGFVFAQNPMVPVPPVVLHPAGQPQR MMGPLSQASRYIGPQNFYQVKDIRRPERRHSDP WGRQDQQQLDRPFNRGKGDRQRFYSDSHHLKR ERHEKEWEQESERHRRDRSQDKDRDRKSREEG HKDKERARLSHGDRGTGDKASRDSRNVDKKPD KPKSEDYEKDKEREKSKHREGEKDRDRYHKDR DHTDRTKSKR
3539	A	157	1769	GSWTVELSLKPSASPSLKWVCLPGAAAVNKHRS GAGGLIRSLIQCTWAPAGPARRGGRGIEDFPYLF FQLTHCQQRICSVTQAGVQWCDHSSLQFPQTPGL NQSSHLSSLSSRDYRMLSSFNWFWDQDFWLPP NVTWTELEDGRVYHPQDLLAALPLALVLLA MRLAFERFIGLPLSRWLGVRDQTRRQVKPNATL EKHFLETEGHRPKPEQLSLLAAQCGLTLQQTQRW FRRRRNQDRPQLTKKFCEASWRFLFYLSFVGGL SVLYHESWLWAPVMCWDYRPNQLTLSCPAADS EASLNVWFTTBLGFYLSLTLPLFDVKKKQCT SSIKPRFHYDPLSTADFKEQVIHHFVAVILMTF&Y SANLLRGLVLLLDSSDYLLLEACKMVNYMQY QQVCDALFLPSFVFFYTRLVLFPTQLYTTYESI SNRGPFPGYYTFNGLMLLQLLHVFWSCLILRML YSFMKKGQMEKDIRSDVEESDSSEAAAAAQEPL QLKNGTAGGPRPAPTDGPRSRVAGRLTNRHITA T
3540	A	267	1397	SPAGYCHSGLLPGCSRSA/CADLAKHQELPGKKL LSEKKLKRYFVDYRRVLVCGGNGGAGASCFHSE PRKEFGGPDGGDGGNGGHVILRVDDQVKSLSSV LSRYQGFGSGEDGGSKNCFGRSGAVLYIRVPVGT VKEGGRVVADLSCVGEYIAALGGAGGKGNRF FLANNNRAPVTCIPGQPGQQRVLHLELKTVAHA GMVGFPNAGKSSLLRAISNARPAVASYPFTTLKP HVGIVHYEGHLQIAVADIPGIIRGAHQNRGLGSA FLRHIERCRFLFVVDLSQPEPWTQVDDLKYELE MYEKGLSARPHAIVANKIDLPEAQANLSQLRDH LGQEVIVLSALTGENLEQLLLHLKVLVYDAYAEA ELGQGRQPLRW
3541	A	1	8008	DTQVSETLKRFAKVTASVKERREILSELGKCV AGKDLPEGAVKGLCKLFCLTLHRYRDAASRRAL QAAIQQLAEAQPEATAKNLLHSLQSSGIGSKAGV PSKSSGSAALLALTWTCLLVRIVFPRAKROGDI WNKLVEVQCLLLLEVLGGSHKHAVDGAVKKLT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				KLWKENPGLVEQYLSAILSLEPNQNYAGMLGLL VQFCTSHKEMDVVSQHKSAALLDFYMKNILMSK VKPPKYLLDSCAPLLRYLSHSEFKDLILPTIQKSL LRSPENVETISSLLASVTLDLSQYAMDIVKGLAG HLKSNSPRLMDEAVLALRNLARQCSDSAMESL TKHLFAILGGSEGKLTVVAQKMSVLSGIGSVSHH VVSGPSSQVLNGIVAEFLFIPFLQQEVHEGTLVHA VSVLALWCNRFTMEVPPKKLTWFKKAFSLKTST SAVRHAYLQCMASYRGDTLLQALDLLPLLIQT VEKAASQSTQVPTITTEGVAAALLLLKLSVADSQA EAKLSSFQWLIVDEKKQVFTSEKFLVMASEDAL CTVLHLTERLFLDHPHRLTGKNVQQYHRALVA VLLSRTWHVRRQAQQTVRKLLSSLGGFKLAHGL LLEELKTVLSSHKVLPLEALVTDAGEVTEAGKAY VPPRVLQEALCVISGVPGLKGDVTDTEQLAQEM LIISHPSLVAVQSGLWPALLARMKIDPEAFITRH LDQIPRMTTQSPLNQSSMNAMGSLSVLSPDRVL PQLISTTTASVQNPALRLVTREEFAIMQTPAGELY DKSIIQSAQQDSIKKANMKRENKAYSFKEQIELE LKEEIKKKKGKKEEVQLTSKQKEMLQAQLDREA QVRRRLQELDGELEAALGLLDIILAKNPSGLTQYI PVLVDSFLPLLKSPLAAPRIKNPFLSLAACVMPSR LKALGTLVSHVTLRLKPECVLDKSWCQBELSV AVKRAVMLLHTHTTTSRVGKGEPGAAPLSAPAFS LVFPFLKMVLTEMPHHSEEEEWMAQILQILTVQ AQLRASNTTPGRVDENGPELLPRVAMLRLLTW VIGTGSPRLQVLASDTLTLCASSSGDDGCAFAE QEEVDVLLCALQSPCASVRETVLRGLMELHMLV PAPDTEKNGNLNLRRLWVVKFDKEEIRKLAEL RLWSMMGLDLQPDLCSLIDDIYHEAAVRQAG AEALSQAVARVQQAEEVETRIETQEKEL PPPVLDALGKVPSPDQWEARCGLALALNLS QYLDSSQVKPLPQFFVPDALNDRHPDVRKCMU AALATLNTHGKENVNSLLPVFEFLKNAPNDAS YDAVRQSVVVLMSGLAKHLDKSDPKVKPIVAKL IALSTPSQQVQESVASCLPPLVPAIKEDAGGMIQ RLMQQLLESCKYAEKGAAYGLAGLVKGLGILS LKQQEMMAALTDAIQDKKNFRRREGALFAFEM LCTMLGKLFEPYVVHVLPHELLCFGDGNQYVRE AADDCAKAVMSNLSAHGVKLVLPSLLAALAEES WRTKAGSVELLGAMAYCAPKQLSSCLPNIVPKL TEVLTDShVKVQKAGQALRQIGSVIRNPEILAI APVLLDALTDPSRKTQKCLQTLDDTKFVHFIDAP SLALIMPVQRAFDQDRSTDTRKMAAQIIGNMYSL TDQKDLAPYLPSTVTPGLKASLLDPVPEVRTVSAK ALGAMVKGMGESCFEDLLPWLMEITLTYEQSSV DRSGAAQGLAEVMAGLVVEKLEKLMPEIVATAS KVDIAPHVRDGYIMMFNYLPITFGDKFTPYVGPII PCILKALADENEFVRDTALRAGQRVISMAYETAI ALLLPQLEQGLFDDLWRIRFSSVQLLDLLFHISG VTGKMTTETASEDDNFGTAQSNKAITALGVERR NRVLAGLYMGRSDTQLVVRQASLHVWKIVVSN TPRTLREILPTLFGLLGFLASTCADKRTIAARTL GDLVRKLGEKILPEIPILEEGLRSQKSDERQGVCI GLSEIMKSTSRDAVLYFSESIVPTARKALCDPLE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EVREAAAKTFEQLHSTIGHQALEDILPFLKQLD DEEVSEFALDGLKQVMAIKSRVVLPYLVPKLTTP PVNTRVLAFLSSVAGDALTRHLGVILPAVMLAL KEKLGTPDEQLEMANCQAVILSVEDDTGHRHIE DLLEATRSPEVGMQRQAAAILNIYCSRSKADYTS HLRSLVSGLRIRLFNDSSPVVLEESWDALNAITKK LDAGNQLALIEELHKEIRLIGNESKGEHVPGFCLP KKGVTSLPVLREGVLTGSPEQKEEAAKALGLVI RLTSADALRPSVVSITGPLIRILGDRFSWNVKAAL LETLSLLAKVGIALKPFLPQLQTTFKALQDSNR GVRLKAADALGKLISHIKVDPLFTLLNGIRAME DPGVRDMLQALRFVIQAGAKVDAVIRKNIVS LLSMLGHDEDNTRISSAGCLGELCAFLTEEELS AVLQQCLLADVSGIDWMVRHGRSLALSAVNV APGRLCAGRYSSDVQEMILSSATADRIPIAVSGV RGMGFLMRHHIETGGGQLPAKLSSLFVKCLQNP SSDIRLVAEKMIWWANKDPLPLDPQAIKPILKA LLDNTKDKNTVVRAYSDAQIVNLLKMRQGEVVF QSLSKILDVASLEVLNEVNRRSLKKLASQADSTE QVDDTILT
3542	A	62	1130	PWNPDFFPGRGLMGVQKGEIGPPGQQGKKGAP GMPGLMGSNSSPGQPGTPGSKGSKGEPGIQGM GASGLKGEPGATGSPGEPGYMGLPGIQGKGD GNQGEKGIQGGQKGENGRQGIPGQGIQGHGAK GERGEKGEPPVGAIGSKGESGVDGLMGPA GPGDPGPQGPGLDGKPGREFSEQFIRQVCTDV IRAQLPVLQSGRIRNCDHCLSQHGSPGIPGPPGI GPEGPRGLPGLPRDGVPLVGVPGRPGVRGLK GLPGRNGEKGSGQGFYGPGEQGGPPGPEGPPGI SKEGPPGDPGLPGKGDHKGPIQGGQPPGICD FSLCTEVIARRDFFRKCETV
3543	A	654	194	PARSLEKMKASVVLSSLLGYL P=SGAYILGRCTV AKKLHDGGLDYFERYSLNWVCLAYFESKFNP AIYENTREGYTGFLFQMRGSDWCGDHGRNRC HMSCSALLNPNELEKTKCAKTIVKGKEGMGA WP TWSRYCQYSDTLARWLDGCKL
3544	A	2	1074	TSRLAAGRLAQWLLRASRSGMLRAGWLRGAAA LALLLAARVVAFAFEPITVGLAIGAASAITG LSYNDIYCRFAECCREERPLNASALKLDLEBK LFGQLATEVTFKALTGFRNNKNPKPLTSLHGW AGTGKNFVSQMGAEHLHPKGLKSNFVHLFVST LHFPHEQKIKLYQDQLQKWIRGNVSACANSV FIFDEM DKLHPGHEVAIKPFLDYEHVERVS YRKAIFIFLSNAGGDLITKTALDFWRAGRKRE DIQLKDLEPVL SVGVFNKXHSGLWHSGLID KNLIDYFIFLPLEYR HVKMCVRAEMRARG SAIDEDIVTRVAEEMTFFPRDEKIYSDKG CKTQVSRDLFH
3545	A	3	273	SAQGRSWGRFYRQIKRHPGIIPMGLICLGM GSAALYLLRLALRSPDVW*SWDRKNNPEP WNRLSPNDQYKFLAVSTDYKKLKKDRPDF
3546	A	23	591	ALSTETRTPDMRLLLVTSLVVVLLWEAGAV PA PKVPIKMVQKHWPSQDPEKAWGARVVEP PEKDDQLVVLFPVQKPKLLTTEEKPRGQGR GPILPGTKAWMETEDTLGRVLSPEPDHDSL YHPPPEEDQG EERPLWVMPNHQVLLGPEED QDHIYHPQ*GSR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GHHCPRPVPRPRLGLGPSLPCPS
3547	A	23	591	ALSTETRTPD MRRLLLVTSLVVLLWEAGAVPA PKVPIKMQVKHWPSEQDPEKAWGARVVEPPEK DDQLVVLFPVQKPKLLTTEEKPRGQGRGPILPGT KAWMETEDTLGRVLSPEPDHDSL YHPPPEEDQG EERPLWVMPNHQVLLGPEEDQDHIYHPQ*GSR GHHCPRPVPRPRLGLGPSLPCPS
3548	A	3	1641	TWLPSPVAEEVQQPEMAAVLNAERLEVSVDGLT LSPDPEERPGAEGAPLAAATAATLATWIRSRP RLRGRTARSPGRRRAAGGAAEEARRLEQRWGFGL ELYGLALRFFKEKDGAFFHTYEEKLKLVALHK QVLMGPYNPDTCPEVGFDFVLGNDRREWAAL GNMSKEDAMVEFVKLLNRCCHLFSTYVASHKIE KEEQEKKRKEEEERRRREERERLQKEEEKRRR EEEEERLREEEERRRIEEERLRLLEQQKQQIMAAL NSQTAVQFQQYAAQQYPGNYEQQQLIRQLQE HYQQYMQQLYQVQLAQQAALQKQEVVAVAG SSLPTSSKVECNCTQVI*CFNRQAKTHTDSSEKE LEPEAAEEALENGPKESLPVIAAPSMWTRPQIKD FKEKIQDADSVITVGRGEVTVRVPTHEEGSYL FWEFATDNYDIGFGVYFEWTDSPNTAVSVHVSE SSDDDEEEENIGCEEKAKKNANKPLLDEIVPVY RRDCHEEVYAGSHQYPGRGVYLLKFDNSYSLW RSKSVYYRVYYTR
3549	A	1837	3593	PAVLVLEPASQSRKQONTASATAQHWSAQIHKE SFLAPVFTKDEQKHRRPYEFEVERDAKARGLEQF SATHGHTPILNGWHGESAMDLSCSSESGPGATS PPFVSASTPKIGAISLQGALGMDLSGILQAGLIHP VTGQIVNGSLRRDDAATRRRRGRRKHVEGGMD LIFLKEQTLQAGILEVHEDPGQATLSTTHPEGP ATSAFEPATLAGEVKEKSPKSLDNL RQQAD SLEVPGFGANSDKPRQRPRCKEPGKLDVSSLS GEERVPAIPKEPILRGFLPENKFNHTLAEPILRDT GPRRRGRRPRSELKAPSIVADSPSGMGPLFMNG LIAGMDLVGLQNMNRNMPGIPLTGLVGFPAGFAT MPTGEEVKSTLSMLPMMLPGMAAVPQMFGVGG LLSPPMATTCTSTAPASLSSTTKSGTAVTEKTAE DKPSSHVDKTDTLAEDKPGPGPFSDQSEPAITTS PVAFPNPLIPGVSPGLIYPSMFLSPGMGMALPAM QQAHRHSEIVGLESQKRKKKTKGDNPNSHPEPA PSCEREPSGDENCAEPSAPLPAEREHGAQAGEGA LKDSNNDTN
3550	A	287	39	QLNLNKIATSQKHRDFVAESVGEKPVGSLAGIGE VMDKKLEEGCFDKAYVVLGQFLVLKKDEDLF*E WLRDTGGARTGRSRE
3551	A	21	3925	GDLLEVGLPPGLEFPRGICLRGLRRTMSLDFGSV ALPVQNEDEEYDEEDYEREKELQQLLTDLPHDM LDDDLSSPELQYSDCEDGTDGQPHHPEQLEMS WNEQMLPKSQSVNGPSCQGLEPYNKVITYKPYQS SAQNGSPAQEITGSDTFEGLQQFLGANENSAE NMQIIQLQVLNKAKEKQLENLIEKLNESERQIRY LNHQLVIKDEKDGLTSLRESQKLFQNGKEREIQ LEAQIKALETQIQALKVNEEQMIKKSRTTEMALE SLKQQLVDLHHSESQRAREQHESIVMGLTKKY EEQVLSLQKNLDATVTALKEQEDICSRDKDHVK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QLERNQEAIKLEKTEIINKLTRSLEESQKQCAHLL QSGSVQVEVAQLQFQLQQAQKAHAMSANMNKA LQEELTELKDEISLYESAAGLGIHPSDSEGELNIEL TESYVDLGIKKVNWKKSKVTSIVQEEDPNEELSK DBFILKLKAEVQRLGNSNMKRHLVSQQLNDLK DCHKKIEDLHQVKKDEKSIEVETKTDTSEKPKNQ LWPESSTSDVVRDDILLKNEIQVLQQQNQELKE TEGKLRNTNQDLNQMQRQMVQDFDHDKQEA VDRCERTYQQHHEAMKTQIRESLLAKHALEKQQL FEAYERTHLQRLSELDKLNKEVTAVQECYLEVC REKDNLELTLRKTTTEKEQQTQEKIKEKLIQQLK EWQSKLDQTIKAMKKKTLDCGSQTDQVTTSDVI SKKEMAIMIEEQKCTIQQNLEQEKDIAIKGAMKK LEIELELKHCEITKQVEIAVQNAHQRWLGELPE LAEYQALVKAQKKWEEQHEVSVNKRISFAVSE AKEKWKSELENMRKNILPGKELEEKIHSLEKELE LKNEEVPVVIRAEALAKARSEWNKEKQEEIHRIOE QNEQDYRQFLDDHRNKINEVLAAAKEDFMKQK TELLQKETELQTCLDQSRREWMTQEAQRIQLEI YQYEEIDTLVLGVLLSDTQKEHISDSEDKQLLEI MSTCSSKWMSVQYFEKLGKCIQKAFQDTPLLV ENADPEWKKRNMAELSKDSASQGTGQGDGP AGHHAQPLALQATEAEADKKKVLEIKDLCCGHC FQELEKAKQECQDLKGKLEKCCRHLQHLERKHK AVVEKIGEENNKVVEELIEENNDMKNKLEELQT LCKTPPRSLSAGAIENACLPSCGGALEELRGQYIK AVKKIKCDMLRYIQESKERAEMVKAENVL*ERQ ETARKMRKYLYICLQQLQDDGKEGAEEKMINA ASKLATMAKLLPETPISSKSQSKTTQSGMSK
3552	A	771	375	ARTRQTSQAREPEKESPAPGGGGLAEIRSROQL SQTGRIPPLAKDQAVEAMFPARGKFLLSFE MYFTREEWGHNLNWGQKDLRYRIEENYKRMV LLVYFQFDAIPLC*TLAHSSWLQLYFRLYF
3553	A	76	72	PGVRGVEAPGGVAPGRNAMRRGERRDAGGPRP ESPVPAGRASLEPPDGPSAGQATGPGEGRSTB SEVYDDGTNTFFWRAHTLTVLFTCTLGYYVTL EETPQDTAYNTKRGIVASILVFLCFGVTAQKDG FSRPHPAYWRFWLCVSVVYELFLIFLQTVQDG RQFLKYVDPKLGVPPLPERDYGGNCLYDPDNET DPFHNIWDKLDGFVPAHFLGWYKTLTMIRDWW MCMISVMFEFLEYSLEHQLPNFSECWWDHWIM DVLVCNGLGIYCGMKTLEWLSLKYKWQGLWN IPTYKGKMKRIAFQFTPYSWVRFEWKPASSLR WLAVCGIILVFLAELNTFYLKFVLWMPPEHYLV LLRLVFFVNVGGVAMREIYDFMDDPKPHKKLGP QAWLVAATATELLIVVKYDPHTLTLSPFYISQC WTLGSLALTWTVWRFLRDITLRYKETRWQK WQNKDDQGSTVGNQDQHLPLGLDEDLLGPGVAE GEGAPTPN*PRGPAPRPLPSAPRAVCGASSRR
3554	A	2	2106	FDEFSALPSPSLQTSWSFGPMSRRALRRLRGEQR GQEPLGPGALHFDLRDDDDAEEEGPKRELGVRR PGGAGKEGVRVNNRFELINIDDEDDPVVNGERS GCALTDVAPGNKGRGQRGNTESKTDGDDTET VPSEQSHASGKLRRKKKKQKNKKSSTGEASENG LEDIDRILERIEDSTGLNRPAPLSSRRKHVLYVE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				HRHLNPDTELKRYFGARAILGEQRPRQRQRYVP KCTWLTPPKSTWPRYSKPGLSMRLLSKKGLSFF AFEHSEYQQAQHKFLVAVESMEPNINIVLLQT SPYHVDSSLQLSDACRFQEDQEMARDLVERALY SMECAFHPLFSLTSGACRLDYRRPENRSFYALY KQMSFLEKRGCPRTALEYCKLILSLEPDEDPLCM LLIDHLALRARNYEYLIRLFQEWVVGASLAHRN LSQLPNFAFSVPLAYFYLLSQQTDLPCEQSSARQ KASLLIQQALTMFPGVLLPLESCSVRPDASVSSH RFFGPNAEISQPPALSQVLNLYLGRSHFLWKEPA TMSWLEENVHEVLQAVDAGDPAVEACENRRKV LYQRAPRNIHRHVILSEIKEAVAALPPDVTTQSV MGFDPLPPSDTIYSYVRPERLSPISHGNTIALFFRS LLPNYTMGERPEEGVAGGLNRNQGLNRLMLA VRDMMANFHLNDLEAPHEDDA*GEGEWD
3555	A	2	2106	FDEFSALPSPSLQTSWSFGPMSRRALRRLRGEQR GQEPLGPGALHFDLRDDDDAEEEGPKRELGVRR PGGAGKEGVRVNNRFELINIDDEDDPVVNGERS GCALTDVAPGNKGRGQRGNTESKTDGDDTET VPSEQSHASGKLRRKKKKQKNKKSSTGEASENG LEDIDRILERIEDSTGLNRPGAPLSSRKHVLYVE HRHLNPDTELKRYFGARAILGEQRPRQRQRYVP KCTWLTPPKSTWPRYSKPGLSMRLLSKKGLSFF AFEHSEYQQAQHKFLVAVESMEPNINIVLLQT SPYHVDSSLQLSDACRFQEDQEMARDLVERALY SMECAFHPLFSLTSGACRLDYRRPENRSFYALY KQMSFLEKRGCPRTALEYCKLILSLEPDEDPLCM LLIDHLALRARNYEYLIRLFQEWVVGASLAHRN LSQLPNFAFSVPLAYFYLLSQQTDLPCEQSSARQ KASLLIQQALTMFPGVLLPLESCSVRPDASVSSH RFFGPNAEISQPPALSQVLNLYLGRSHFLWKEPA TMSWLEENVHEVLQAVDAGDPAVEACENRRKV LYQRAPRNIHRHVILSEIKEAVAALPPDVTTQSV MGFDPLPPSDTIYSYVRPERLSPISHGNTIALFFRS LLPNYTMGERPEEGVAGGLNRNQGLNRLMLA VRDMMANFHLNDLEAPHEDDA*GEGEWD
3556	A	3388	1650	KTRGTMFYYPNVLQRHTGCFATIWLAAATRGSR VKREYLRVNVVKTCEEILNYVLVRVQPPQGLP RPRFSLYLSAQLQIGVIRVYSQQCYLVEDIQHIL ERLHRAQLQIRIDMETELPSLLLPNHLAMMETLE DAPDPFFGMMMSVDPRLSPFDIPQIRHLLAAIPE RVEEIPPEVPTPEPREPERIPVTVLPPEAITLEAEP MLBIEGERELPEVSRRELDLIAEEEEAILLEIPRL PPAPAE*GQELLDQVGCQCWEGSPHFSCFPLR VEGMGEALGPEELRLTGWEPGALLMEVTPPEEL RLPAPPSPERRPPVPPPPRRRRRRLLFWDKETQI SPEKFQEQQLQTRAHCWECMVQPPERTIRGPAEL FRTPTLSGWLPEELLGLWTHCAQPPPKALRREL EEAAAEERRKIEVPSEIEVPREALEPSVPLMVS EISLEAAEBEKSRLIPPEERWAWPEVEAPEAPA LPVVPELPEVPMEMPLVLPPELELLSLEAVHRAV ALELQANREPDFSSLVSPLSPRRMAARVFYLLLV LSAQQLHVKQEKPYGRLLIQGPRFH
3557	A	3388	1650	KTRGTMFYYPNVLQRHTGCFATIWLAAATRGSR VKREYLRVNVVKTCEEILNYVLVRVQPPQGLP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RPRFSLYLSAQLQIGVIRVYSQQCQYLVEDIQHIL ERLHRAQLQIRIDMETELPSLLLPNHLAMMETLE DAPDPFFGMMSVDPRLPSPFDIPQIRHLEAAIPE RVEEIPPEVPTEPREPERIPVTVLPEAITLEAEPIR MLEIEGERELPEVSRRELDLLIAEEEEAILLEIPRL PPPAPAE*GQELLDQVGCQCWEGSPHFSCFFPLR VEGMGEALGPEELRLTGWEPGALLMEVTPPEEL RLPAPSPERRPPVPPPPRRRRRRRLFWDKETQI SPEKFQEQLQTRAHCWECMVQPPERTIRGPAEL FRTPTLSGWLPELLGLWTHCAQPPPKALRREL EEAAAEERRKIEVPSEIEVPREALEPSVPLMVSL EISLEAAEEKSRISLIPPEERWAWPEVEAPEAPA LPVVPELPEVPMEMPLVLPPELELLSLEAVHRAV ALELQANREPDFSSLVSPSPRRMAARVYLLLV LSAQQLHVQKQEPYGRLLIQPGPRFH
3558	A	489	2360	IRPRPRGRRRRALDSPNAAAPPVYVCRSPGEPTSL VNMAEDIAKLAETLAKTQVAGGQLSFKGKSLK LNTAEDAKDVIKEIEDFDSLEALRLEGNVTGV ARVIAKAL*KKSELKRCHWSMDMFTGRLRTEIPPA LISLGEGLITAGAQLVELDLSDNAFGPDGVQGF ALLKSSACFTLQELKLNNCGMGIGGGKILAAALT ECHRKSSAQGKPLALKVVFVAGRNRLNDGATAL AEAFRVIGTLEEVHMPQNGINHPGITALAQAFV NPLLRVINLNDNTFTEKGAVAMAETLKTLRQVE VINFGDCLVRSGAVAIADAIRGGLPKLKELNLS FCEIKRDAALAVAEAMADKAELEKLDLNGNTLG EEGCEQLQEVLEGFNMAKVLASLSDDEDEEEEEE EGEEEEEEAEEEEEEDEEEEEEEEEEEEEEPQQRG QGEKSATPSRKILDPNTGEPAPVLSPPPADVSTF LAFPSPEKLLRLGPKSSVLIAQQTDTSDPEKVSA FLKVS SVFKDEATVRMAVQDAVDALMQKAFNS SSFNSTNITLRLVHMGLLKSEDKVKALANLYGP LMALNHMVQQDYFPKALAPLLAFVTKPNAL SCSFARHSLLOTLYKV
3559	A	489	2360	IRPRPRGRRRRALDSPNAAAPPVYVCRSPGEPTSL VNMAEDIAKLAETLAKTQVAGGQLSFKGKSLK LNTAEDAKDVIKEIEDFDSLEALRLEGNVTGV ARVIAKAL*KKSELKRCHWSMDMFTGRLRTEIPPA LISLGEGLITAGAQLVELDLSDNAFGPDGVQGF ALLKSSACFTLQELKLNNCGMGIGGGKILAAALT ECHRKSSAQGKPLALKVVFVAGRNRLNDGATAL AEAFRVIGTLEEVHMPQNGINHPGITALAQAFV NPLLRVINLNDNTFTEKGAVAMAETLKTLRQVE VINFGDCLVRSGAVAIADAIRGGLPKLKELNLS FCEIKRDAALAVAEAMADKAELEKLDLNGNTLG EEGCEQLQEVLEGFNMAKVLASLSDDEDEEEEEE EGEEEEEEAEEEEEEDEEEEEEEEEEEEEEPQQRG QGEKSATPSRKILDPNTGEPAPVLSPPPADVSTF LAFPSPEKLLRLGPKSSVLIAQQTDTSDPEKVSA FLKVS SVFKDEATVRMAVQDAVDALMQKAFNS SSFNSTNITLRLVHMGLLKSEDKVKALANLYGP LMALNHMVQQDYFPKALAPLLAFVTKPNAL SCSFARHSLLOTLYKV
3560	A	2	1198	FVRELPRPRPGAATAAIMVSVINTVDTSHEDMIH DAQMDYYGTRLATCSSDRSVKIFDVRNGGQILIA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				DLRGHEGPVWQVAWAHPMYGNILASCSYDRKV IIWRENGTWEKSHEHAGHDSSVNSVCWAPHDY GLILACGSSDGAISLLTYTGEGQWEVKKINNAHT IGCNAVSWAPAVVPGSLIDHPSGQKPNYIKRFAS GGCDNLIKWLKEEEDGQWKEEQKLEAHSDWVR DVAWAPSIGLPTSTIASCSQDGRVFIWTCDDASS NTWSPKLLHKFNDVVWHVSWSITANILAVSGGD NKVTLWKESVDGQWVCISDVNKGQGSVSASVT EGQQNEQ*QDRWGLAPHPAPGLPLPGPTNQTT GKSPQLQDDYFPRRSYRCSHRLICLNVIDDAL
3561	A	540	86	WRVKEMTSTLPKALGRKTASRSHTTLQGGSCCP VLWTAKLRCRKLRFPLPPPPSSSAWPWQGWGI RGEQEAEGPLGETGPPVGPESGLRQWRKLIKGR YGEWRGSGQKTGQPS*TTMQGGETEENRTETTT GNKQRESEAPWVRHTYTT
3562	A	1920	242	PMMAMPFFERFKSSIQRPSPVLVLSQNTKRESGR KVQSGNINAAKTIADIHRTCLGPKSMMKMLLD MGGIVMTNDGNAILREIQVQHPAAKSMIEISRTQ DEEVGDGTTSVILAGEMLSVAEHFLEQQMHPTV VISAYRKALDDMISTLKKISIPVDISDSMMLNIIN SSITTKAISRWSSLACNIALDAVKMVQFEENGRK EIDIKKYARVEKIPGGIIEDSCVLRGVMNKDVTH PRMRYIKNPRIVLLDSSLEYKKGESQTDIEITRE EDFTRILQMEEYIQLCEDIIQLKPDVVITEKGIS DLAQHYLMRANITAIRRVKTDNNRIARACGARI VSRPEELREDDVGTGAGLLEIKKIGDEYFTFITDC KDPKACTILLRGASKEILSEVERNFDAMQVCRN VLLDPQLVPGGGASEMAVAHALTEKSKAMTGV EQWPYRAVAQALEVIPRTLQNCGASTIRLLTSLR AKHTQENCETWGVNGETGTLVDMKELGIWEPL AVKLQTYTAVETAVALLLRIDDVSGHLYKGGD QSRQGGAPDAGQE
3563	A	1571	560	GPSLLGTRGTPNPARTLQIFFLIIGRRLTGRMAAV DDLQFEFGNAATSLTANPDATTVNIEDPGETPK HQPGSPRGSGREEDDELLGNDDSDKTELLAGQK KSSPFWTFEYYQTFDVTYQVFDRIKGSLLPIPG KNFVRLYIRSNPDLYGPFWICATLVFAIAISGNLS NFLIHLGEKTYHYVPEFRKVSIAATHIYAYAWLVP LALWGFLMWRNSKVMNIVSYSFLEIVCVYGYSL FIYIPTAILWIIPKAVRWILVMIALGISGSLAMT FWPAVREDNRRVALATTIVTIVLLHMLLSVGCLA YFFDAPEMDHLPTTATPNQTVAAAKSS
3564	A	1	328	NSRVDDFVAHLQRPLGPASCLGILRPAMTAHSF ALPGIIFTFWGLVGIAGPWFPKGPNGVITIML VATAVCCYLFWLIALAQLNPLFGPQLKNETIYW VRFLWE
3565	A	2	1081	FVTDFFPARSMAATSLMSALAAARLLQPAHSCSLRL RPFHLAAVRNEAVVISGRKLAQQIKQEVQRQVEE WVASGNKRPHLSVILVGENPASHSYVLNKTRAA AVVGINSETIMKPASISEEELLNLINKLNDDNDVD GLLVQLPLPEHIDERRICNAVSPDKDVGPHVIN VGRMCLDQYSMLPATPWGVWEIKRTGIPTLGK NVVVAGRSKNVGMPIAMLLHTDGAHERPGGDA TVTISHRYTPKEQLKKHTILADIVISAAGIPNLITA DMIKEGAVIDVGINRVHDPVTAKPKLVGDVDF

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EGVRQKAGYITPVPGGVGPMTVAMLMKNTIIAA KKVLRLEEREVLKSKELGVATN
3566	A	3	1130	SCRRGRQQRRNVSLSSQFAHTMAAPAQQTTP GGGKRKGKAQYVLAKRARRCDAGGPRQLEPGL QGILITCNMNERKCVEEAYSLLNEYGDDMYGPE KFTDKDQQPSGSEGEDDDAEAAALKKEVGDIKAS TEMRLRRFQSVESGANNVVFIRTLGIEPEKLVHHI LQDMYKTKKKKTRVILRMLPISGTCKAFLEDMK KYAETFLFPWFKAPNKGTFQIVYKSRNNSHVNR EEVIRELAGIVCTLNSENKVDLTNPQYTVVVEIK AVCCLSVVKDYMLFRKYNLQEVVKSPKDPSQLN SKQNGKEAKLESADKSDQNNTAEGKNNQQVP ENTEELGQTKPTSNPQVVNEGGAPELASQATE GSKSNENDFS
3567	A	248	3498	GKKDSSPWTCPFHPPLQLFFVIRNTRQLGDFHLA KIKVRNYWTADGDLDIGAKNVKLYVNRNLIFNG KLDKGDREAPADHSILVDQKNEKSEQLEAMNA HSEBSKGTHEMAGASGDKEGLGCSPPAETLAD AKLSSQGNVSGKRKNSTNCRKDSLSQLEEYLRLS AVPTSMGDMPSAPATSPPVKCPPVHEEPSLIQQL ENLMGRKICEPPGKTPSWLQPSPTGKDRKQGG KPKPLWLSPEKPLAWKGRLPSSDDVIGEGPGETEA RDKGLRHEPGWGTSRSVNTKERPQRATTKVHSD DSDIFNQPPNRERPASGRRGSRKDAGSSSHGDDQ PASREDTWSSRTPSRSRWRSEQEHTLHESWSSLS AFDRSHRGRISNTELPDILDELLQKSSRHSCLP PSKKGEQPLSRGQDGYSGETDAGGDFKIPVLPY GQRLVIDIKSTWGDRIYVGLNGIEIFSSKGPEVQI SNIKADPPDINILPAYGKDPVVNTLIDGVNRTQ DDMHVWLAFTTRGRSHSITDFTHPCHVALRIW NYNKSRHIEFGVYDITMLLEDTGTTGEIAKASG TAGAPEHFGDTILFTTDDDEALFTDEMFDLD VGSLSLSQDEEAMRRPSTADGGERPFTQAGL GADERIPELELPSSSPVPQVTIPEPCITGICLQLN FTASWGDLHYLGLTGLEVVGKEGQALPEHLHQIS ASPRDLNELPEYSDDSRITDKLIDGTNITMEDEH MWLIPFSPGLDHVVITRLDRAESLAGLRFWYNK SPEDTYRGAKIVHVSVDGLCVSPPEGFLIRKGP NCHFDFAQEILFVDYLRAQLLPQARRLDMSLE CASMDYEAPLMPCGFIFQFQLLTSWGDPPYYIGLT GLELYDERGEKIPLSENNIAAFPDSVNSLEGVGG DVRTPDKLIDQVNDTSDGRHMLAPILPGLVNR VYVIFDLPTTVSMIKLWNYAKTPHRGVKEFGLL VDDL VYNGILAMVSHLVGGILPTCEPTVPYHTI LFTEDRDIRHQEKHTTISNQAEDQDVQMMNENQ IITNAKRKQSVVDPALRPKTCISEKETRRRRRC
3568	A	50	1724	AQGGTLSAASRFCRGGLLGPWLHPASEMAATLD LKSKEEKDAELDKRIEALRRKNEALIRRYQEIEE DRKKAELEGVAVTAPRKGRSVEKENVAVESEKN LGPSRRSPGTPRPPGASKGGRTPPQQGGRAGMG RASRSWEGSPGEQPRGGGAGGRGRGRGRGSPH LSGAGDTSISDRKSKEWEERRRQNEKMNEEME KIAEYERNQREGVLEPNPVRNFLDDPRRRSGPLE ESERDRREESRRHGRNWGGPDFERVRCGLEHER QGRRAGLGSAGDMLTSMGTGRERSEYLRWKQER

386

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				EAAAETRKLEKQEKRLKKEKKRLAALALASS ENSSSTPEECEETSEKPKKKKKQKPQEVPOENGM EDPSISFSKPKKKKSFSKEELMSSDLEETAGSTSIP KRKKSTPKEETVNDPEEAGHRSRSKKKRKFSKEE PVSSGPEEAVGKSSSKKKKKFKASQED
3574	A	284	2032	CGNERTARLWVQPVVSTMPQASEHRLGRTREPP VNIQPRVGSKLFPAPRARSKERRNPASGPNPMLR PLPPRPLPDERLKKLELGRGRTSGPRPRGPLRA DHGVPLPGSPPTVALPLSRTNLARSKSVSSGDL RPMGIALGGHRTGELGAALSRLALRPEPTLRR STSLRRLGGFPGPPTLSIRTEPPASHGSFHMISAR SSEPFYSDDKMAHHTLLGSGHVGLRNLGNTCF LNAVLCQLSSTRPLRDFCLRRDFRQEVPGGGRRA QELTEAFADVIGALWHPDSCEAVNPTRFRAVFQ KYVPSFSGYSQQDAQEFLKLLMERLHLEINRRGR RAPPILANGPVPSPPRRGGALLEEPELSDDDRANL MWKRYLEREDSKIVDLFGVQLKSLCKQACGY RSTTFEVFCDSLPIPKKGFAGGKVSLRDCFNLT KEEELESENAPVCDRCRQKTRSTKCLTVQRFPR LVLHLNRFASRSIKKSSVGVDPLQLSLGDF ASDKAGSPVYQLYALCNHSGSVHYGHYTALCR CQTGWHVYNDNRVSPVSENQVASSEGYVLFYQL MQEPPRCL
3575	A	1	2408	RELDLADLPERIKPPYANGLSTSHLRSSSVEDVK LIISEGRPTIEVRRCSMPVICEHTKQFQTISEESN QGSLLTVPGDTSPSPKPEVFSNVPERDLSNVSNH SSFATSPGTGASNSKYVSADRNLKNTAPVNTVMD SPVHLEPSSQGVQIQNKSWEMPVDRLETLSTRDF ICPNNSNIPDQESSLQSFCSNENKVLKENADFLSLR QTELPGNSCAQDPASFMPQPCSFPSQSI SDAES IKKPMISI.SYVANQEFJLQOZNAVQIIS...DTT NESTKDTENTFVLGLVQK...FVPVYSESTIQEA SPNFKA YTLVLPSEKDFNGSDASTQLNTHYAF SKLTYKSSSGHEVENSTTDTQVISHEKENKLES VLTHLSRCDSDLCMNAGMPKGNLNEQDPKHC PESEKCLLSIEDEESQOSILSSLENHSQOSTQPEM HKYQGLVKVLEENAEEDDKTENQIPQRMTRNK ANTMANQSKQILASCTLLSEKDSSESSPRGRIRLT EDDDPQIHHPKRKRVSRVPQVQVSPSLLQAKEK TQQSLAAIVDSLKLDEIQPYSSERANPYFEYLHIR KKIEEKRLKLLCSVIPQAPQYYDEYVTFNGSYLLD GNPLSKICIPTITPPPSLSDPLKELFRQEQEVVRMKL RLQHSIEREKLIVSNEQEVLRVHYRAARTLANQT LPFSACTVLLDAEVYNVPLDSQSDSDKTSVRDRF NARQFMSWLQDVDDKFDKLTCLLMRQQHEA AALNAVQRLEWQLKLQELDPATYKSISIYIEIQEF YVPLVDVNDDEFELTPI
3576	A	5	1421	LRLAWHDGARWPLGTPRAAATRREAAALPPVT LALLCLDGVLSSAENDFVHRIQEELDRFLQKQ LSKVLLFPPLSSRLRYLIHRTAENFDLLSSFSVGE GWKRRTVICHQDIRVPSSDGLSGPCRAPASCP YHGPRISNQGAAAVPRGARAGRWRGRKPDQ PLYVPRVLRQEEWGLTSTSVLKREAPAGRDP PGDVGAGDPNSDQGLPVLMTQGTEDLKGPGR CENEPLLDVPGPEPLGPESQSGKGDMVEMATRF

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GSTLQLDLEKGKESLLEKRLVAEEEEDEEEVEED GPSSCEDDYSELLQEITDNLTKKEIQIEKIHLDTS SFMEELPGEKDLAHVVEIYDFEPALKTEDLLATF SEFQEGFRIQWVDDTHALGIFPCRASAAEALTR EFSVLKIRPLTQGTKQSKLKALQRPKLLRLVKER PQTNATVARRLVARALGLQHKKKERPAVRGPLP P
3577	A	102	1998	DTRTPGSLEMGPLQFRDVAIEFSLEEWCHLDTAQ RNLYRNVMLNYSNLVFLGIVVSKPDIAHLEQG KKPLTMKRHEMVANPSGPVICSHFAQDLWPEQN IKDSFQKVILRRYEKRGHGNLQLIKRCESVDECK VHTGGYNGLNQCSTTTQSKVFQCDKYGKVFHK FSNSNRHNIRHTEKKPFKCECGKAFNQFSTLITH KKIHTGEKPYICEECGKAFKYSSALNTHKRIHTG EKPYKCDKCDKAFIASSTLSKHEIHTGKKPYKCE ECGKAFNQSSSTLTKHKKIHTGEKPYKCEECGKAF NQSSSTLTKHKKIHTGEKPYVCEECGKAFKYSRIL TTHKRIHTGEKPYKCNKCGKAFIASSTLSRHEFIH MGKKHYKCEECGKAFTWSSVLTRHKRVHTGEKP YKCEECGKAFKYSSTLSSHKRSHTGEKPYKCEEC GKAFVASSTLSKHBIHTGKKPYKCEECGKAFNQ SSSLTKHKKIHTGEKPYKCEECGKAFNQSSSLTK HKKIHTGEKPYKCEECGKAFNQSSSLTIKHKKIHT REKPYKCEECGKAFHLSTHLTTHKILHTGEKPYR CRECGKAFNHSATLSSHKKIHSGEKPYECDKCG KAFISPSLSRHEIHTGEKP
3578	A	1725	445	RPRRRGTHHFSCVLGSFRVSAMFPRVSTFLPLRP LSRHPLSSGSPETSAAAIMLLTVRHGTVRYRSSA LLARTKNNIQRIFYGTNSVICSKDKQSVRTEETS KETSESQDSEKENTKKDLLGIKGMKVELSTVNV RTTKPPKRRPLKSLEATLGRLRRATEYAPKKRIEP LSPELVAAASAVADSLPFDKQTTKSELLSQLQQH EESRAQRDAKRPKISFSNISDMKVARSAVARV RSRPELRJQFDEGYDNYPGQEKTDLLKKRKNIFT GKRLNIFDMMAVTKEAPETDTSPLWDVEFAKQ LATVNEQPLQNGFEEELIQWTKGKLWEFFINNEA GFDDDGSEFHEHIFLEKHLESFPKQGPRIHFMEV TCGLSKNPYLSVKQKVEHIEWFRNYFNEKKDILK ESNIQFKLRPWKFLFRNN
3579	A	1725	445	RPRRRGTHHFSCVLGSFRVSAMFPRVSTFLPLRP LSRHPLSSGSPETSAAAIMLLTVRHGTVRYRSSA LLARTKNNIQRIFYGTNSVICSKDKQSVRTEETS KETSESQDSEKENTKKDLLGIKGMKVELSTVNV RTTKPPKRRPLKSLEATLGRLRRATEYAPKKRIEP LSPELVAAASAVADSLPFDKQTTKSELLSQLQQH EESRAQRDAKRPKISFSNISDMKVARSAVARV RSRPELRJQFDEGYDNYPGQEKTDLLKKRKNIFT GKRLNIFDMMAVTKEAPETDTSPLWDVEFAKQ LATVNEQPLQNGFEEELIQWTKGKLWEFFINNEA GFDDDGSEFHEHIFLEKHLESFPKQGPRIHFMEV TCGLSKNPYLSVKQKVEHIEWFRNYFNEKKDILK ESNIQFKLRPWKFLFRNN
3580	A	3673	1619	LYCVAPYSRHLGRMSHLPMLLRKKIEKRNK LRQRNLKFQASNLTLSETQNGDVSEETMGSRK VKKSKQKPMNVGLSETQNGGMSQEA VGNKVT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				KSPQKSTVLTNGEAMQSSNSESKKKKKKKKRK MVNDAEPDTKKAKTENKKGKSEESAETTKETEN NVEKPDNDEDESEVPSPSLPLGLTGAFEDTSFASLC NLVNENTLKAIKEMGFTNMTEIQHKSIRPLLEGR DLLAAAKTGSGKTLAFLIPAVELIVKLRFMPRNG TGVLLSPTRELAMQTFGVKELMTHHVHTYGLI MGGSNRSAAEQKLGNGINIIVATPGRLLDHMQN TPGFMYKNLQCLVIDEADRILDVGFEELKQIIKL LPTRQTMFLFSAQTQTRKVEDLARISLKKEPLYVG VDDDKANATVDGLEQGYVVCPEKRFLLLFTFL KKNRKKKLMVFFSSCMSVKYHYELLNYIDLPLV AIHGKQKQNKRTTTFQFCNADSGTLLCTDVAA RGLDIPEVDWIVQYDPPDDPKEYIHRVGR TARGL NGRGHALLLRPEELGFLRYLKQSKVPLSEFDFS WSKISDIQSQLEKLEKNYFLHKSQAQYKSYTRA YDSHSLKQIFNVNINLNPQVALSFGFKVPPFVDL NVNSNEGKQKKRGGGGGFGYQKTKKVEKSKIF KHISKSSDSRQFSH
3581	A	23	453	LCRCICIKNITPHCLWDKVLQSQFTYILDNLSNFMS HHPHSLRNSCLIRMDLLYWQFTIYTITFCFSLSG RLTLSAQHISHRPCLLSYLLFWKVHHLFLEGFPC SPRLDEMSFHQPQHPVHVSVVHLPVYKGSMT QVSPH
3582	A	3	950	TRGCGNKMAGKKNVLSSLA VYAEDSEPESDGEA GIEAVGSAAEEKGGLVSDAYGEDDFSRLGGDED GYEEEEEDENSQSEDDDDSETEKPEADDPKDNT AEKRD PQELVASFSEVRNMSPEIKIPPEPPGRC SNHLQDKIQKLYERKIKEGMDMNYIIQRKKEFRN PSIYEKLIQFCAIDELGTNYPKDMFDPHGWS YEEALAKAQKIEMDKLEKAKKERTKIEFVTG KGTTTATSTTTTASTAVADACKRKS PVI TIAQPTILTTTATLPAVVTTTSASGSKTTVIS AVGTIVKKAQ
3583	A	3	950	TRGCGNKMAGKKNVLSSLA VYAEDSEPESDGEA GIEAVGSAAEEKGGLVSDAYGEDDFSRLGGDED GYEEEEEDENSQSEDDDDSETEKPEADDPKDNT AEKRD PQELVASFSEVRNMSPEIKIPPEPPGRC SNHLQDKIQKLYERKIKEGMDMNYIIQRKKEFRN PSIYEKLIQFCAIDELGTNYPKDMFDPHGWS YEEALAKAQKIEMDKLEKAKKERTKIEFVTG KGTTTATSTTTTASTAVADACKRKS PVT TIAQPTILTTTATLPAVVTTTSASGSKTTVIS AVGTIVKKAQ
3584	A	3	1139	PGSTISSRADRLGAPVLAHPKMAERQEEQRGSP LRAEGKADA EVKLILYHWTHSFSSQKVRVIAE KALKCEBHDVSLPLSEHNPEWFMRLNSTGEVPV LIHGENIICEATQIIDYLEQTFDERTPRLMPDKES MYYPVQHYRELLDSLPMDAYTHGCILHPELTV DSMPAYATTRIRSQIGNTESELKKLAEENPDLOE AYIAKQKRLKSKLLDHDNVKYLKILDELEKVL DQVETELPRRNEETPEEGQQPWLCGESFTLADVS LAVTLHRLKFLGFARRNWGNGKRPNLETYYERV LKRKTFNKVLGHVNNILISAVLPTAFRVAKKRAP KVLGTTLVVGLLAGVGYAFMLFRKRLGSMILA LRPRPNYF

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
3585	A	1	1777	RRHSPGSPAFAPSSRATAICPRAARAPATLLLAG AVLWPAAGAWELTILHTNDVHSRLEQTSDDSK CVNASRCMGGVARLFTKVQQIRRAEPNVLLDA GDQYQGTIWFVYKGAEVAFHFMNALRYDAMA LGNHEFDNGVEGLIEPLLKEAKFPILSANIKAKGP LASQISGLYLPYKVLPGDEVVGIVGYTSKETPF LSNPGTNLVFEDEITALQPEVDKLTNLNVNKIAL GHSGFEMDKLIAQKVRGVDVVVGHSNTFLYT GNPPSKEVPAGKYPFIVTSDGRKVPVVQAYAF GKYLGYLKIEFDERGNVISSHGPNILLNSSIPEDPS IKADINKWRIKLDNYSTQELGKTTIVLDGSSQSC RFRECNMGNLICDAMINNLRHTDEMFWNHVS MCILNGGGIRSPIDERNNGTITWENLAAVLPGG TFDLVQLKGSTLKKAFEHVHRYGQSTGEFLQV GGIHVVYDLSRKPGDRVVKLDVLCTKCRVPSYD PLKMDDEVYKVLNPNFLANGGDGFQMIKDELLRH DSGDQDINVVSTYISKMKVTPA VEGRIKFSTGS HCHGSFSLIFLSLWAVIFVLYQ
3586	A	1399	881	LSNKDVLSPQLKDENSEKLRRKLNEVQSFSEAQTE MVRTLERKLEAKMIKEESDYHDLESVVQQVEQN LELMTKRAVKAENHVVKLKQEISLLQAQVSNFQ RENEALRCGQGASLTVVKQADVALQNLRVVM NSAQASIEQLVSGAETLNLVAEILKSIDRISEVKD EEEDS
3587	A	88	1639	GCVGRGLPLPPRHPTPPSSSSSPFVLLAFLLLVRL DPAVSGKMAAPRPPPARLSGVMVPAPIQDLEAL RALTALFKEQRNRETAPRTIFQRLDILKKSSHA VELACRDPSQVENLASSLQLITECFRCLRNACIEC SVNQNSIRNLDTIGVAVDLILLFRELRLVEQESLT AFRCGLQFLGNIA SRNEDSQSIVVWHAFPELFSL CLPFDKTIAYSELT TNLNHERMKF LTL LN IADVIDAYQKHPESWPTLTDLFLKSPELVQA MFPKLNNQERVTLLEMIAKITSDEPLTKDDIPVF LRHAELIASTFVDQCKTVLLEA SEEPDDEEALA TIRLLDVLCEMTVNTTELLGYLQVFPGLLERVIDL LRVIHVAGKETTNIFSNCGCVRAEGDISNVANGF KSHLIRLIGNLCYKNKDNQDKVNELDGIPLLDN CNISDSNPFLTQWVITYAIRNLTEDNSQNQDLIAK MEEQGLADASLLKKVGFEVEKKGEKLILKSTRD TPKP
3588	A	3	1462	DSPRNRFEILGRPTRTPTRPGRPAMEDLDALLSD LETTTSHMPRSGAPKERPAEPLTPPPSYGHQPQT GSGESSGASGDKDHLYSTVCKPRSPKPAAPAAPP FSSSSGVLGTGLCELDRLQLNATQFNITDEIMS QFPSSKVASGEQKEDQSEDKKRPSLPSSPSPGLPK ASATSATLELDRLMASLSDFRVQNHLPASGPTQP PVVSSSTNEGSPSPPEPTGKGSLDTMLGLLQSDLSR RGVPTQAKGLCGSCNKPIAGQVVTALGRAWHPE HFVCGGCSTALGSSSFFEKDGAPFCPECYFERFSP RCGFCNQPIRHKMVTALGTHWHPEHFCCVSCGE PFGDEGFHEREGRPYCRRDFLQLFAPRCQGCQGP ILDNYISALSALWHPDCFVCRECFAPFSGGSFEEH EGRPLCENHFHARRGSLCATCGLPVTGRCVSAL GRRFHPDHFTCTFCLRPLTKGSFQERAGKPYCQP CFLKLFQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
3589	A	226	6793	SPKKSRKCNLSFRLISAERWRFFLLILMEMPRKP RLTLFVQRRINIATEREFDPEEFYLLLEAAEGHA KEGQGIKTIDPRYIISQLGLNKDPLEEAMHLGNY DSGTAETPETDESVSNSASLKLRRKPRESDFETI KLISNGAYGAVYFVRHKESRQRFAMKKINKQNL ILRNQIQQAFVERDILTFAENPFVVS MYCSFETR HLCMVMEYVEGGDCATLMKNMGPLPVDMMARM YFAETVLALAYLHNYGIVHRDLKPDNLLVTSMG HIKLTDGLSKVGLMSMTNL YEGHIEKDAREFL DKQVCGTPEYIAPEVILRQGYGKPDVWWAMGII LYEFVGCVPFFGDTPEELFGQVISDEINWPEKDE APPPDAQDLITLLLRQNPLERLTGGGAYEVKQHR FFRSLDWNLSLRQKAEFIPQLESEDDTSYFDTRSE KYHHMETEEEDDTNDEDFNVEIRQFSSCSHRFSK VFSSIDRITQNSAEKEDSVDKTKSTTLPSTETLS WSSEYSEMQLSTSNSSDTESNRHKLSSGLLPKL AISTEGEQDEAASCPGDPHEEPGKPALPPECAQ EEPEVTTTASTISSSTLSVGSFSEHLDQINGRSECV DSTDNSSKPSSEPASHMARQRLSTTEKKKISGV TKSLASALSMLPGDMFAVSPLGSPMSPHSLSSD PSSSRDSSPRDSSAASASPHQPIVHSSGKNYGFT IRAIRVYVGDSDIYTVHHIVWNVEEGSPACQAGL KAGDLITHINGEPVHGLVHTEVIELLLKSGNKVSI TTTFFENTSIKTGPARRNSYKSRMVRRSKSKKK ESLERRRSLFKKLAKQPSPLHTSRFSCLNRSLS SGESLPGSPHSLSPRSPTPSYRSTPDFSPGTNSSQ SSSPSSAPNSPAGSGHIRPSTLHGLAPKLGGQRY RSGRRKSAGNIPLSPLARTPSPTQPTSPQRSPL LGHSLGNSKIAQAFPSKMHSPPTIVRHIVRPKSAE PPRSPLLKRVQSEELSPSYGSDKKHLCSRKHSL EVTQFEVQREQEETAPIQSDENITMTTSLRA RPVIAQULKRPVSRKVGRQESVDESDRERKAK VVVKKADGFPEKQESHQKFHGPGLINFEALFK LEEREKKVYPKAVERSSTFENKASMQEAPLGS LKDALHKQASVRASEGAMSDGPVPAEHRQGGG DFRRAPAGTILQDGLCHSLDRGISGKGEGTEKSS QAKELLRCEKLDKLANIDYLRKKMSLEDKEDN LCPVLKPKMTAGSHECLPGNPVRPTGGQEQEPPA SESRAFVSSTHAAQMSAVSFVPLKALTGRVDSGT EKPGLVAPESPVRKSPSEYKLEGRSVSCLEPIEGT LDIALLSGPQASKTELPSESAQSPSPSGDVRSV PPVLPSSSGKKNDTTSARELSPSSLMNKSYLEP WFLPPSRGLQNSPAVSLPDPEFKRDRKGPHPTAR SPGTVMESNPQQREGSSPKHQDHTTDPKLLTCLG QNLHSPDLARPRCPLPPEASPSREKPLRESSERG PPTARSERSAARADTCREPSMELCFPETAKTSDN SKNLLSVGRTHPDFYTQTQAMEKAWAPGGKTN HKDGPGEARPPPRDNSSLHSAIPCEKELGKVRR GVEPKPEALLARRSLQPPGIESEKSEKLSFPLQ KDGAKEPERKEQPLQRHPSSIPPLTAKDLSSPA ARQHCSSPSHASGREPGAKPSTAEPSSSPQDPPKP VAAHSESSSHKPRPGPDGPPKTKHPDRSLSSQK PSVGATKGKEPATQSLGGSSREGKGHSKSGPDVF PATPGSQNKASDGIQGGGSPVPLHTDRAPLDA KPQPTSGGRPLEVLEKPVHLPRPGHPGPSEPADQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * =Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				KLSAVGEKQTLSPKHPKPSTVKDCPTLCKQTDN RQTDKSPSQPAANTDRRAEGKKCTEALYAPAE DKLEAGLSFVHSENRLKGAERPAAGVGKGFPEA RGKGGPGQKPPTADKPNGMKRSPSATGQSSFRS TALPEKSLSCSSFPETRAGVREASAASSDTSSAK AAGGMLELPAPSNRDRKAQPAGEGRTHMTKS DSLPSFRVSTLPLESHHPDPNTMGGASHRDRALS VTATVGETKGKDPAPAQPPARKQNVGRDVTKP SPAPNTDRPISLSNEKDFVVRQRRGKESLRSSPHK KAL
3590	A	3	935	RATTPKNEVQDYVSVEYLSPHMGGTDPFKYSY PPLVDDDFQTPLCENGPIITSEDETSSKEDIESDGK ETLETISNEEQTPLLKKINPTSTSKAEENEKVDS KVKAFFKKPLSVFKGPLLHISPAEELYFGSTESGEK KTLIVLTNVTKNIVAFKVRITTAPEKYRVKPSNSS CDPGASVDIVVSPHGGLTVSAQDRFLIMAAEME QSSGTGPAELTQFWKEVPRNKVMEHRLRCHTVE SSKPNTLTLKDNAFNMSDKTSEDICQLSRLLS NRKLEDQVQRCIWFQQLLSLTMLLAFVTSFFY LLYS
3591	A	303	2	GGSWGPLCPVSPAMSLSDPGLGYHPTCWTLRWP PLCSLHALHVHCLFSSRLGTPVSPRLAMPNCS CEAGGCACAGSCKCKCKCTSCCKSCCSCPL
3592	A	1052	1779	GKTMMRKMLLAAALSVTAMTAHADYQCSVTP RDDVIVSPQTVQVKGNGNLVITPDGNNVMYNGK QYSLNAAQREQAKDYQAEIRSTLPWIDEGAKSR VEKARIAIDKIIQEMGESSKMRSRLTKLDAQVK EQMNRHETRSDDLTFHYKAIDQVRAEGQQLVNQ AMGGILQDSINEMGAKAVLKSGGNPLQNVLGSL GGLQSSIQTWKKQEKDFQFGKDVCSRVTLE DSRKAALVGNLK
3593	A	3	1837	LSFEKVDIQTNDLTKEMYBKG \VSFELQRDFS QETDFSEASLLEKQEQEVHSAGNIKKEKSNTIDGT VKDETSPVEECFFSQSSNSYQCHTITGEQPSGCTG LGKSISFDTKLVKHEIINSEERPFKCEELVEPFRCD SqliQHqENNTeeKPYQCSECGKAfSINEKLIW QRLHSGEKPFKCVCEGKSFSYSSHYITHQTHSGE KPYQCKMCGKAfSVNGSLSRHQRIHTGEKPYQC KECGNGfSCSSAYITHQRVHTGEKPYECNDCGK AFNGNAKLIQHQRHTGEKPYECNECGKGfRCSS QLRQHQSHTGEKPYQCKEKGfGNNTKLIQH QRIHTASLAEQLFKASGNHPNWGCCLTISSPGPS VYGPKMNMrgAPNSRLAGGREKRTQDfDfGQC SFLPSHSPSCfEPWNVTdySSWYRQKQVLsgv WSSPLSILKLPRTLIRISIHQEMDTPGEMLMtGR GSLGPTLTTEAPAAAPGKQGPPGTGRCLQAPGT EPGEQTPEGARELSPLQESSPPGGVKAEEEEQRAG AEPGTRPSLARSDDNDHEVGALGLQQKSPGAG NPEPEQDCAARAPVRAEAVRRMPPGAEGSVVL DD
3594	A	39	261	RAAMMDTSRVQPIKLAIVIKVLGRTGSQGQCTQ VRVEFMDDTSRSIIRSVKGPVREGDVLTLLESERE ARRLR
3595	A	973	68	GRVGTKHQMADDAGAAGGPGGGPGGMGNRG GFRGGFGSGIRGRGRGRGRGRGRGARGGKAE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				DKEWMPVTKLGRLVKDMKIKSLEETLFLSLPIKE SEIDFFLGASLKDEVLKIMPVQKQTRAGQRTF KAFVAIGDYNHGVGLGVKCSKEVATAIRGAILA KLSIVPVRRGYWGNIKIGKPHTPVCKVTGRCSV LVRLIPAPRGTGIVSAPVPPKLLMMAGIDDCYTS ARGCTATLGNFAKATFDAISKTYSYLTPDLWKE TVFTKSPYQEFTHLVKTHTRVSVQRTQAPAVA TT
3596	A	106	2960	DERRVGAADMFGRSRSWVGGGHGKTSRNIHSL DHLKYLHYVLTKNNTVTEQNRNLLVETIRSITEL IWGDQNDSSVDFDFLEKNMFVFFLNILRQKSGRY VCVQLLQTLNLFENISHETSLYLLSNVYVNSII VHKFDFSDEEIMAYYISFLKTLKLNHTVHFF YNEHTNDFALYTEAIKFFNHPESMVRIAVRTITL NVYKVSLDNQAMLHYIRDKTAVPYFSNLVWFIG SHVIELDDCVQTDDEHRNRGKLSDLVAEHLDDL HYLNDILIINCEFLNDVLTDHLLNRLFLPLVYVSL ENQDKGGERPKISLPVSLYLLSQVFLIIHHAPLVN SLAEVILNGDLSEMYAKTEQDIQRSSAKPSIRCFI KPTETLERSLEMNKHKGKRRVQKRPYKNVGE EDEEKGPTEDAQEDAEEKAGTEGGSGIKTSGES EEIEMVIMERSKLSLAASVQEQNTTDEEKSA AATCSESTQWSRPFLDMVYHALDSPDDDYHALF VLCLLYAMSHNKGMDPEKLERIQLPVPNAEKT TYNHPLAERLIRIMNNAAPQDGKIRLATLESLCL LLKQQVLSAGCIMKDVHLACLEGAREESVHLV RHFYKGEDIFLDMFEDEYRSMTMKPMNVEYLM MDASILLPPTGTPLTGIDFVKRLPCGDVEKTRRAI RVFFMLRSLSLQLRGEPETQLPLTREEDLIKTDV LDLNNSDLIACVITKDGGMVQSRSLAVDIYQMS LVEPVSRLG...VKTALLO...VEDES R...TIHKPASSPHSKPPILQA...PS...RCHIAK QLAKGRIQARRMKMQRIAALLD...IQTTEVLG FGLGSSTSTQHLPRFYDQGRRGSSD...T...QRSVF ASVDKVPGFAVAQCINEHSSPSLSSQSPSSAGSP SGSGSTSHCDSSGTTSSSTPSTAQSPAGIGHVTQ
3597	A	427	277	GVRRIQHHWAQMHECNVHTYASLFLCLLHTG KLCCCLNSHRHFHCIKYSK
3598	A	1	503	FRPRTKKATAMYLEHYLDSIENLPCELQRNFQL MRELDQRTEDKKAIDILAAEYISTVKTLSPDQR VERLQKIQNAYSKCKEYSDDKVQLAMQTYEMV DKHIRRLDADLARFEADLKDKMEGSDFESSGGR GLKKGRGQKEKGRSGRGRRTSEEDTPKKKKH KGG
3599	A	2	3907	KTITALAFSPDGKYLVTGESGHMPAVRVWDVAB HSQVAELQEHKYGVACVAFSPSAKYIVSVGYQH DMIVNVWAWKKNIVVASNKVSSRVTAVSFSED CSYFVTAGNRHIKFWYLDSDSKSVNATVPLLG RSGLLGELRNNLFTDVACGRGKKADSTFCITSSG LLCEFSDRRLLDKWVELRVYPEVKDSNQACLPP SSFITCSSDNTRLWNTESGVHGSTLHRNLSDDL IKIYVDGNTQALLDTLPGGDKADASLLDPRVGI RSVCVSPNGQHLASGDRMGTLRVHELQSLSEML KVEAHDSEILCLEYSKPDGLKLLASASRDRLH VLDAGREYSLQQTLDHSSSITAVKFAASDGQVR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				MISCGADKSIYFRTAQKSGDGVQFTRTHHVVRK TTL YDMDVEPSWKYTAIGCQDRNIRIFNISSGKQ KKLFGKSQGEDGTLIKVQTDPSGIYIATSCSDKNL SIFDFSSGECVATMFGHSEIVTGMKPSNDCKHLIS VSGDSCIFVWRLSSEMTISMQRQLAELRQRQRGG KQQGPSPQRASGPNRHQAPSM LSPGPALSSSDSD KEGEDEGTEELPALPVLAKSTKKALASVPSPAL PRSLSHWEMSRAQESVGFDPAPAAPNGPRRRG RWVQPGVELSVRSM LDRQLET LAPSLQDPSQD SLAIPSGPRKHGQEALETSLTSQNEKPPRPQASQ PCSYPHIIRLLSQEEGVFAQDLEPAPIEDGIVYPEP SDNPTMDTSEFQVQAPARGTLGRVYPGSRSEK HSPDSACSVDYSSSCLSSPEHPTEDSESTEPLSVD GISSDLEEPAEGDEEEEEEGMGYPYGLQEGSPQ TPDQEQFLKQHFETLASGAAPGAPVQVPERSES SISSRFLQVQTRPLREPSSSSLALMSRPAQVPQ ASGEQPRGNGANPPGAPPEVEPSSGNPSPPQAA VLLPRCRLNPDSSWAPKRVATASPFSGLQKAQS VHSLVPQERHEASLQAPSPGALLSREIEAQDGLG SLPPADGRPSRPHSYQNPTTSSMAKISRISIVGEN LGLVAEPQAHAPIRVSPLSKLALPSRAHLVLDIPK PLPDRPTLAASFVPTKGRAPGEAEKPGFPVGLGK AHSTTERWACLGEGETTPKPRTECOAHGPGSSPCA QQLPVSSLFQGPENLQPPPPEKTPNPMECTKPGA ALSQDSEPAVSLEQCEQLVAELRGSVRQAVRLY HSVAGCKMPSAEQSRIAQLLRDTFSSVRQELEAV AGAVLSSPGSSPGAVGAEQTQALLEQYSELLRA VERRMERKL
3600	A	1688	916	IPGSTISCSMALCEAAGCGSALLWPRLLLFGDSIT QFSFQGGWGASLADRLVRKCDVLNRGFSGYN TNYAKILFLES'GNSLDIPVA'VTFGANDSAL KDLKQKHIPLEEYAAANLKSVMQYLSVDI NLITPTPLCETAWEEOCHQCKLNRLNSVVGEY ANACLOVAQDCGTDVLDLWTLMQDSQDFSSYL SDGLHLSPKGNBFLFSLWPLIEKKVSSLPLLLPY WRDVAEAKPELSLLGDGDH
3601	A	44	223	VHFLIPQLAKCFWTMNAARNKSEKRYYSEFL QIAHLFNYGLSSFLREFIIFLIKLLQ
3602	A	37	1124	VPKPASGKRRLEFRPQDSKACAA TPHSPGRITSR TRGSQKVRSPVPRLPWAQASASTDWEGLRGVPG PALRRENFLAAAASGRSGRTPTGGVGFRDVGGP HFPIFAAHFLWCNLHTPRRPACNAPWHSPVGEI SPPPRESQLRRDPEVHFESPAHPLGFRLLPGRGLP ANAVTVETAAMAAPRQIPSHIVRLKPSCSTDSSF TRTPVPTVSLASREL PVSSWQVTEPSSKNLWEQI CKEYEAEPFPEGYKVKQEPVITVAPVEEMLFH GFSAEHYFPVSHFTMISRTPCPQDKSETINPKTCS PKEYLETFIFVLLPGMASLLHQAKKEKCFEVVL QMTPSGGKACVWGHL PSSHTI
3603	A	286	587	NISNKAEVSSHPSVISHMSDFSQPRPEDNQSVLR RMQKKYWKTKQVFIKATGKKEDEHLVASDAEL DAKLEV FHSVQETCTELLKIEKYQLRLNGMKS
3604	A	103	2440	QPRRRVFPAAGRGPGRKCSQWGRQASVSFEDVT VDFSKEEWQHLDPAQRRLYWDVTLENYSHLLS VG YQIPKSEAAFKLEQGEGPWMLEGEAPHQSCS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				GEAIGKMQQQGIPGGIFFHCERFDQPIGEDSLCSI LEELWQDNDQLEQRQENQNLLSHVKVLIKERG YEHKNIEKIIHVTTKLVPISIKRLHNCDTILKHTLN SHNHNRRNSATKNLGKIFGNGNNFPHSPSTKNEN AKTGANSCEHDHYEKHLSHKQAPTHHQKIHPEE KLYVCTECVMGFTQKSHLFEHQRIHAGEKSREC DKSNKVFQKPQVDVHPSVYTGEKPYLCTQCCK VFTLKSNIHQIHTGQKPYKCECGKAFFQRS DLFRHLRIHTGEKPYECSECGKGFSONSDLSIHQ KTHTGEKHYESNECGKAFTRKSAALRMHQRIHTG EKPYVCADCGKAFIQKSHFNTHQRIHTGEKPYEC SDCGKSFTKKSQHLVHQRIHTGEKPYICTECGKV FTHRTNLTHQKTHTGEKPYMCAECGKAFTDQS NLIKHKQTHTGEKPYKCNCGKAFIWKSRLLKIH QKSHIGERHYECKDCGKAFIQKSTLSVHQRIHTG EKPYVCECGKAFIQKSHFIAHRIHTGEKPYECS DCGKCFTKKSQLRVHQIHTGEKPNICAECGKAF TDRSNLTHQKIHTREKPYECGDCGKTFTWKSRL NIHQKSHTERHYECSKCGKAFIQKATLSMHQII HTGKKPYACTECQKAFTDRSNLIKHKMHSGEK RYKASD
3605	A	3	322	SFRMSGRGKGGKGLGKGGAKRHRKVLDRDNIQGI TKPAIRRLARRGGVKRISGLIYEETRGVLKVFLN VIRDAVTYTEHAKRKTVTAMDVVYALKRQGR LYGFGG
3606	A	1	1749	VPVTAEAKLMGFTQGCVTTFEDVAIYFSQEEWGL LDEAQRLLYRDVMLENFALITALVCWHGMEDE ETPEQSVSVEGVVPQVTPASPSTQKIQSCDMCV PFLTDILHLTDLPGQELYLTGACAVFHQDQKHHS AEKPLESDMDKASFVQCCLFHESGMPFTSSEVG KDFLPLGLQCCMANYEKPKN'SKCEAFHVGE SHYKWSQCEKESLHTFFHPRVCTGKRLYESS KCGKACCCECLVQLQRVHPGERPYECSECGKS FSQTSHLNDHKKHTHTGERPYVCGQCGKSFSQRAT LIKHHRVHTGERPYECGCGKSFSQSSNLIEHCRI HTGERPYECDECGKAFGSKSTLVHRHQRTHTGEK PYECGECGKLFRQSFLVVHQRIHTTARPYECGQ CGKSFSCLKGLIQHQLHSGARPFECDECGKSFSQ RTTLNKHKKVHTAERPYPVCGECGKAFMFKSKL VRHQRTHTGERPFECSECGKFFRQSYTLVEHQKI HTGLRPYDCGQCGKSFIQKSSLIQHQQVVHTGERP YECGKCGKSFTQHSGLILHRKSHTVERPRDSSKC GKPYSPRSNIV
3607	A	92	331	AMAGPGPGGDPDEQYDFLKLVLVGDA SVGKT CVVQRFKTGAFSERQGSTIGVDFTMKLEIQGKR VKLQIWDTAGQER
3608	A	545	379	AIKGYIHLSAPRNRMYHTTASNGRMLFMKVMTM YMRRGVQIMGWSVRMAFMACFTQ
3609	A	118	873	VWMAWQVSLLELEDRLQCPICLEVFKE SMLQC GHSYCKGCLVSLSYHLDTKVRCPMCWQVVDGS SSLPNVSLAWVIEALRLPGDPEPKVCVHHRNPLS LFCEKDQELICGLCGLLGSHQHHPVTPVSTVCSR MKEELAALFSELKQEQKKVDELI AKLVKNRTRIV NESDVFSWVIRREFQELRHPVDEEKARCLEGIGG HTRGLVASLDMQLEQAQGTRERLAQAECVLEQF

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				GNEDHHEFTWKFHSMASR
3610	A	2	987	DPRVRPPLLQPPPPLLPRLVILKMAPLDLDKYVEI ARLCKYLPENDLKRCLDYVCDLLLEESNVQPV TPVTVCGLDIHQFYDLCELFRTGGQVPDTNYIFM GDFVDRGYYSLETFTYLLALKAKWPDRITLLRG NHESRQITQVYGFYDECQTKYGNANAWRYCTK VFDMLTVAALIDEQILCVHGGSPDIKTLTDQIRTI ERNQEIPIHKGAFCDLVWSDPEDVDTWAI SPRGA GWLFGAKVTNEFVHINNKLICRAHQLVHEGYK FMFDEKLVTVWSAPNYCYRCGNASIMVFKDVN TREP KLFRAVPDSERVIPRTTTPYFL
3611	A	2459	869	AEKMTAELREAMALAPWGPVKVKKEEEEENF PGQASSQQVHSENIKVWAPVQGLQTGLDGSEEE EKGQNISWDMAVVLKATQEAAPASTLGSYSLPG TLAKSEILETHGTMNFLGAETKNLQLLVPKTEIC EEAEKPLIISERIQKADPQGPPELGEACEKGNMLK RQRIKREKKDFRQVIVNDCHLPESFKEEENQCKK KSGGKYSLSNGAVKNPKTQLGQKPFTCSVCGKG FSQSANLVVHQRIHTGEKPFECHECGKAFIQSAN LVVHQRIHTGQKPYVCSKCGKAFTQSSNLTVHQ KIHSLEKTFKCNECEKAFFSYSSQLARHQKVHTE KCYECNECGKTFTRSSNLIVHQRIHTGEKPFACN DCGKAFTQSANLIVHQRSHTGEKPYECKECGKA FSCFSLIVHQRIHTAEKPYDCSECGKAFFSLSCL IVHQRIHSGDLPYVCNECGKAFTCSSYLLIHQRIH NGEKPYTCNECGKAFTQRSSLTVHQRIHTGEK YECEKCGAFAFISNSHLMRHHRTHLVE
3612	A	318	2245	SPMAEALVNTQPIMVTEEFVKPSQGHVTFEDI AVYFSQEEWGLLDEAQRCLYHDVMLENFSLMA SVGCLHGIEAEEAPSEQTLQAQGVSAQRTPKLGP SIPNAH3CEM'CT VMKDILZLSEH'GTLPWQKPY TSVASGKWFSFGSNLQQHQNQD3GEKKEEES ALLNSCKIPLSDNLFCKDVEKDFPTILGLLQHQ TTHSRQEYAHRSRETFQORRYKCEQVFNEKVHV TEHQRVHTGEKAYKRREYKSLNSKYLVEHQ THNAEKPYVCNICGKSLHKQTLVGHQQRHTRE RSYVCIECGKSLSSKYSLEHQRTTHNGEKPYVCN VCGKSFRHKQTFVGHQQRHTGERPYVCMCEGK SFIHSYDRIRHQRVHTGEGAYQCSECGKSFIYKQ SLLDHHRHTGERPYECKECGKAFFHKRLLEHQ RIHTGEKPYVCIICGKSFISSDYMRHQRIHTGER AYECSDCGKAFISKQTLKHHKIHTREPYECSE CGKGFYLEVKLLQHQRIHTREQLCECNECGKVF SHQKRLEHQKVHTGEKPCCECGKCFRHRSTS LIHQKQVHSGERPYNCTACEKAFTYKNKLVEHQ RIHTGEKPYECGKCGKAFNKRYSLVRHQKVHIT EEP
3613	A	817	3345	NQSHPDSETVTVEGGRRKMKSQERSNECLPPK KREIPATSRSEEKAPTLPSDNHRVEGTAWLPGN PGGRGHGGRRHGPAAGTSVELGLQQGIGLHKALS TGLDYSPPSAPRSVPVATTLPAAYATPQGPVSP VQYAHLPHTFQFIGSSQYSGTYASFIPSQLIPTAN PVTSASASAAGATTPSQRSQLEAYSTLLANMGS LSQTPGHKAEQQQQQQQQQQQQQQQQQQQQQ QQHQQQQQQQQQQQQQQQQQHLSRAPGLITPGSP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				PAQQNQYVHISSSPQNTGRTASPPAIPVHLHPHQ TMIPHTLTGPPSQVVMQYADSGSHFVPREATK KAESSRLQQAIAKEVLNGEMEKSSRYGAPSSA DLGLGKAGGKSVPHPYESRHVVVHPSPSDYSSR DPSGVRASVMVLPNSNTPAADLEVQQATHREAS PSTLNDKSGHLHGKPGHRSYALSPHTVIQTTHSA SEPLPVGLPATAFYAGTQPPVIGYLSGQQQAITY AGSLPQHLVIPGTQPLLIPVGSTDMEASGAAPAI TSSPQFAAVPHTFVTTALPKSENFNPEALVTQAA YPAMVQAQIHLPPVQSVASPAAPPTLPPYFMK GSIIQLANGELKKVEDLKTEDFIQSAEISNDLKIDS STVERJEDSHSPGVAVIQFVGEHRAQVSVEVLV EYPFFVFGQGWSGCCPERTSQLFDLPCKLSVGD VCISLTILKNLKNKSGVKKGGQPVDPASVLLKHSKA DGLAGSRHRYAEQENGINEQGSQAQMLSENGELKF PEKMGLSAAFPFLTKIEPSKPAATRKRRWSAPESR KLEKSEDEPPLTLPKPSLIPQEVKICIEGRSNVKG
3614	A	3	114	FFESRLRCKCCEPRGSGWARFGCWRLQPEFKPKQ LEG
3615	A	3	1603	DAWALTNQFSDSKQHIEVLKESLTAKEQRAAILQ TEVDALRLRLEEKETMLNKKTKQIQDMAEEKGT QAGEIHDLDKMDLVKERKVNVLQKKIENLQEQ RDKEKQMSSLKERVKSQADTTNTDTALTLEE ALAEKERTIERLKEQRDRDEREKQEEIDNYKKDL KDLKEKVSLQGDSEKEASLLDLKEHASSLASS GLKKDSRLKLEIALEQKKEECKMESQLKKAH EAALEARASPEMSDRIQHLEIRETRYKDESSKAQ AEVDRILLEILKEVENEKNDKDKKIAELESLSRQ VKDQNKKVANLKHKEQVEKKKSAQMLEEARRR EDNLNDSSQQLQDSLRRKKDDRFEIEALRESVQ ITAEREMVLAQEESARTNAEKQ... KQELSMKAKLSST...SLAEKETHLTNLAERR KHLEEVLEMKEALLAAISEKDANIALELSSSK KKTQEEVAALKREKDRLVQQLKQQTQNRMKLM ADNYEDDHFKSSHSNQTNHKPSPDQDEEEGIWA
3616	A	244	1420	RRRWRARGGLVPTLAWAEATGAYVPGRDKPD PTWKRNFRLNRKEGLRLAEDRSKDPHPHKI YEFVNSGVGDFSQPDTSPTDNGGGSTSDTQEDIL DELLGNMVLAPLPDPGPPSLAVAPEPCPQLRSPS LDNPTFPNLPSENPLKRLLVPGEEWFEVTA YRGRQVFQQTISCPEGLRLVGSEVGDRTLP VTLDPDGMSTDRGVMSYVRHVLSCLGGLAL WRAGQWLWAQRLGHCHTYWAVSEELLPSNGH GPDGEVPKDKEGGVFDLGPFFVGSGLGPPDLITFE GSGRSPRYALWFCVGESWPQDQPWTKRLVMVK VVPTCLRALVEMARVGGASSENTVDLHISNSHP LSLTSDQYKAYLQDLVEGMDFGQPGES
3617	A	852	304	RGGLLSKMARVLKAAAANAVGLFSRLQAPIPTV RASSTSQPLDQVTGSVWNLGRNLHVAIAVPDLE KAAAFYKNILGAQVSEAVPLPEHGVSVVFNLG NTKMELLHPLGRDSPAGFLQKNKAGGMHHICIE VDNINAAVMDLKKKKIRSLSEEVKIGAHGKPVIF LHPKDCGGVLVELEQA
3618	A	3	5992	DNIDETYGVNVQFESDEEEGEDVDYGEVREEAS DDDMEGDEAVVRCTLSANMYVDEILVWCASEL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, /-possible nucleotide insertion)
				<p> NIPEFFPLESPHKKVG YGLSSRTWLQGGGKVIEA GRDLLVASGELMSSKKKDLHPRDIDAFWLQRQL SRFYDDAIVSQKKADEVLEILKTASDDRECENQL VLLLGFNFTDFIKVLRQHRMMILYCTLLASAQSE AEKERIMGKMEADPELSKFLYQLHETEKEDLIRE ERSRRERVRQSRMDTDLETMDLDQGGALAPRQ VLDLEDLVFTQGSFHMANKRCQLPDGSFRRQRK GYEEVHPALPKPKPGSSEEQLLPVEKLPKYAQA GFEGFKTLNRIQSKLYRAALETDENLLLCAPTGA GKTNVALMCMLEIGKHINMDGTINVDDFKIYI APMRSVLQEMVGSFGKRLATYGITVAELTGDHQ LCKEEISATQIVCTPEKWDIITRKGGERTYTQLV RLIILDEIHLHDDRGPVLEALVARAIRNIEMTQE DVRLIGLSATLPNYEDVATFLRVDPKGLFYFDN SFRPVPLEQTYVVGITEKKAIRKFOIMNEIVYEKIM EHAGKNQVLVPVHSRKETGKTARAIKDMCLEKD TLGLFLREGSASTEVLRTEAEQCKNLEKDLLPY GFAIHAGMTRVDRTLVEDLFGDKHIQVLVSTA TLAWGVNLPHTVIKGTQVVSPEKGRWTELGA LDILQMLGRAGRPQYDTKGEGILITSHGELQYYL SLLNQQLPIESQMVSKLPDMLNAEIVLGNVQNA KDAVNWLGYAYLYIRMLRSPTLYGISHDDLKGD PLLDQRRDLVHTAALMLDKNNLVKYDKKTGN FQVTELGRIASHYYITNDTVQTYNQLLKPTLSEIE LFRVFSLSSEFKNITVREEKLELQKLLERVPVVK ESIEEPSAKINVLLQAFISQLKLEGFALMADMVY VTQSAGRLMRAIFEIVLNRGWAQLTDKTLNLCK MIDKRMWQSMCPLRQFRKLPEEVVKKIEKKNFP FERLYDLNHNEIGELIRMPKMGKTIHKYVHLFPK LELSVHLQPIRSTLKVLTITPDFQWDEKVHGSS EAFWVVERNTSTLHHEITLTAQAQDEHLA TFFVPVFEHLPQTLVSDRWLSCETQLPVSFR HLILPEKYPPFELLDLQPLPVSALRNSAFESLYQ DKFPFENPIQTQVNTVYNSDDNVFVGAPTGSBK TICAEFAILRMLLQNSETRCVYITPMRLWQEQVY MDWYEKFQDRLNKKVLLTGETSTDLKLLGKG NIIISTPEKWDILSRRWKQRKNVQNNLFFVDEV HLIGGENGPVLEVICSRMRYISSQIERPIRIVALSSS LSNAKDV AHWLGCSATSTFNHFNVRPVPLELHI QGFNISHTQTRLLSMAKPVFHAITKHSPKKPVIVF VPSRKQTRLTAIDILTTCAADIQRQFLHCTEKDL IPYLEKLSDSLKETLLNGVGYLHEGLSPMERRL VEQLFSSGAIQVVVASRSLCWGMNVAHLVIIM DTLYYNGKIHAYVDYPIYDVLQMVGHANRPLQ DDEGRVCVIMCQGSKKDFKKFLYEPLPVESHLD HCMHDHFNAEIVTKTIENKQDAVDYLTWTFLYR RMTQNPNNYNLQGISHRHLSHLSSELVEQTLSDL EQSKCISIEDEMDVAPLNLGMIAAYYYINYTTIEL FSMSLNAKTKVRGLIEISNAEYENIPIRHHEDN LLRQLAQKVPKLNPKFNDPHVKTNLLQAHN SRMQLSAELQSDTEILSKAIRLIQACVDVLSNG WLSPALAAMELAQMVTQAMWSEDSYLRRLPPF PSGLFKRCTDKGVESVFDIMEMEDEERNALLQLT DSQIADVARFCNRYPNIELSYEVVDKDSIRSGGP VVVLVQLEREEVETGPVIAPLFPQKREEGWVVV </p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				IGDAKSNSLISIKRLTLQOKAKVKLDFVAPATGG RHNTLYFMSDAYMGCDQEYKFSVDVKEAETDS DSD
3619	A	3	5992	DNIDETYGVNVQFESDEEEGDEDVYGEVREEAS DDDMEGDEAVVRCTLSANMYVDEILVWCASEL NIPEFFPLESPHKKVGYGLSSRTWLQGGGKVIBA GRDLLVASGELMSSKKKDLHPRDIDAFWLQRQL SRFYDDAIVSQKKADEVLEILKTASDDRECENQL VLLGFNTFDIKVLRQHRMMILYCTLLASAQSE AEKERIMGKMEADPELSKFLYQLHETEKEDLIRE ERSRRERVRQSRMDTDLETMDLDQGGALAPRQ VLDLEDLVFTQGSFMANRCQLPDGSFRRQRK GYEEVHVPALPKPFGSEEQLLPVEKLPKYAQA GFEGFKTLNRIQSKLYRAALETDENLLLCAPTGA GKTINVALMCMLREIGKHINMDGTINVDDFKIYI APMRSLVQEMVGSFGKRLATYGITVAELTGDHQ LCKEISATQIIVCTPEKWDITRKGGERTYTQLV RLIILDEIHLHDDRGPVLEALVARAIRNIEMTQE DVRIGLSATLPNYEDVATFLRVPKGLFYFDN SRFPVPLEQTYVGITEKKAIRFQIMNEIVYEKIM EHAGKNQVLVVFVHSRKETGKTARAIKDMCKED TLGLFLREGSASTEVLRTEAEQCKNLEKDLLPY GFAIHAGMTRVDRTLVEDLFGDKHIQVLVSTA TLAWGVNLPAAHTVIKGTQVVSPEKGRWTELGA LDILQMLGRAGRPQYDTKGEGILITSHGELQYYL SLLNQQLPIESQMVSKLPDMLNAEIVLGNVQNA KDAVNWLGYAYLYIRMLRSPTLYGISHDDLKGD PLLDQRRDLVHTAALMLDKNNLVKYDKKTGN FQVTELGRIASHYYITNDTVQTYNQLLKPTLSEIE LFRVFSLSSEFKNITVREEKLELQKLLERVPVVK ESIEEPSAETVLCAANSOITETETAMADMVY FQTSAGRLMRAIFEIVLNRCMAQLPKTLNLCK MIDKRMWQSMCPLRQFRKLEAVVKKIEKKNFP FERLYDLNHNEIGELIRMPKMGKTAKYVHLFPK LELSVHLQPIRSTLKVLTITPDFQWDEKVVHGSS EAFWILVEDVDSEVILHHEYFLKAKYAQDEHLI TFFVPVFEPLPPQYFIRVVS DRWLSCETQLPVSR HLILPEKYPPTELLDLQPLPVSA LRNSAFESLYQ DKFPFFNPIQTQVFNTVYNSDDNVFVGAPTGS GK TICAFAILRMLLQNSEGRCVYITPMRLWQE QVY MDWYEKFQDRLNKKVVLLTGETSTD LKLLGKG NIIISTPEKWDILSRRWKQRKNVQNNL FVVDEV HLIGGENGPVLEVICS RMRYISSQIERPIRIVALSSS LSNAKDVAHWLGCSATSTFNHFPNVRPVLELHI QGFNISHTQTRLLSMAKPVFHAITKHSPKKPVTVF VPSRKQTRLTAIDILTTCAADIQRQFLHCTEKDL IPYLEKLSDSLKETLLNGVGYLHEGLSPMERRL VEQLFSSGAIQVVVASRSLCWGMNVA AHLVIIM DTLYYNGKIHAYVDYPIYDVLQMVGHANRPLQ DDEGRCVIMCQGSKKDFFKKFLYEPLPVESHLD HCMHDFHNAEIVTKTIENKQDAVDYLTWTFLYR RMTQNPNYYNLQGISHRHLSDHLSSELVEQTLSDL EQSKCISIEDEMDVAPNLGMIAAYYYNYTTIEL FSMSLNAKTKVRGLIEISNAAEYENIPIRHHEDN LLRQLAQKVPHKLNPNPKFNDPHVKTNLLQAH

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SRMQLSAELQSDTEELSKAIRLIQACVDVLSSNG WLSPALAAMELAQMVTQAMWSEDSYLRLPPF PSGLFKRCTDKGVESVFDIMEMEDEFERNALLQLT DSQIADVAFRCNRYPNIELSYEVVDKDSIRSGGP VVVLVQLEREEVETGPVIAPLPQKREEGWWVV IGDAKSNSLISIKRLTLQKAKVKLDFVAPATGG RHNTLYFMSDAYMGCDQEYKFSVDVKEAETDS DSD
3620	A	1205	323	VIKMALAARLLPQFLHSRSLPCGAVRLRTPAVAE VRLPSATLCYFCRCRLGLGAALFPRSARALAASA LPAQGSRWVPLSSPGLPAAFASFPACQRSYSTE EKPQQHQKTKMIVLGFSNPINWVRTRIKAFLIWA YFDKEFSITEFSEGAQAFAHVSKLLSQCKFDLL EELVAKEVLHALKEKVTSLPDNHKNALANIDEI VFTSTGDISIYDEKGRKFVNILMCFWYLTANIP SETLRGASVFQVKLGNQNVETKQLLSASYEFQR EFTQGVKPDWTIARIEHSLLE
3621	A	2	2995	SSSRSRHSSISPVRLPLNSSLGAELSRKKKRAAAA AAAAKMDGKESYERSGSYSGRSPSPYGRSSSS PFLSKRSLRSPLPSRKSMKSRSRSPAYSRHSSSH SKKKRSSSRSRHSSISPVRLPLNSSLGAELSRKKK ERAAAAAAKMDGKESYERSGSYSGRSPSPYGR RRSSSPFLSKRSLRSPLPSRKSMKSRSRSPAY RHSSSHSKKKRSSSRSRHSSISPVRLPLNSSLGAEL SRKKKRAAAAAAAKMDGKESKGSVPFLPRKE NSSVEAKDSGLESKKLPRSVKLEKSAPDTELNVN THLNTEVKNSSDTGKVKLDENSEKHLVKDLKAQ GTRDSKPIALKEEIVTPKETETSEKETPPPLPTIASP PPPLPTTTTPPQTPPLPPLPPIPALPQQPPLPPSQPA FSQVPASSTSTLPPSTHKTSAVSSQANSQPPVQV CVKTQVSVTAAPHLKTEGPIPIPIPLPCBDM DSPKETLPSKWKKEKEQTRHLLTDLPLPPLP GDLSPDSPEPKAITPPQPYKKRPKICCPRYGUR RQTESDWGKRCVDKFDIIGIGEGTYGQVYKAKK KDTGELVALKKVRLDNEKEGFPITAREIKILRQL IHRSVVNMKEIVTDKQDALDFKKDKGAFYLVFE YMDHDLMLGLESGLVHFSEDHIKSFMKQLMEGL EYCHKKNFLHRDIKCSNILLNNSGQIKLADFGLA RLYNSEESRPYTNKVITLWYRPPKLLGEERYTP AIDVWSCGILGELFTKKPIFQANLELAQLELISR LCGSPCPAVWPDVIKLPYFNTMKPKKQYRRRLR EEFSFIPSAALDLLDHMLTLDPSKRCTAEQTLQSD FLKDVELSKMAPDLPWHQDCHELWSKKRRRR RQSGVVVEEPPPSKTSRKETTSGTSTEPVKNSSPA PPQPAPGKVESGAGDAIGLADITQQLNQSELAVL LNLLQSQTDLSPQMAQLLNHNSNPEMQQLEAL NQSISALTEATSQQQDSETMAPEESLKEAPSAPVI LPSAEQTTLEASSTPADMQNILAVLLSQLMKTQE PAGSLEENNSDKNSGPGPRRTPTMPQEEAAGRS NGGNAL
3622	A	16	390	TPERGSAYPETAAVRRPAGECPITMSDLEAKLST EHLGDKIKDEDIKLRVIGQDSSEIHFVKMTTPLK KLKKSQCQRQGVVNSLRLFEQQRADNHTPEE LGMEEDVIEVYQEIQIGHSTV
3623	A	2	1544	PPPAPGPDGLNEGCLHRLSMPHQRPRTCAMNPE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				LTMESLGLHARGGGGGGGGGGGGGGGGGGGGGP GHEQELLASPSPHHARRGPRGSLRGPPPPPTAHQ ELGTAAAAAASRSAMVTSMASILDGGDYRPE LSIPLHHAMSMSCDSSPPGMGMSNTYTTLTPLQP LPISTVSDKFHHPHPHHHHHHHHHHQRLSGN VSGSFTLMRDERGLPAMNNLYSPYKEMPGMSQS LSPLAATPLGNGLGGLHNAQQSLPNYGPPGHDK MLSPNFDAAHTAMLTRGEQHLRGLGTPPAAM MSHLNLGHHHPGHTQSHGPVLAPSRERPPSSSSGS QVATSGQLEEINTKEVAQRITAELKRYSHIPQAIFA QRVLCRSQGTLSDLLRNPKPWSKLKS GRETFRR MWKWLQEPEFQRM SALRLAACRKEQEPNKDR NNSQKKSRLVFTDLQRRTLFAIFKENKRPSKEMQ ITISQQLGLELTTVSNFFMNAARRRSLEKWQDDLS TGGSSSTSSTCTKA
3624	A	27	2152	SARKAEAATSGTAARDGSVGRNLVPPPSASAPK AEVESNEKDNRPEEEEQVIHEDDERPSEKNEFSR RKRSKSEMDNVQSKRRRYMEEYEA EFQVKIT AKGDINQKLQKVIQWLLEKLCALQCAVFDKTL AELKTRVEKIECNKRHKTVLTELQAKIARLTKRF EAAKEDLKKRHEHPPNPVSPGKTVNDVNSNNN MSYRNAGTVRQMLESKRNVSESAPPSFQTPVNT VSSTNLVTPPAVVSSQPKLQTPVTSGLTATSVP APNTATVVATTQVPSGNPQTISLQPLPVILHVPV AVSSQPQLLQSHPGTLVTNQPSGNVEFISVQSPT VSGLTKNPVSLPSLPNPTKPNVPSVSPSIQRNP TASAAPLGTTLAVQA VPTAHSIVQATRTSLPTVG PSLYSPSTNRGPIQMKIPISAFSTSSAAEQNSNTT PRIENQTNKTIDASVSKKAADSTSQC GKATGSDS SGVIDLTMDDESGASQDPKKNHTPVSTMSSSQ FVSRLQPIQAPPLQPSGVTSQFSCITITL LPTA PNTVNVTHRPVTQVITRLPVPRAPANE PTT LPAPPAQAFLRGTMQAPAVRQVNPQNSVTVRV PQTTTYVNNGLTLGSTGPQLTVHHRPPQVHTEP PREVHPAPLPEAPQPQLPPEAGSTSRPSEATLEV SHAFRVKMAIVLVMCEPGGSKLCHC
3625	A	210	1115	ASFLRPQGHDSGEREPFSQTPGLMQFFSIPVQIT LQGSRRRQGRTAFFASGKKRETDYSDGDPDLDVH KRLPSSTGEDRAVMLGFAMMGFSVLMFFLLGTT ILKPFMLSQREESTCTAIHTDIMDDWLDCAFTCG VHCHGQGGKYPCLQVFVNL SHPGQKALLHYNEE AVQINPKCFYTPKCHQDRNDLLNSALDIKEFFDH KNGTPFSCFYSPASQSEDVILIKKYDQMAIFHCLF WPSLTLLGGALIVGMVRLTQHLSLLCEKYSTVV RDEVGKVPYIEQHQFKLCIMRRSKGRAEKS
3626	A	9	921	SSVVEFSALSVSMACLSPSQLQKFQDGFVLVLEG FLSAEECVAMQQRIGEIVAEMDVPLHCRTEFSTQ EEEQLRAQGSTDYFLSSGDKIRFFFEKGVFDEKG NFLVPPEKSINKIGHALHAHPVFKSITHSFKVQT LARSLGLQMPVVVQSMYIFKQPHFGGEVSPHQD ASFLYTEPLGRVLGVWIAVEDATLENGCLWFIPG SHTSGVSRRMVRAPVGSAPGTSFLGSEPARDNSL FVPTPVQRGALVLHGEVVHKSQNLSDRSRQA YTFHLMASGTTWSPENWLQPTAELPFPQLYT
3627	A	231	644	INSSPRTGRDHQELNLHTERDSRSQRAVLKIPRQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				NPGIFYWIFLPSRSHSASHGSRQRQVSCQGTQDEI LKMRNTFAELKNSLEALSSRMDQAERIGTQAG VQWRDHGSLQPQPPEFKQCFHLSLPSSWDYRAC LS
3628	A	2	810	GCKHLLQNSWYDPRVREADRVGQARRPRAAM DWLMGKSKAKPNGKKPAAEERKAYLEPEHTKA RITDFQFKELVVLPREIDLNEWLASNTTFFHHIN LQYSTISEFCTGETCQTMVAVCNTQYYWYDERGK KVKCTAPQYVDFVMSSVQKLVTDEDVFTKYG REFPSSFESLVRKICRHLFHVLAHTYWAHFKETLA LELHGHNLNTLYVHFILFAREFNLDPKETAJMDL LTVLCSGGRRGSTVGA VGMGPAAGAPGAQNH VKER
3629	A	699	1604	CSHGSSAVSAWSPLFQASEVERQLSMQVHALRE DFREKNSSTNQHIIRLESQAIEKMLSDRKRELEH RLSATLEENDLLQGTVEELQDRVLILRQGHDKD LQLHQSLQLQEVRLSCRQLQVKVEELTEERSLQ SSAATSTSLLEIEQSMEAELEQEREQLTLLSVE MTALKEERDRLRVTSDEKPEQLQKAIRDRE AIAKKNAVELELAKCRMDMMSLNSQLLDAIQQ KLNLSQLLEAWQDDMHRVIDRQLMDTHLKERS QPAAALCRGHSAGRGDEPSIAEGKRLFSFRKI
3630	A	423	1	PAKVLTDIYLSKTEGAQVDEPVVITPRAEDCGD WDDMEKRSSGRRSGRRRSGSKTSDSPGADALP ESAARDDAVFDDEVAPNAASDNASAEKKVKSPR AALDGGVASAASPEKSPSGTKGQLRGESDRSK QPPPASSP
3631	A	2082	674	WSGFWQLPGVRGVGSAPGGDGAFTSRRGSSRR PGAACPGCRGAGSERAPGGMGRRRAPELYRAPF PLYALQVDPETGLLIAAGGGGAAGTGIKNGVHF LOETLINGRIALLNHLTETLALAGD LALGQDAHCQLLRFAQHQQC KAENLGSKEQ GPRQRGAAPAEKKCGAETQHEGLRVENLQA VQIDFSSDPLQKVVCFNHDNTLLATGCTDGYVR VWKVPSLEKVFLEKAHEGEIEDLALGPDGLVT VGRDLKASVWQKDQLVTQLHWQENGPTFSSTP YRYQACRFGQVPDQAGLRLFTVQIPHKRLRQPP PCYLTAWDGSNFLPLRTKSCGHEVVSLDVSES GTFLGLGTVTGSVAIYIAFSLQCLYYVREAHGIV VTDVAFLPEKGRGPPELLGSHETALFSVAVDSRCQ LHLLPSRRSVPVWLLLLLCVGLIIVTILLQSAFFG FL
3632	A	942	40	PWCQRVEVRSCGSSKRSCSRWSGSSWDGSRSLG RGLNHTSLNRSPPFTPDTHCCSPCCQPTCCRT TCCRTTCWKPTTVTCSSTPCCQPSCCVPSCCQP CCHPTCCQNTCCRTTCCQPTCVASCCQPSCCSTP CCQPTCCGSSCCGQTSCGSSCCQPCGSSCCQPC HPTCYQTICFRTTCCQPTCCQPTCCRNSTCQPTCC GSSCCQPCCHPTCCQPTICRSTCCQPSCVTRCCSTP CCQPTCGGSSCCSQTCCNESSYCLPCCRPTCCQTT CYRTTCCRPSCCCSPCCVSSCCQPSCC
3633	A	605	3004	GPEGYRGRRARHPSLGGSTTGHC GGGRGAEGTGT DPAAPAARLNV DGLLVYFPYDYTYPEQFSYMRE LKRTLDAKGHVLEMPSGTGKTVSLLALIMAYQ RAYPLEVTCLIYCSRTVPEIEKVIEELRKLNFYB

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KQEGEKLPFLGLALSSRKNLCHPEVTPPLRFQKD VDGKCHSLTASYVRAQYQHDTSPLPHCRFYEEFD AHGREVPLPAGIYNLDDLKALGRRQGWCPYFLA RYSILHANVVVSYHYLLDPKIADLVSKELARK AVVVFDEAHNIDNVCIDSMSVNLTRRTLDRCQG NLETQKTVLRKETDEQRLRDEYRRLVEGLREA SAARETDAHLANPVLPEVLQEA VPGSIRTAEHF LGFLRRLLEYVKWRLRVQHVVQESPPAFLSGLA QRVCIQRKPLRFCAERLRSLLHTLEITDLADFSL TLLANFATLVSTYAKGFTIIEPFDDRTPTIANPIL HFSCMDASLAIKPVFERFQSVIITSGTSLPLDIYPK ILDFHPVTMATFTMTLARVCLCPMIIGRNDQVA ISSKFETREDIAVIRNYGNLLEMSAVVPDGIVAF FTSYQYMESTVASWYEQGILENIQRNKLFIETQ DGAETSVALEKYQEACENGRGAILLSVARGKVS EGIDFVHHYGRAVIMFGVPYVYTQSRILKARLEY LRDQFQIRENDFLTFDAMRHAAQCVGRAIRGKT DYGLMVFADKRFARGDKRGKLPRIQEHLTDA NLNLTVDEGVQVAKYFLRQMAQPFHREDQLGL SLLSLEQLESEETLKRIEQIAQQL
3634	A	159	384	LKMSSKTASTNNIAQARRTVQQLREASIERIKV SKASADLMSYCEEHARSDPLLIGIPTSENPFKDKK TCIL
3635	A	5	409	TELSQLEKAHPPADMGRKSKRPPPKKMTGT LETQFTCPFCNHEKSCDVKMDRARNTGVISCTV CLEEFQTPITCILGNLGGFQRVGRGLESGPCSSGP LCALVQGGQSRPEEQVPPSDFCGVRRRCRAGFQCQ
3636	A	48	282	DHLKSCYQDSHEDPTKMKRFLFLLTISLLVMVQ IQTGLSGQNDTSQTSSPSASSSMSGGIFLFFVANAI IHLFCFS
3637	A	1	1248	ARAGSVVGGAAAPGPPAGC...ARA*RI*SSPAA RRRCDWVEDGAG...EILMTVSKFASICTMGAA ASALEKEIGPEQFPVNEHYFGLVNFNTCYCNSV LQALYFCRPFREKGLAYKSQPRKKESLLTCLADL FHSIATQKKKVGVIPPKKFITRLKENELFDNYM QQDAHEFLNYLLNTIADILQEERKQEQNGRLPN GNIDNENNNSTPDPTWVHEIFQGTLTNETRCLTC ETISSKDEDFLDLSVDVEQNTSITHCLRGFSNTET LCSEYKYCYCECRSKQEAHKRMKVKKLPMILAL HLKRFKYMDQLHRYTKLSYRVVFPLELRLFNST GDATNPDRMYDLVA VVHCGSGPNRGHYIAIV KSHDFWLLFDDDDIVEKIDAQAIEEFYGLTSDISKN SESGYILFYQSRD
3638	A	11	630	PAGIPVSTISSDRRASTDLTRKMKPDETPMFDPNL LKEVDWSQNTATFSPAISPHPEGGLVLRPLCTA DLNRGFFKVLGQLTETGVVSPEQFMKSFEHMKK SGDYVTVVVEDVTLGQIVATLIEHKFIHSCAK RGRVEDVVVSDECRGKQLGNLLSTLTLLSKKL NCYKITLECLPQNVGFYKKFGYTVSEENYMCRR FLK
3639	A	2	1200	PRVRLRPSRSRSGRLLSTRAPGPSFRLHSSPL LPHAMKSPFYRCQNTTSVEKGNSAVMGGVLFST GLLGNLLALGLLARSGLGWCSRRLRPLPSVFY MLVCGLTVTDLGKCLSPVVLAAAYAQNRSRLRV LAPALDNSLCQAFAMSFGLSSTLQLLAMALE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				CWLSLGHPPFFYRRHITLRLGALVAPVVSFAFLAF CALPFMGFGKFVQYCPGTWCFIQMVHEEGSLSV LGYSVLYSSLMALLVLATVLCNLGAMRNLYAM HRRLQRHPRSCSTRDCAEPRADGREASQPLEELD HLLLLALMTVLFTMCSLPVIYRAYYGAFKDVKE KNRTSEEAEDLRALRFLSVISIVDPWIFIFRSPVFR IFFHKIFIRPLRYRSRCSNSTNMESSL
3640	A	930	182	PLPPTLAMFLTRSEYDRGVNTFSPEGRLFQVEY AIEAIKLGSTAIQITSEGVCLA VEKRTSPLMEPS SIEKIVEIDAHIGCAMSGLIADAKTLIDKARVETQ NHWFTYNETMTVESVTQAVSNLALQFGEEDADP GAMSRPFGVALLFGGVDEKGPQLFHMDPSGTFV QCDARAIGSASEGAQSSLOQVYHKSMTLKEAIKS SLILKQVMEEKLNATNIELATVQPGQNFHMFTE EELEEVIKDI
3641	A	2	1254	PTGQGGRRAEARSCLLSKAMLRSGYRALPLGD FDRFQSSFGFLGSQKGCLSPERGCVGTGADVPO SWPSCCLCHGLISFLGFLLLVTFPISGWFAKIVPT YERMIVFRLGRIRTPQGPGMVLLLPFIDSFQRVDL RTRAFNVPPCKLASKDGAVLSVGADVQFRIWDP VLSVMTVKDLNTATRMATAQNAMTKALLKRPLR EIQMEKLIKSDQLLEINDVTRAWGLEVDRELA VEAVLQPPQDSPAGPNLDSTLQQLALHFLGGSM NSMAGGAPSPGPADTVEMVSEVEPPAPQVGARS SPKQPLAEGLLTALQPFLEALVSQVGACYQFNV VLPSGTQSA YFLDLTTGRGRVGHGVDPDGIPDVV VEMAEADLRALLCRELRPLGAYMSGRLKVKGDL LAMAMKLEAVLRALK
3642	A	1	237	RRGEIDMATEGDVELELETETSGPERPPEKPRKH DSGAADLERVTDYAEKEIQSSNLETAMSVIGDR RSPFQKAKQER
3643	A	94	541	RLRRRRRRRMEAVVVFVSLLDCCALIFLS TLSDLECDYINARSCSKLNKWVPELIGHTIVTV L LMSLHWFI LLNLPVATWNIYRYIMVPSGNM GVFDPTEIHNRGQLKSHMKEAMIKLGFHLLCFF MYLSMILALIND
3644	A	95	2808	TSCRHFPITSEDPLNYLLLTVERIYAYQALPLGFL FCSRDPVPEYLNHCGVKYVLISDRASFCALHIFFS PFRNVFRPAAGGGIAPPPRLWFQPSLSDAEMEIPK LLPARGTLOQGGGGGGIPAGGGRVHRGPDSPAGQ VPTRRLLLPRGPQDGGPGRRREEASTASRGPGPS LFAPRPHQPSGGGGGGGDDFFLVLLDPVGGDVE TAGSGQAAGPVLREEAEEGPGLQGGESGANPAG PTALGPRCLSAVPTPAPISAPGPAFAAGTVTIHN QDLLLRFENGVLTLATPPPHAWEPGAAPAQQPG CLAPQAGFPHAAHPGDCELPDPLLAEPAPAP APAPEEAEGPAAALGPRGPLGSGPGVVLYLCPE ALCGQTFAKKHQLKMHLTHSSSQGQRPFKCPL GGCGWTFITTSYKLKRHLQSHDKLRPFGCPAEGC GKSFTTVYNLKAHMKGHEQENSFKCEVCEESFP TQAKLGAHQRSHFEPERPYQCAFSGCKKTFTITS ALFSHNRAHFREQLFSCSFGCSKQYDKACRLK IHLRSHTGERPFLCDFDGCWNFTSMKLLRHKR KHDDDRRFMCPEGCGKSFTRAHLKGHSITHL STKPFVCPVAGCCARFSARSSLYHSHKKHLQDQVD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TWKSRCPISSCNKLFTHSKHSMKTHMVKRHKVGO DLAQLEAANSLTPSELTSQRQNDLSDAEIVSLF SDVPDSTSAALLDTALVNSGILTIDVASVSTLAG HLPANNNNSVGQAVDPPSLMATSDPPQSLDLSL FGTAATGFQQSSLNMDEVSSVVGPLGSLDSL MKNSSPEPQALTPSSKLTVDITDITLTPSSTLCENSV SELLTPAKAEWSVHPNSDFFGQGETQGFNPAA GNHGSQKERNLITVTGSSFLV
3645	A	2194	1707	TVSFHKTMASLKCSTVVCICLEKPKYRCPACRV PYCSVVCFRKHKEQCNPETRPVEKKIRSAIPKT VKPVENKDDDDSIADFLNSDEEDRVSLQNLKN LGESATLRSLLLNPRLRLQMLVNLQGEDKAKLM RAYMQEPLFVEFADCCLGIVEPSQNEES
3646	A	85	1948	ERGGGKAAAAAAAAAARALAASGQDPRPHPR APPWDDSGDDDEATTPADKSELHHTLKNLSLKL DDLSTCNLDIAKHGAALQRSITELDGLKIPSESG EKLKVVNERATLFRITSNAMINACRDFLELAEIHS RKWQRALQYEQEQRVHLEETIEQLAKQHNSLER AFHSAPGRPANPSKSFIEGSLTPKGEDSEDEDT EYFDAMEDSTSFITVITEAKEDSRKABGSTGTSSA DWSSADNVLDGASLVPKGSSKVKRRVRIPNKP YSLNLWSIMKNCIGRELSRIPMPVNFNEPLSMLQ RLTEDLEYHLLDKAVHCTSSVEQMCLVAAFSV SSYSTTVHRIAKFPNPMLETFELDRLLDDMGLRS LCEQVSHPPSAAHYVFSKHGWSLWQEITISSKF RGKYISIMPLGAHLEFQASGNHYVWRKSTSTVH NIIVGKLWIDQSGDIEIVNHKTNDRCQLKFLPYSY FSKEAARKVTGVVSDSQGAHYVLGSGWDEQM ECSKVMHSSPSSSDGKQKTVYQTLAKLLWK KYPLPENAMMYFSELALTNHHEEGVAPTDS RLRFDQJAEKGRWDEANTVQJEEKRLSD RRRLEACGPGSSSSEE
3647	A	46	5007	PTGDACVSTSCELASALSHLDASHLTENLPKAAS ELGQQPMTELDSSSDLISSPGKKGAHPDPKTS VDTGQVSRPENPSQPASPRVTKCKARSPVRLPHE GSPSPGEKAAAPPDYSKTRSASETSTPHNTRRVA ALRGAGPGAEGMTPAGAVLPDPLTSQEQRQGA PGNHSKALEMTGIHAPESSQEPSLLEGADSVSSR APQASLSMLPSTDNTEACGHVSGHCCPGGSRE SPVTDIDSFIELDASAARSPSSQTGDSGSQEGSA QGHPPAGAGGGSSCRAEPVPGGQTSSPRRAWAA GAPAYPWASQPSVLDINPDKHFTVNKNFLSN YSRNFSSFHEDSTSLSGLDSTEPSLSSMYGDAE DSSSDPESLTEAPRASARDGWSPPRSRLVSLHKED PSESEEEQIEICSTRGCPNPPSSPAHLPTQAAICPAS AKVLSLKYSTPRESVASPREKVACLPGSYTSGPD SSQPSSLLEMSSQEHETHADISTSQNHRPSCAET TEVTSASSAMENSPLSKVARHFHSPPIILSSPNMV NGLEHDLDDDETNLQYETSINAAASLSSFSVDVP KNGESVLENLHISESQDLDDLQPKPMIARRPIM AWFKEINKHNQGTILRSKTEKEQPLMPARSPDS KIQMVSSSQKKGVTVPHSPPQPKTNLENKDLSSK SPAEMLLTNGQKAKCGPKLRSLKGAQVNSE APAANAVKAGGTDHRKPLISPQTSHKTLKAVS QRLHVADHEDPDRNTTAAPRSPQCVLESKPPLAT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SGPLKPSVSDTSIRTFVSPLTSPKPVPEQGMWSRF HMAVLSEPDRCPTTPKSPKCRABGRAPRADSG PVSPAASRNGMSVAGNRQSEPRASHVAADTAQ PRPTGEKGGNIMASDRLENTNQLKIVEISAEAVSE TVCGNKPAESDRRGGLAQGNCQEKSEIRLYRQ VAESSTSHPSLPSHASQAEQEMSRFSMAKLAS SSSSLQTAIRKAEYSQKSSLMDSRQVPRNSIPG GPSGEDHLYFTPRPATRTYSMPAQFSSHFGREGH PPHSLGRSRDSQVPVTSSVVPEAKASRGGLPSLA NGQGIYSVKPLLDTSRNLPTDEGDIISVQETSCL VTDKIKVTRRHICYEQNWPHSTSFSSVKQRIKS FENLANADRPVAKSGASPFSLVSSKPPIGRRSSGS IVSGSLGHPGDAAARLLRRSLSSCSENQSEAGTL LPQMAKSPSIMTLTISRQNPPTSSKGSSELKKS LGPLGIPTMTLASPVKRNKSSVRHTQPSPVRS KLQELRALSMPLDLKLCSEDYSAGPSAVLFKTEL EITPRRSPGPPAGGVSCPEKGGNRACPGGSGPKT SAAETPSSASDTGEAAQDLFRRSWSVNLDQLLV SAGDQQRQLQSVLSSVGSKSTILTIQEAKAQSENE EDVCFIVLNRKEGSGLGFSVAGGTDVEPKSITVH RVFSQGAASQEGTMNRGDFLLSVNGASLAGLAH GNVLKVLHQAQLHKDALVVIKKGMDQPRPSAR QEPPTANGKGLLSRKTIPLPGIGRSVAVHDALC VEVLKTSAGLGLSLDGGKSSVTGDGPLVIKRVY KGGAAEQAGIEAGDEILAINGKPLVGLMHFDA WNIMKSVPEGPVQLLRKHRNSS
3648	A	337	1564	KSRLSVTLMPVQLSEHPWNEHMSLRISVGGGLP VLASMTKAADPRFRPRWKVVLTFFVGAAILWLL CSHRPAPGRPPTHNAHNWRLGQAPANWYNDTY PLSPQRTAGIRYRIAVIADLDTESRAQEENTWF TYLTKGYLTGEGGDKVAVEWDKTHGVLEJHL AEKGRGNEUSDELFTNGKLYSVDDRTGVVYQIE GSKAVPWVLTSDGDGTVEKGFKAEWLAVKDER LYVGGLGKEWTTTGDVVNENPEWVKVVGYYK GSVDHENWVSNYNALRAAGIQPPGYLIHESAC WSDTLQRWFFLPRRASQERYSEKDDERKGANLL LSASPDFGDIAVSHVGAVVPTHGFSFKFIPNTDD QITVALKSEEDSGRVASYMAFTLDGRFLLPETKI GSVKYEGIEFI
3649	A	1	775	PTRPGSGSAGGARVGSGEFGVEMAALAPLPLPA QFKSIQHHLRTAQEHDKRDPVVAAYYCRLYAMQ TGMKIDSKTPECRKFLSKLMDQLEALKKQLGDN EAITQEI VGCAHLENYALKMFLYADNEDRAGR HKNMIKSFYASLLIDVITVFGEITDENVKHRKY ARWKATYIHNCLKNGETPQAGPVGIBEDNDIEEN EDAGAASLPTQPTQPSSTYDPSNMPSGNYTGI QIPPGAHA PANTPAEVPHSTGVAK
3650	A	20	963	KMAATLGPLGSWQQWRRCLSARDGSRRLLLL LLGSGQGPPQVQVAGQTFEYLKREHLSKPYQGE APRPCFLRDWELQVHFQKIHGQKKNLHGDGLAI WYTKDRMQPGPVFGNMDKFGVGLGVFDYTPNE EKQQERVFPYISAMVNNGSLSYDHERDGRPTL GGCTAIVRNHLYDTFLVIRYVKRHLTIMMDIDGK HEWRDCIEVPGVRLPRGYFGTSSITGDLSDNHD VISLKL FELTVERTPEEEKLHRDVFLPSVDNMKL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				PEMTAPLPPLSGLALFLIVFFSLVFSVFAIVIGILY NKWQEQRKRFY
3651	A	1	1218	RSWAYVKKCKNNMCPNRGLHDGPEPCWLHHA AGTVSAVQARGLQPSQSRSRPRVPLATALAYG PAHTPPLSRIGWAMQPPPGPLGDCRLDWEDLQ QDFQNIQVSAAADAGSPPSRVSLAQGGSGSGPGC KPSLPAAEAEGAAQELNQMKERQGLFFDMEAYL PKKNGLYLSLVLGNNVTLLSKQAKFAYKDEYE KFKLYLTILLISFTCRFLNLSRVTDAAFNFLVW YYCTLTIRESILINNGSRIKGWVVFHHYVSTLSG VMLTWPDLGMYQKFRNQFLSFSMYQSFVQFLQ YYYQSGCLYRLRALGERHTMDLTVEGFQSWMW RVLTFLLPFLFFGHFWQLFNALTLFNLAQDPQCK EWQVLMCGFPFLLFLGNFTTLRVVHHKFHSQ RHGSKKD
3652	A	640	164	VTTSCIPFAFGLGVRASERLAEIDMPYLLKYQPM MQTIGQKYCMDPAVIAGVLSRSPGDKILVNMG DRTSMVQDPGSQAPTSWISQSQVFQTTTEVLTTRI TELQRRFPTWTPDQYLRGGLCAYSGGAGYVRSS QDLSCDFCNDVLARAKYLKRHGF
3653	A	2	909	IVRRDWQEVSDIHLAMANCKMTKSIRFPALHC YTGGEVVLPKDQEEWKRRRTGLLLYENYQSETG LICATYWGMKIKPGFMGKATPPYDVQFHMEASV ENCIIVSMNTADPGSQGITHSLLLQVIDDKGSILPP NTEGNIGIRIKPVRPVSLFMCYEGDPEKTAKVEC GDFYNTGDRGKMDEEGYICFLGRSDDIINASGYR IGPAEVESALVEHPAVAESAVVGSPDPIRGEVVK AFIVLTPQFLSHDKDQLTKELQQHVKSVTAPYKY PRKVEFVSELPKTITGKIERKELRKKETGQM
3654	A	2	909	IVRRDWQEVSDIHLAMANCKMTKSIRFPALHC YTGGEVVLPKDQEEWKRRRTGLLLYENYQSETG LICATYWGMKIKPGFMGKATPPYDVQFHMEASV ENCIIVSMNTADPGSQGITHSLLLQVIDDKGSILPP NTEGNIGIRIKPVRPVSLFMCYEGDPEKTAKVEC GDFYNTGDRGKMDEEGYICFLGRSDDIINASGYR IGPAEVESALVEHPAVAESAVVGSPDPIRGEVVK AFIVLTPQFLSHDKDQLTKELQQHVKSVTAPYKY PRKVEFVSELPKTITGKIERKELRKKETGQM
3655	A	2	2364	SPGPSLPESAESLDGSQEDKPRGSCAEPFTDTG MVAHINNSRLKAKGVGQHDNAQNFGNQSFEE RAACLRKGELFEDPLFPAEPSSLGFKDLGPNSKN VQNISWQRPKDIIINPLFIMDGISPTDICQGILGDC WLLAAIGSLTTCPKLLYRVVPRGQSFKKNYAGIF HFQIWQFGQWVNVVDDRLPTKNDKLVFVHST ERSEFWASALLEKAYAKLSGSYEALSGGSTMEGL EDFTGGVAQSFQLQRPQNLLRLLRKAVERSSL MGCSIEVTSDESLESMTDKMLVRGHAYSVTGLQ DVHYRGKMETLIRVRNPWGRIEWNGAWSDSAR EWEEVASDIQMQLLHKTEDGEFWMYSYQDFLNN FTILLEICNLTPDTLSGDYKSYWHTTFYEGSWRTG SSAGGCRNHPGTFTWNPQFKISLPEGDDPEDDAE GNVVVCTCLVALMQKNWRHARQQAQLOQTIGF VLYAVPKEFQNIQDVHLKKEFFTKYQDHGFSEIF TNSREVSSQLRLPPGEYIIPSTFEPHRDADFLLRV FTEKHSSESWEDEVNYAEQLQEEKVSEDDMDQ

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				DFLHLFKIVAGEGKEIGVYELQRLNRMIAIKFKS FKTKGFGGLDACRCMINLMDKDGSGKLGLEFKI LWKKLKKWMDIFRECDQDHSGTLNSYEMRLVIE KAGIKLNNKVMQVLVARYADDDLIIDFDSFISCF LRLKTMFTFFLTMDPKNTGHICLSLEQVLGEGW EGICRIAPACPSTPPPPSSDVPGPASCPRLFPWDL LPVSTVAADDHVGIEAL
3656	A	3	174	PLCTHYLLPELPEKSSRTSPRSRPGNMLSGDPHLP QPLCHCLDHCPCCFSGKRLVA
3657	A	1	444	DTRSTYHNAHSLPTYVKSPACQMTYIKSPACQ TQTCYVQGASPCQSYVQAPASGSTSQCVDTP CSAPCSTSYCCLAPRTFGVSPLRRWIQRPNCNT GSSGCCENSGSSGCCSGGCGCGSCGCGSSGCCCL GIIPMKSRSPALL
3658	A	92	1537	SEAPVQPQPYTMTSFYSTSSCPLGCTMAPGARNV FVSPIDVGCQPVAEANAASMCLLANVAHANRVR VGSTPLGRPSLCLPPTSHTACPLPGTCHIPGNIGIC GAYGKNTLNGHEKETMKFLNDRLANYLEKVRQ LEQENAELETTLLERSKCHESTVCPDYQSYFRTIE ELQQKILCSKAENARLIVQIDNAKLAADDFRIKL ESERSLHQLVEADKCGTQKLLDDATLAKADLEA QQESLKEEQSLKSNHEQEVKILRSQLEKFRIEL DIEPTIDLNRVLGEMRAQYEAMVETNHQDVEQ WFQAQSEGISLQAMSCSEELQCCQSEILELRCTV NALEVERQAQHTLKDCLQNSLCEADRYGTELA QMQLISNLEEQLSEIRADLERQNEQYQVLLDVK ARLENEIATYRNLTPLQSLFHACLLYFLSKLWPC HRWVSLWPWSQHGEMILKARVRRRLRLVALGSG VPSPCPVFLQD
3659	A	2	402	DLLQCLNQLYSASTEMSCQSQCCQPPPKCTP KCFPKC.IK.LTTCPPKCTPQYSA.CPPKCTSCG SSSGGCCSSEGGGCCCHHRPRQSLRRRPQSSSC CGSGSGQQSGGSSCCHSSGGSGCCHSSGGGCC
3660	A	26	710	CSAVEVKMAARTAFGAVCRRLWQGLGNFSVNT SKGNTAKNGGLLLSTNMKWVQFSNLHVDVPKD LTKPVVTISDEPDILYKRLSVLVKGHDKAVLDSY EYFAVLAAKELGISIKVHEPPRKIERFTLLQSVHI YKKHRVQYEMRTLYRCLELEHLTGSTADVLEY IQRNLPQGVAMEVTKFCFFIFLDTIRTVTRTHQGA NLGNTIRRKRRKQVIKQGGHFCLNLK
3661	A	2	370	DVSVAASEPTVYRNPTKMSCQQNQCCQPPPKC PIPKYPPKCPKSCASSCPPPISSCCGSSSGGCCSSG GCGCCSSEGGGCCLSHHRHHRSHCHRPKSSNCY GSGSGQQSGGSGGCCSGGGCC
3662	A	205	1277	RKSLPHNPQKMLKKPLSAVTWLCIFIVAFVSHP AWLQKLSKHKTPAQPLKAANCCEEVKELKAQ VANLSSLLSELNKKQERDWSVVMQVMELESN SKRMESRLTDAESKYSEMNNQIDIMQLQAAQTV TQTSAGKETSPLRERGVPPLQHCFYIPDDFLGS PELEVFCDMETSGGGWTTIQRKSGLSVSYRDW KQYKQGFSGIRGDFWLGNENHRLSRQPTRLRVE MEDWEGNLRVAYSHFVLGNELNSYRLFLNGY TGNVGNDAQYHNNTAFSTKDKDNDNCLDKCA QLRKGGYWYNCCTDSNLNGVYYRLGEHNKHL GITWYGWHGSTYSLKRVEMKIRPEDFKP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
3663	A	64	1456	LSSAKETLAQMYNTVWNMEDLDLEYAKTDINC GTDLMFYIEMDPPALPPKPPKPTTVANNGMNNN MSLQDAEWYWGDISREEVNEKLRDTADGTFVLV RDASTKMHGDYTLTLRKGGNNKLIKIFHRDGKY GFSDPLTFSSVVELINHYRNESLAQYNPKLDVKL LYPVSKYQQDQVVKEDNIEAVGKKLHEYNTQFQ EKSREYDRLYEBYTRTSQEIQMKRTAIEAFNETIK IFEEQCQTQERYSKYEIEFKREGNEKEIQRMHN YDKLKSRISEIIDSRRRLBEDLKKQAAEYREIDKR MNSIKPDLIQLRKTRDQYLMWLTQKGVRRQKKL NEWLGNENTEDQYSLVEDDEDLPHHDEKTNWV GSSNRNKAENLLRGKRDGTFVRESSKQGCYAC SVVVDGEVKHCVINKTATGYGFAEPYNLYSSLK ELVLHYQHTSLVQHNDLSLNVTLAYPVYAQQRR
3664	A	944	406	GATVEDQSCNFGSLRWVVSVPHISARSCPDPLLS RTGRVPGGRGAGLPRHHSRCCLOVFFNGANVR QVDVPTLTGAFGILAAHVPTLQVLRPGLVVVHA EDGTTSKYFVSSGSIAVNADSSVQLLAEAAVTL MLDLGAAKANLEKAQAEVGTADEATRAEIQIR IEANEALVKALE
3665	A	98	1388	ASQLAFGGKLTSTPSRDFQCGCRGAVTCCSFHEH RHQSGRCLSTGMAPNLKGRPRKKKPCPQRDSF SGVKDSNNNSDGKAVAKVKCEARSALTKPKNN HNCKKVSNEEKPKVAIGEECRADEQAFLVALYK YMKERKTPIERIPYLGFKQINLWTFQAAQKL GYETITARRQWKHIYDELGGNPGSTSAATCTRR HYERLILPYERFIKGEEDKPLPIKPRKQENSSQE NENKTKVSGTKRIKHIEPKSKKEKENAPKPDAA EVSSEQEKEQETLISQKSIPEPLPAADMKKKIEGY QEFSAPLASRVDPEKDNETDQGSNSEKVAEEA GEKGF... ALVDSKQESKLCCFTES... HRWQTRMRRRMTNCPPW...ILPTAP
3666	A	113	1492	LLQEMCTKTIPVLWGCFLWNLTVSSSQITYPGI KARITQRALDYGVAQGMKMEQMLKEKKLPDL SGSESLEFLKVDYVYNFNSNIKISAFSPNTSLAF VPGVGKALTNHGTANISTDWGFESPLFVLVNSF AEPMEKPILKNLNEMLCPIIASEVKALNANLSTLE VLTKIDNYTLLDYSLISSPEITENYLDNLKGIFY PLENLTDPPFSPVPFVLPERSNSMLYIGIAEYFFKS ASFAHFTAGVFNVTLSLEEISNHFVQNSQGLGNV LSRIAETIYLSQPFMVRIMATEPPIINLQPGNFTLDI PASIMMLTQPKNSTVETIVSMDFVASTSVGLVIL GQRLVCSLSLNRFLALPESNRSNIEVLRFENILSS ILHFGVLPANAKLQOGFPLPNPHKFLFVNSDIEV LEGFLLISTDLKYETSSKQQPSFHVWEGNLISRQ WRGKSAP
3667	A	1	181	FRRLGSGRNGGGSMNAPPAFESFLLFEGEKITIN KDTKVPNACFLTINKEDHTLGNLIK
3668	A	212	431	VAGEAVPFFPMYSEPLKPSYLALVLWYFLLTG YCITKPEVIFKIEQGEWPWLEKGFPSQCHPAKYL WCLHD
3669	A	458	1056	FSGVCFAGIAGSMATLLHDAVMNPAEVVKQRLQ MYNSQHRSAISCIRTVWRTEGLGAFYRSYTTQLT MNIPFQSIHFITYEFLQEQVNPHTYNPQSHISGG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LAGALAAAATTPLDVCKTLLNTQENVALSLANIS GRLSGMANAFRTVYQLNGLAGYFKGIQARVIYQ MPSTAISWSVYEFFKYFLTKRQLENRAPY
3670	A	145	298	RNPCPLTFLPSTLMVLLLSLTFFSALTFHSICQLRN TGVEVDIVFQRVSF
3671	A	3	462	ILKVAKKERTMSSLPVPYKLPVSVLSVSGSCVIKGT PIHSFINDPQLQVDFYTDMDSDIAFRFRVHFG NHVVMNRREFGIWMLBETTDYVPFEDGKQFELC IYVHYNEYEIKVNGHHLRALSHRIPPSFVEDGC KCPRRYLPWTSVCVCN
3672	A	1	1028	HYAKLGTRPRLKFMSSPSLSDLGKREPAAAAD RGTQQRACANATWNSIHNGVIAVFQRKGLPDQ ELFSLNEGVRQLLKTGSGFTYELQNQLLTGKM VILRDKIRFYEGQKLLDSLAEWDFFFSDVLPML QAIFYPVQGKEPSVRQLALLHFRNAITLSVKLED ALARAHARVPPAIVQMLLVLGQVHESRGVTEDY LRLTLVQKVVSPLYLGTGLHSSEGPFTHSCELEK RLLRRSRSGDVLAKNPVVRSKSYNTPLNPNVQE HEAEGAAAGGTSIRRHVSSEMTSCPEPQGFSDPP GQGPTGTFRSSPAPHSGPCPSRLYPTTQPPEQGLD PTRS
3673	A	2	712	RPPRVWYPELRELSAAAPRWSHRTAPGIMVFYF TSSSVNSSAYTIYMGKDKYENEDLIKHWGPEDI WFHVDKLSSAHVYLRHLKGENIEDIPKEVLMDC AHLVKANSIQGCKMNNVNVVYTPWSNLKKTAD MDVGQIGFHRQKDVKIVTVEKKVNEILNRLEKT KVERFPDLAAEKECRDREERNEKKAQIQEMKKR EKEEMKKKREMDLRSYSSLMKVENMSSNQDG NDSDEFM
3674	A	2	712	RPPRVWYPELRELSAAAPRWSHRTAPGIMVFYF TSSSVNSSAYTIYMGKDKYENEDLIKHWGPEDI WFHVDKLSSAHVYLRHLKGENIEDIPKEVLMDC AHLVKANSIQGCKMNNVNVVYTPWSNLKKTAD MDVGQIGFHRQKDVKIVTVEKKVNEILNRLEKT KVERFPDLAAEKECRDREERNEKKAQIQEMKKR EKEEMKKKREMDLRSYSSLMKVENMSSNQDG NDSDEFM
3675	A	921	1321	VTLAKMRVHISSCLKVQEQMANCPKFVPVPTS QPIPSNIPNRSTFACPYCGARNLDQQLVKHCVE SHRSDPNRVVCPICAMPWGDPSYKSANFLQHL LHRHKFSYDTFVDYSIDEEAAAFQAALALSSEN
3676	A	3	1856	TLGRWLLGVYETVAPTACLPRPRLRRRRRRRR RRMISRYTRKAVPQSLELKGITKHALNHPPPEK LEEISPTSDSHEKDTSSQSKSDITRESSFTSADTGN SLSAFPSYTGAGISTEGSSDFSWGYGELDQNATE KVQTMFTAIDELLYEQKLSVHTKSLQEECQQWT ASFPHLRILGRQITPSEGYRLYPRSPSAVSASYET TLSQERDSTIFGIRGKKLHFSSSYAHKASSIAKSSS FCSMERDEEDSIIVSEGIIEEYLAFDHIDIEGFHG KKSEAAATEKQKLGYPPIAPFYCMKEDVLAYVFD SVWCKVVSCEQLTRSHWEGFASDDESNAV RPDSESSCVLSELHPLVLPVPSKVLVYITSPMS LCQASRHQPNVNDLLVHGMPQLPRNLSLMDKLL DLDDKLLMRPGSSTILSTRNWPNAVFEFTSSLS YTVQSTRRRNPPRTLHPISTSHSCAETPRSVEEIL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RGARVPVAPDSLSSPSPTPLSRNNLLPPIGTAEVE HVSTVGPQRQMKPHGDSSRAQSAVVDEPNYQQ PQERLLLPDFFPRPNTTQSFLDQTQYRRSCAVEYP HQARPGRGSAGPQLHGSTKSQSGGRPVSRTRQGP
3677	A	246	757	MRLQGAIFVLLPHLGPIVWLFTRDHMSGWCEG PRMLSWCPFYKVLVLLVQTATYSVVGYSYLWVK DLGGGLGWPLALPLGLYAVQLTISWTVLVLFPT VHNPGALALLHLLL YGLVVSTALIWHPINKLAAL LLPYLAWLTVTSALTYHLWRDSLCPVHQPPTEKSD
3678	A	20	1508	RGKAEFFLAMAGTNALLMLENFIDGKFLPCSSYI DSYDPSTGEVYCRVPNSGKDEIEAAVKAAREAFP SWSSRSPOERSRVLNQVADLLEQSLEEFQAESK DQGKTLALARTMDIPRSVQNRFFASSSLHHTSE CTQMDHLGCMHYTVRAPVGVAGLISPWNPLFY LLTWKIAPAMAAGNTVIAKPSELTSVTAWMLCK LLDKAGVPPGVVNVFVGTGPRVGEALVSHPEVPL ISFTGSQPTAERITQLSAPHCKKLSLELGGKNPAI FEDANLDECIPATVRSSFANQGEICLTSRIFVQK SIYSEFLKRFVEATRKWKVGIPSDPLVSGALISK AHLEKVRSYVKRALAEGAQIWCGEVDKLSLPA RNQAGYFMLPTVITDIKDESCMTEEIFGPVTCV VPFDSEEEVIERANNVKYGLAATVWSSNVGRVH RVAKKLQSGLVWTCWLIRELNLPGGMKSSGI GREGAKDSYDFFTEIKTITVKH
3679	A	1862	502	MAGTKPYMEIQTITREYYEHL YANKLENLEEMD KFLDTYTLPRNLNQEVEESLNRPIGSEIEAINSLP TKKIPGPDRTAKFYQRYKEELSNIHYLGLSHH LLALNFIIVSFGKKSAWSSAQVKVTDITDFDGEV RVFEGDNTTTPPKRQVYTHCGGWALASAKIT YDELSTAMAEELNAVIVSIEYRLVPKVYFPEQIH DVVRAITNYFLKPEVLQKYMVDPGRICISGDSAG GNLAAALCCQFTQDASLKNKLKLQALITYPVLQA LDFNTPSYQQNVNTPIPRYVMVKYVVDYFKG NYDFVQAMIVNNHTSLDVEEAAAVRARLNWTS LLPASFTKNYKPVVQTTGNARIVQELPQLLDARS APLIADQAVLQLLPKTYILTCEHDVLRDDGIMYA KRLESAGVEVTLDFEDGFHGCMTFSWPTNFSV GIRTRNSYIKWLDQNL
3680	A	249	2146	RSWGAPWFWRMRLRRRHMPRLAMVGCFAV LFLFLLHRDVSSREEATEKPWLKSLVSRKDHVLD LMLEAMNNLRDSMPKLQIRAPEAQQTLSINQSC LPGFYTPAELKPFWERPPQDPNAPGADGKAFQK SKWTPLETQEKEEGYKKHCFNAFASDRISLQSL GPDTRPPECVDQKFRRCPLATTSVIIVFHNEAWS TLLRTVYSVLHTTPAILLKEILLVDDASTEHLKE KLEQYVKQLQVVRVVRQEERKGLITARLLGASV AQAEVLTFLDAHCECFHGWLEPLLARIAEDKT VVSPIVITIDLNTFEFAKPVQGRVHSGNFDWS LTFGWETLPPHEKQRRKDETYPKSPITFAGGLFSI SKSYFEHIGTYDNQMEIWGGENVEMSFVRVWQC GGQLEIPCSVVGHVFRTPKSPHTFPKGTSVIARNQ VRLAEVWMDSYKKIFYRRNLQAAKMAQEKSGF DISERLQLREQLHCHNFSWYLHNVPPEMFVPDL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				TPTFYGAIKNLGTNQCLDVGENNRGGKPLIMYS CHGLGGNQYFEYTTQRDRLRHNIKQLCLHVS KGLGLGSCHFTGKNSQVPKDEEWELAQDQLRNS GSGTCLTSQDKKPAMAPCNPSDPHQLWLFV
3681	A	2982	1869	LKDTLKSQMTQEASDEAEDMKEAMNRMIDELN KQVSELSQLYKEAQAELEDYRKRKSLDVDTA EYIHKAHEKLMQLTNVSRKAEDALSEMKSQY SKVLNELTQLKQLVDAQKENSVSITEHLQVIT TLRTAAKEMEELKISNLKEHLASKEVEVAKLE KQLLEEKAAMTDAMVPRSSYEKLQSSLESEV SVLASKLKESVKEKEKVHSEVVQIRSEVSQV KREKENIQTLTKSKEQEVNELLQKFQQAQEE LAEMKRYSESSSKLEEDKDKKINEMSKEVTK LKEALNSLSQLSYSTS SSKRQSQQLALQQV KQLQNQLAECKKQHQE VISVYRMHLLYAVQ GQMDVDVQKVLKQILTMC KNQSQKK
3682	A	447	1024	AQALTAGRQLALAAPFIAPISPLPRLNPPS QSWNSTPFFKVKLPPQKEVITSDELMAHLG NCLLSIKPQEKSEGLQLNFQQNVDDAMTVLP KLATGLDVNVRFTGVSDFEYTPECSVFDLL GIPLYHGWLVDPQQSPEAVRAVGKLSYNQL /VGEDHHLQTLQ*HQP RDRKPCRAVPGD HRGPSDLPRTV
3683	A	2	942	LEIKQEEKFVGQCIKEELMHGECVKEEKDF LKKEIVDDTKVKEEPPINHPVGCKRKLAM SRCETCGTEBAKYRCPRCMRYSCSLPCVKK HKAELTCNGVRDKTAYISIQQFTEMNLLSD YRFLEDVARTADHISRDAFLKRPISNKYMY FMKNRARRQGINKLLPNGFTKRKENSTFFD KKKQQFCWHVKLQFPQSQAIST*KKRVPD DKTINEILKPYIDPEKSDPVIRQLKAYIRS QQTGVQILMKIEYMQQNLVRYEYELDPYKSI !DNLRIJEVETYPILHVVKGEENNDM... VHLQVSTKNVGNEN
3684	A	119	153	SLQENVQEKVRVCPGLGGLLPNGTPSITAAA AAPQVLWRHVQPGCSHHLHACVIRAAACRA GEGHADRHAGPPET/PVTLPSWPWSSPWER QCPMHL*APGHAFRPVPTHEHRRGWAALGH HRAAGPLREPASGSQPAPASC*PECHHGCP EQTRQCQDLLREAVVAPEQRG*PCAHLQT* ATATTLCPQVPAGRVWQPGHSCHELLPHR HDGSH*HHCAAHRRPVTRRQAAHGVPLPD ACYSPIHTLPAAPPPATRPAGHTATHPE* GGDLTPVPDGPDPDPRDVGQIPGAGGGSQ LAPCCPPFFAAPVSVQGTQGLGPKNVLH*Q WEGIRWQKEPE/PGPPPEVELKRGAKCRIGD HGLGAVLGQGEYAS*SPSIPW*ASSACPLH PTP/TVYTQSPAAA PGWTRPPSP/PPPGL YPGP/PASHAPGVRGGISHQLYSLP*LCRE CCSCP/PPPAHGGRCPSLLPPEALAK LLL
3685	A	101	438	AWVLQCKINTELQTEVVMLKSMVLWLGEQV QS LQLQQQLHCHFNHTHICVTNLEYN*KE YPWDLVKAHLQGA STNITFDIGELQKKIL DLNKQTQEFQPSL*AWTEFQGGLE
3686	A	105	845	VSDVVKNNQLVEVQCRQDGCDAVENVHQM FMFNWFTDCLWTLFLSNYQPSVESSSPGGS ATSDDHEFDPSADMLVHDFDDERTLEEEEM MEGETNFSSEIEDLAREGDMPIHELLSLYGY GSTVRLPEEDEEBE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EEEEGEDEDEDADNDDNSGCSGENKEENIKDSS GQEDETQSSNDDPSQSVASQDAQEIRPRCKYF DTNSEVEEESEDEDEDYIP/SIISFTQSSDGI*SSSSSE DWKKEIMVGS
3687	A	49	1225	PVLVTSLRMREADTLRPPQLMEVSADIISTVEFN HTGELLATGDKGGRVVIFQREPESKNAPHSQGE YDVYSTFQSHEPEFDYLSLEIEEKINKIKWLPQQ NAAHSLSTNDKTIKLWKITERDKRPEGYNLKDE EGKLKDLSTVTSLQVPVLKPMDLMEVSPRRIFA NGHTYHINSISVNSDCETYMSADDLRLNLWHLAI TDRSFTP\NIVDIKPANMEDL TEVITASEFHPHHC NLFVYSSSKGSLRLCDMRAAALCDKHSKLFEEPE DPSNRSFFSEIISVSDVKFSHSDRYMLTRIDYLT VKVWDLNMEARPIETYQVHDYLRSLKCSLYEND CIFDKFECAWNGSDR/IIMTGA YNNFFRMFDRNT KRDTVLEASRGSSKPRAVL
3688	A	1	401	KKVPGRLSEMSFSLNFTLPANTTSSPVTDCGPSL GLAAGIPLL VATALLVALLFTLIHRRRSSIEAMEE SDRPECISEIDNPKISENPRRSPTHEKNTMGAQE AHYVKT VAGSEEPVHDIRPTIEMERRR
3689	A	698	889	GRVLVHCAMGVSRSATLVLAFLMIYENMTLVEA IPDGAGPPQISALTQAFVRQLQVLDNRLGRE
3690	A	61	153	MG AHLVRRYLGDASVEPDPLQMPTFPDYGF
3691	A	61	153	MG AHLVRRYLGDASVEPDPLQMPTFPDYGF
3692	A	3	2831	PLVRRLLRQTLRRVGGARAVREAVMRAVL TWR DKAEHCINDIAFKPDGTQLILAAGSRLLVYDTS GTLLQPLKGHKDTVYCVAYAKDGKRFASGSAD KSVIIWTSKLEGILKYTHNDAIQCVSYNPITHQLA SCSSSDFLWSPEQKS VSKHKSSSKICCSWTNDG QYLALGMFNGHSIRNKNKEEKVKIERPGGS' SPI WICWNPSSESTVARE... TVNR IQ EESITLKS AVYSSQGS EAE EEE EEDDS PRDDNL EERN DILAVADWGQKVSFYQLSKQIGKDRAL NFDPCISYFTKGEYILLGGSDKQVSLFTKDGVR LGTVGEQNSWVWTGQAKPDSNYVVGCCQDGTI SFYQLIFSTVHGLYKDRYAYRDSMTDVIVQHLIT EQKVRICKELVKKIATYRNRLAIQLPEKILIYEL SEDLSDMHYRVKEKIKKFECNLLVVCANHILC QEKRLQCLSFSGVKEREWQMESLIRYK VIGGPP GREGLLVGLKNGQILKIFVDNLFAIVLLKQATAV RCLDMSASRKKLAVVDENDTCLVYDIDTKELLF QEPNANSVAWNTQCEDMLCFSGGGYLNIAKSTF PVHRQKLQGFVVGYNKSKIFCLHVFSISAVEVPQ SAPMYQYLDRLKFKEAYQIACLGVIDTDWRELA MEALEGLDFETAKKBRKKRGETNNDLFLADVFS YQGKFHEAAKLYKRSGHENLALEMYTDL CMFE YAKDFLGSGDPKETKMLITKQADWARNIKEPKA AVEMYISAGEHVKAIEICGDHGWVDM LIDIARK LDKAEREPLLLCATYLLKKLDSPGYAAETYLKMG DLKSLVQLHVETQRWDEAFALGEKHPEFKDDIY MPYAQWLAENDRFEEAQAFHKAGRQREAVQV LEQLTNNAVAESRFNDAA YYYWMLSMQCLDIA QDPAQKD
3693	A	3	1099	SSFPTCMRTVFHSNTSVSSLLHRPGHVTPQLTIHG GWRHHRDHTAIDEWDFNPSKFLIYTCLLLFSVLL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				PLRLDGIHQWSYWAVFAPLWLWKLVLVAGASVG AGVWARNPRYRTEGEACVEFKAMLIAGVGHLL LMFEVLVCDRVERGTHFWLLVFMPLFFVSPVSV AACVWGRHDSLELEILCSVNILQFIFALKLDRI IHWPWL VVFVPLWILMSFLCLVVLYYIVWSLLFL RSLDVVAEQRRTHVTMAISWITIVVPLLTFEVL VHRLDGHNTFSYVSIFVPLWLSLLTLMATTFRRK GGNHWWFAIRRD/CQDQLPQPTGKPPPPPLTDH HGEKALPLQNKDRGSWPASRGSPRL
3694	A	483	761	PRSLIDYKSYM DTKLLVARFLEQSSCTMTDPDIHE LVENIKSVLKSDEEHMEEAITSASFLEQIMAH SX QHIAHKLXPETAGLXTSELRLXITP
3695	A	483	761	PRSLIDYKSYM DTKLLVARFLEQSSCTMTDPDIHE LVENIKSVLKSDEEHMEEAITSASFLEQIMAH SX QHIAHKLXPETAGLXTSELRLXITP
3696	A	456	733	LSAALWEEPILSLWSETKELNTRGKMNYPIQIPH RPHVGLRVRPGPGTILSNAPKSLCPGMSNSDRGI HGGEGQGPGRAGHLGRGGGMSFL
3697	A	877	1873	VWL*TLS*HTCALMTVCRSCLVKYLEENNTCPT CRIVHQSHPLQYIGHDRMTQDIVYKLVGLQEA EMRKQREFYHKLGMVPGDIKGETCSAKQHLD S HRNGETKADDSSNKEAAE
3698	A	1	572	KQCGIPHEVVRDENSSVYAEVSRLLLATGHWKR LRRDNPRFNLMLGERNRLPFGRLGHEPGLVQLV NYYRGADKLCRKASLVKLIKTSPELAESCTWFPE SYVIYPTNLKTPVAPAQNGIQPPISNSRTDEREFL ASYNRKKEGEGNVWIAKSSAGAKVWVQW*M TDLEEDIPSPVGLGLESEWPL
3699	A	2008	2432	LHCKMGALETQTHPCSQNMLRSLQKCCCKVEE HHLQPVQVLQTLHSATAGTCRRPARPPAPPT MTPWESKQSCNDESERASARGGRYGLCA GGRALGGSFPPPLPGETLFSGCKHRRRRR AAPGEEAGT
3700	A	32	1318	GYQIGMALASGPARRALAGSGQLGLGGFGAPRN GAYEWGVRSTRKSEPPPLDRVYEIPGLEPITFAG KMHFVPWLARPIFPWDRGYKDPFRFYSPLHE HPLYKDQACYIFHHRCLLEGVKQALWLTKT KL IEGLPEKVLSLVDDPRNHENQDECVLNVISHARL WQTTEEIPKRETYCPVIVDNLQLCKSQILKHPSL ARRICVQNSTFSATWNRESLLLQVRGSGGARLST KDPLPTIASREEIEATKNHVLETFYPISPIDLHECN IYDVKNDTGFQEGYPYPHTLYLLDKANLRPH RLQPDQLRAKMILFAFGSALAQARLLYGND AKV LEQPVVVQSVGTDGRVFHFLVFQLNTDLD SNE GVKNLAWVDSDDLQLYQHFCLPVIKKRVVVEP VGPVGFKPETFRKFLALYLHGAA
3701	A	86	465	WTLCPGEAGMVGYDPKPDGRNNTKFQVAVAGS VSLVTRALISPFVVIKIRFQLQHERLSRSDPSAK YHGILQASRQILQEEGPTAFWKGHVPAQILSIGY GAVQFLSFEMLTTELVRHGSVVDARE
3702	A	166	814	GFWEKTNQSSHSMDPLGAPSQFVDVDTLPSWGD SCQDELNSSDTTAEIFQEDTVRSFFLYNKDVNGK VVLWKGDVALLNCTAIVNTSNESLTDKNPVSESI FMLAGPDLKEDLQKLKGCRTEAQLTKGFNLAA RFIHTVGPKYKSRYRTAAESSLYSCYRNVQLA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KEQSMSSVGFVCVNSAKRGYPLKDATHIALRTVR RFLEIHGETIEKV
3703	A	128	1255	SLGPSFKSATIPCCGDTMAPEEDAGGEALGGSFW EAGNYRRTVQRVEDGHRLCGDLVSCFQERARIE KAYAQQLADWARKWRGTVEKGPQYGILEKAW HAFFTAERLSALHLEVREKLQGGQDSERVRAWQ RGAFHRPVLGGFRESRAAEDGFRKAQKPWLKRL KEVEASKKSYHAARKDEKTAQTRESHAKADSA VSQEQLRKLQERVERCAKEAEKTKAQYEQTILAE LHRYTPRYMEDMEQAFETCQAAERQRLFFKD MLLTLHQHLDLSSSEKFHELHRDLHQGIEAASDE EDLRWWRSTHGPGMAMNWPQFEWSLDTQRTI SRKEKGGRSPDEVTLTSIVPTRDGTAPPPQSPGSP GTGQDEEWSDEESP
3704	A	1	271	ARGEDLALATGGGPDVTVTHSNMPCPNLSVYDC WLNKECSVGEHTFEDLGLCPGRNQREKKRSYK DFLREEEKIAAQVRNSSKKKLKDSE
3705	A	170	1318	LNWANLVIMWPREEEKEKVQDYSLGGLSPDLRI DVSRRKKILKAYDEDEDEDLYPDHPPPSLPLPG QFTCPQCRKSFTRRSFRPNLQLANMVQIRQMCP TPYRGNRSDQGMCFKHQEAALKLFCEVDKEAIC VVCRESRSHKQHSVLPLEEVVQEYKAKLQGHVE PLRKHLEAVQKMKAKEERRVTELKSQMKSELA AVASEFGRLTRFLAEEQAGLERRLREMHEAQLG RAGAAASRLAEQAQLSRLAEAQERSQQGGLR LLQDIKETFNRCCEVQLQPPEVWSPDPCQPHSHD FLTDAIVRKMSRMFCQAARVDLTLPDTAHPAL MLSPDRRGVRLAERRQEVADHPKRFSADCCVLG AQGFRSGRHYWEVCMGP
3706	A	204	1996	SRERQTTWMDHNFAPAPPEMQSHGAPGPGTSFS EHVLCGRPRCPGPGGSFLTPVLRKTHLDTFP QEHIPQTSSRLGLGARTSVPPQETGIALVLSLP LPTSSLVPRKLSSISLTLHQNSQARSLDRPLSHWE EIPTPGKKAAPHEGGRVSSPGSPVTLVPGGRVH SEGPGNPGLTKSNRMLATEKPLVSSYLALPFQSR LAQSAPVLAEPGSLGQGHVSVTDHMPTRASPG KGKPRARGIPRPRGRQLQRANTVNLAMDRTRD AARHLATMATNRPSLAINLATPNTSQLDTGTEFP ALDIKLTARDLSSVGTVKSGKTVNLATAGTIKP GTAMNLTIVGTTKPGMVMDLIASEPDKLGKAM ATRSTAKPDMTTEGIAMDSATSDPVKPDITITATV GTSRLETAMALARVNRAKLGTAKNLSALDTSR MGTA VGSVVPVTPDPATGKTTLGSVNNLTISDV ATCLLMPSRSTDLALDNTNAAMDRAPEASLDL ATEYKKGKCRNLVGDGLGCREGEVCELGDGSMK PMSINSNLLGYIGIDTIEQMRKKTMTKGFDFNIM VVGTEGCGAAAGLVAGSTKDPISFPQ
3707	A	3	549	SSSISRDFLGQAACASGTMRLWLRDFVLPTAACQ DAEQPMRYETLFQALDRNGDGVVDIGELQEGLR NLGIPLGQDAEEKIFTTGDVNKDGKLDFFEFMKY LKDHEKKMKLAFKSLDKNNDGKIEASEIVQSLQ TLGLTISEQQAELILQSIDVDGMTVDWNEWDRD YFLFNPVTDIEEIR
3708	A	1	1866	EFRGAGRANMLAPRGAAVLLLHLVLQRWLAAG AQATPQVFDLLPSSSQRLNPGALLPVLTDPALND

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LYVISTFKLQTKSSATIFGLYSSTDNSKYFEFTVM GRLSKAILRYLKNKGK VHLVVFNNLQLADGRRH RILLRLSNLQRGAGSLELYLDCIQVDSVHNLPR FAGPSQKPETIELRTFQRKPQDFLEELKL VVRGSL FQVASLQDCFLQQSEPLAATGTGDFNRQFLGQM TQLNQLLGEVKDLLRQEVNETSFLRNTITECQAC GPLKFQSPSTPVPPASPAPPTRPRRCDSNPCF RGVQCTDSRDGFQCGPCPEGYTGNITCIDVDEC KYHPCYPGEHCINLSPGFRCDCPVGFTGPMVQ GVGISFAKSNKQVCTDIDECRNGACVPNSICVNT LGSYRCGPCKPGYTGQIRGCKAERNCRNPELN PCSVNAQCIERQGDVTCVCGVGWAGDGYICGK DVDIDSYPDEELPCSARNCKKDNCKYVPNSGQE DADRDGIGDADDEDADGDGILNEQDNCVLHNV DQRNSDKDIFGDACDNCLSVLNNDQKDTGDG RGDADDDMDGDGDIKNILDNCPKFPNRDQRDK DGDGVGDACDSCPDVSNPNQ
3709	A	144	417	TQAMEGLLHYINPAHAISLLSALNEERLKGQLCD VLLIVGDQKFRAHKNVLAASSEYFQSLFTNKENE SQTVFQLDFCEPDAFDNVLYNY
3710	A	245	688	FGMLKNKGHSSKKDNLAVALQDHILHDLO LRNLSVADHSKTQVQKKENKSLKRDTKAIDTGL KKTTCQPKLEDSEKEYVLDPKPPPLTLAQKGLI GPPPPPLSSDEWEKVKQRSLQGDVQPCPICKE EFELRPQVFSIRG
3711	A	3	773	SLEMSSDGEPLSRMDESDSISSTIMDVSTISSGRS TPAMMNGQGSTTSSSKNIAYNCCWDQCQACFNS SPDLADHIRSIHVDGQRGGVFVCLWKGCKVYNT PSTSQSWLQRHMLTHSGDKPFKCVGGCNAFSA SQGGLARHVPTTHFSQNSKVSQPKAKESPSK AGMNTTRKLYKRRRSLALHDEFTLQTLAAS HRAICFNLSAEESLKGHSVVFHSTVSILLQIK YKTLQKNISTIISKSLKI
3712	A	2	344	RATWHNAGKEREAVQLMAGAEKRVKASHSFLK GLFGGNTRIEEACEMYTRAANMFKMAKNWSAA GNAFCQAAKLHMQLQSKHDSATSFVDAGNAYK KADPQGKTARHVACYLCV
3713	A	20	974	GAAATACSSSSSSGAPATWAAHGPBKDVASPS SVSLSPRRSRLVLRCGLRRNPERPSSPALRLL LLLLLLLLLLGFLSPGPERGVGGFRGRRAL LWAAALGHVVSQKVMRRAPGSRLSSGGGGGG TNYRSWWDWQPRTDASADPGNLKYSSSRDRG GSSSYGLQPSNSAVVSRQRHDDTRVHADIQND KGGYSVNGGSGENTYGRKSLGQELRVNNVTSPE FTSVQHGSRALATKDMRKSQERSMSYCDERLS YLLRRITRENDRRRLATVKQLKEFIQPENKLV LVKQLDILAAVHVDLNER
3714	A	237	458	IFALKSPSYLLPCCTPEGKMDHKQLCWSHPQKSG QSSRSCCICSNQHGLIWKYSNLMLQCCHQYVK DIGFIKL
3715	A	970	1524	LCTLSPGISGTAGSCLTTEPGTELGTSAQNGFYH EAVVLFQALKLNPQDHRLEGNRSFCHERLGQP AWALADAQVALTLRPGWPRGLFRLGKALMGLQ RFREAAAVFQETLRGGSQPDAAARELRSCLLHLTL QGQRGGICAPPLSPGALQPLPHAEAPSGPLSLRC

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				PRSTALRSPGLSPLLH
3716	A	85	308	QGLPSTMVKLGCSFSGKPGKDPGDQDGAAMDS VPLISPLDISQLQPPLPDQVVIKTQTTEYQLSSPDQQ NYTKSR
3717	A	58	618	GAGCTSPGLWARKAAARCLPTYPSPRAQPSNVGR RRRRRPGLGALAAGVPAMAESVERLQQRVQELE RELAQERSLQVPRSGDGGGGRVRIEKMSSEVVD SNPYSRLMALKRMGIVSDYEKIRTFAVAIVGVGG VGSVTAEMLTRCGIGKLLLFYDKVELANMNRLL FFQPHQAGLSKVQAAGHTPEE
3718	A	3	593	RGAGGRAGGRADGQPNMADQQRQSLSTSGESL YHVLGLDKNATSDDIKKSyrKLALKYHPDKNP NPEAADKFKEINNAHAILTDAKRNIDYKYGSLG LYVAEQFGREENVNTYFVLSSWWAKALFVFCGLL TCCYCCCCCCCCFNCCKGCKPKAPEGEETEFY VSPEDLEAQLQSDEREATDTPIVIQPASATEP
3719	A	2	2173	SGGVRMGSRADGPRTSghvtgkMAVFPWHSRN RNYKAEFASCRLEAVPLEFGDYHPLKPITVTESK TKKVNRKGSTSTSSSSSSSVVDPLSSVLDGTDPL SMFAATADPAALAAAMDSSRRKRDRDDNSVVG SDFEPWTNKRGEILARYTTTEKLSINLFMGSEKG KAGTATLAMSEKVRTRLEELDDFEEGSQKELLN LTQQDYVNRIEELNQSLKDAWASDQVKVAPKN VHPGKLVYERIFSMCVDSRSVLPDHFSPENANDT AKETCLNWFFKIASIRELIPRFYVEASILKCNKFLS KTGISECLPRLTCMIRGIGDPLAGSVYARAYLASRV GMEVAPHLKETLNKNFFDFLLTFKQIHGDTVQN QLVVQGVVELPSYLPYPPAMDWIFQCISYHAPEA LITEMMERCKKLGNNALLNSVMSAFRAEFAT RSMDFIGMIKECDESGFPKHLLFRSLGLNLALAD SELDLQCLNEINWELKKNFODILNCAEVWV EYTCKHFTKREVTVLAEVIKHMTPDRAFEDSY PQLQLIKKVIAHFHFSVLFSVEKFLPFLDMFQK ESVRVEVCKCIRTPLSKSPPRTRSS*MPFCMF ARPCMTL/CNALTLEDEKRLSYLINGFIKMVSF GRDFEQQLSFYVESRSMFCNLEPVLVQLIHSVNR LAMETRKVMKGNHSRKTAAFVRSWGAYWFTTTP SLAGIFTRLNLYLHSG
3720	A	24	296	ENLFRAGFAFSLRSSFYISKTYCSWFSNLISGSL ADFNKSGTRDYSRQMAVRE/KVFDVIIRCFKRH GAEVIDTPVFELKVRNGQEETW
3721	A	2	310	PSCLTCVGHCSIGGSCTMIGIMMPECHCSLHMTG PRCEEHVFIQQPGHIASILIPLLVLLLALVAGVV FWHKRRVQGAQGFQHQRTMGAMNVEIGNPTY K
3722	A	75	722	MELVAGCYEQVLFGFVHPEPEACGDHEQWTL VADFTTHAHTASLSAVAVNSRFVVTGSKDETHI YDMKKKIEHGALVHHS GTTICLFYGNRHLISGA EDGLICTWDAKKWECLKSIKAHKGQVTFLSIHP GKLALSVGTDKTLRTWNLVEGRSAFIKNIKQNA HIVEWSPRGEQYVVIIQNKIDYQLDTASISGTTN EKRISSVKFLSES
3723	A	110	316	MELSDNRRSGGLEGLAEKCPNLTYLNLSGNKIK DLSTVEALVSGTVLSLDLLFLVKFSEICLCLLISI
3724	A	3	406	VDRGTEAWQRDPAFSGLQRVGGVDVSFVKGDS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				VRACASLGVLSPFELEVYYEESRMVSLTAPYVSG FLAFREVPFLELVQQLREKEPGLMPQVLLVDGN GVLHHRGFGVACHLGVLTDLPVGVAKKLLQV DG
3725	A	3	406	VDRGTEAWQRDPAFSGLQRVGGVDVSFVKGDS VRACASLGVLSPFELEVYYEESRMVSLTAPYVSG FLAFREVPFLELVQQLREKEPGLMPQVLLVDGN GVLHHRGFGVACHLGVLTDLPVGVAKKLLQV DG
3726	A	1	433	SSDDRSIFRRLKLNIAIFDEGHMLKNMGSIRYQ HLMTINANNRLLLTGTPVQNNLELMSLLNFVM PHMFSSSTSEIRRMFSSKTKSADEQSIYEKERIAH AKQIKPFILRRVKEEVLKQLPPKKDRIELCAMSE KQEQLYL
3727	A	6	383	RIPRGKACXTVLGRSTGELEGFASSRLPPQPCGW GQSSDLLSRIDLDELMMKDEPPLDFPDLEGFEY AFNEKGQLRHIKTGEFFVFNYREHLHRWNQKRY EALGEITKYVYELLEKDCNSKKVS
3728	A	3	2452	EIAGAAAENMLGSLCLPGSGSVLLDPCTGSTISE TTSEAWSVEVLPSDSEAPDLKQEERLQELSCSG LGSTSDDTDVREVSSRPSTPGLSVVSGISATSEDP NKIEDLRSECSSDFGGKDSVTSPDMDEITHDFLYI LQPKQHFQHIEAEADMRIQLSSSAHQLTSPPSQSE SLLAMFDPLSSHEGASAVVRPKVHYARPSHPPD PPILEGAVGGNEARLPNFGSPMF*LPAEMEAFKQ RHS/YTPERLVRSSSDIVSSVRRPMSDPSWNR PAGNEERELPAAAIGATSLVAAPHSSSSSPSKDSS RGETEERKDSDEKSDRNRPWWRKRFVSAMPK APIPFRKKEKQEKDKDDLGPDRFSTLTDDPSRSL AQAQVAEDILDKYRNAIKRTSPSDGAM/NVEST EAMDCGSAHDEPRDEALQPSADDLP... AHPQDSAFSYRDAKKLRLP... HSTRNGLPDHTDPEDNEIVCFKLVQIAEAINLQD KNLMAQLQETMRCVCRFDNRCTCKLLASIAEDY RKRAPYIAYLTRCRQGLQTTQAHLELLQRVLR DKEVANRYFTTVCVRLLESKEKKIREFIQDFQK LTAADDKTAQVEDFLQFLYGAMAQDVIWQNAS EEQLQDAQLAIERSVMNRIFKLAFYPNQDGDILR DQVLHEHIQRLSKVVTANHRALQIPEVYLREAP WPSAQSEIRTISAYKTPRDKVQCILRMCSTIMNLL SLANEDSVPGADDFVPVLVFLIKANPPCLLSTV QYISSFYASCLSGEESYWWMQFTAAVEFIKTIDD RK
3729	A	3	2452	EIAGAAAENMLGSLCLPGSGSVLLDPCTGSTISE TTSEAWSVEVLPSDSEAPDLKQEERLQELSCSG LGSTSDDTDVREVSSRPSTPGLSVVSGISATSEDP NKIEDLRSECSSDFGGKDSVTSPDMDEITHDFLYI LQPKQHFQHIEAEADMRIQLSSSAHQLTSPPSQSE SLLAMFDPLSSHEGASAVVRPKVHYARPSHPPD PPILEGAVGGNEARLPNFGSPMF*LPAEMEAFKQ RHS/YTPERLVRSSSDIVSSVRRPMSDPSWNR PAGNEERELPAAAIGATSLVAAPHSSSSSPSKDSS RGETEERKDSDEKSDRNRPWWRKRFVSAMPK APIPFRKKEKQEKDKDDLGPDRFSTLTDDPSRSL AQAQVAEDILDKYRNAIKRTSPSDGAMANYEST

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EVMGDGESAHDSRDEALQNISADDLPDSASQA AHPQDSAFSYRDAKKKLRLALCSADSVAFVLT HSTRNGLPDHTDPEDNEIVCFLKVQIAEAINLQD KNLMAQLQETMRCVCRFDNRTCRKLLASIAEDY RKRAPYIAYLTRCRQGLQTTQAHLERLLRVLR DKEVANRYFTTVCVRLLESKEKKIREFIQDFQK LTAADDKTAQVEDFLQFLYGAMAQDVIWQNAS EEQLQDAQLAIERSVMNRIFKLAFFYPNQDGDILR DQVLHEHIQRLSKVVTANHRALQIPEVYLREAP WPSAQSEIRTISA YKTPRDKVQCILRMCSTIMNLL SLANEDSVPGADDFVPVLVFLIKANPPCLLSTV QYISSFYASCLSGEESYWWMQFTA AVEFIKTIDD RK
3730	A	3	2452	EIAGAAAENMLGSLCLPGSGSVLLDPCTGSTISE TTSEA WSVVEVLPSDSEAPDLKQBERLQELESCG LGSTSDDTDVREVSSRPSTPGLSVVSGISATSEIDP NKIEDLRSECSSDFGGKDSVTSPDMDEITHDFLYI LQPKQHFQHIEAEADMRIQLSSSAHQLTSPPSQSE SLLAMFDPLSSHEGASAVVRPKVHYARPSHPPPD PPILEGAVGGNEARLPNFGSPMF*LPAMEAFKQ RHS/YTPERLVRSSSDIVSSVRRPMSDPSWNRR PGNEERELPPAAIGATSLVAAPHSSSSSPSKDSS RGETEERKDSDEKSDRNRPWWRKRFVSAMPK APIPFRKKEKQEKDKDDLGPDRFSTLTDDPSRLS AQAQVAEDILDKYRNAIKRTSPSDGAMANYEST EVMGDGESAHDSRDEALQNISADDLPDSASQA AHPQDSAFSYRDAKKKLRLALCSADSVAFVLT HSTRNGLPDHTDPEDNEIVCFLKVQIAEAINLQD KNLMAQLQETMRCVCRFDNRTCRKLLASIAEDY RKRAPYIAYLTRCRQGLQTTQAHLERLLRVLR DKEVANRYFTTVCVRLLESKEKKIREFIQDFQK LTAADDKTAQVEDFLQFLYGAMAQDVIWQNAS EEQLQDAQLAIERSVMNRIFKLAFFYPNQDGDILR DQVLHEHIQRLSKVVTANHRALQIPEVYLREAP WPSAQSEIRTISA YKTPRDKVQCILRMCSTIMNLL SLANEDSVPGADDFVPVLVFLIKANPPCLLSTV QYISSFYASCLSGEESYWWMQFTA AVEFIKTIDD RK
3731	A	1	1305	VNTAMHEAKLMECEDELVEIQQRKQMIQIAVKIK ETKVMKLRKLAQQVANCRCQCLERSTVLINQAEH ILKENDQARFLQSAKNIAERVAMATASSQVLIPDI NFNDAFENFALDFSREKKLLEGLDYLTA PNPPSIR EELCTASHDTTIVHWISDDEFSISSYELQYTFTGQ ANFISLYNSVDSWMIVPNIKQNHYTIVHGLQSGTR YIFIVKAINQAGSRNSEPTRLKTNSQPFLDPKMT HKKLKISNDGLQMEKDESSLKKSHTPERFSGTGC YVYGVLHNSDNS*MFISLSFPLSHRYAIGIAYKSA PKNEWIGKNASSWVFSRCNSNFVVRHNNKEML VDVPPHLKRLGVLLDYDNY/NMFSFYDPANSLAH LHTFDVTFVLPVCPTFTIWNKSLMILSGLPAPDFI DYPERQECNCRPOESPYVSGMKTCH
3732	A	127	2832	LGQRLSLVPRPSLKRRLGKRLSLGLRERMMSLW WS/GPKVRTQATTGARPKTETKSVPAARPKTEAQ AMSGARPKTEVQVMGGARPKTEAQGITGARPKT DARAVGGARSKTDAKAIPGARPKDEAQA WAQS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				<p> EFGTEAVSQAEGVSQTNAVAWPLATAESGSVTK SKIACLWIEN*SMWM/PETFPGTQGGKGIQWFG PGEETNMGSWCYSRPRAREEASNESGFWSADET STASSFWTGEETSVRSWPRESNTRSRHRAKHQT NPRSRPRSKQEA YVDSWSGSEDEASNPFSSWVG ENTNNLFRPRVREEANIRSKLRTNREDCFESESED EFYKQSWVLPGEEANIDSGTETKKILLPWKLRA QKDVDSDRVKQEPFEEVHGSWFWAEKEASLE GGASAICESEPGTEEGAIGGSA YWAEKSSLGAV AREEAKPESEEEAIFGSWFWDRDEACFDLNPCPV YKVSDFRDAAEELNASSRPQTWDEVTVEFKPG LFHGVGFRSTSPFGIPEEASEMLEAKPKNLELSPE GEEQESLLQPDQPSPEFTFQYDPSYRSVREIREHL RARESAESESWSCSCIQCELKIGSEEFEEFLLMD KIRDPFIHEISKIAMGMRSASQFTRDFIRDSGVVS LIETLLNYPSSRVTSFLENMIHMAPPYPNLNMI TFICQVCEETLAHSVDSLEQLTGKNGCFRHLTMT IDYHTLIAN*YGPGFLLF*PQAQCGETKPHVLK MLLNLSENPA VAKKLFSAKALSIFVGLFNIEETN DNIQIVIKMFQNISNIKSGKMSLIDDDFSLEPLISA FREFEELAKQLQAQIDNQNDPEATGTTAFVGKG NNPSANRERLSPSVFCPGAQEAESLPARRVRGEE QRLLEEVGARTADGIPEGW </p>
3733	A	2	3274	<p> DVPLIRIEEDTGEIFTTGARIDREKLCAGIPRDEHC FYEVEVAILPDEIFRLVKIRFLIEDINDNAPLPAT VINISIPENSAINSKYTLPAAVDPDVGINGVQNYE LKSQNIIFGLDVIETPGGDKMPQLIVQKELDREEK DTYVMKVKVEDGGFPQRSSTAILQVSVTDNDN HPVFKETEIEVSIPENAPVGTSTVLHATDADIGE NAKHFSFNLVSNIARRLFHLNATTGLITIKPLD REEEENKILV*ASDCCCT*RAM/LV*VTV NDN*PSIDIRYIVNPVNDIV*SENIPLNKIALIT VTDKDADHNGRVTCFLLHEIPFRLRPVFSNQFL ETAAYLDYESTKEYAIKLLA* DAGKPPLNQSAM LFIKVKDENDNAPVFTQSFVTVEEPENNSPGIQLT KVSAMDADSGPNAKINYLLGPDAPPEFSLDCRT GMLTVVKKLDREKEDKYLFTILAKDNGVPPLTS NVTVFVSIIDQNDNSPVFTHNEYNFYVPENLPRH GTVGLITVTDPDYGDNSAVTSLDENDDFDIDSQ TGVIRPNISFDREKQESYTFYVKAEDGGRVSRSSS AKVTINVVDVNDNKPVFVPPSNCSYELVLPSTN PGTVVFQVIAVDNDTGMNAEVRYIVGGNTRDL FAIDQETGNITLMEKCDVTDGLHRLVKANDL GQPDLSFVIVNLFVNESVNTATLINELVPQKH LKHQ*PQILEIADVSSPTS DYVKILVA AVAGTTTV VVVIFTTAVVRCRQAPHLKAAQKNMQNSEWATP NPENRQMIMMKKKKKKKHSPKNLLN VVTIEE TKADDVDSGDNRVTLDLPIDLEEQTMGKYNWV TTPTTFKPDSPDLARHYKSASPQAFQIQPETPLN LKHIIQELPLDNTFVACDSISNCSSSSDPYSVSD CGYPVTTFEVPVSVHTRPPVDLEVGGAGSQGQVAI LTSSLMELLCLMVA AFLPLELRPLGQQNVMSW EQEAKILLVGYWGDGEWCHFHHLIPGPVNPNG YERKQYHILDSSEDTQPSGELCPVVRPFILSIQ LLQDDGEHCGTKQGFQPAVQLGLLPHKTLK </p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3734	A	1	840	GTRPGHLPAPSDGFCV/HL*SIPSWGSE*GESL/EM QLITSLGLQEFDIARNVLELIYAQTLVWIGIFFCPL LPFIQMIMLFIMFYSKNISLMMNFQPPSKAWRAS QMMTFFIFLLFFPSFTGVLCTLAITWRLKPSADC GPFRGLPLFIHSIYSWIDTLSTRPGYLWVWWTYRN LIGSVHFFFILTLIVLITYLYWQITEGRKIMIRLLH EQIINEGKDKMFLIEKLIKLDMEKKANPSSLVLE RREVEQQGFLHLGEHDGSLDLRSRRSVQEGNPR A
3735	A	2	432	VEVCRRYLWKMTVDASQNVQCCVIFSHFFIFN NLSKIKLLHTDTLLKIESKKHKA YLRSAAIEEERE SEFALRPTFDLTVRRNHLIEDVLNQLSQFENEDL RKELWVSFSGEIGYDLGGS/VKKEIFYCLFAEMIQ PEYGMFMY
3736	A	1542	343	KGAPSFVRLYQYPNFAGPHAALANKSFFKADKV TMLWNKKATAVLVIASDVKTGASYYGEQTL HYIATNGESAVVQLPKNGPIYDVVWNSSTEFCA VYGFMPAKATIFNLKCDPVDFGTGPRNAAYYS PHGHILVLAGFGNLIQI*AD/IMKVWNVKNYKLI SKPVASDSTYFAWCPDGEHILTATCAPRLRVNN GYKIWHYTGSIHKYDVPSNAELWQVSWQPFLLD GIFPAKTTITYQAVPSEVPNEEPKVATAYRPPALRN KPITNSKLHEEEPPQNMKPQSGNDKPLSKTALKN QRKHEAKKAAKQEARSDKSPDLAPTPAQSTPR NTVSQSISGDPEDKKIKNLKKKLKAIEQLKEQAA TGKQLEKNQLEKIQKETALLQLEDLELGI
3737	A	3190	664	VAMGTPRAQHPPPPQLFLILLSCPWIQGLPLKEE EILPEPGSETPTVASEALAEHLHGALLRRGPENG YLPGPPLGPEGGEEETTTTITTTTITTTTITTTT NNNISEGEGYVESPDLGSPVSRITGLLDCTYSIHV YGGYHIEIQVQTLNLSQBEELLVLAGCCPGLAP RLLAISSMLGEGQVLRSPNRLLLHFQSPRVTS GGFRIHYQAYLLSCGFPPRAHGDVSVTDLHPGG TATTCDSGYQLQGEETLICLNGTRPSWNGETPS CMASCGGTIHNATLGRIVSPEPGGAVGPNLTCR WVIEAABGRRLHLHFERVSLDEDNDRMLMVRSG SPLSPVIYDSMDDDVPERGLISDAQSLYVELLSET PANPLLLSLRFEAFEDRCFAPFLAHGNVTTTDP YRPGALATFSCLPGYALEPPGPPNAIECVDPTEPH WNDTEPACKAMCGGELSEPAGVVLSPDWPQSY SPGQDCVWGVHVQEEKRILLQVEILNVREGDML TLFDGDGPSARVLAQLRGPQPRRLLSSGPDILT QFQAPPGPPNPGLGQGFVLHFKEVPRNDTCPELP PPEWGWRTASHGDLIRGTVLTYQCEPGYELLGS DILTCQWDLSSWAAPPACQKIMTCADPGELANG HRTASDAGFPVGSVHQRCLPGYSLEGAAMLTC YSRDTGTPKWSDRVPKCALKEYECLNPGVPENG YQTLYKHHYQAGESLRFFCYEGFELIGEVTITCV PGHPSQWTSQPPLCKVTQTTDPSRQLEGGNLAL AILLPLGLVIVLGSVYTYTKLQKSLFGFSGSH SYSPITVESDFSNPLYEAGDTREYEVSI
3738	A	3190	664	VAMGTPRAQHPPPPQLFLILLSCPWIQGLPLKEE EILPEPGSETPTVASEALAEHLHGALLRRGPENG YLPGPPLGPEGGEEETTTTITTTTITTTTITTTT NNNISEGEGYVESPDLGSPVSRITGLLDCTYSIHV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				YPGYGIEIQVQTLNLSQEEELLVLAGGGSPGLAP RLLANSSMLGEGQVLRSPNRLLLHFQSPRVPRG GGFRIHYQAYLLSCGFPPRAHGDVSVTDLHPGG TATFHCDSGYQLQGEETLCLNGTRPSWNGETPS CMASCGGTIHNA TLGRIVSPEPGGAVGPNLTCTCR WVIEAAEGRRLHLHFERVSLDEDNDRMLMVRSGG SPLSPVIYDSMDVPERGLISDAQSLYVELLSET PANPLLLSLRFEAFEDRCFAPFLAHGNVTTTDP YRPGALATFSCLPGYALEPPGPPNAIECVDPTEPH WNDTEPACKAMCGGELSEPAGVVLSPDWPOSY SPGQDCVWGVHVQEEKRILLQVEILNVREGDML TLFDGDGPSARVLAQLRGPQRRRLSSGPDLT QFQAPPGPPNPLGQGFVLHFKEVPRNDTCPELP PPEWGWRTASHGDLIRGTVLTYQCEPGYELLGS DILTCQWDLWSAAPPACQKIMTCADPGEIANG HRTASDAGFPVGS HVQYRCLPGYSLEGAAMLTC YSRDTGTPKWSDRVPKCALKYEPCLNPGVPENG YQTLYKHHYQAGESLRFFCYEGFELIGEVTITCV PGHPSQWTSQPPLCKVTQTTDPSRQLEGGNLAL AILLPLGLVIVLGSVYIYYTKLQGKSLFGFSGSH SYSPITVESDFSNPLYEAGDTREYEVS
3739	A	734	445	LLEPEPAEEYTEQSEVEST/EGMILI*CCLYFAAFQ TNVSNIFYALQYVNRQFMAETQFTSGEKEQVDE WTVETVEVRVLCIAKLLSLSSVS NFYLY
3740	A	2	1578	MAHYITFLCMVLVLLQNSVLAEDGEVRSSCRT APTDLVFILDGYSYVGPNFEIVKKWLVNITKNF DIGPKFIQGVVQYSDYPVLEIPLGSYDSGEHLTA AVESILYLGGNKTGKAIQFALDYLFKSSRFLT KIAVVLTDGKSQDDVKDAAQAARDSKITLFAIG VGSETEDAELRAIANKPSSTYVFYVEDYIAISKIR EVMKQYLCEESVCPVPAARDEPGFVFLD VNKKKIKIQLSPKIKGYEVTSKVISELNV FPEGLPPSYVVFVSTQRFKVKKIWDLWRLTIDG/* PQIAVTLNGVDKILLFTTTSVINGSQVVFAPQV KTLFDEGWHQIRLLVTEQDVTLYIDDQDIENKEL HPVLGILINGQTQIGKYSKKEETVQFDVQKLRIY CDPEQNNRETACBPGFCLNGPSDVGSTPAPCICP PGKPGQLQGPKGDPGLPGNPGYPGQPGQDGKPV TESLVISGISGITGYQGIAGTPGVPGSPGIQGARGL PGYKGEPGRDGDK
3741	A	5048	1236	MSAPAGSSHPAASARIPPKFGGSAVSGAAAPAGP GAGPAPHQQNGPAQNQMVPQSGYGLHHQNYIA PSGHYSQGPQKMTSLPLDTQCGDYYSALYTVPT QNVTPNTVNQQPGAQQLYSRGPAPHIVGSTLGS FQGAASSASHLHTSASQPYSSFNHYNPAMYS ASSSVASQGFSTCGHYAMSTVSNAAYPSVSYPS LPAGDTYQGMFTSQNAPTVPVKDNSFSGQNTA ISHPSPLPPLPSQQHHQQSLSGYSTLTWSSPGLP STQDNLIRNHTGSLAVANNNPITTVADSLSCPVM QNVQPPKSSPVVSTVLSGSSGSSSTRTPPTANHPV EPVTSVTQPSSELLQKGVQYGEYVNNQASSAPT PLSSTSDDEREEDEEAGVDSSTTSSASPMFNS YDALEGGSYDMLSSASSAPDPAPEDPASAP APASAPAPVVPQPSKMAKPLAMAIQHFSLVIRML QHHLFLEYSPSNPVYSGFQQYPQQYPGVNQLSSS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=/possible nucleotide insertion)
				IGGLSLQSSQPESLRPVNLTQERNILPMTVPVWAP VPNLNADLKKLNCSPDSFRCTLTNPQTQALLNK AKLPLGLLLHPPFRDLTQLPVITSNTIVRCRSCRTYI NPFVVSFIDQRR*KCNLCYRVNDVPBEFMYNPLT RSYGEPHKRPEVQNSITVEFIASSDYMLRPPQPAV YLFVLDVSHNAVEAGYLT/LWCQSLLENLDKLP GDSRTARIGFMTFD*STYSFLQFTQEGLSQPQMLI VSDIDDVFLPTPDSLLVNLYESKELIKDLLNALPN MFTNTRETHSALGPALQAAFKLMSPTGGRVSVF QTQLPSLGAGLLQSREDPNQRSSTKVQHLGPAT DFYKKLALDCSGQQTAVDLFLLSSQYSDLASLA CMSKYSAGCIYYPSPFHYTHNPSQAEKLQKDLK RYLTRKIGFEAVMRIRCTKGLSMHTFHGNFFVRS TDLLSLANINPDAGFAVQLSIEESLTDTSLVCFQT ALLYTSSKGERRIRVHTLCLPVVSSLSDDVYAGVD VQAAICLLANMAVDRSVSSSLSDARDALVNADV DSLSAYGSTVSNLQHSALMAPSSLKLFPLYVLAL LKQKAFRTGTSTRLDDRVIAMCQIKSQPLVHLM KMIHPNLYRIDRLTDEGAVHVNDRIVPQPPLQKL SAEKLREGAFLMDCGSVFYIWWGKGCDDNNFIE DVLGYTNFASIPQKMTHELPELDTLSSERARSFIT WLRDSRPLSPILHIVKDESPAKAEFFQHLIEDRTE AAFSYYEFLHVVQQQICK
3742	A	934	68	SMLASQGVLLHPYGVPMIVPAAPYLPGLIQGNQE AAAAPDTMAQPYASAQFAPPQNGIPAETAPHP HPAPEYTGQTTVPEHTLNLYPPAQTHSEQSPADT SAQTVSGTRNKQD*RTSDGWSPKQTQS*KHGK QVSSPSGLHVSNIPIFRFRDPDLRQMFAGQFGKILD VEIFNERGSKGFGVTFENSADADRAREKLHGT VVEGRKIEVMNATARVMNKKTVNPYTNQWK INPVVG.IVYSPETYAGTVLLCQA PSTDFRGAKLH/TSRFL/SC3
3743	A	3	1456	QFQQAWMQNKVPIAPNEVLNDRKEDIKLEBKK KTQAEIEQEMATLQYTNPQLEQLKIERLAQKQV EQIQPPSSGTPLLGPQFPQGGPMSQIPQGF/PTA PSISADANEHGS/KGPPGPQGGQFRPPGPQGGQMG QGPPLHQGGGGPQGFMGPPQGGPQGGPGLPRQD MHGPQGMQRHPGPHGPLGPQGGPQGGSSGPQ HMGPPQGGPQGHIGPQGGPQGHLPQGGPPT QGMQGGPPGRGMQGGPPHGHIGGGPGSQGIQGP VSQGPLMGLNPKGMQGGPPGPRENQGPAPQGM MGHPPQEMRGPHPPGGLLGHGPQEMRGPPQEI MQGPPPGQSMGLGPPQELRGPPGSQSGQGGPQGS GPPPGGGMQGGPPGPQGGQNPARGPHPSQGGPIFFQ QKQTPLLGDGPRAFPNQEQQSTGPPPLIPGLGQQ GAQGRIPPLNPGQGGPKNKVS/ERGAPPRHEGRA PPRGRDGFPGPMKTLV
3744	A	1571	652	PLTGRKCPGWTHSGSRRSPRIAEVPGFPKRAEA SRQFSETADRLLELLRRAVMAAARATTPADGEEP APEAEALAAARERSRFLSGLELVKQGAEARVFR GRFQGRAAVVKHFRPKGYRHPALEARLGRRTV QEARALLRCRRAGISAPVVFFVDYASNCLYMEEI EGSVTVRDVFSPLWRLKKTQGLSNLAKTIGQVL ARMHDEDLIHGDLTTSNMLLKPPLEQLNIVLIDF GLSFISALPEDKGVDLYVLEKAFLSTHPNTETVFE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				AFLKSYSTSSKKARPVLKKLDEVRLRGKKRSMV G
3745	A	127	1433	GSHRFSLASPLDPEVGPYCDTPTMRTLFLNLLWLA LACSPVHTTSLKSDAKKAASKTLLKSQFSDKPV QDRGLVVTDLKAESVLEHRSYCSAKARDRHFA GDVLGYVTPWNHGYDVTKVFGSKFTQISPWWL QLKRRGREMFVETGLHDVDQGWMAVRKHAK GLP*CLGSCLRTGLTMISG/YVLDSEDEIEELS VQVAKNQHFDDGFVVEVWNQLLSQKRVGLIHM LTHLAEALHQARLLALLVIPPATPGTDQLGMFT HKEFEQLAPVLDGFSMLTYDYSTAHPGPNAPL SWVRACVQVLDPKSKWRSKILLGNFYGMDYA TSKDAREPVVGARYIQTLKDHPRPMVWDSQVSE HFFEYKKSRSGRHVVFPYTLKSLQVRLELARELG VGVSIVELGQGLDYFYDLL*VGIAASAVDVFFSK PWSE
3746	A	1	898	IDRAAECRTKPLPMAVSIRGNADSIACLVLMVL YLIKKRLVACAAVFYGFVHMKIYPETYILPITL HLLPDRDNDKSLRQFRYTFQAEL*ELLKRLCNR TALMFVAVAGLTFFALSFGFYEYGEWFLEHTYF YHLTRDIRHNHNSPYFYMLYLTAESKWSFSLGIA AFLPQLILLSAVSFAYYRDVFCWFLHTSIFVTFN KVCTSQYFLWYLCLLPLVMPLVRMPWKRAVVL LMLWFIGQAMWLAPAYVLEFQGKNTFLFIWLA GLFFLLINCSILIQISHYKEEPLTERIKYD
3747	A	1	2325	MVISFQGLVTFGDVAVDFSQEEWEWLNPIQRNL YRKVMLENYRNLASLGLCVSKPDVISSLEQKKEP WTVKRKMTRAWCPDLKAVWKIKELPLKKDFCE GKLSQAVITERLTSYNLEYSLLGEHWDYDALFET QPGLVTIKNLAVDFRQQLHPAQKNFCKNGIWEN NSDLGSAGI/TAEDL/TLIDG/ET/MVKREL TQELPSGQRS/VHETQELFPKQ/VAEG/IDRTSN TKLDCSSFRENWSDSYVFGKRL/GOETQFRQE PITHNKTLSKERERTYNKSGRWFLDDEEEKVH NRDSIKNFQKSSVVIKQTGIYAGKKLFLKMECKK TFTQSSSLTVHQRIHTGEKPYKCNECGKAFSDGS SFARHQRCHTGKKPYECIECGKAFIQNTSLIRHW RYYHTGEKPFDCIDCGKAFSDHIGLNQHRRIHTG EKPYKCDVCHKSFRYGSSTLVHQRIHTGEKPYE CDVCRKAFSHHASLT/QHQRVHSGEKPFCKKEC GKAFRQNIHLASHLRIHTGEKPFCEAECGKSFIS SQLATHQRIHTGEKPYECKVCSKAFTQKAHLAQ HQKTHHTGEKPYECKEBCGKAFSQTTHLIHQRVH TGEKPYKCMECGKAFGDNSSCTQHQRLLHTGQRP YECIECGKAFKTKSSLICHRRSHTGEKPYECSVC GKAFSHRQSLSVHQRIHSGKKPYECKECKRTFIQI GHLNQHKRVHTGERSYNYKKSRKVFRQTAHLA HHQRIHTGESSTCPSLPSTSNPVDLFPKFLWNPSS LPSP
3748	A	823	1	GGYTKSGYDSACKDFVPHDLEVQIPGRVFLVTG GNSGIGKATALEIAKRGGTVHLVCRDQAPAE RGEIIRE/SGNQNIHLHIVDLSDPKKIWKFFVENFKQ EHKLHVL/VNNAGCMVKNKREAHKKMDFEKNFG CQYSGVCTFLTTRPDPLCWRKNTDPRVTIVSSG GMLVQKLNNQ*SPVRKNTIWMGMTMVYAQNKVS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ERQQVVLTERWGPAPGVHFFSSMHPGWAIDTPG VRQAMPGFHVQASGYRLRSEAQGADTMLWLAL SSARSRTAQR
3749	A	1939	715	GFLRLSQATRQRLSIPVMVLTLDPTRD\QCFGDR FSRLLLDDEFLGYDDILMSSVKGLAENEENKGFRL NVVSGEHYRFVSMWMARTSYLAAFANHGQSF TLSVSHACCGYSHHQIFVFVDLLQMLEMNMAIA FPAAPLLTVILALVGMEAIMSEFFNDTTTAFYILI VWLADQYDAICCHTSTSKRHWLRFFLYLHFAFY AYHYRFNGQYSSLALVTSWLFIOHSMIYFFHHYE LPAILQHVRIQEMLLQAPTLGPGTPTALPDDMN NNSGAPATAPDSAGOPPAGPVSPGASGSPGPV AAAPSSLVAAAASVAAAAGGDLGWMAETAIIIT DASFLSGLSASLLERRPASPLGPAGGLPHAPQDS VPPSDSAASDITPLGAAVGGPSPASMAPTEAPSE VGS
3750	A	2	844	GLLEPFSKLLSFVIONAVFTLAYLVELCGLCYRA FTKERDKFYLSRSVVLELLQALKLKSPLPDTNLL LLVQFICADAGTKLAESTILSKQMIASVPGCGTA AMECVRQYINEVLDFMADMHTLTKLKSHMKTC SQPLHEDTFGGHLKVGLAQIAAMDISRGNHDRN KAVIRYLPWLYHPPSAMQQGPKEFIECVSHIRLL SWLLGSLTHNAVCLKWPPLPGLPIPLDAGSHV ADHLIVILIGFPEQSKTSVLHMCSLFHAFSLAQL WDSLLARQSGRW
3751	A	431	2	AFTRKCEETAFTVPQCEIIPTE/WVCCRIPGTSSLER NPGVKEGCEFCPPKVEFFKDDANHDPQWSRQ QLIAAKFGFAALGI/QTEVDIMSHAT*AVFEIPEKS RL/PQNCTPVDMKIEFGVHVTSKEILTDVIDNDS* RHSPS
3752	A	131	1278	AVEGSGP...TAS...PMSVLCN...FLWQNEG PGGRW...YISRR...SSDPAWAVEWIELPRGLSLSS LGSARTLR...WSRSSRPSSVDSQDLPEVNVGDTV AMPLPKSRRAL...FJAALARSSLHGISQVVKDHV TKPTAMAQGRVAHLIEWKGWSKPSDSPAALESA FSSYSDLSEGEQEARFAAGVAEQFAIAEAKLRA WSSVDGEDSTDDSYDEDFAGGMDTDMAGQLPL GPHLQDLFTGHRFSRPVRQGSVEPESDCSQTVP DTLCSSLCLEDGLLGSAPLAPSCWAMSCFSPN CPPAGKVPSAAW/APLEAQDSLINSPLTESCLSP AEEBPAPCKDCQPLCPPLTGSWERQRQASDLASS GVVSLDEDEAEPEEQ
3753	A	3	1138	YYSSVRQRTCEPRFRECAALIEGSATEVYAG EWRADRRSGFGVSQRSNGLRYEGEWLGNRRHG YGRTRPDGSREEGKYKRNLVHGGVRVRSLLPL ALRRGKVKVDRAVEGARAVSAARQREQEIA AARAADALLKAVAASSVAEKAVEAARMALIA QDLQPMLEAPGRRPRQDSEGSDTEPLDEDSGV YENGLTPSEGSPELPSSPASSRQPWRPPACRSPLP PGGDQGPFSPPKAWPEEWGGAGAQAEELAGYE AEDEAGMQGPGPRDGSPLLGCSDSGSLREEE GEDEEPLPLRAPAGTEPEPIAMLVLRGSSSRGPD AGCLTEELGEPATERPAQPGAANPLVVGAVAL LDLSLAFLFSQLLT
3754	A	2	3338	SSLLEKMTSSDKDFRFRMATSDLMSELQKDSIQLD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				EDSEKVVVKMLRLLEDKNGEVQNLAVKWLGV PLGAFHASLLHCLLPQLSSPRLAVRKRAVGALGH LATACSTDLFVELADHLLDRLPGPRVPTSPTAIRT LIQCLGSVGRQAGHRLGAHLDRVLVDFCNL DDELRESCLQAFEAFLRKCPKEMGPHVFNVT LCLQYIKHDPNPNYDSDEDEEQMETEDSEFSEQE SEDEYSDDDDMSWKVRRAAAKCIAALISSRPDL LPDFHCTLAPVLIRRFKEREENVKADVFTAYIVL LRQTRPPKGWLEAMEEPTQTGSNLHMLRGQVPL VVKALQRQLKDRSVRARQGCFSLLTELAVLPG SLAEHMPVLVSGIIFSLADRSSSSTIRMDALAFLO GLLGTEPAEAFHPLPILLPPVMACVADSFYKIA AEALVVLQELVRALWPLHRPRMLDPEPYVGEMS AVTLARLRATDLDQEVKERAISCMGHLVGHLD RLGDDEPTLLLLLDRLRNEITRLPAIKALTLVAV SPLQLDLQPIAEALHILASFLRKNQRLRLATLA ALDALAQSQGLSLPPSAVQAVLAELPALVNESD MHVAQLAVDFLATVTQAQPASLVEVSGPVLSEL LRLRSPLLPAGVLAAGFLQALVGTRPPCVDY AKLISLLTAPVYEQAVDGGPGLHKQVFHSLARC VAALSAACPQVEASTASRLVCDARSPHSSTGVK VLAFLSLAEVGQVAGPGHERELKAVLLEALGSPS EDVRAAASYALGRVGAGSLPDFLPFLLEQIEAEP RRQYLLHSLKEALGAAQPDLSKPYAEDIWALL FQCEGAEETRGVVAECIGKLVVNPSFLLPRL RKQLAAGRPHTRSTVITAVKFLISDQPHPIDPLK SFIAVHNKPSLVRLDLDLPLLYQETKIRRDILRE VEMGPFKHTVDDGLDVRKAAFECMYSLLESCLG QLDICEFLNHVEDGLKDHYDIRMLTFIMVARLAT LCPAPVLQRVDRLEIPLRATCTAKVKAGSVKQBF EKQDELKRSATRAVAALLTPEVETSPMADESS QIRSNPELAALFESIQETTSAPSNDMSMELS
3755	A	2	3338	SSLEKMTSSDKDFRFRMATSDLMSELQKDSIQLD EDSEKVVVKMLRLLEDKNGEVQNLAVKWLGV PLGAFHASLLHCLLPQLSSPRLAVRKRAVGALGH LATACSTDLFVELADHLLDRLPGPRVPTSPTAIRT LIQCLGSVGRQAGHRLGAHLDRVLVDFCNL DDELRESCLQAFEAFLRKCPKEMGPHVFNVT LCLQYIKHDPNPNYDSDEDEEQMETEDSEFSEQE SEDEYSDDDDMSWKVRRAAAKCIAALISSRPDL LPDFHCTLAPVLIRRFKEREENVKADVFTAYIVL LRQTRPPKGWLEAMEEPTQTGSNLHMLRGQVPL VVKALQRQLKDRSVRARQGCFSLLTELAVLPG SLAEHMPVLVSGIIFSLADRSSSSTIRMDALAFLO GLLGTEPAEAFHPLPILLPPVMACVADSFYKIA AEALVVLQELVRALWPLHRPRMLDPEPYVGEMS AVTLARLRATDLDQEVKERAISCMGHLVGHLD RLGDDEPTLLLLLDRLRNEITRLPAIKALTLVAV SPLQLDLQPIAEALHILASFLRKNQRLRLATLA ALDALAQSQGLSLPPSAVQAVLAELPALVNESD MHVAQLAVDFLATVTQAQPASLVEVSGPVLSEL LRLRSPLLPAGVLAAGFLQALVGTRPPCVDY AKLISLLTAPVYEQAVDGGPGLHKQVFHSLARC VAALSAACPQVEASTASRLVCDARSPHSSTGVK VLAFLSLAEVGQVAGPGHERELKAVLLEALGSPS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EDVRAAASYALGRVGAGSLPDFLPFLLEQIEAEP RRQYLLHSLKEALGAAQPSLKPYAEDIWALL FQRCEGAEEGTRGVVAECIGKLVLVNPSFLLPRL RKQLAAGRPHTRSTVITAVKFLISDQPHPIDPLK SFIAVHNKPSLVRDLDLPLLYQETKIRRDIRE VEMGPFKHTVDDGLDVRKAAFECMYSLLESCLG QLDICEFLNHVEDGLKDHDIRMLTFIMVARLAT LCPAPVLQRVDRLIEPLRATCTAKVKAGSVKQEF EKQDELKRSAMRAVAALLTIPEVGKSPIMADFSS QIRSNPELAALFESI QKDST SAPSTDSMELS
3756	A	112	1361	SLEEQQGRHPSFAPK CASQILGRIMITLITEQLQK QTLDELKCTRFSSISLPLPDHADISNCGNSFQLVSE GASWRGLPHCSAEFQ/DQPQLQLPSLRPEPAPQ TTHRGNSPKEQPFSQVLRPEPPDPEKLPVPPAPPS KRHCRSLSPVVDLSRWQPVWRPAPSKLWTPIKH RGSGGGGGPQVPHQSPPKRVSSL/SVPSSQCLFS MCPSSHTLQPSFLQPGPGPDSSRPCAASFQSGSW ESDAESLSPCPPQRRFSLSPSLGPAQSRFLPSARSS PASSPELPWRPRLNLPRLSRSPQCDLDARKTGV KRRHEEDPRRLRPSLDFDKMNQKPYSGGLCLQE TAREGSSISPPWFMACSPPLSASCPTGGSSQVL SESEEEEGAVRWGRQALSKRTLQCRDFGDLDL NLIEEN
3757	A	413	1	PKPMLQQDFT/SLPDQGLDHIAE/NSYFDARSLCA AELVCKEWQQVTSE*MLWKKLIERMVHAYPLW KGLSEKVV/DQHLFKNRPTDGPPNSFHRSLYPKII QVIETIESNWQCG*HTLQRIQCHSEKSKGVYCLQ YDDEK
3758	A	2	613	FVSGSPWRMDGSTERLEARRPAGRLPWSSRQEM TRRPSLMAGRQHGWSAQQSATVANPVPGANPD LLPHFLGEPEE VYTVKNKPVLLVCKAYATQIFF KCNGEWVRQVDHVIERSIDGSSGLI DEVRINV SRQQVEKVFGLLEEYWCQCVAWSSSGTTKSQKA YIRIAYLRKNFEQEPLAKEVSLEQGIVLPCRPEGI PPAE
3759	A	1	561	ADDTLHLWNLRQKRPAIHLHLKFCRERVTFCHLP FQSKWLYVGTERRGNIHIVNVEFTLSGYVIMWN KAIELSSKSHPGPVVHISDNPMDEGKLLIGFESGT VVLWDLKSKKADYRYTYDEAIHSVAWHHEGKQ FICSHSDGTLTIWNVRSPAKPVQTITPHGKQLKD GKKPEPCKPILKVEFXTTR
3760	A	1	824	LPACRCGCVAGCPSNHGICRCLRASERQVCVMH LKHLRTLSPQDGAAKVTCAWSONNAKFAVC TVDRVLLYDEHGERRDKFSKPADMKYGRKS YMKGMASFDPSTKIAIGQTDNIYVYKIGEDWG DKKVICNKFIQTVKFRPVPGTLG*TNIQYTYL*IQ PGVAFLTSECDFSYCKDGASWLFMVICCLP*SPA VSFIGD*SAVTCLQWPAEYIIVFGLAEGKVRLS NTKTNKSSTYGTESYVVSLLTNCSGKGILSGHA DGYQR
3761	A	2253	320	PVIQRCSQPYGFSLLISFFLKCVSETSQQPPSRKVF QLLPSFPTLTRSKSHESQLGNRIDDVSSMRFDLSH GSPQMVRDGLSVTHRFSTKSWLSQVCHVCQK SMIFGVKCKHCRLKCHNKCTKEAPACRISFLPLT RLRRTESVPSDINNPDRAAEPHFGTLPKALTKK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EHPPAMNHL DSSSNPSSTTFSTPSSPAPFTSSNPS SATTPNPSPIGQRADSRFNFPSC/AYFIHHR/QQFI FPDISAFAHAAPLPEAADGTRLDDQPKADVLEAH EAEAEPEAGKSEAEDDEDEVDLPSRRPWGR PISRKASQTSVYLQEWDPFEQVELGEPGQGRW GRVHRGRWHGEVAIRLLEMDGHNQDHLKLFKK EVMNYRQTRHENVVLFMGACMNPPLAITSFC KGRTLHSFVRDPKTSLDINKTRQIAQEIIKGMGYL HAKGIVHKDLKSRNVFYDNG\KVVTIDFGLFAGIS GVVPEGRRENQLKLSHDWLCYLAPEIVREMTGP KDEDQLPFSKAADVYAFGTWVYELQARDWPLK NQAAEASIWQIGSGEGMKRVLTSSVLGKEVSEN LSACWAFDLQERPSFSLMDMLEKLPKLNRRLS HPGHF*KSADINSSKVVPRFERFGLGVLESSNPK M
3762	A	2	1578	MAHYITFLCMVLVLLQNSVLAEDGEVRSSCRT APTDLVFILDGSSVGPENFEIVKKWLVNITKNF DIGPKFIQVGVVQYSDYPVLEIPLSYDSGEHLTA AVESILYLGNTKTGKAIQFALDYFAKSSRFLT KIAVVLTDGKSQDDVKDAAQAARDSKITLFAIG VGSETEDAELRAIANKPSSTYVFYVEDYIAISKIR EVMKQKLCEESVCPTRIPVAARDERGF DILLGLD VNKKVKKRIQLSPKKIKGYEVTSKVDLSELT SNV FPEGLPPSYVVFVSTQRFKVKKIWDLWRILTIDG/* PQIAVTLNGVDKILLFTTTSVINGSQVVTANPQV KTLFDEGWHQIRLLVTEQDVTLYIDDQIENKPL HPVLGILINGQTQIGKYSGKEETVQFDVQKLRIY CDPEQNNRETACEIPGFCLNGPSDVGSTPAPCICP PGKPGLOGPKGDPGLPGNPGYPGQPGQDGKPV TESLVISGISGITGYQGIAGT*GVPGSPGIQGARGL GKYKG*PGRDGP
3763	A	3	1267	CKVWRNPLNLRGALNRYTWVTGREPLTYD MNL SAQDHQTFCTCDSDHLRPADAIMQKAWRE RNPQARISAAHEALNECATAYILLAEBAATTIA EAEKLFKQALKAGDGCYRRSQQLQHHSQYEA QHSVLVPLQVTRHQCLGVHQQKASNVCQKTRE DQGSSENDERFNEGVPSEYVQYP*KPFAKALLEL QAYADVQAVLAKYDDISLPKSA TICYTAALLKA RAVSDKFSPEAASRRGLSTAEMNAVEAIHRAVEF NPHVPKY LLEMKSLILPPEHILKRGDSEAIAYAFF HLAHWKRVEGALNLLHCTWEGTFRMIPYPLEKG HLFPYPICETETADRELLPSFHEVSVPKKEPFFI LFTAGLCSFTAMLALLTHQFPELMGVFAKAVSV CLEGGLGEWMGKAKGIKAA
3764	A	25	1032	RSADGLCGNKDRERGNEFTRNQAAQEVVNP KKMKKKKYVNSGTVTLLSFAVESECTFLDYIKG GTQINFVAIDFTASNGNPSQSTSLHYMSPYQLN AYALALTAVGEIIQHYDSDKMFALGFGAKLPPD GRVSHEFFLNGNQENPSCCGIDGILEAYHRSRLT VQLYGPTNFAPVTVHVARNAAAVQDGSQYSVL LIITDGVISDMAQTKAIVNGSKLPMSIIVGVGQ AEFNAMVELDGDVRISSRGKLAERDIVQFVPPR DYVDRTGNHVL SMARLARDVLAEPDQLVSYM KAQGIRPRSPPAAPTHSPSQSPARTPPACPLHTI
3765	A	172	3456	LGMMDSPKIGNGLPVIGPGTDIGISSLHMVG YLG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KNFDSA KVP SDEYCPACKEKGK LKALKTYRISFQ ESIFLCEDLQCIYPLGSKSLNNLISPDL EECHTPHK PQKRKSLESSYKDSLLLANSKKTRNYIAIDGGKV LNSKHNGEVYDETSSNLPDSSGQONPIRTADSLE RNEILEADTVDMATTKDPATVDVSGTGRPSPQN EGCTSKLEMPLESKCTSFPAALCVQWKNAALC WLD CILSALVHSEELKNTVTGLCSKEESIFWRL TKYNQANTLLYTSQ LSGVKDGDCKKLTSEIFAEL ETCLNEVRDEIFISLQPLRCTLGDMESPVFAFPL LLKLETHIEKLFYLSFSDFECSQCGHQYQNRH MKSLVTFTNVIPEWHPLNAAHFGPCNNCNSKSOI RKMVLEKVSPIFMLHFVEGLPQNDLQHYAFHFE GCLYQITSVIQYRANNHFITWLDADGSWLECD LKGPCSERHKKFEVPASEIHIVTWERKISQVTDKE AACLP LKKTNDQH ALSNEKPVSLTSCSVGDAAS AETASVTHPKDISVAPRTLSQDTAVTHGDHLLSG PKGLVDNLP LTL EETIQKTASVSQ LNSEAFLEN KPVAENTGILKTNTLLSQESLMASVSAPCNEKLI QDQFVDISFSPQVNTNMQSVQLNTEDTVNTKS VNNTDATGLIQGVKSVEIEKDAQLKQFLTPKTEQ LKPERVTSQVSNLKKKETTADSQTTSKSLQNS LKENQKKPFVGSWVKGLISR GASFMPLCVSAHN RNTITDLQPSVKGVNNFGGFKTKGINQKASHVSK KARKSASKPPPI SKPPAGPPSSNGTAAHPHAHAA SEVLEKSGSTSCGAQLNHSSYNGGISSANHEDLV EGQIHKLRLKLRKKLKA EKKKLAALMSSPQSRT VRSENLEQVPQDGPNDCE SIEDLLNELPYPIDIA NESACTTVPGVSLYSSQTHEELAE LLSPTPVSTE LSENGEGDFRYLGMGD SHIPPPVPSEFNDVSQNT HLRQDHNYSPTKKNPCEVQPD SLTNNACVRTL NLES PMKT DIFDEFESSA LNALANTL DLF YLFENY
3766	A	3	1662	AQQIVYRNVMLENYKNLVSLGYQLTKPDVILRL EKGEFPWLVEREIQETHPDSETA FEIKSSVSSRSI FKDKQSCDIKMEGMARNDLWYLSLEEVWKCRD QLDKYQENPERHLRQVAFTQKKVLTQERVSESG KYGGNCLLPAQLVLREYFHKRDSHTKSLKHDLV LNGHQDSCASNSNECGQTF CQNIHLIQFARTHTG DKSYKCPDNDNSLTHGSSLGISKGIHREKPYECK ECGKFFSWRSNLTRHQLIHTGEKPYECKECGKSF SRSSH LIGHQKTHTGEEPYECKECGKSFSWFSHL VTHQRTHTGDKLYTCNQCGKSF/VHSSRLIRHQR THTGEKPYECKEPCGKSFRQSTHLILHQRTHVRVR PYECNECGKSYSQSRSHLVVHHRIHTGLKPFECKD CGKCFSRSSHL YSHQRTHTG EKPYECHDCGKSFS QSSALIVHQRIHTGEKPYECCQCGKAFIRKNDLIK HQRIHVGEETYKCNQCGHIFSQNSPFIVHQIAHTG EQFLTCNQCGTALVNTSNLIGYQTNHIRENAY
3767	A	3	1622	AQQIVYRNVMLENYKNLVSLGYQLTKPDVILRL EKGEFPWLVEREIQETHPDSETA FEIKSSVSSRSI FKDKQSCDIKMEGMARNDLWYLSLEEVWKCRD QLDKYQENPERHLRQVAFTQKKVLTQERVSESG KYGGNCLLPAQLVLREYFHKRDSHTKSLKHDLV LNGHQDSCASNSNECGQTF CQNIHLIQFARTHTG DKSYKCPDNDNSLTHGSSLGISKGIHREKPYECK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ECGKFFSWRSNLTRHQLIHTGEKPYECKECGKSF SRSSHLIGHQKTHGTGEEPYECKECGKSFSWFSHL VTHQRTHTGDKLYTCNQCGKSF/VHSSRLIRHQR THTGEKPYECPECGKSFRQSTHLILHQRTHVRVR PYECNECGKSYSQSRSHLVVHHRHTGLKPFECKD CGKCFSRSSHLVSHQRTHTGEKPYECHDCGKSFS QSSALIVHQRIHTGEKPYECCQCGKAFIRKNDLIK HQRIHVGEETYKCNQCGIIFSQNSPFIVHQAHTG EQFLT CNQCGTALVNTSNLIGYQTNHIRENAY
3768	A	185	2258	SIHKMSRKISKESKKVNISSSLESDISLETTVPTD DISSEEREKGVRITRQLIERKELLHNIQLLKIELS QKTMIDNLKVDYLTKEBELEEKLNDAHQKQL LTLRLDNQLAFQKQDASKYQELMKQEMETILLR QKQLEETNLQLEKAGDVRRSLRDFELTEEQYIK LKAFPEDQLSIPEYVSVRFYELVNPLRKEICELQV KKNILABELSTNKNQLKLTETYEEDRKNYSEV QRCQRLALELADTKQLIQQGDYRQENYDKVKS ERDALEQEVIELRRKHEILEASHMIQTKERSELSK EVTLEQTVILLQKDKEYLNRQNMELSVRCAHE EDRLERLQAQLEESKKAREEMYEKYVASRDHY KTEYENKLHDELEQIRLKTNQEIDQLRNASREMY ERENRNLREARDNAVAEKERAVMAEKDALEKH DQLLDYREVLQLSTESKVTEFLHQSKLKSFESE RVQLLEETARNLTQCQLECEKYQKKLEVLTK FYSLQASSEKRITELQANSEHQARLDIYEKLEK ELDEIIMQTAEIENEDEAERVLFSGYGANVPTT AKRRLKQSVHLARRVLQLEKQNSLI/LKRSQTSK GPSNTAFTRSLTEANSLNQTQPPYRYLIESVRQ RDSKIDSLTESIAQL/ERKDVSNLNKEKSALLQTN GKMALADLADQLLNHP
3769	A	3	2297	DAAEFRVVDAMEVDFKPEEYTFKILANILII EGNLKFVVQDGTPLIENGKVSIAEELSTKIDM VEKALLYRTVATGRDIIDKQHTQEASYGRDAF AKAIYERLFCWIVTRINDIIEVKNUTTHGKNTV IGVLDIYGFEIFDNNSEFCINYCNEKLQQLFIQL VLKQEQEFYQREGIPWKHIDYFNNQIIVDLVEQQ HKGIIAILDDACMNVGKVTDEMFLAALNSKLGK HAHFSSRKL CASDKILEFDRDFRIRHYAGDVVYS VIGFIDKNKDTLFDKRLMYNSSNPVLKNMWP EGKLSITEVTKRPLTAATLFKNSMIALVDNLASK BPYYVRCIKPNDKKSPQIFDDERCRHQVEYLGLL ENVRVRRAGFAFRQTYEKLHRYKMISEFTWPN HDLPSDKBAVKKLIERCGFQDDVAYGKTKIFIRT PRTLFTLEELRAQMLIRIVLFLQKVWRGTLARMR YKRTKAALTIIRYRRYKVKSYTHEVARRFHGK TMRDYGKHVKWSPPKVLRFFEEALQTIFNRWR ASQLIKSIPASDLQVRAKVA AVEMLKGQRADL GLQRAWEGNYLASKPDTPQTSQTFVPVANELKR KDKYMNVLFSCHVRKVNRFKVEDRAIFVTDRH LYKMDPTKQYKVMKTIPLYNLTGLSVSNGKDQL VVFHTKDNKDLIVCLFSKQPTHESTRIGELVGVLV NHFKSEKRHLQVNVNTPVQCSSLHGKKCTVSVE TRLNQPPDFTKNRSGFILSVPGN
3770	A	3	6276	HKVAAPDVVVPTLDTVRHEALLYTWLAEHKPL VLCGPPGSGKMTLFSALRALPDMEVVGLNFSS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				ATTPELLLKTDFHYCEYRRTPNGVVLPVQLGK WLVLFCDEINLPDMDKYGTQQRVISFIRQMVEHG GFYRTSDQTWVKLERIQFVGACNPPTDPGRKPLS HRFLRHVPVVYVDYPGPASLTQIYGTFNRAMLR LIPSLRTYAEPLTAAMVEFYTMSQERFTQDTQPH YIYSPREMTRWVRGIFEALRPLETLPEGLIRIWA HEALRLFQDRLVEDEERRWTDENIDTVALKHFP NIDREKAMSRPILYSNWLSDYIPVDQEELRDYV KARLKVFYEEELDVPLVLFNEVLHDVLRIDRIFR QPQGHLLIGVSGAGKTTLSRFVAVMNGLSVYQ IKVHRKYTGEDFDEDLRTVLRSSGCKNEKIAFIM DESNVLD SGFLERMNTLLANGEVPGLFEGDEYA TLMQTCKEQAQKEGLMLDSHEELYKWFTSQVIR NLHVVFMTNPSSEGLKDRAATSPALFNRCVLNW FGDWSTEALYQVGKEFTSKMDLEKPNYITVPDYM PVVYDKLPQPPSHREAIIVNSCVFVHQTTHQANA RLAKRGGRTMAITPRHYLDFINHYANLFHEKRSE LEEQMHNLVGLRKIKETVDQVEELRRDLRIKS QELEVKNAAANDKLKKMVKDQQAEEKKKVMS QEIQEQLHKQQEVIADKQMSVKEDLDKVEPAVI EAQNAVKSIIKQHLVEVRSMANPPAAVKLALES ICLLLGESTTDWKQIRSIIMRENFIPTIVNFSABEIS DAIREKMKKNYMSNPSSYNYEIVNRASLACGPMV KWAIAQLNYADMLKRVEPLRNELOKLEDDAKD NQKANEVEQMIRDLEASIARYKEEYAVLISEAQ AIKADLAAVEAKVNRSTALLKSLSAERERWEKT SETFKNQ MSTIAGDCLLSAAFIAYAGYFDQQMR QNLFTTWSHHLQANIQFRDIA RTEYLSNADER LRWQASSLPADDLCTENAIMLKRFRNYPLIDPS GQATEFIMNEYKDRKITRTSFLDDAFRKNLESAL RUCNPI LVQDVESYDPVL EEN E V P R T G C E E LITLGDQDID E E S V I F L S T R D P T V E F P P D L C A V T F V N F T V T R S S L Q S Q C L N E V L K A E R P D V D E K K S O L L K L Q G E F Q L R L R Q L E K S L L Q A L N E V K G R I L D D C T I T T L E N L K R E A A E V T R K V E E T D I V M Q E V E T V S Q Q Y L P L S T A C S S I Y F T M E S L K Q I H F L Y Q Y S L Q F F L D I Y H N V L Y E N P N L K G V T D H T Q R L S I I T K D L F Q V A F N R V A R G M L H Q D H I T F A M L L A R I K L K G T V G E P T Y D A E F Q H F L R G N E I V L S A G S T P R I Q G L T V E Q A E A V V R L S C L P A F K D L I A K V Q A D E Q F G I W L D S S S P E Q T V P Y L W S E E T P A T P I G Q A I H R L L L I Q A F R P D R L L A M A H M F V S T N L G E S F M S I M E Q P L D L T Q I V G T E V K P N T P V L M C S V P G Y D A S G H V E D L A A E Q N T Q I T S I A I G S A E G F N Q A D K A I N T A V K S G R W V M L K N V H L A P G W L M Q L E K K L H S L Q P H A C F R L F L T M E I N P K V P V N L L R A G R I F V F E P P P G V K A N M L R T F S S I P V S R I C K S P N E R A R L Y F L L A W F H A I Q E R L R Y A P L G W S K K Y E F G E S D L R S A C D T V D T W L D D T A K G R Q N I S P D K I P W S A L K T L M A Q S I Y G G R V D N E F D Q R L L N T F L E R L F T R S F D S E F K L A C K V D G H K D I Q M P D G I R R E E F V Q W V E L L P D T Q T P S W L G L P N N A E R V L L T T Q G V D M I S K M L K M Q M L E D E D D L A Y A E T E K K T R T D S T S D G R P A W M R T L H T T A S N W L H L I P Q T L S H L K R T V E N I K D P L F R F F E R E V K M G A K L L Q D V R Q D L A D V V Q V C E G K K K Q T N Y L R T L I N E L V K G I L P A R S W S H Y

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TVPAGMTVIQWGVPIARRIKQLQNISLAAASG GAKELKNIHVCLGGLFVPEAYITATRQYVAAQAN SWSLEELCLEVNVTTSSQGATLDACSFGVTGLKL QGATCANNKLSLSNAISTALPLTQLRWVKQTNT EKKASVVTLPVYLNFTADLIFTVDIEATKEDPR SFYERGVAVLCTE
3771	A	1	2043	LPLLHAGFNRRFMENSSIIACYNELIQIEHGEVRS QFKLRACNSVFTALDHCHEAIEITSDDHVIQYVN PAFERMMGYHKGELLGKELADLPKSDKNRADL LDTNTCIKKGKEWQGVYARRKSGDSIQQHVKI TPVIGQGGKIRHFVSLKKLCCTDNNKQIHKIHR DSGDNSQTEPHSFYKNNRKESIDVKSISRRGSDA PSLQNNRYPSMARIHSMTEAPITKVINIINAAQEN SPVTVAEALDRVLEILRTTEL YSPQLGTKDEDPH TSDLVGGMLMTDGLRRLSGNEYVFTKNVHQSHSH LAMPITINDVPPCISQLLDNEESWDFNIFELEAITH KRPLVYLGLKVFSRFGVCEFLNCSETTLRAWFQ VIEANYHSSNAYHNSTHAADV LHATAFFLGKER VKGSLDQLDEVAALIAATVHDVDHPGRTNSFL/C NAGSELAVLYNDTAVLESHTALAFQLTVKDT KICNIFKNID/RGNHYRTLRAIDMVLATEMTKH FEHVNFVNSINKPMAAEIEGSDCECNPAGKNFP ENQILIKRMMIKCADVANPCRPLDLCEWAGRIS EEYFAQTDEEKRQGLPVVMPVFDRTNCSIPKSI SFIDYFITDMFDAWDAFAHLPALMQHLADNYKH WKTLDLCKCKSLRLPSDRLKPSHRGGLLTDKGH CESQ
3772	A	1013	50	TLVHADGPPSLHITETCLAYREKRIGIDLVDHDTVE HELKEAEIIQGIMALLTRTLEEASEQIRMNRSK YNLEKDLKDKFVALTIDDICFSLNNNSPNIRYSEN AVKLEPNSVSI EDWLDFSSTNVFKADKORNNSL LKLALVDRLSQTANYLRKQCDVVHTEANGL KDTKDARDQLADHLAKVMEEIASQEKNTALEK AILDQEGPAKVAHTRLETRTHRPVVELCRDVAQ YRLMKEVQEITHNVARLKETLAQAQAEKGLH RRQLALQEEIQVKENTTYIDEVLCMQMRKSIPLR DGEDHGVWAGGLRPDAVC
3773	A	1	955	AAARESERQLRLRLCVLNEILGTERDYVGTLRFL QSAFLHRIRQNVADSVEKGLTEENVKVLFSNIEDI LEVHKDFLALEYCLHPEPQSQHELGNVFLKFK DKFCVYEEYCSNHEKALRLVELNKIPTVRAFL SCMLLGGRKTTDIPLEGYLSPIQRICKYPLLLKE LAKRTPGKHPDHPAVQSAQAMKTVCSNINETK RQMEKLEALEAAA/QSHIEGWEGSNLTDICTQLL LQGTLLKISAGNIQERAFFLDNLLVYCKRKS RV TGSKKSTKRTKSINGSLYIFRGRINTEVMEVENVE DGTGSPSPSLA
3774	A	4254	2061	ELQGD FSVDPVPSMAWCENSICVGFKR DYLI RVDGKGSIKELFPTGKQLEPLVAPLADGKVAVG QDDLTVVLNEEGICTQKCALNWTDPVAMEHQ PYIAVLPRYVEIRTFEPRLLVQSIELQRPFITSGG SNIIYVASNHFWRLIPVPMATQIQQLQDKQFE LALQLAEMKDDSDSEKQQQIHHIKNLYAFNLFC QKRFD ESMQVFAKLGTDPHVMGLYPDLLPTDY RKQLQYPNPLPVLSGALEKAHLALIDYLTQKRS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QLVKKLNDSDHQSSSTPLMEGTPTIKSKKKLLQII DTLLKCYLHTNVALVAPLLRLNNHCHIEESEH VLKKAHKYSELILYEKKGLHEKALQVLVDQSK KANSPLKGHERTVQYLQHLGTENLHLIFSYSVW VLRDFPEDGLKIFTEDLPEVESLPRDRVLGFLIEN FKGLAIPYLEHHHVWEETGSRFHNCLIQLYCEKV QGLMKEYLLSFPAGKTPVPAGEEEGELGEYRQK LLMFLEISSYDPRGLICDFPFDGLLEERALLGR MGKHEQALFIYVHILKDTMRMAEEYCHKHYDRN KDGNDVYLSLLRMYLSPPSIHCLGPIKLELLEPK ANLQAALQVLELHHSKLDTTKALNLLPANTQIN DIRIFLEKVEENAQKKRFNQVLKNLLHAEFLRV QEBRLHQQVKCIITEEKVCMVCKKKIGNSAFAR YPNGVVVHYFCSKEVNPADT
3775	A	1832	839	MSRARGALCRACLALAAALALLPLPRAP APARTPAPAPRAPPSPRAPPSLRPDDVFIAVKTTT KNHGPRLRLRLRTWISRARQQTFITDGDDEPELE LQGGDRVINTNCSAVRTRQALCCKMSVEYDKFI ESGRKWFCHVDDDNVYNARSLHLLSSFSQSOD VYLGRPSLDHPIEATERVQGGRTVTTVKFWFAT GGAGFCLSRGLALKMSPWASLGSFMSTAEQVRL PDDCTVGIVVEGLLGARLLHSPLFHSHLENLQRL PPDTLLQQVTLSHGGPENPQNVNVAAGGFSLHQ DPTRFKSIHCLLYPDTDWCPROKQGAPTSR
3776	A	3	796	PRAKLGTRARNMAGQDAGCGRGDDYSEDEGD SSVSRAAVEVFGKLKDLNCPFLEGLYTEPKTIQE LLCSPSEYRLEILEWMCTRVWPSLQDRFSSLKGV PTEVKIQEMTKLGHELMMLCAPDDQELLKGCACA QKQLHFMDQLDITRSLTIGCSSCSLMEHFEDT REKNEALLGELFSSPHLQMLLNPECDFWPLDMQ PLNKO'DDWQWASASAE'EEFKI'EL'GQ BSAAKLHAL'EL'FAQHEQGAAGAANTS
3777	A	3	413	SEEDVIEGKTAVIEKRRKRSSAGVVED/IGGA NMLEGVGVNDINKALLAKRKRLEMYTKASLRIS QKIEHVWKTQQDQROKLNQEYSQQFLTFFQQW DLDMQKAEQEKEKLVGIMIRFIINQVSSRNGQPS LLL
3778	A	132	788	SRLPPPPPHLADGRAGARVPR SARLSRWVWQD WTHGPVIRPPAAARTMWVNPEEVLLANALWITE RANPYFILQRRKGHAGDGGGGGGLAGLLVGTLD VVDSSARVAPYRILYQTPDSL VYWTIACGGSR KEITEHWLEWLEQNLLQTL SIFENENDITTFVRGKI QGIIAEYNKINDVKEDDDTEKFKEAIVKFHRLFG MPEBEKLVNYSCSYWKG
3779	A	2	934	CKSCTLFPQNPNLPPSTRERPPGCKTVFVGGLPE NATEEIIQEVFEQCGDITAIRKSKKNFCHIRFAEEF MVDKAIYLSGYRMRLGSSTDKKDSGRLHVDFA QARDDFYEWCKQRMRAREERHRRKLEEDRLR PPSPPAIMHYSEHEAALLAEKLDKDSKFSEAMQ VLLSWIERGEVNRASANQFYSMVQSANS HVRL MNEKATHEQEMEEAKENFKNALTGILTQFEQIV AVFNASTRQKAWDHFSKAQRKNIDIWAKHSEE LRNAQSEQLMGIRREEEMEMSDDENCDSPTKKM RVDESALGAP
3780	A	1	2535	AAQAEREELAAGRMPGGGPQGAAPAAAGGGGVS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				HRAGSRDCLPPAACFRRRRLARRPGYMRSSSTGP GIGFLSPAVGTLFRFPGGVSGEESHSESRRARQC GLDSRGLLVRSPVSKSAAAPTVTSVRGTSAHFGI QLRGGTRLPDRLSWPCGPGSAGWQQEFAMDS SETLDASWEAACSDGARRVRAAGSLPSAELSSNS CSPGCGPEVPPTPPGSHSAFTSSFSIRLSLGSAGE RGEAEGCPPSREAESHCSQSPQEMGAKAASLDGP HEDPRCLSQPFSLLATRVSAADLAQAARNSSRPER DMHSLPDMDPGSSSSLDPSLAGCGGDGSSGSGD AHSWDTLLRKWEPVLRDCLLRNRRQMBVISLRL KLQKLQEDAVENDDYDKAETLQQRLEDLEQEKI SLHFQLPSRQPALSSFLGHLAAQVQAALRRGATQ QASGDDTHTPLRMEPRILLEPTAQDSLHVSITTRD WLLQEKQQLQKEIALQARMFVLEAKDQQLRRE IEEQEQQQLQWQCGDLTPLVGQLSLGQLQEVSKA LQDTLASAGQIPFHAEPETIRSLQERIKSLNLSLK EITTKVCMSEKFCSTLRKKVNDIETQLPALLEAK MHAISGNHFWTAKDLTEEIRSLTSDREGLEGLLS KLLVLSSRNVKKLGSVKEDYNRLRREVEHQETA YETSVKENTMKYMETLKNKLCSCCKPLLKGVW EADLEACRLLIQCLQLQEARGSLSVEDERQMDD LEGAAPPPIPRLHSEDKRKTPLKESYILSAELGEK CEDIGKKLLYLEDQLHTAIHSHDEDLIQSLRRELQ MVKETLQAMILQLQPAKEAGEREAAAASCMTAG VHEAQA
3781	A	3	995	GRRRAGPAHSARMYNMMETELKPPGPQQTSGG GGGNSTAAAAGGNQKNSPDRVKRPMNAFMVW SRGQRRKMAQENPKMHNSEISKRLGAEWKLLSE TEKRPFIDEAKRLRALHMKHEPDYKYRPRRKT TLMKKDKYTLPGGLLAPGGNSMASGVGVGAGL GAGVNGEMTSYAEENGVSNGSYSMKQDQI YPQINSLNAGAAQMOPMHRVDVSALQYNSM TSSQTYMNG/SRPTYSMSYSQQGTPGMAPGSMG SVVKSEAESPPVVTSSSHSRAPCQAGDLRDMIS MYLPGAEVFEPAPPSRLHMSQHYQSGPVPGTAI NGTLPLSHM
3782	A	1	2649	FRVPDSCPVVLSFTQLDPLDPRPESSTQEIGEELI NGVIYSISLRKVQLHHGGNKGQRWLGYESAL NLYETCKVRTVKAGTLEKLVEHLVPAFQGSCLS YVTIFLCTYRAFTTTQQVLDLLFKRYGRCDALTA SSRYGCILPYSDEDDGGPDQLKNAISSILGTWLD QYSEDFCQPPDFPCKQLVAYVQLNMPGSDLER RAHLLLAQLEHSEPIEAEPGEEDWALSPVPALK PTPELELALTPARAPSPVPAPAPEPEPAPTAPGSE LEVAPAPAPELQQAPEPAVGLESAPAPALELEPA PEQDPAPSQTLELEPAPAPVPSLQPSWSPVVAEN GLSEEKPHLLVFPPDLVAEQFTLMDAELFKKVVP YHCLGSIWSQRDKKGKEHLAPTIRATVTQFNSV ANCVITTCNGNRSTKAPDRARVVEHWIEVAREC RILKNFSSLYAILSALQNSIHLKKTWEDVSRDS FRIFQKLSEIFSDENNYSLSRELLIKEGTSKFATLE MNPKRAQKRPKETGIIQGTVPYLGTFITDLVML DTAMKDYLGRINFEKRRKEFEVIAQIKLLQSA CNNYSIAPDEQFGAWFRAVERLSETESYNLSCEL EPPSESASNTLRTKNTAIVKRWSRQAPSTELS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				TSGSSHSKSCDQLRCGPYLSSGDIADALSVHSAG SSSSDVEEINISFVPESPDGQEKFWESASQSSPET SGISSASSSTSSSSASTTPVAATRTHKRSVSGLCNS SSALPLYNQVGDCCIRVSLDVDNGNMYKSILV TSQDKAPAVIRKAMDKHNLEEEEPEDYELLQILS DDRKLKIPENANVFYAMNSTANYDFVLKKRTFT KGVVKVKGASSTLPRMKQKGLKIAKGIF
3783	A	3	869	RSGQGKVYGLIGRRRFQQMDVLEGLNLLITISGK RNKLRVYYLSWLRNKLHNDPEVEKKQGWTTV GDMEGCGHYRVVKYERIKFLVIALKSSVEVYAW APKPYHKFMAFKSFADLPHRPLLVDLTVEEQQR LKVIYTGSSAGFHAVDVDSGNSYDIYIPVHIQSQT PHAIIFLPNTDGMEMLLCYEDEGVVYNTYGRHK DVVLQWGEEMPTSVAYICSNQIMGWGEKAEIRS VETGHLDGVMHKRAQRLKFLCERNDKVFFASV RSGSSSQVYFMTLNRNCIMNW
3784	A	1213	457	LSPRQVDGLAGLQKGLSLSLYQFLMNGIRLGTY GLAEAGGYLHTAEGTHSPARSAAGAMAGVMG AYLGSPIMVKTHLQAQAASEIAVGHQYKHQ MFQALTEIGQKHGLVGLWRGALGGLPRVIVGSS TQLCTFSSTKDLSQWEIFPQSWKLALVAAMM SGIAVVLAMAPFDVACTRLYNQPHRCTGQGPLY RGILDALLQTARTEGIFGMKGIGASYFRLGPHIT LSLFFWDQLRSLYYTDTK
3785	A	193	813	RRRGRHSLCGGKMLAYCVQDATVVDVEKRRNP SKHYVYIINVTWSDSTSQTYYRRYASKFFDLQML LDKFPNESGQKDPKQRIIPFLPGKILFRRSHIRDV AVKRLKPIDEYCRALVRLPPHISQCDEVFRFFEAR PEDVNPPEQGPSPPDAVLPGVGNKGKQELKAG PNWPGRTHHVNCVTQKCLFVFHFKFSSSGNKE SK
3786	A	3785	1632	EFVGRAASTTVVTRWRMAADAGIRRVVPSDLY PLVLGFLRDNLSEVANKFAKATGATQQDANAS SLLDIYSFWLNRSKVPENELQANGPVAKKAKK KASSSDSEDSSEEEVQGPAAKKAAPAKRVGL PPGKAAAKASESSSSESSDDDEEDQKKQPVQ KGVKPAKAGQAPPKAKSSSDSDSSSEDEPP KNQKPKITPVTVKAQTKAPKPARA\APKIANGK AASSSSSSSSSSDDSEEEKAAATPKKTPVKKQV VAKAPVKAATTPTRKSSSEDSSSDEEEQKKPM KNKPGPYSSVPPSAPPPKSLGTQPPKKA\VEKQ QPVESSEDSSDESSESEEEKPPTKAVVSKATTK PPPAKKA\ESSSDSSSDSSEDEAPSKPAGTTK NSSNKP\AVTTKSPAVKPAAAPKQPVGGGQKLLT RKADSSSEBESSSEEEKTKKM\AVTTKPKATAK AALS\PAKQAPQGSRDSSSDSSSSEEEKTSK SAVKKKQKVAGGAAPSKPASAKKGA\ESSNSS SSDDSSSEEEKLGKGS\PRQAPKANGTSALTA QNGKAAKNSEEEEEKKAAVVVSKSGSLKKR KQNEAAKEAETPQAKKIKLQTPNTFPK\KKGEK RASSPFRRVREEEIEVDSRVADNSFDAKRGAAAGD WGERANQVLKFTKGKSF\REKTKKKRGSYRGG SISVQVNSIKFDSE
3787	A	3	5078	IPEG/RALSAEHTSSLVPSLHITTLGQEQAILSGAV PASPSTGTADFPSILTFLQPTENHASPSVPPEMPTL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				<p>PAEGSDGSPPATRDLLSSKVPNLLSTSWTFPRW KKDSVTAILGKNEEANVTIPLQAFPRKEVLSLHT VNGFVSDFTGVSSSIPITAPRTNPLPSGPPLPSILS IQATQTVFPSELLAFSSTKPEVYAAAVDHSGLPAS APKQVRASPSSMDVYDSLITGDMKKPATTDVFW SSLSAETGSLSTESIISGLQQQTNYDLNGHTISTTS WETHLAPTAPPNGLTSAADAISQDFKDTAGHS VTAEGFSIQDLVLGTSIEQPVQQSDMTMVGSHID LWPTSNNNHSRDFQTAEVAYYSPTTRHSVSHPO LQLPNQPAHPLLLTSPGPTSTGSLQEMLSDGTD GSEISSDINSSPERNASTPFQNLGYHSAEASSISTS VFPTSSRVLRASQHPKKWTADTVSSKVQPTAA AAVTLFLRKSSPPALSAALVAKGTSSSPLAVASG PAKSSSMTTLAKNVTKAASGPKRTPGAVHTAF PFTPTMYARTGHTTSTHTA/IARKHGHCLWPVV YNLP/PP/GKPQAMHTGLPNPTNLEMPRSTPRPL TVTAALTSITASVKATRLPPLRAENTDAVLPAAS AAVVTGKMASNLECMSSKLLVKTVLFLTQRR VQISESLKFSIAKGLTQALRKAFHONDVSAHVDI LEYSHNVTVGYYATKGLVYLPVAVIEMLGYY GVSNTADLKQHTPHLQSVAVLASPWNQPAQ YFQLKTVLQFVSQADNIQCKFAQTMEQRLQKA FQDAERKVLNTKSNLTQIVSTSNASQAVTLVYV VGNQSTFLNGTVASSLLSQLSAELVGFYLTYPPL TIAEPLEYPNLDISETTRDYWVITVLQGVNSLV GLHNQSFARVMEQRLAQLFMMSQQQGRFRKRA TTLGSYTVQMVKMQRVPGPKDPAELTYTLYN GKPLLGTAALKILSTIDSQRMALTLHHVLLQAD PVVKNPPNWLWIAAVLAPVAVTVIIIITAVLCR KNKNDFKPDTMINLPQRAKPVQGFYAKQHLG QQCADEFVPTQETVVLPLTADPQEDVACD GSTIKTAKSTETPDRSPSENGSVISNESGKPS RSPQNVMAQQKVTKEEARKRNPASDEEEGAV LFDNSSKVAAEPDFTSSGVSQLIAIKPTALPMVPP TSDRSQESSAVLNGEVNKKALKQKSDIEHYRNKL RLKAKRKGYDYDFPAVETSKGLTERKKMYEKAP KEMEHVLDPDSELCAPFTESKNRQQMKNSVYRS RQSLNSPSPGETEMDLLVTRERPRRGIRNSGYDT EPEIIBETNIDRVPEPRGYRSRQVKGHSETSTLSS QPSIDEVRQQMHMLLEEAFSLASAGHAGQSRHQ EAYGSAQHLPYSEVVTAPGTMTRPRAGVQWVP TYRPEMYQYSLPRPAYRFSQLPEMVMGSPPPVP PRTGPVAVASLRRSTSDIGSKTRMAESTGPEPAQ LHDSASFTQMSRGPVSVTQLDQSALNYSNGTVP AVFAIPAANRPGFTGYFIPTPSSYRNQAWMSYA GENELPSQWADSVPLPGYIEAYPRSRYPQSSPSRL PRQYSQPANLHPSLEQAPAPSTAASQQSLAENDP SDAPLTNISTAALVKAIREEVAKLAKKQTMDFEF QV</p>
3788	A	2	1737	<p>MKGLYTDAMKSDNVKDKDAKISFLQKVIDV VMVSGEPLAKPARIVAGHEPERTNELLQIGKC CLNKLSSDDAVRRVLAGEKGEVKGRASLTSRSQ ELDNKNVREEESRVHKNTEDRGDAEIKERSTSRD RKQKEELKEDRMPREKDKDKEKAKENGGRNHR EGERERAKARAPDNERQKDRGNRERDRDSEK</p>

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KETERKSEGGKEKERLRDRDRERDRDKGKDRDR RRVKNGEHSWDLDRNNREHDKPEKKSASSGE MSKKLSDGTFKDSKAETETEISTRASKSLTTKTS KRRSKNSVEGDSTSDAEGDAGPAGQDKSEVPET PEIPNELSSNIRRIIPRPGSARPPRVRQDSMEAL QMDRSGSGKTVSNVITESHNSDNEEDDQFVVEA APQLSEMSEIEMVTA VELEEEKHGGLVKKILET KKDYEKLQQSPKPGEKERSLFESA WKKEKDVS KEIEKLRTSIQTLCKSALPLGKIMDYIQEDVDAM QNELQM\YHSENRQHAEALQQEQRITDCAVEPL KAELA\VELEQLIKD\Q\QDKICAVKANILKNEEKIQ KMOVYSINLTSRR
3789	A	1	4369	MRTLGTCLATLAGLLTAAGETFSGGCLFDEPYS TCGYSQSEGGDDFNWEQVNTLTKPSTDPWMPSGS FMLVNASGRPEGQRAHLLLPQLKENDTHCIDFH YFVSSKSNPPGLLN\YVKVNNGLGNPIWNISG DPTRTWNRAELAISTFWPNFYQVIFEVITSGHQG YLAIDEVKVLGHPCTRTPHFLRIQNVENAGQFA TFQCSAIGRTVAGDRLWLQGIDVRDAPLKEIKVT SSRRFIASFNVVNTTKRDAGKYRCMVRTEGGVGI SNYAE\VVKEPPVPIAPPQLASVGATYLVWQLN ANSINGDGPVAREVEYCTASGSWNRQPV DSTS YKIGHLDPDTEYEISVLLTRPGEGGTGSPGALRT RTKCADPMRGPRKLEVVEVKSRQITRWEPPGY NVTRCHSYNLTVHYCYQVGGQEQVREEVSWDT ENSHPQHTITNLSPYTNVSVKLILMNPEGRKESQ ELIVQTDDELPGA\PTESIQQSTFEEKIFLQWREP TQTYGVITLYEITYKAVSSFDPEIDL SNQSGRVSK LGNETHFLFFGLYPGTTYSTIRASTAKGFGPAT NOFTTKISAPSPMPAYELETPLNQTDNTVTVMKLP AIIIRAD\VS\YQIVVEETERRTKITTEV\KCYP VEMIFQNASLLNSQ\VAALPADSLQAAQPFTIG DNKTYNGYWNTPLL\KSYET\FQAASRANGET KIDCVQVATKGAATPKP\VEPEKQTDHTVKIAG VIAGILLFVIFLGVVLVMKKRKLAAKKRKETMSS TRQEIDLWIGELNGPRSYAEQGTKLATRAFSMD THNLNGRSVSSPSSFTMKTNTLSTSVPSYYPDE THTMASDTSSLVQSHTYKKREPADV\PYQTGQLH PAIRVADLLQHITQMKCAEGYGFKEEYESFFEGQ SAPWDSAKK DENRMKNRYGNIIAYDHSRVRLQT IEGDTNSDYINGNYIDGYHRPNHYIATQGPMQET IYDFWRMVWHENTASIIMVTNLVEVGRVKCKK YWPDDTEIYKDIKVTLIETELLA EYVIRTFAVEKR GVHEIREIRQFHFTGWPDPHGVPYHATGLLGFVR QVKS\KSPPSAGPLVVHCSAGAGRTGCFIVIDIML DMAEREGVVDIYNCVRELRSRRVNMVQTEEQY VFHDAILEACL CGDTSVPASQVRSLYYDMNKLD PQTNSSQIKEEFRTLNMVTPTLRVEDCSIALLPN HEKNRCMDILPPDRCLPFLITIDGESSNYNAALM DSYKQPSAFIVTQHPLPNTVKDFWRLVLDYHCTS VVMLNDVDPAQLCPQYWPENGVHRHGPIQVEF VSADLEEDIISRIFRIYNAARPQDGYRMVQQFQFL GWPMYRDTPVSKRSFLKLIRQVDKWQEEYNGG EGRTVVHCLNGGGRSGTFCASIVCEMLRHQRTV DVFHAVKTLRNNKPNMVDLLDQYKFCYEVVALE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				YLNSG
3790	A	261	485	EEQTPLHIASRLGKTEIVQLLQHMAHPDAATTN GYTPLHISAREGQVDVVASVLLGRQGAHFSRLT KVRRTMS
3791	A	1	5874	LPPVTMSGKYIMEEHDSDQVWSIDELPSKQG YYLQGNLYRCVAEVGSFEHNLTTDLLNHLVVFQ KVFMKEVNEVIQKVSGGEQPIPLWNEHDGTADG DKPKILLYSLNLQFKGIQVTATTPSMRAVRFETG LIELELSNRLQTKASPGSSSYLKLFGKCQVDLNL ALGQIVKHQVVEEAGSDFHQVAYFKTRIGLRNA LREEISGSSDREAVLITLNRPIVYAQPVAFDRAVL FWLNYKAAAYDNWNEQRMALHKDIHMATKEVV DMLPGIQQTSAQAFGTPFLQLTVNDLGICLPITNT AQSNHTGDLDTGSALVLTISTLITACSSSLVSK GHFKNFCIRFADGFETSWDDWKPEIHGDLVMNA CVVPDGTVEVCSRTTGQAAAESSAGTWTLNVL WKMCGIDVHMDPNIGKRLNALGNLTTLTGEED IDDIADLNSVNIADLSDEDEVDTMSPTIHTAATDY RRQAASASQPGELRGRKIMKRIVDIRLNEQAKV IDDLKKGASEGTINQEIQRYYQLESVAVNDIR DVRKKLRRSSMRAASLKDKWGLSYKPSYSRSKS ISASGRPPLKRMERASSRVGETEELPEIRVDAASP GPRVTFNIDTFPEETELDLLSVTIEGPSHYSSNSE GSCSVFSSPKTPGFGSPGIPFQTEGRRDDLSSTS EDSEKDEKDEHERERFYIRKPSHTSRKKATGF AAVHQLFTERWPTTPVNRSLSGTATERNIDFELD IRVEIDSGKCVLHPTTLQEHDDISLRRSYDRSSR SLDQDSPSKKKKFQNTYASTTHLMTGKKVPSSL QTKPSDLETTVFYIPGVDVKLHYNSTLKTESP ASRGSSLPRTLKSKESKLYGMKDSATSPSPPI PST VQKNTLTLPPQPPPIPAAKGKSGCCVKTAK WVALQSLPEEMVISPCLLDFLE NYTAVSSQDEDMGHFEIPDPMEBSVTSLSVSSSTS AYSSFPVDVVVYVRVQPSQIKFSCLPVSRVECM KLPSLDLVFSSNRGELETGTTYPATLSPGGNA TQSGTKTSASKTGIPGSSGLGSPLGRSRHSSSQSD LTSSSSSSGLSFTACMSDFSLYVFHPYGAGKQIT AVSGLTPGSGGLGNVDEEPTSVTGRKDSLINLE FVKVSLSRIRSGGASFFESQSVSKSASKMDTTLI NISAVCDIGSASFKYDMRRLSEILAFPRAWYRRSI ARRLFGLDQTINLPTSGPGTPDSIEGVSQLHSPSS RKAYCKTWEQPSQSASFTHMPQSPNVFNEHMTN STMSPGTVGQSLKSPASIRSRVSDSSVPRRDSLS KTSTPFNKSNAASQGGTPWETLVVFAINLKQL NVQMNMSNVMGNTTWTTSGLKSQGRLSVGSNR DREISMSVGLGRSQLDSKGGVVGGTIDVNALEM VAHISEHPNQPSHKIQTMTGSTEARVDYMGSSIL MGIFSNADLKLQDEWKVNLNTLDSSITDKSEIF VHGDLDKWDIFQVMISRSTTDLIKIGMKLQEFFT QQFDTSKRALSTWGPVYPPLPKTMTSNLEKSSQE QLLDAAHHRHWPGVLKVVSCHISLFQIPLPEDG MQFGGMSMLHGNHMTLACFHGPNFRSKSWALF HLEEPNIAFWTEAQKIWEDGSSDHSTYIVQTLDF HLGHNTMVTKPCGALESPMATITKIRRRHENPP HGVASVKEWFNYVTATRNEELNLLRNVDANNT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ENSTTVKNSSLLSGFRGGSSYNHETETTFALPRM QLDFKSIHVQEPQEPSLQDASLKPKECSVVTEF TDHICVTMDAELIMFLHDLVSAYLKEKEKAIFPP RILSTRPGQKSPIIIHDDNSSDKDREDSITYTTVDW RDFMCNTWHLEPTLRLISWTGRKIDPVGVYILQ KLGFIHARTTTPKWLRQGVMDPLDKVLSVLIKK LGTALQDEKEKKGKDKEEH
3792	A	1	364	QNGSTPLHHAASKNRHEIALMLLEGGANPDGKD HYEATAKHQATAKGNFKMIHLLYYKASTIIQDT EGNTPPHLVCDARVEEAKLLVSQGA/SIYIENKEE KDP/LQVAKGALGLVLKRMVEG
3793	A	2	340	DIVPNPKMAPLGDEAPTEKVLTPELSEEBVSTR DDIQFHHFSSEEALQKVYFVAKEDPSSQEEAHT PEAPPPQPPSSERCLGEMKCTLVRGDSSPRQAE KSGPASRPAL
3794	A	421	158	SYWVGEDYTYKFFEVLIDPFHKAIRRNPDQTQWI SKAVYKHREMCGLTSTGRKSHGLEKDRMFPHAI GGSCRAA*RRRKTLLQPCYH
3795	A	24	592	GGMDSRVSGTTSNGETKPVYPVMEKKEEDGTLE RGHWNNKMEFVLSVAGEIIGLGNVWRFYLYCYK NGGGAFFIPYLVFLFTCGIPVFLLETALGQYTSQG GVTAWRKICPIFEGIGYASQMIVILLNVYYIVLA WALFYLFSSFTIDLPWGGCYHEWNTHECMFEQK TNGSLNGTSENATSPVIEFW
3796	A	3	592	KPASTYSTSQPSMAPLLPIRTLPLILILLALLSPGA ADFNISLLSGLLSPALTESLLVALPPCHLTGGNAT LMVRRANDSKVVTSSFVVPPCRGRRELVSVDSD GAGFTVTRL SAYQVTNLVPGTKFYISYLVKKGT ATESSREIPMFTLPRRNMESIGLGMARTGGMVVI TVLLSVAMFI/VLGFIIALALGSRK
3797	A	1	1556	ATRLRGSGS...GAYPN I...LPGVKPVFATVDGQEK...TKVTLDNGL RVASQNKFGQFCTVGILNSGSRYEAKYLSGIAH FLEKLAFSSTARFDSKDEILLTLEKHG...DCQTS RDTTMYAVSADSKGLDTVALLADVVLQPRLT DEEVEMTRMAVQFELEDNLNRPDPEPLITEMIHE AAYRENTVGLHRFCPTENVAKINREVLHSYLRN YYTPDRMVLAVGVGVEHEHLVDCARKYLLGVQP AWGSAAVDIRSVAQYTGGAIKLERDMSNVSL GPTPIPELTHIMVGLESCSFLEEDFIPFAVLNMMM GGGGSFSAGGPGKGMFSRLYLNVNLRHHWMYN ATSYHHSYEDTGLLCIHASADPRQVREMVEITK EFILMGGTVDTVELERAKTQLTSMMLMNLESRP VIFEDVGRQVLATRSRKLPHLCTLRNVKPEDV KRVASKMLRGKPAVAALGDLTDLPTYEHIQTAL SSKDGRLPRTYRLFR
3798	A	73	759	KRLVEAGVPRTFDGIVGEGGAQSRSCWPWGVTA QTPAFSADSLNCLKNCMSITMGSVRPSVEQFHKY LPWFLNDRPNIKCPKGGAAYSTSVNLTSDGQV LASRFMAYHKPLKNSQDYTEALRAARELANIT ADLRKVPGTDPAFEVFPYTTINVFYEQYLTILPEG LFMLSLCLVPTFAVSCLLLGLDLRSGLLNLLSIV MILVDTVGFMALWGISYNAVSLNVLVS
3799	A	73	759	KRLVEAGVPRTFDGIVGEGGAQSRSCWPWGVTA QTPAFSADSLNCLKNCMSITMGSVRPSVEQFHKY

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				LPWFLNDRPNKCPKGGGLAAYSTSVNLTSDGQV LASRFMAYHKPLKNSQDYTEALRAARELAANIT ADLRKVPGTDPFAFEVFPYITINVFYEQYLTLPEG LFMLSLLCLVPTFAVSCLLLGLDLRSGLLNLLSIV MILVDTVGFMALWGISYNAVSLINLVS
3800	A	250	1032	GIFRSLRVLFPLFSVGRPQFARSLSAAPQLSDTAD TMGFGDLKSPAGLQVLNDYLADKSYIEGYVPSQ ADVAVFEAVSSPPADLCHALRWYNHIKSYEKE KASLPGVKKALGKYGPADVEDTTGSGATDSKD DDDIDLFGSDDEESEEAERLREERLAQYESKKA KKPALVAKSSILLDVKPWDEETDMAKLEECVRS IQADGLVWGSSKLVPVGYGIKKLQIQCVVEDDK VGTDMLEEQITAFEDYVQSMDDAAFNKI
3801	A	155	656	SREMELVTFRDVAIEFSPPEWKCLDPAQQNLYR DVMLENYRNLVSLGFVISNPDVTCLEQIKEPCN LKIHETAAPPAICSPFSQDLSPVQGIEDSFHKLIL KRYEKCCHENLQLRKCKRVNECKVQKGVNNG VYQCLSTTQSKIFQCNTCVRVFSTSSHKNHK
3802	A	1	1428	VTVSPETHMDLTGCVTFEDIAIFYSQDEWGLLD EAQRLLYLEVMLENFALVASLGCGHGTEDEETP SDQNVSVGVSSQSKAGSSTQKTQSCMCVPVLKD ILHLADLPQKPYLVGECTNHHQHQQHHSACKS LKRDMDRASYVKCCLFCMSLKPFRKWEVGKDL PAMLRLLRSLVFPGGKKPGTITECGEDIRSQKSH YKSGECGKASRHKHTPVYHPRVYTGKKLYECSK CGKAFRGKYSLVQHQRVHTGERPWECNECGKF FSQTSHLNDHRRHTGERPYECSECGKLFQNSS LVDHQKIHTGARPYECSQCGKSFSQKATLVKHQ RVHTGERPYKCGECGNSFSQSAILNQHRRHTGA KPYECGQCGKSFSQKATLIKHQRVHTGERPYKC GDCGYSFSQSAILQHHTPVHTGARPYECGQC SOKSGLIQHVHTGERPYECGNSFSQSAIL IHHQKCHNT
3803	A	193	617	LPFLGSESKNGEADSSDKEMKHGQKSPTGKQTS QHLKRLKKSGLGHLKWTKAEDIDIETPGSILVNT NLRALINKHTFASLPQHFQYLLLLPEVDRQMG SDGILRLSTSALNNEFFAYAAQGWKQRLAEGKF VFSIIM
3804	A	197	479	SSSRASPPEHPSSQAHCGLPLVLSHACPEVTNKWS TGSSSPNSSWVSSPLQPEGLSGSSRMKGGSATKI LLETLLAAHMTADQGIASSQRCLL
3805	A	1	385	QSADTLFPGDINFNVSGLSAVTLQDQTVSDRLAS EELPSTAVPTPATTAPAPAPAPATAPALVSAAT KERTSEVPPRPASPKVTRSPPEAAPVEDMARR SELAVGQEGTEGGRGEGTGSPMSSY
3806	A	47	1033	LQGDTWHLSTLSHFSRLHGGVPGRGLLEGNNLQ PQAPGHDMTSIPFPGDRLLQVDGVLCGLTHKQA VQCLKGPQVARLVLERRVPRSTQQCPSANDSM GDERTAVSLVTALPGRPSSCVSVTDGPKF*SSN* KRIANGLGFSFVQMEKESCSHLKSDLVRIKRLFP GHPAEENGAIAGDIILGREWEGPRKASSRCRG SWAMQLSVQAGPSFASYYPAAVEVLHLLRGAPQ EVTLLLCRPPPGALPELEQEWQTPELSADKEFTR ATCTDSCTSPILGSRGQLGGTVPPQMKGAWGL RPSSQKAIREGTMGAKTERDLGPVP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
3807	A	656	1238	RCPSLLPPSWPLPTLQTLTRTPGNKAIAGGAGLW AVLWGSERTPPYR*GN*NQRGAVPCLRPRLRP QDKFLVLASDGLWDMLSNEDVRLVVGHLAEA DWHKTDLAQRPANLGLMQSLLQRKASGLHEA DQNAATRLIRHAIGNNEYGEMEAERLAAMLTL EDLARMYRDDITVTVVYFNSESIGAYYKGG
3808	A	26	2195	SQYSESVAGRQASPERLLGSYHAMASTVEGGDT ALLPEFPRGPLDAYRARASFSWKELALFTEGEG MLRFKKTIFSALENDPLFARSPGADLSLEKYREL NFLRCKRIFEYDFLSVEDMFKSPLKVPALIQCLG MYDSSLAAYLLHSLVFGSAVYSSGSRHLTYIQ KIFRMEIFGCFALTELSHGSNTKAIRTTAHYDPAT EEFIHSPDFEAAKFWVGNMGKTATHAVVFAKL CVPGDQCHGLHPFIVQIRDPKTLTPMPGVMVGGDI GKKLGQNGLDNGFAMFHKVRVPRQSLNRMGD VTPEGTYVSPFKDVRQRFASLGSLSGRVSIVSL AILNLKLAIAIALRFSATRRQFGPTEEEIPVLEY PMQQWRLLPYLAAYALDHFSKSLFLDLVELQR GLASGDRSARQELGREIHALASASKPLASWTT QQGIQECREACGGHGYLAMNRLGVLRDDNDPN CTYEGDNNILLQQTSNYLLGLLAHQVHDGACFR SPLKSVDFLDAYPGILDQKFEVSSVADCLDSAVA LAAWKWLVCYLLRETYQKLNQEKRSSSDFEAR NKCQVSHGRPLALAFVELTVVQRFHEHVHQPSV PPSLRAVLGRLSALYALWSLSRHAALLYRGGYF SGEQAGEVLESAYLALCSQLKDDAVALVDVIAP PDFVLDSPIGRADGELYKNLWGAVLQESKVLER ASWWPEFSVNKPVIGSLKSKL
3809	A	117	830	CFGIMERVGCITLTTTYAHPRPPTNPLPAISTMAS SYDRFPHSNI.THSLSLPWRPSTYYKVASNSPSV APYCTRSQRVWNTMIPFVSNPTTTPDDEW YRDELNYQESNTSRHNSKLRVTSKLDNDKYQ QTRKTQADTTQNLGERVNDIGFWYHEIHELDEM IGETNALTDVKKRLERALMETEAPLCVARECLF HREKRMGIDLHDEVEAQLLTVNVGEMEQSQA A
3810	A	3	518	VIQELGGSGADLGEHSCR PASQPRFRPAEARS HPATRRPASGPAMGKTNSKLAPVLEDLVQNT FSEQELKQWYKGLKDCPSGILNLEEFQQLYKF FPYGDASKFAQHAFRTFDKNGDGTIDFREFICAL SVTSRGSFEQKLNWAFEMYDLGDGGRITRLEML EIE
3811	A	81	1147	GCGYGC SGAGGAJGEPMAKWGEGDPRWIVEE RADATNVNNWHWTERDASNWSTDKLKTLFLAV QVQNEBGKCEVTEVSKLDGEASNNRKGKLIFFY EWSVKLNWTGTSKSGVQYKGHVEIPNLSDENSV DEVEISVSLAKDEPDNLVALMKEEGVKLLREA MGIYISTLKTEFTQGMILPTMNGESVDPVGPAL KTEERKAKPAPSKTQARPVGVIPTCKITLKETFL TSPEELYRVFTTQELVQAFTHAPATLEADRGKF HMDVGNVSGEFTDLVPEKHIVMKWRFKSWPEG HFATITLTFIDKNGETELCMEGRGIPAPEERTRQ GWQRYFFEGIKQTFGYGARLF
3812	A	20	558	PCGTAASTHAYDRRAKCRQQQQQQQNGGQNKV RPAKKKTSPAREVSSSESGTSGQFTPPSSSTVPTIAS

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SSAPVSIWSPASISPLSDPLSTSSSCMQRSYPMTYT QASGYSQGYAGSTSYFGGMDCGSYLTPMHQQL PGPGATLSPMGNTNAVTSHLNQSPASLSTQGYGAS KLWGFNFNH
3813	A	1	1016	CTEPPRRSTRTPAALASLRPYTDYVVVSDQILQES EDFFTLIESHEGKPLKLMVYNSKSDSCREVTVP NAAWGGEGSLGCGIGYGYLHRIPTQPPSYHKKPP GTPPPSALPLGAPPDALPPGPTPEDSPSLETGSRQ SDYMEALLQAPGSSMEDPLPGPGSPSHSAPDPDG LPHFMETPLQPPPPVQRVMDPGFLDVSGISLLDN SNASVWPSLPSSTELTTTAVSTSGPEDICSSSSHE RGGEATWSGSEFEVSFLDSPGAQAQADHLPQLT LPDSL TSAASPEDGLSAELLEAQAEPEASTEGLD TGTEAEGLDSQAQISTTE*HPGL*QGP
3814	A	2	884	VFWQVRNAGSSPLSAACPLFRTPAPQPCGSWGR CCIPHASTGCRPMAERGELDLTGAKQNTGVWL KVPKYLSSQWAKASGRGEVGLRIAKTQGRTE VSFTLNEDLANIHDIGGKPASVSAPREHPFVLQSV GGQTLTVFTSSSDKLSLEGIVVQRAECRPAASE NYMRLKRLQIESSKPVRLSQQLDKVVTNYKP VANHQYNIEYERKKKEDGKRARADKQHVLDML FSAFEKHQYYNLKDLDITKQPVVYLKEILKEIG VQNVKGHIKNTWELKPEYRHYQGEEKSD
3815	A	17	411	NIGDWEDIGKSPERIIQYYGPATWAQDGSRGYCT PIYMLNHIURLQAVLEIMNERANALDLLAQOTT MRNANYQNRLALDYLLAHEGGV*GKFSLTNCC LEIDDNGKAIMEITARMRKLHIPVQTWER
3816	A	3	1172	SHWQRRDRRCVRNMAERGRKRCGPGEGHQRI EWRKWKQKKEEKKKWKDLKLMKKLERQRAQ BEQAKRLEEEEAEEKEDRGRPYTL SVALPGSIL DNAQSPELNTYL AGQIAKACATC VDEFEDEE EQDAKTVEGEFTGVGKKGQACVQLA...EQYLEC PQYLKRAFFPKHQDLQFAGLLNPLDSPHMRQD EESEFREGVVVDRPTRPGHGSFVNCGMKKEVKI D...NLEPLRVTVRLNQQQHPDCKTYHGKVVS QDPRTKAGLYWGYTVRLASCLSAVFAEAPFDG YDLTIGTSESGSDVASAQLPNFRHALVFGGLQG LEAGADADPNLEVAEPSVLFDLVYVNTCPGQGS TIRTEEAILSLAALQPGLIQAGARHT
3817	A	246	1197	FLSAGMSNFTHYAYLLMESLMLGKVPHPVPSH HFIFHDDGSARQKGESDYKVIIQQWFSKSGPWTT SSNVTWGLLELQQSISESAVLTIPPGDSGAGSNLI TMFLRNKEDLCSGRSKVNRGWSGRCKQRG KTEQPGEPLEHVYVTIKHVALESRHQKGELQC LIKMCIPLSKPLQMFSPPHWEAWLQRVQQLAK NTRYFRQRLQEMGFITYGNENASVVPLLYMPG KVAAFARHMLEKKIGVVVVGFATPLAEARARF CVSAAHTREMLDTVLEALDEMGDLLQLKYSRH KKSARPELYDETSFELED
3818	A	215	789	NPQSSSSEGSSEIFQVNGHNRLLVQRSEVTQAPG QYTVDVEGHGCTFIQATLKYNVLLPKKASGFSLS LEIVKNYSSTAFDLTVTLKYTGIRNKSSMVVIDV KMLSGFTPTMSSIELENKGQVMKTEVKNDHVL FYLENVFGRA DSFTFSVEQSNLVFNIQAPGMVY DYEEKEEYALAFYHINSSSVSE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3819	A	1	1483	RIPDSIISRGVQGLPRDTASLSTTPSESRAQATSR LSTASCPITPKVQSRCSSKENILRASHSAVDITKVA RRHRMSPFPLTSMDFAFITVLEMTPTVLGTEINR DGMGRVLAQDVYAKDNLPFPASVKDGYAVRA ADGPGDRFTIGESQAGEQPTQTVMPPGQVMRVTT GAPIPCGADAVVQVEDTELIRESDDGTEELEVRIL VQARPGQDIRPIGHDIKRGECVLAKGTHMGPSI GLLATVGVTEVEVNKFPVAVMSTGNELLNPED DLLPGKIRDSNRSTLLATIQEHGYPTINLGIVGDN PDDLNLALNEGISRADVITSGGVSMGEKDYLKQ VLIDDLHAQIHFGFVFMKPGLPPTFATLIDIGVR KIIFALPGNPVSAVVTCNLFVVPALRKMQGILDP RPTIHKARLSCDVKLDPRPEYHRCILTWHHQEPLP WAQSTGNQMSSRLMSMRSANGLMLPPKTEQY VELHKGEVVDVMVIGRL
3820	A	2216	487	PQEPALKSEFSQVASNTIPLPLPQNTCKDNGPCK QVCSTVGGSAICSCFPGYAIMADGVSCEDQDECL MGAHDCSRRQFCVNTLGSFYCVNHTVLCADGYI LNAHRKCVDINECVTDLHTCSRGEHCVNTLGSF HCYKALTCEPGYALKDGECEVDDECAMGTHTC QPGFLCQNTKGSFYCQARQRCMDGFLQDPEGNC VDINECTSLSEPCRPFGFSCINTVGSYTCQRNPLIC ARGYHASDDGTKCVDVNECETGVHRCGEGQVC HNLPGSYRCDCKAGFQDAFGRGCIDVNECWAS PGRLCQHTCENTLGSYRCSASCGLLAADGKRC EDVNECEAQRCSEQECANTYGSYQCYCRQGYQLA EDGHTCTDIDECAQAGILCTFRCLNVPGSYQCA CPEQGYTMTANGRSCKDVDECALGTHNCSEAET CHNIQGSFRCLRFECPPNYVQVSKTKCERTTCHD FLECQNSPARITHYQLNFQTGLLVPAHIFRIGPAP AFTGDTIALNIKGNEEGYFGTRRLNAYTGVVYL QRAVLEPRDFALDVEMKLWRQGSVTTFLAKMHI FFTTFAL
3821	A	2216	487	PQEPALKSEFSQVASNTIPLPLPQNTCKDNGPCK QVCSTVGGSAICSCFPGYAIMADGVSCEDQDECL MGAHDCSRRQFCVNTLGSFYCVNHTVLCADGYI LNAHRKCVDINECVTDLHTCSRGEHCVNTLGSF HCYKALTCEPGYALKDGECEVDDECAMGTHTC QPGFLCQNTKGSFYCQARQRCMDGFLQDPEGNC VDINECTSLSEPCRPFGFSCINTVGSYTCQRNPLIC ARGYHASDDGTKCVDVNECETGVHRCGEGQVC HNLPGSYRCDCKAGFQDAFGRGCIDVNECWAS PGRLCQHTCENTLGSYRCSASCGLLAADGKRC EDVNECEAQRCSEQECANTYGSYQCYCRQGYQLA EDGHTCTDIDECAQAGILCTFRCLNVPGSYQCA CPEQGYTMTANGRSCKDVDECALGTHNCSEAET CHNIQGSFRCLRFECPPNYVQVSKTKCERTTCHD FLECQNSPARITHYQLNFQTGLLVPAHIFRIGPAP AFTGDTIALNIKGNEEGYFGTRRLNAYTGVVYL QRAVLEPRDFALDVEMKLWRQGSVTTFLAKMHI FFTTFAL
3822	A	2502	1540	MAAATRGCRPWGSLGLLGLVSAWWDLAS LRCTLGAFCECDFRPDLPGLECDLAQHLAQHL AKALVVKALKAFVRDPAPTKPLVLSLHGWTGTG KSYVSSLLAHYLFQGGRLSPRVHHFSPVLHFPHP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SHIERYKKDLKSWVQGNLTACGRSLFLFDEMMDK MPPGLMEVLRPFLGSSWVVYGTNYRKAIFIFISN TGGEQINQVALEAWRSRRDREEILLQLEPVISR AVLDNPHHGFNSNGIMEERLLDAVVPFLPLQRHH VRHCVLNELAQLGLEPRDEVVQAVLDSTTTFFPE DEQLFSSNGCKTVASRIAFFL
3823	A	1	3174	YGCEKTTEGRIPLKNIYRLFSADRKRVTALEAC SLPSSRNDSIPQEDFTPEVYRVFLNNLCPRPEIDNI FSEFGAKSKPYLTVQMMDFINLKQRDPRLNEIL YPPLKQEQVQVLEKYEPNNSLARKGQISVDGFM RYLSGEENGVSPEKLDLNEEDMSQPLSHYFINSS HNTYLTAGQLAGNSSVEMYRQVLLSGCRCVELD CWKGRTAEEEPVITHGFTMTTEISFKEVIEAIAEC AFKTSPPILLSFENHVDSPKQQAQMAEYCRIFG DALLMEPLEKYPLESGVPLPSPMDLMYKILVKN KKKSHKSSESGGKKKLEQASNTYSDSSSMFEP SPGAGEADTESDDDDDDDDCKKSSMDEGTAGSE AMATEEMSNLVNYIQPVKFESFEISKKRNKSFEM SSFVETKGLBQLTKSPVEFVEYNKMQLSRIYPKG TRVDSSNYMPQLFWNAGCQMVAlNFQTMDLA MQINMGMYEYNGKSGYRLKPEFMRRPDKHFD FTEGIVDGIANTLSVKIISGQFLSDKKVGTYVEV DMFGLPVDTRRKAFKTKTSQGNVNPVWEEPI VFKKVVLPTLACLRIVYEEGGKFIGHRILPVQAI RPGYHYICLRNERNQPLTLPVVFVYIEVKDYVPD TYADVIEALSNPIRYVNLMEQRAKQLAALTLEDE EEVKKEADPGETPSEAPSEARTTPAENGVNHTTT LTPKPPSQALHSQAPGSVKAPAKTEDLIQSVLTE VEAQTIIEELKQKSFVKLQKKHYKEMKDLVKR HHKKTDLIKEHTTKYNEIONDYLRRRAALEKS AKKDSKKKSEPPSPDHGSSIEQDI AALDAENTG KLIDLKDKQQQANLNRQEYQYSEKYQKRE LLIQKLTDAEBCQNNQLKKLEICEKEKKELK KMDKKRQEKITEAKSKDKSQMEBEKTEMIRSYI QEVVQYIKRLBEAQSKRQEKLEKHKEIRQQILD EKPKLQVELEQYQDKFKRLPLEILEFVQEAMKG KISED SNHGSAPLSLSSDPGKVNHKTPSSEELGGD IPGKEFDTP
3824	A	1	426	ILHWFVHRWVSGRNNREKIGVHVGFEEILNMEPY CCRETLKSLRPECFTYDLSAVVMHHGKGFGSGH YTAYCYNSEGGFWVHCNDSKLSMCTMDEVCKA QAYILFYTQRTVTEGH SKLLPPELLLSQHPNED ADTSSNEILS
3825	A	3	364	GIRAKFPNKIPVVVERYPRETFLPPLDKTKFLVPQ ELTMTQFLSIIRSRMVLRAEAFYLLVNNKSLVS MSATMAEITYRDYKDEDFVYMTYASQETFGCLE SAAPRDGSSLEDRLPLHL
3826	A	1	1237	PEKKFERECREAEKAQQSYERLDNDTNATKADV EKAKQQLNLRTHMADENKNEYAAQLQNFNGEQ HKHFYVVPQIYKQLQEMDERRTIKLECYRGFA DSERKVIPIISKCLEGMILAAKSVDERRDSQMVV DSFKSGFEPPGDFPFEDYSQHIYRTISDGTISASKQ ESGKMDAKTTVGAKGKLWLFGKKPKGPALED FSLPPEQRRKKLQQRIDELNRELQKESDQKDAL NKMKDVEKNPQMGDPGSLQPKLAETMNNIDR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				LRMEIHKNEAWLSEVEGKTGGRGDRRHSSDINH LVTQGRESPEGSYTDDANQSEVRGPPQHQHGHNE FDDEFEDDDPLPAIGHCKAIYFPDGHNEGTLAMK EGEVLVIIIEEDKGDGWTRARRQNGEEGYVPTS YI DVTLEKNSKGS
3827	A	2	1584	INPVSSAVNGEAHSSHETRGQNSNALPSVLELL SQSCLIPAMSSYLNRNDSVLDMARHVPLYRALLEL LRAIASCAAMVPLLLPLSTENGESEEEQSECQTS VGTLAKMKTCVDYTNRLRSKRENVKTGVKP DASDQEPEGLTLLVPDIQKTAIEIVYAATTSRQA NQEKKLGEYSKKAAMKPKPLSVLKSLEEKYVAV MKKLQFDTFEMVSEDEDGKLGFKNYHYMSQV KNANDANSAARARRLAQEA VTLSTSLPLSSSSSV FVRCDEERLDIMKVLITGPADTPYANGCFEFDVY FPQDYPPSPPLVNLETTGGHVSFRNPNLYNDGKV CLSILNTWHGRPEEKWNPQTSSFLQVLVSVQSLI LVAEPYFNEPGYERSRGTSGTQSSREYDGNIRQ ATVKWAMLEQIRNPSPCFKEVIHKHFYLRVEM AQCEEWIADIQYSSDKRVGRTMSHHAALKRH TAQLREELLKLPCEGLDPDTDDAPEVCRA TTGA EBTLMHDQVKPSSSKELPSPDFQL
3828	A	1415	845	PRVPATLVSLDPWHCFPTAGRLAGSTWVPACT LQLGPSSEHLDNHRAPLLSLPSQESLSFTPWYLV ACKPLFHIFCPLFACFMQEGKVQYFLHLHSHMRL LNYYFFPFLAPESLMQALEDLDYLAALDNDGNL SEFGIIMSEFPLDPQLSKSILASCEFDVDEVLTIA AMVTGILNDYSFSSFFANLH
3829	A	199	683	VDHTPVLSKPPQCFSSVKWGATLSARSQKTSGIGR LMVHVIEATELKACKPNGKSNPYCEISMGSQSYT TRTIQDTLNPKNWFNCQFFIKDLYQDVLCLTLFD NDQFSDDFLCSEVAKIEQEQKCPNTRRL LHEVPTGEVVRKDLQFEQKTL
3830	A	1747	404	RKMMEESGIEI PPGTPPPNPAGLAATAMSSTPV PLAATSSSFSPNVSESEFPPLAYSTPQPPLPPVRP SAPLPFVPPPAVPSVPLVTSMPPPVSPSTAAAFG NPPVSHFPSTAPNTLLPAPPSGPPISGFSVGSTY DITRGHAGRAPQTPLMPSFSAPSGTGLLPTPITQQ ASLTSLAQGTGTTSAITFPEEQEDPRITRGQDEAS AGGIWGFIKGVAGNPMVKSVDKTKHSVESMIT TLDPGMAPYIKSGGELDIVVTSNKEVKVA AVR AFQEVFGLAVVVGAEAGQSNIAQPVGYAAGLKG AQERIDSLRRTGVIHEKQTA VSVENFIAELL PDK WFDIGCLVVEDPVHGHILETFTQATPVPLEFVQQ AQLTPQDYNLRWSGLLVTVGEVLEKSLNVS TDWHMAFTGMSRRQMIYSAARAIAAGMYKQRLP PRTV
3831	A	5	674	FWTRSAWHEGLQOMKANDPSLQEVNLYNIKNIP IPTLREFAKALETNTHVKKFSLAATRSNDPVAIAF ADMLKVNTTLTSLNIBSHFITGTGILALVEALKEN DTLTEIKIDNQRQQLGTAVEMELAQMLENSRIL KFGYQFTKQGPRTVAAAITKNNDLA WQKDTQ EQTSIWQVVSQSIAGFNPQFEVQGNARSWMEE LGKAFHQFVRRELKQTEGKLP
3832	A	164	782	EPWVPMDVAESPERDPHSPEDDEEQPQGLSDDDDIL RDSGSDQDLDGAGVRASDLEDEESAARGPSQEE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				EDNHSDEEDRASEPKSQDQDSEVNELSRGPTSSP CEEEDGEGEDRTSDLRDEASSVTRELDEHLDY DEEVPEEPAPAVQDEAEKAGAEDDDEEKGEFTP REEGKAGVQSVGEKESLEAAKEKKKEDDDGEID DEEMY
3833	A	122	1676	SQPHFTQKMENKDTDSKKSEEYEDDFEKDLE WLINENEKSDASIIEMACEKEENINQDLKENETV MEHTKRHSDPDKSLQDEVSPRRNDIISVPGIQLD PISDSSENSFQESKLESQKDEEEDEEVRYYIM EKIVQANKLLQNQEPVNDKRERKLKFKDQLVDL EVPPLEDTTTSKNYFENERNMFGKLSQLCISNDF GQEDVLLSLTNGSCENKDRITLVERDGKFELN LQDIASQGFLLPPINNANSTENDPQQLPRSSNSSV SGTKKEDSTAKIHAVTHSSTGEPLAIIAQPLNR KTCPSAVNSDRSKGNKSNHRTQSAHISPTST YCLSPRQKELQKLEEKREKLKREERRKIEEEK EKKRENDIVFKAWLKKKREQVLEMRRIQRAKEI EDMNSRQENRDPQQAFLWLKKKHBEQMKERQ TEELRKQBECLFFLKGTEGRERAFKQWLRKRKM EKMAEQQAVRERTRQLRLEAKRSKQLQHHLYM SEAKPFRFTDHYN
3834	A	575	774	RSRTEELSNGSKLAMS KDLVTFGDVA VNFQS QEE WEWLNPAQRNLYRK VMLENYRSLVSLGKDMSP
3835	A	2	100	ASDFYLRYVGHKGKFGHEFLEFEFRPDGVVYV
3836	A	91	749	RPTPGHGFWMQPLTKDAGMSLSSVTLASALQV RGEALSEEEIWSLLFLAAEQLEDLRNDSSDYV CPWSALLSAAGSLSFQGRVSHIEAAPFKAPELLQ GQSEDEQPDASQMHVYSLGMTLYWSAGFHVPP HQPLQLCEPLHSILLTMCEDQPHRRCTLQSVLEA CRVHEKEVSVYPAPAGLHRRRLVGLVLTISEVS TEPCFSS3SCW3CVAIK
3837	A	3	1214	SLGCTNSARGKGQD...RTLMANGAPFTTDWFS KLRVSCGYIGDNCKNGADVNAKDMKMTALH WATERHHRDVVELLIKYGADVHAFSKFDKSAFD IALEKNNAEILVILQEAMQNQVNVNPERANPVD PVSMAAPFIFTSGEVVNLASLISSTNTKTTSGDPH ASTVQFSNSTTSVLATLAALAEASVPLSNHRAT ANTEBIEGNSVDSSIQQVMGSGGQRVITIVTDGV PLGNIQTSIPTGGIGHPPFIVTVQDGGQVLTVPAGK VAEETVIKEEEEEKLPLTKKPRIGEKTNSVEESKE GNERELLQQQLQEANRRAQEYRHQLLKKEQEAE QYRLKLEAIARQQPNGVDFTMVEEVAEVDVAVV VTEGELEERETKVTGSAGATGPPTRVSMATVSS
3838	A	1	1332	MIEDNKENKDHSLERGRASLIFSLKNEVGGLIKA LKIFQEKHVNLLHIESRKSRRNSEFEIFVDCDIN REQNDIFHLLKSHTNVLSVNLDPNFTLKEDGME TVPWFPPKISDLHDHCANRVLMYGSELDADHPGF KDNVYRKRRKYFADLAMNYKHGDPKVEFTEE BIKTWGTVFQELNKLYPHACREYLNPLLSKY CGYREDNIPQLEDVSNFLKERTGFSIRPVAGYLS RDFLSGLAFRVFHTQYVRHSSDPFYTPEDTCH ELLGHVPLLAEPFAQFSQEIQLASLGASEEAVQ KLATCYFFTVEFGLCKQDQQLRVFGAGLLSSISE LKHLSGHAKVKPFDPKITCKQECLITTFQDVYF VSESFEDAKEKMREFTKTIKRPFGVKYNPYTRSI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				QILKDTKSITSAMNELQHDLDVVSDALAKVSRKPSI
3839	A	3093	520	MVNFTVDQIRAIMDKKANIRNMSVIAHVDHGKSTLTDLSLVCKAGIIASARAGETRFTDTRKDEQERCITIKSTAIISLFYELSENDLNFIKQSKDGAGFLINLIDSPGHVDFSSEVTAALRVTDGALVVVDCVSGVCVQTETVLRQAIAERIKPVLMMNKMDRALLELQLEPEELYQTFQRIVENVNVIISTYGESES GPMGNIMIDPVLGTVGFGSGLHGWAF TLKQFAEMYVAKFAAKGEGQLGPAERAKKVEDMMKKLWGDYFDPANGKFSKSATSPEGKKLPRTFCQLILDPIFKVFDAIMNFKKEETAKLIEKLDIKL DSEDKDKEGKPLKAVMRRWLPAGDALLQMITIHLPSVTAQKYRCELLEYGPPDDEAAMGIKSCDPKGPLMMYISKMPVTSDKGRFYAFGRVFSGLVSTGLKVRIMGPNYTPGKKEDLYLKPIQRTILMMGRYVEPIEDVPCGNIVGLVGVDQFLVKTGTITTFEHAHNMRVMKFSVSPVVRVAVEAKNPADLPKLVEGLKRLAKSDPMVQCIIEESGEHIIAGAGELHLEICLKDLEEDHACIPKKS DPVVS YRET VSEESNVLC LSKSPNKHNR LYMKA RPPFDGLAEDIDKGEVSARQELKQRRARYLAEKY EWDVAEARKIWCFGPDGTGNILTDITKGVQYLNEIKDSVVAGFQWATKEGALCEENMRGVRFDVHDVTLHADAIHRGGGQIPTARRCLYASVLT AQPLMEPIYLVEIQCEQVVGGIYGV LNRKRGHVFEESQVAGTPMFVVKAYLPVNESFGFTADLRSNTGQAFPPQCVDHWQILPGDPFDNSSRPSQVVAETRKRKGLKEGIPALDNFLDKL
3840	A	2	753	SSTRSRDFCCSEATQGS LTRRERRASGVRRRSQSSAMASKILLNVQEEVTCPICLELLTEPLSLDCGHSICRACTIVSNKEAVTSM:30KSSCPVCGISYSFELQANQHLANIVERLKEVKLSPONGK...LDCDHGEKLLLFCKEDRKVICWLCERSQEHRRGHHTVLT EEFVKCEQEKLOAVLKRLKKEEEAEKLEADIRSEKTSWKYQVQTERQRIQTEFDQLRSILNNEBQRELQRL EEEBEKKT
3841	A	2	405	GKAFSCFTYLSQHRRTHMAEKPYECKTCKKAFSHFGNLKVHERIHTGEKPYECKECKRAFSWLTCLLRHERIHTGKKS YECCQCGKAFTRSRFLRGHEKTHTGEKMHECKECCGKALSSSLHRHKRTHWRDTL
3842	A	311	88	AVLKNMAPMTALGLLDLHILNLILFLSAGEDFTSVVSEIMMYILLVFLTLWLLIEMIYCYRKVSKAEAAQENA
3843	A	3	1175	APIRNSRIDDFVRRVESKATSARCGLWGS GPRRRPASGMFRGLSSWLGLQQPVAGGGQPNGDAPPEQPSETVAESAEEELQQAGDQELLHQAKDFGNLYLNFASAA TKKITESVAETAQTIKKSVEEGKIDGIIDKTIIGDFQKEQKKFVEEQHTKKSEAAVPPWVDTNDEETIQQILALSADKRNFLRDPAGVQFNFDQDQMPVALVMLQEDEL LSKMRFALVPKLVKEEVFWRNYFYRVSLIKQSAQLTALAAQQQAAGKEEKSNGREQDLPLAEAVRPKTPPVVKSQ LKTQEDEEISTSPGVSEFVSDAFDACNLNQEDLRKEMEQLVLDKKQEETA VLEEDSADWEKELQQELQEYEV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				VTSEKRDENWDKEIEKMLQEEN
3844	A	798	148	LPPAQIPEAWLLANVVVVLILVPLKDRLLDPLLL RCKLLPSALQKMALGMFFGFTSVIVAGVLEMER LHYIHHNETVSQQIGEVLYNAAPLSIWWQIPQYL LIGISEIFASIPGLEFAYSEAPRSMQGAIMGIFCLS GVGSLGSSLVALLSLPGGWLHCPKDFGNINNCR MDLYFFLLAGIQA VTALLFVWIAGRYERASQGP ASHSRFSRDRG
3845	A	3	1934	PEDSAPQYSRLFPNASQHITPSYNYAPNPDKH MRYTGPMKPIHMEFTNMLQRKRLQTLMSVDDS METTYNMLVETGELDNTYIVYTADHG YHIGQFG LVKGKSMPIYEFDIRVPFYVRGPVNEAGCLNPHIV LNIDLAPTILDIAGLDIPADMDGKSILKLLDTERP VNRFHLLKKKMRVWRDSFLVERGKLLHKRDNDK VDAQEENFLPKYQRVKDL CQRAEYQTACEQLG QKWQCVEDATGKLLKHKCKGPMRLGGSRALSN LVPKYYGQGSEACTCDSDGYKLSLAGRRKKLFK KKYKASYVRSRSIRSVAIEVDGRVYHVGLGDAA QPRNLTKRHWP GAPEDQDDKDGGDFSGTGGLP DYSAANPIKVTHRCYLENDTVQCDDL DYKSLQ AWKDHKLHIDHEIETLQNKIKNLREVRGHLKKK RPEECDCHKISYHTQHKGRLLKHRGSSLHPFRKGL QEKDKVWLLREQKRKKKLRKLLKRLQNNDTCS MPGLTCFTHDNQHWQTAPFWTLGPFCACTSAN NNTYWC MRTINETHNLFCEFATGFLEYFDLNT DPYQLMNAVNTLDRDVLNQLHVQLMELRSCKG YKQCNPRTRNMDLGLKDGGSYEQYRQFQRRKW PEMKRPSSKSLGQLWEGWEG
3846	A	3	1934	PEDSAPQYSRLFPNASQHITPSYNYAPNPDKH MRYTGPMKPIHMEFTNMLQRKRLQTLMSVDDS METTYNMLVETGELDNTYIVYTADHG YHIGQFG LVKGKSMPIYEFDIRVPFYVRGPVNEAGCLNPHIV LNIDLAPTILDIAGLDIPADMDGKSILKLLDTERP VNRFHLLKKKMRVWRDSFLVERGKLLHKRDNDK VDAQEENFLPKYQRVKDL CQRAEYQTACEQLG QKWQCVEDATGKLLKHKCKGPMRLGGSRALSN LVPKYYGQGSEACTCDSDGYKLSLAGRRKKLFK KKYKASYVRSRSIRSVAIEVDGRVYHVGLGDAA QPRNLTKRHWP GAPEDQDDKDGGDFSGTGGLP DYSAANPIKVTHRCYLENDTVQCDDL DYKSLQ AWKDHKLHIDHEIETLQNKIKNLREVRGHLKKK RPEECDCHKISYHTQHKGRLLKHRGSSLHPFRKGL QEKDKVWLLREQKRKKKLRKLLKRLQNNDTCS MPGLTCFTHDNQHWQTAPFWTLGPFCACTSAN NNTYWC MRTINETHNLFCEFATGFLEYFDLNT DPYQLMNAVNTLDRDVLNQLHVQLMELRSCKG YKQCNPRTRNMDLGLKDGGSYEQYRQFQRRKW PEMKRPSSKSLGQLWEGWEG
3847	A	1	1257	MVFSAVLTAFTGTSTNTTFVVYENTYMNITLPPP FQHPDLSPLLRYSFETMAPTGLSSLTVNSTAVPTT PAAFKSLNPLQITLSAIMIFILFVSFLGNLVVCLM VYQKAAMRSAINILLASLAFADMLLAVLNMPPA LVTLTTRWIFGKFFCRVSAMFFWLFVIEGVAILL IISDRFLIIVQRQDKLNPYRAKVLIASWATSFVCV AFPLAVGNPD LQIPSRAPQCVFGYTTNPGYQAYV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				ILISLISFFIPFLVILYSFMGILNTRLRHNLRIHSYPE GICLSQASKLGLMGLQRPFFQMSIDMGFKTRAFTT ILILFAVFIVCWAPFTTYSLVATFSKHFFYQHNF EISTWLLWLCYLKSALNPLIYYWRIKKFHDACLD MMPKSFKFLPQLPGHTKRRIRPSAVYVCGEHT VV
3848	A	3	2827	SSAVAARRRRSWASLVLAFLGVCLGITLAVDRS NFKTCBESSFCKRQRSIRPGLSPYRALLDSLQGP DSLTVHLIHEVTKVLLVLELQGLQKNMTRFRIDE LEPRRPRYRVPDVLVADPPIARLSVSGRDENSVE LTMAEGPYKIILTARPFRLDLEDRLLSVNARG LLEFEHQRAPRVSQGSKDPABGDGAQPEETPRD GDKPEETQGKAEKDEPGAWEETFKTHSDSKPYG PMSVGLDFSLPGMEHVYGIPEHADNLRKLVTEG GEPYRLYNLDVFQYELYNPMALYGSVPVLLAHN PHRDLGIFWLNAAETWVDISSNTAGKTLFGKMM DYLQGSGETPQTDVRWMSETGIIDVFLLLGPSISD VFRQYASLTGTQALPPLFSLGYHQSRWNYRDEA DVLEVDQGFDDHNLPCDVIWLDIEHADGKRYFT WDPSRFPQPRTMLERLASKRRKLVAIVDPHIKVD SGYRVHEELRNGLYVKTRDGSDEYGCWCPGS AGYPDFTNPTMRAWWANMFSYDNYEGSAPNLF VWNDMNEPSVFNGPEVTMLKDAQHYGGWEHR DVHNIYGLYVHMAADGLRQSRSGMERPFVLA RAFFAGSQRFQAVWTGDNTAEWDHLKISIPMCL SLGLVGLSFCGADVGGFFKNPEPELLVRWYQMG AYQPFRAHAHLDTGRREPWLLPSQHNDIIRDAL GQRYSLLPFWYTLTYQAHREGIPVMRPLVWQYP QDVTTFNIDDQYLLGDALLVHPVSDSGAHGVQV YLPQGQGEVWYDIQSYQKHHGPQTLVPLTLSSIP VFQREGTEVPRWMRVRESECMYDDPITTFVALS TQGTAAQGEFLDDGHTFNYQTRQEFLEKESFG NTLVSSADPEGHFETPIWIERVVIIAGKPAVV LQTKGSPESRSLSFQHPETSVLVLRKPGINVASD VSEHLR
3849	A	1	1717	RARNARGCWGVCRRSGFSSAVCGAARMEQVAEG ARVTAVPVSAADSTEELAEVEEGVGVVGEDNDA AARGAEAFGDSEEDGEDVFEVEKILDMKTEGGK VLYKVRWKGYTSDDDTWEPIHLEDCKEVLLEF RKKIAENKAKAVRKDIQRLSLNNDIFEANSDDQ QSETKEDTSPKKKKKKLRQREEKSPDDLKKKKA KAGKLKDKSKPDLESSLESVFDLRTKKRISEAK BELKESKKPKKDEVKETKELKKVKKGEIRDLKT KTREDPKENRKTKEKFVESQVESESSVLNDSPF PEDDSEGLHSDSREEKQNTKSARERAGQDMGLE HGFEKPLDSAMSAEEDTDVRGRKKKTPRKAED TRENKLENKNAFLEKKTVPKKQRNQDRSKSAA ELEKLMPVSAQTPKGRRLSGEERGLWSTDSABE DKETKRNESKKPKKDEVKETKELKKVKKGEIRD LKTKTREDPKENRKTKEKFVESQVESESSVLND SPFPEDDSEGLHSDSREEKQNTKSARERAGQDM GLEHGFEKPLDSAMSAEEDTDVRGRKKKTPRK AEDTRENKLENKNAFLEKKTVPKKQRNQDRSK SAAELEKLMPVSAQTPKGRRLSGEERGLWSTDS AEEDKETKRNESKKPKKDEVKETKELKKVKKGE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				IRDLKTKTREDPKENRRTKKEKFVESQVESESSV LNDSPFPED/RQ*RAFRQQREEKSPDDLKKKKA KAGKLKDKSKPDLESSLESVDFDLRTKKRISEAK EELKESKKPK
3850	A	1113	3975	PAAAAAAAAAAAAAAGRGPSFTPCFSPSLAVEPS RRTRLGSDPAQAMAGNVKKSAGGGSGSGGS GSGGLIGLMDAFQPHHHHHHLSHPHPPGTVDK KMVEKCKWKLMDKVVRCLQNPKLALKNSPPYIL DLLPDTYQHLRILSRYEGKMETLGENEYFRVF MENLMKKTQITISLFKEGKERMIEENSQPRRL TKLSLIFSHMLAELKGIFPSGLFQGDIFRITKADA AEFWRKAFGEKTIIVPWKSFRQALHEVHPISGLE AMALKSTIDLTCNDYISVFEDIFTRLFQPWSSLL RNWNSLAVTHPGYMAFLTIDEVKARLQKFHHP GSYIFRLSCTRLGQWAIGYVTADGNILQTIHPNKP LFQALIDGFREGFYLFDPGRNQNPDLTGLCEPT QDHIKVTQEYELCYCEMGSTFQLCKICAENDKD VKIEPCGHLMTSCLTSWQSESGQGCPCFRCCEIK GTEPIVDPFDPGRSGSLLRQGAEGAPSPNYDDD DDERADDTLMMKELAGAKVERPPSPFMAPQA SLPPVPPRLDLLPQRCVPSASALGTASKAASGS LHKDKPLVPPTLRDLPPPPPPDRPYSVGAESRPQ RRPLPCTPGDCPSRDKLPPVPSRLGDSWLPPIP KVPVSAPSSSDPWTGREL TNRHSLPFLPSQMEP RPDVPRLGSTFSLDTSMSMNSSPLVGPECDHPKI KPSSSANAIYSLAARPLVPKLPPGEQCEGEEDTE YMTPTSSRLRPLDTSQSSRACDCDQIDSCITYEA MYNIQSQAPOSITESSTFEGENLAAAHANTGPES ENEDDGYDVPKPPVPAVLARRTLSDISNASS/FG LFVLERDP*PONVTEGSQVPERPPKFPRRINSER KAGSCQQAASAATAISPOLSTENLMQC DYQDIQKLVIAQNNIEMANVILSVSISSPAH VAT
3851	A	2	2781	GRVGSMDGAMGPRGLLLCMYLVSLILQAMPA LGSATGRKSSEKQAVDTAVDGVIRSLKVCN KVTSRFAHYVVTSSQVNTANEAREVAFDLEIPK TAFISDFAVTADGNAFIGDIKDKVTAWKQYRKA AISGENAGLVRASGRTEQFTIHLTVNPQSKVTF QLTYEEVLKRNMQYEIVIKVKPKQLVHHFEIDV DIFEPQGISKLDAQASFLPKELAAQTIKKSFGKK GHVLFRTVSQQQSCPTCSTSLNGHFKVTYDVS RDKICDLLVANNHFAHFFAPQNLTMNKNVVFV IDISGSMRGQKVKQTKBALLKILGDMQPGDYFD LVLFGRVQSWKGSVQASEANLQAAQDFVRGF SLDEATNLNGGLLRGIEILNQVQESLPESLNHASI LIMLTDGDPTEGVTDRLQILKNVRNAIRGRFLY NLGFGHNVDNFLEVMSENNGRAQRIYEDHD ATQQLQGFSYQVAKPLLVDVDLQYPQDAVLALT QNHKKQYYEGSEIVVAGRIADNKQSSFKADVQA HGEGQEFSTCLVDEEEMKKLLRERGHMLENHV ERLWAYLTIQELLAKRMKVDREVRANLSSQALR MSLDYGFVTPLTSMIRGMADQDGLKPTIDKPSE DSPPLEMLGPRRTFVLSALQPSPTHSSNTQRLPD RVTGVDTDPHFIHVPQKEDTLCFNINEEPGVILS LVQDPNTGFSVNGQLIGNKARSPGQHDGTYFGR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LGIANPATDFQLEVTQNTITLNPFGGPPVFSWRD QAVLRQDGVVVTINKKRNLVVSVDGDTAEVVA LHRVWKGSSVHQDFLGLLMCWDKSGMSSPGR KGCWGGQFFHPIRFLKVS*HPPPGSDPQKAQMPT MVVRNPPGLTVTRGLQKDYSKDPWHGAEVSC WFIHNGA*ITDCAITDYIVPDI
3852	A	39	1735	TQVAEAGRGEGVVAEATGRPQSAGMNLLELES FGQNYPEEADGTLDCISMALCTFNWGTLLAV GCNDGRIVWDFLTRGIA*NKFSAHHPVCSLC WSRDGHKLVSASTDNIVSQWDVLSGDCDQRF PSPILKVQYHPRDQNKVLVCPMKSAVMLTSD SKHVVLVDDSDNLNVASFDRRGEYTYTGNK GKILVLKTDSDLVASFRTTGTSTNTAKSIEFA RKGSCFLINTADRIIRVYDGREILTCGRDGEPEPM QKLQDLVNRTPWKKCCPSGDGEYIVAGSARQH ALYIWEKSGNLVKILHGTGELLDDVAWHPVRP IIASISSGVVSIWAQNQVENWSAFAPDFKELDEN VEYEERESEFDIEDEDKSEPEQTGADAAEDBEVD VTSVDPIAAFCSSEDELEDSKALLYLPIAPEVEDP EENPYGPPDAVQTSMLDEGASSEKKRQSSADG SQPPKKPKKTINIELQGVNDEVHPLLGVKGDG KSKKKQAGRPKGSKGKEKDSPFKPKLYKGDRL PLEGSAKGKVAELSPLTAGGAISELL
3853	A	45	2603	PLLFTCGREVRARDPEKEGTIVVAGLKVQVQPRF LWILCFSMEETQGELTSSCGSKTMANVSLAFRDV SIDLSQEEWECLDAVQRDLYKDVMLNYSNLVS LDLEYKYITKNLLSEKNVCKIYLSQLQTGEKSKN TIHEDTIFRNLQCKHEFERQERHQMGCVSQMLI QKQISHPLHPKIHAREKSYECKEKRKAFRQVSYLI QHLRIHTGERPYKCMCEGKAFCRVGDLRVHHTI HAGERPYTCKECCGKAFLR... PYECKEKGK...RDLRVHQTIHAGERPYE... ECGKAFLRHYQLTEHQRIHTGERPYECKVCGKT FRVQRHISQHQIHTGVKPYKNECGKAFSHGS YLVOHQIHTGEKPYECKEKGKSFSAELARH RRIHTGEKPYECRECGKAFLQTELTRHRTHTG EKPYECKEKGKAFICGYQLTLHLRHTHTGEIPYC KECGKTFSSRYHLTQHYRIHTGEKPYICNECGKA FRLQGELTRHHRIHTCEKPYECKEKGKAFIHSNQ FISHQRIHTSESTYICKECGKIFSRRYNLTQHFKIH TGEKPYICNECGKAFRFQTELQHHRIHTGEKPY KCTECGKAFIRSTHLTQHRIHTGEKPYECTECG KTFSRHYHLTQHRGHTGEKPYICNECGNAFICS YRLTLHQRIHTGELPYECKEKGKTFSSRYHLTQH FRLHTGEKPYSCKECGNAFRLQAELTRHHIVHTG EKPYPCKEKGKAFSVNSELTRHHRIHTGEKPYQC KECGKAFIRSDQLTLHQKILVRNPMHNVKRIR WPLENAL*QRICNLNRLFLVTEHVGIPTSCSQFI RNYFVC
3854	A	108	894	LQSCWVPGPWPSPVSWLSWLKDLPSCEIHSASLS AVLQGPQCSEMLWPKNLTSWDDSSSVSSGISDTI DNLSTDDINTSSSISSYANTPASSRKNLDVQTD KHSQVERNLSWGGDDVKSDGGSDSGIKMEPGS KWRNRPSPDVSDSDKSTSGKKNPVISQTGSWRR GMTAQVGITMPRTKASAPAGALKTPGTGKRPG

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				SGGGAPTPAAPPQLARMAWAFSLSAASTPAVSP STSPSAVEGSPATILPLASSPPPTTP*LPLSELTV* RPQELVRGRGCLGPGAPTAPPPQLARMAWAFS LSAASTPAVSPSTSPSAVEGSPATILPLASSPPPT TP
3855	A	1	772	FRGGDGAPGVLPKPGNPLPFPLPPLQYPPSTLSHS DNLAMTSRSTARPNQGPQASKICQFKLVLLGESA VGKSSLVLRVKGQFHEYQESTIGAAFLTQSVCL DDITVKFEIWDTAGQERYHSLAPMYRGAQAAI VVYDITNQETFARAKTWVKELQRQASPSIVVGL AGNKADLANKRMVEYEEAQA YADDNLLFMET SAKTAMNVNDLFLAIA*EVAKRVNPQNLG/GVA AGRSRGVDLHEQS\QQNKSQCCSN
3856	A	2815	352	LGLEAAARPRPGGPAAMQDGNFLLSALQPEAGV CSLALPSDLQLDRRGAEQPEAERLRAARVQEQV RARLLQLGQQRHNGAAEPEPEAETARGTSRGQ YHTLQAGFSSRSQGLSGDKTSGRPIAKPAYSPA SWSSRSVVDLSCSRRLSSAHNGGSAFGAAGYGG AQPTPPMPTRPVSFHERGGVGSRADYDTLSRLSL RLGPGLDDRYSLVSEQLEPAATSTYRAFAYER QASSSSSRAGGLDWPEATEVSPSRTRAPAVRTL QRFQSSHRSGVGGAVPGAVLEPVARAPSVRSL LSLADSGHLDPVHGFSYSGHRTLQRLSSGFDDI DLPSAVKYLMA SDPNLQVLGAAYIQHKCYSDAA AKKQARSLQAVPRLVKLFNHANQEVQRHATGA MRNLIYDNADNKLALVEENGIFELLRTLREQDDE LRKNVTGILWNLSSSDHLKDRLAKKTPLEQLTD LGV*APLSGAGOPPLIQQNAEAEIFYNATGFPR NLSSASQATRQKMRECHGLVDALVTSINHALDA GKCEDKSVENAVCVLRNLSYRLYDEMPPSALQR FGRGRDLA GAPPGEVVGCTTFQERRLEPLA ALALTFAEVSKDPKGLEWLWSPQIVGLY*LLQ RCELNRHTTEAAAGALQNTGGDPRGPGLSRL AF EQERILNPLLDVRTADHHQLRSLTGLIRNLS RNARNKDEMSTKV VSHLNEKLPGSVGEKSPPAE VLVNNAVFNNLGLASPI/ALARDLLYFDGLRK LIFIKKRSDPDSEKSSRAASSLLANLWQYNKLH RDFRAKGYRKEDFLGP
3857	A	1034	204	VAVTLLSQLPSAIQRTAAWEMRAPLTFRVPLALD LIKPEHCTVNVDNSLSIPVIAELVVRKPSEKGM QKKKKTKDLGFRAGKESKTEWRK*GLQDMASQ MFALPLK*PVTAAFHDSSMPSSLLQIEMEQLFLE ARLO/PDSKSEARRNQCD SMLLRNQCLCSTCQE MKMVQPRMTKIPDDPKASFENCMSYRMSLHQP KFQTTPEPFHDDIPTENIHLQNL/PILGPRTAVFHG LLTEAYKTLKERQRSSLPRKEPIGKTTEAVSGRSS SPPRLPERK
3858	A	203	3469	SHQIEQNSAMAPRKRGGRGISFIFCCFRNNDHPE ITYRLRNDNSNFALQTMEPALPMPVVEELDVMFSE LVDELDLTDKHEAMFALPAEKKWQYCSKKK DQEENKATSWPEFYIDQLNSMAARKSLLALEK EEEEEESKTIESLKTALRTKPMRFVTRFDLDGLS CILNFLKTM DYETSESRIHTSLIGCIKALMNNNSQG RAHVLAHSESIN VIAQSLSTENIKTKVAVLEILGA VCLVPGGHKKVLQAMLHYQKYASERTRFQTLIN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				DLDKSTGRYRDEVSLKTAIMSFINAVLSQGAGVE SLDFRLHLRYEFLMLGIHPVMDKLRKHENSTLD RHLDDFFEMLRNEDELEFAKRFELVHIDTKSATQM FELTRKRLTHSEAYPHFMSILHHCLQMPYKRSGN TVQYWLLLDRIQQIIVIQNDKGQDPDSTPLENFI KNVVRMLVNENEVKQWKEQAEKMRKEHNEHQ QKLEKKERECDAKTQKEEMMQTLNKMKEKLE KETTEHKQVKQVADLTAQLHELSSRAVCASIP GGPSGAPGGPFPSSVPGSLPPPPPPPLPGGMLPP PPPLPPGGPPPPPPPPPLGAIMPPPGAPMGLALK KKSIPQPTNALKSFNWSKLPENKLEGTVWTEIDD TKVFKILDLEDLERTFSA YQRQQDFFVNSNSKQK EADAIDDTLSSKLKVKELSVIDGRRRAQNCNILLS RLKLSNDEIKRAILTMDEQEDLPKDMLEQLLKVF PEKSDIDLLEEHKHELDRMAKADRFLFEMSRIH YQRLQSLYFKKKFAERVAEVKPKVEAIRSGSEE VFRSGALKQLLEVVLAFGNMKNKGQRGNAYGF KISSLNKIADTKSSIDKNITLLHYLITTVENKYPSV LNLNEELRDIPQAAKVNMTELDKEISTLRSGLKA VETELVYQKSQPPQPGDKFVS VVSQFITVASFSFS DVEDLLAEAKDLFTKAVKHFGEEAGKIQPDFF GIFDQFLQAVSEAKQENENMRKKKEEEERRARM EAQLKEQRERERKMRKAKENSEESGEFDDL VSA LRSGEVFDKDL SKLKRNRKRITNQMTDSSRERPI TKLNF
3859	A	1279	141	RVEHLSEFLVDIKPSLTFDVIPLDPYGPAGSDPS LEFLVVSEETYRGGMAINRFRLENDLEELALYQI QLLKDLRHTENEEKVSSSSFRQRMGLNLLRPY ERPELPTCLYVIGLTGISGSGKSSIAQLKGLGAF VIDSDHLGHRA YAPGGPAYQPVVEAFGTDILHK DCENRVLGSRVFGNKKC REEMDKAVAEVCVIDAAVLEAGWQNEVH EVWTAVIPETEAVRRIVERDGLSEAAQSRLLQ MSGQQLVEQSHVVLSTCGSRISPNAWRKPGPS CRSAFPRLIRPSTEFKSVGPDWLELTSDPVVRRN GGLDAHPSGSGPEVQAILCRTWPGLVDTGSLPNTL VFGQH
3860	A	1	3881	MGQKSVGASYVQIPLVPPLSRHPKGLGHEDRWS SYCLSSLAQNICTSKLHCPAAPEHTDPSEPRGSV SCCSLLRGLSSGWSSPLLPAVCPNPNKAIFTVDA KTTEILVANDKACGLLGYSQDLIGQKLTQFFLR SDSDVVEALSEEHMEADGHAAVVFQTVVDIISRS GEKIPVSVWMKMRMQERRLCVVVLEPVERVST WVAFQSDGTVTSCDSLFAHLHGYSVSGEDVAGQ HITDLIPSVQLPPSGQHHPKLNKIQRVSVGRARDGT TFPLSLKLKSPSSEEATTGEAAPVSGYRASVWV FCTISGLITLLPDGTHGINHSFALTFLGYGKTELL GKNITFLIPGFYSYMDLAYNSSQLPDLASCLDV GNESGCGERTLDPWQGDPAEGGQDPRINVLVA GGHVVPREIRKLMESQDIFTGTQTELIAGGQLL SCLSPQAPAGVDNVPESLFPVHGEQALPKDQIT ALGREEPVAIESPGQDLLGESRSEPVDPKPFASCE DSEAPVPAEDGGSDAGMCGLCQKAQLERMVGS GPSGSDLWAGAAVAKPQAKGQLAGGSLLMHCP CYGSEWGLWWSQDLAPSPSGMAGLSFGTPTLD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				EPWLGVENDREELQTCLIKEQLSQLSLAGALDVP HAELVPTECQAVTAPVSSCDLGGRLCGGCTGS SSACYALATDLPGGLEAVEAQEVDVNSFSWNLK ELFFSDQTDQTSSNCSCATSELRETPSSSLAVGSDP DVGSLQEQGSCVLDDRELLLTGTCVDLGGQRR FRESCVGHDPTEPLEVCLVSSEHYAASDRESPGH VPSTLDAGPEDTCPSAEEPRLNVQVTSTPVIMR GAAGLQREIQEGAYSGSCYHRDGLRLSIQFEVRR VELQGPTPLFCCWLKDLLHSQRDSAARTLFL ASLPGSTHSTAELTGPSLVEVLRARPWFEEPPK AVELEGLAACEGEYSQKYSTMSPLGSGAFGFVW TAVDKEKNKEVVVKFIKKEKVLEDCWIEDPKLG KVTLEIAILSRVEHANIIVLDIFENQGGFFQLVME KHGSGLDLFAFIDRHPRLDEPLASYIFRQVRAG/Q SRLVSAVGYLRLKDIHRDIKDENVIAEDFTIKLI DFGSAAYLERGKLFYTFCGTIEYCAPEVLMGNPY RGPELEMWSLGVTLTYTLVFEENPFCELEETVEAA IHPPYLVSKEMLMSLVSGLLQVPERRTTLEKLV DPWVTQPVNLADYTWEEVFRVVKPESGVLSAAS LEMGNRSLSDVAQAQELCGGPVPGAPNGQGCL HPGDPRLTS
3861	A	1	3881	MGQKSVGASYVQIPLVPPLSRHPKGLGHEDRWS SYCLSSLAQNICTSKLHCPAAPEHTDPSEPRGSV SCCSLLRGLSSGWSSPLLPAVPCNPKNKAIFTVDA KTTBILVANDKACGLLGYSSQDLIGQKLTQFFLR SDSDVVEALSEEHMEADGHAAVVFQTVVDIISRS GEKIPVSVWMKMRQERRLCCVVLEPVERVST WVAFQSDGTVTSCDSLFAHLHGYSVSGEDVAGQ HITDLIPSVQLPPSGQHHPKNLKIORSVGRARDGT TFPLSLKLKSQPSSEEATTGEAAPVSGYRASVWV FCTISGLTLPDGTIHGPHFALTLFGYGF TFL GKNITLCTYMDLAYNSSLQLPDLASCLDV GNESGCGEITLDPWQQQDPAEGGQDPRINVLA GGHVVPRLKXLMESQDIFTGTQTELIAGGQLL SCLSPQPAPGVINTEGSLPVHGEQALPKDQKIT ALGREEPVAIESPGQDLLGESRSEPVVKPFASCE DSEAPVPAEDGGSDAGMCGLCQKAQLERMGVS GPSGSDLWAGAAVAKPQAKGQLAGGSLLMHCP CYGSEWGLWWSQDLAPSPSGMAGLSFGTPTLD EPWLGVENDREELQTCLIKEQLSQLSLAGALDVP HAELVPTECQAVTAPVSSCDLGGRLCGGCTGS SSACYALATDLPGGLEAVEAQEVDVNSFSWNLK ELFFSDQTDQTSSNCSCATSELRETPSSSLAVGSDP DVGSLQEQGSCVLDDRELLLTGTCVDLGGQRR FRESCVGHDPTEPLEVCLVSSEHYAASDRESPGH VPSTLDAGPEDTCPSAEEPRLNVQVTSTPVIMR GAAGLQREIQEGAYSGSCYHRDGLRLSIQFEVRR VELQGPTPLFCCWLKDLLHSQRDSAARTLFL ASLPGSTHSTAELTGPSLVEVLRARPWFEEPPK AVELEGLAACEGEYSQKYSTMSPLGSGAFGFVW TAVDKEKNKEVVVKFIKKEKVLEDCWIEDPKLG KVTLEIAILSRVEHANIIVLDIFENQGGFFQLVME KHGSGLDLFAFIDRHPRLDEPLASYIFRQVRAG/Q SRLVSAVGYLRLKDIHRDIKDENVIAEDFTIKLI DFGSAAYLERGKLFYTFCGTIEYCAPEVLMGNPY

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RGPELEMWSLGVTLTYTLVFEENPFCELEETVEAA IHPPYLVSKEMLMSLVSGLLQVPERRTTLEKLV DPWVTQPVNLADYTWEEVFRVKNKPESGVLSAAS LEMGNRSLSDVAQAQELCGGPVPGAPNGQGCL HPGDPRLTTS
3862	A	399	2069	TMDRSKRNSIAGFPFPRVEARLEEFEGGGGGEGNV SQVGRVWPSSYRALISAFFRLTRLDDFTCEKIGSG FFSEVFKVRHRASGQVMALKMNTLSSNRANML KEVQLMNRLSHPNILRYINSGNLEQLLDSNLHLP WTVRVKLAYDIAVGLSYLHFKGIFHRDLTSKNC LIKRDENGYSVVADFGLAEKIPDVSMGSEKLA VVGSPFWMAPEVLRDEPYNEKADVFSYGILCEII ARIQADPDYLPRTENFGLDYDAFQHMVGDCPPD FLQLTFNCCNMDPKLRPSFVEIGKLTLEELSRLOE EEQERDRKLQPTARGLLEKAPGVKRLSSLDKIP HKSPCPRRTIWLRSQSDFSRKPPRTVSVLDPYY RPRDGAARTPKVNPFSARQDLMGGKIKFFDLPSK SVISLVFDLDAPGPGTMPLADWQEPLAPPIRRWR SLPGSPEFLHQEACPFVGREESLSDGPPRLSSLK YRVKEIPFFRASALPAAQAHEAMDCSILQEENG GSRPQGTSPCPAGASEEMEVEERPA GSTPATFSTS GIGLQTQKQDG
3863	A	399	2069	TMDRSKRNSIAGFPFPRVEARLEEFEGGGGGEGNV SQVGRVWPSSYRALISAFFRLTRLDDFTCEKIGSG FFSEVFKVRHRASGQVMALKMNTLSSNRANML KEVQLMNRLSHPNILRYINSGNLEQLLDSNLHLP WTVRVKLAYDIAVGLSYLHFKGIFHRDLTSKNC LIKRDENGYSVVADFGLAEKIPDVSMGSEKLA VVGSPFWMAPEVLRDEPYNEKADVFSYGILCEII ARIQADPDYLPRTENFGLDYDAFQHMVGDCPPD FLQLTFNCCNMDPKLRPSFVEIGKLTLEELSRLOE EEQERDRKLQPTARGLLEKAPGVKRLSSLDKIP HKSPCPRRTIWLRSQSDFSRKPPRTVSVLDPYY RPRDGAARTPKVNPFSARQDLMGGKIKFFDLPSK SVISLVFDLDAPGPGTMPLADWQEPLAPPIRRWR SLPGSPEFLHQEACPFVGREESLSDGPPRLSSLK YRVKEIPFFRASALPAAQAHEAMDCSILQEENG GSRPQGTSPCPAGASEEMEVEERPA GSTPATFSTS GIGLQTQKQDG
3864	A	3	911	SWNMDSDSCAAAFHPEEYSPSCKRRRTVEDFNK FCTFVLAYAGYIPYKPEELPLRSSPSPANSTAGTI DSDGWDAGFSDIASSVPLPVSDRCFSLQPTLLQ RAKPSNFLDRKKTDKLLKKKKRKRSDAPGK EGYRGGLLKLEAADPYVETPTSPTLQDIPQAPSD PCSGWSDTPSSGSCATVSPDQVKEIKTEGKRTI VR/QEAQLMARNDGNFSSLESIFPS\DDDSWDLV TCFCMKPFAGRPMIECNECHTWIHLSCAKIRKSN VPEVFVCQKCRDSKFDIRRSNRSRTGSRKFLD
3865	A	3	3573	QERLRSRSPDRAAREAGSARGQPKRTERVEQ FLTIARRRGRRSMPVSLEDSGEPTSCPATDAETAS EGSVESASETRSGPQSASTAVKERPASSEKVKGG DDHDDTSDSDGLTLKELQNRLRRKREQEPT RPLKGIQSRLRKKRREEGAETVGSEASDTVEGV LPSKQEPENDQGVVSQAGKDDRESKLEGKAAQD IKDEEPGDLGRPKPECEGYDPNALYCICRQHNN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RFMICCDRCEEWFHGDVGISEARGRLERNGE DYICPNTILQVQDETHSETADQOEAKWRPGDA DGTDCSTIGTIEQKSSDQGIKGRIEKAANPSGKK KLKIFQPGPGPVPTQLPVLWQVLEIAVSRISAF LLHCISCKVIEAPGASKCIGPGCCHVAQPDSVYCS NDCILKHAAATMKFLSSGKEQKPKPEKMKMK PEKPSLPKCGAQAQGIKISSVHKRPAPEKKETTVK KAVVVVPARSEALGKEAACESSTPSWASDHNYNA VKPEKTAAPSPSLLYKSTKEDRRSEEKAAATAAS KKTAPPGSTVGKQPAPRNLVKKSSFANVAAAT PAIKKPPSGFGKTIPKRPWLSATPSSGASAARQAG PAPAAATAASKKFGSAALVGA VRKPVVPSVPM ASPAPGRLGAMSAAPSQPNQIRQNRSLKEIL WK/RFLFFILFRVNDSDDLMTENEVGKIALHIEK EMFNLQVTDN/RA YKSKYRSIMFNLKDPKNQG LFHRVLREEISLAKLVRKPEELVSKELSTWKER PARSVMESRTKLHNESKKTAPRQEAIPDLEDSP VSDSEEQESARAVPEKSTAPLLDVFSMLKDDTT SQHRAHLFDLNCICTGQVPSAEDEPAPKKQKLS ASVKKEDLKSKHDSSAPDPAPDSADEVMPEAVP EVASEPGLESASHPNVDRTYFPGPPGDGHPEPSPL EDLSPCPASCGSGVVTTTVTSGRDPRTAPSSCT AVASAASRPDSTMVEARQDVPKPVLTSVMVPK SILAKPSSSPDPRYLSVPPSPNISTSESRSPEGDDTT LFLSRLSTTWKGFNMQSVAKFVTKAYPVSGCFD YLSDELPTIHIGGRIAPKTVWDYVGKLKSSVSK ELCLIRFHPATEEEVAYISLYSYFSSRGRFGVVA NNNRHVKDLYLPLSAQDPVPSKLLPFEGPGKRR LSGWR
3866	A	2	3181	AQQPVGRRGGASGAGGGRRGTTPRAGAGPGF QVSSGGCRISTMRRLRPGHDEVERIFRNLFO FNKTVEHGFPHQPAALGYSPSLRLAIGTRSGAIK LYGAPGVFEMGLHQENNAVTQIHLPGQCQLVT LLDDNSLHLWSLKVKGGASELQEDSFILRGPP GAAPSATQITVVLPSSCELLYLGTESGNVVFVQ LPAFRALEDRTISSDAVLQRLPEEARHRRVFEMV EALQEHPRDPNQILIGYSRGLVVIWDLQGSRLY HFLSSQQLENTWWQRDGRLLVSHSDGSYCQWP VSSEAQQEPEPLRSLVPYGPFPCKAITRILWLTRQ GLPFTIFQGGMPRASYGDRHCISVIHDGQQTAFD FTSRVIGFTVLTEADPAATFDDPYALVVLAEEL VVIDLQTAGWPPVQLPYLASHCSAJTCSHHVSN IPLKLWERIJAAGSRQNAHPSTMWEPIDGGTSLTP APPQRDLLLLTGHDGTVRFDASGVCLRLLYKL STVRVFLTDTDPNENLSAQGEDEWPPLRKVGSF DPYSDDPRLGIQKIFLCKYSGYLA VAGTAGQVLV LELNDEAAEQAVEQVEADLLQDQEGYRWKGHE RLAARSGPVRFEFGFQPFVLVQCQPPAVVTSAL HSEWRLVAFGTSHGFLFDHQRRQVFVKCTH PSDQLALEGPLSRVKSLSKSLRQSFRMRRSRV SRKRHPAGPPGEAQEGSAKAERPGLQNMELAPV QRKIEARSAEDSFTGFVRTLYFADTYLKDSSRHC PSLWAGTNGGTIYAFSLRVPPAERRMDEPVRAE QAKEIQLMHRAPVVGILVLDGHSVPLPEPLEVAH DLSKSPDMQGGSHQLLVVSEEQFKVFTLPKVS AK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LKLKLTALEGSRVRRVSVAHFGSRRRAEDYGEHH LAVLTNLGDIQVVSLPLLKPQVRYSCIRREDVSGI ASCVFTKYGGFYLLISPSEFERFSLSTKGLVEPRC LVDSAETKNHRPGNGAGPKKAPSRARNSTQSD GEEKQPLVMERALLSDERAATGVHIEPPWGA ASAMAEQSEWLSVQAAR
3867	A	2	3181	AQQPVGRRGGASGAGGGRRGTTPRAGAGPGF QVSSGGCRLSKMRRFLRPGHDPVRERLKRDLFQ FNKTVEHGFPHPQSALGYSPSLRILAIOTRSGAIK LYGAPGVFEMGLHQENNAVTQIHLLPGQCQLVT LLDDNSLHLWSLKVKGGASELQEDSFLLRGPP GAAPSATQITVVLPHSSCELLYLGTESGNVFFVQ LPAFRALEDRTISSDAVLQRLPEARHRRVFEMV EALQEHPRDPNQILIGYSRGLVVIWDLQGSRLVY HFLSSQQLENIWWQRDGRLLVSHSDGSYCQWP VSSEAQQPEPLRSLVPYGPFPCKAITRILWLTRQ GLPFTIFQGGMPRASYGDRHCISVIHDGQQTAFD FTSRVIGFTVLTEADPAATFDDPYALVVLAEEL VVIDLQTAGWPPVQLPYLASLHCSAITCSHHVSN IPLKLWERIJAAGSRQNAHFSTMEWPIDGGTSLTP APPQRDLLLTGHEDGTVRFWDASGVCLRLLYKL STVRVFLTDTPNENLSAQGEDEWPPLRKVGFSF DPYSDDPRLGIQKIFLCKYSGYLAVAGTAGQVLV LELNDEAAEQAVEQVEADLLQDQEGYRWKGHE RLAARSGPVRFEFGFQPFVLVQCQPPAVVTSAL HSEWRLVAFGTSHGFGLFDHQRRQVFVKCTH PSDQLALEGPLSRVKSLLKSLRQSFRRMRRSRVS SRKRHPAGPPGEAQEGSAKAERPGLQNMELAPV QRKIEARSAEDSFTGFVRTLYFADTYLKDSSRHC PSLWAGTNGGTTYAFSLRVPPAERRMDEPVRAE QAKKQLMHRAPVVGHTDGHSVNPFPEVVAH DLKSPDMQGSQHLVVCMLDFKVFILPKVSAK LKLKLTALEGSRVRRVSVAHFGSRRRAEDYGEHH LAVLTNLGDIQVVSLPLLKPQVRYSCIRREDVSGI ASCVFTKYGGFYLLISPSEFERFSLSTKGLVEPRC LVDSAETKNHRPGNGAGPKKAPSRARNSTQSD GEEKQPLVMERALLSDERAATGVHIEPPWGA ASAMAEQSEWLSVQAAR
3868	A	1	2497	GDGGGLVCEBPSGRFFLAGIVSWGICAEARRP GVYARVTRLRDWLEATTKASMLAPTMAPAPA APSTAWPTSPESPVVSTPTKSMQALSTVPLDWVT VPKLQECGARPAMEKPTRVVGFGAASGEVPW QVSLKEGSRHFCGATVVGDRWLLSAAHCFNHT KVEQVRAHLGTASLLGLGGSPVKIGLRRVVLHP LYNPGILDFDLAVLELASPLAFNKYIQPVCLPLAI QKFPVGRKCMISGWGNTQEGNATKPELLQKASV GIIDQKTCVLYNFSLTDRMICAGFLEGKVDSCQ VSGIKALYESELADARRVLDETARERARLQIEIG KLRAELDEVNKSARKKREGELTVAQGRVKDLESL FHRSEVELAAALSDDRGLSDVAELRAQLAKAE DGHAVAKKQLEKETLMRVDLENRCQSLQEELDF RKSVEEEVRETRRRHERRLVEVDSSRQOEYDFK MAQALEELRSQHDEQVRLYKLELEQTYQAKLDS AKLSSDQNDKAASAAREELKEARMRLESLSYQL SGLQKQASAAEDRIRELEEAMAGERDKFRKMLD

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=-possible nucleotide deletion, \=possible nucleotide insertion)
				AKEQEMTEMRDVMQQQLAEYQELLDVVKLALD MEINAYRKLLGEEERLKLSPSPSSRVTVSRATSS SSGSLSATGRLGRSKRKR\WRWRSPW\QRPKRP HGHGWQRWLPPGAPGLGLGQRHIEIDLEGKFV QLKNNSDKDQSLGNWRIKRVLEGEIEAYKFTP KYILRAGQMVTVWAAAGAGVAHSPSTLVWKGQ SSWGTEGESFRTVLVNADGEEVAMRTVKKSSVM RENENGEEEEEEAEFGEDLFHQQQGDPRTTSRGC YVM
3869	A	1	1942	RYRAGIPGDGRKDYIRLTRPGLTLPGRAMFARG RRRRSGRAPPEAEDPDRGQPCNSCREQCPGFLH GWRKICQHCKCPREEHAVHVPVDLERIMCRLIS DFQRHSISDDSGCASEEYAWVPPGLKPEQVYQ FFSCLPEDKVYPVNSPGEKYRIKQLLHQLPPHDS EAQYCTALIEEEKKELRAFSQQRKRENLG/RLG IVRIFPVITVGAANCECGKQIGGGDIQVAFASRL GLLGQPSCFVCTTCQELLVDLIYFYHVGVKYYC GRHHAECRLPRCQACDEIIFPECTEAEGRHWHM DHFCCFECEASLGGQRYVMRQSRPHCCACYEAR HAEYCDGCGEHIGLDQGGQMA YEGQHWASDRC FCCSRCGRALLGRPFLPRRGLIFCSRACSLGSEPT APGPSRRSWSAGPVTA PLAASTASFSA VKGASET TTKGTSTELAPATGPEEPSRFLRGAPHRHSMPEL GLRSVPEPPPESPGQPNLRPDDSAFGRQSTPRVSF RDPLVSEGPRRTLSAPPAQRRRPRSPPPRAPSRR RHHHHNNHHHHNNRHPSSRRRHYQCDAGSGSDSE SCSSSPSSSSSESSEDDGFFLGERIPLPPLCRPMP AQDTAMETFNPSLSLPRDSRAGMMPRQARDKNC IVA
3870	A	2	3485	FVWRVIFYVHASCMPPRARSWEGAHAPVGMHV AEAHACSSOQQQMPPAQFWMLEWLLHLCAFLS TSPFPHWCCSNPHGSIADKPEEIVI...SRAAE NMAVEPRVATIKQRPSSRCFPAGSDMNSVYERQ GIAVMTPTVPGSPKAPFLGIPRGTMRRQKSIDSRI FLSGITEBERQFLAPPMLKFTRSLMPDTSIEDIPP PQSVPPSPPPSPPTTYNCPKSPTPRVYGTIKPAFNQ NSAAKVSPATRSDTVATMMREKGMVFRRELD YSLDSEDLYSRAGPQANFRNKRGMQMPENPYSE VGKIASKAVYVPAKPARRKGMVVKQSNVEDSPE KTCSEIPTIIVKEPSTSSSGKSSQGSSMEIDPQAPE PPSQLRPDES LTVSSPFAAAIAGAVRDREKRLEA RRNSPAFLSADLGDEHVGLGPPAPRTRPSMFPEE GDFADEDSAEQLSSPMPSATPREPENHFVGGAEA SAPGEAGRPLNSTSKAQGPRESSPAVPSASSGTAG PGNYVHPLTGRLLDPSSPLALALSARDRAMKES QQGPKGEAPKADLNKPLYIDTKMRPSLDAGFPT VTRQNTRGPLRRQETENKYETDLGRDRKGDDK KNMLIDIMDTSQQKSAGLLMVHTVDA TKLDNA LQEEDEKAEVEMKPDSSPSEVPEGVSETEGALQI SAAPEPTTVPGRTIVAVGSMEEAVILPFRIPPPPLA SVDLDEDFIFTEPLPPPLEFANSFDIPDDRAASVPA LSDLVKQKSDTPQSPSLNSSQPTNSADSKKPAS LSNCLPASFLPPESFDAVADSGIEEVDSSSSDH HLETTSTISTVSSISTLSSEGGENVDTCTVYADGQ AFMVDKPPVPPKPKMKPIHKSNALYQDALVEE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				DVDSFVIPPPAPPPPPGSAQPGMAKVLQPRTSKL WGDVTEIKSPILSGPKANVISELNSILQQMNREKL AKPGEGLDSPMGAKSASLAPRSPEIMSTISGTRST TVTFTVRPGTSQPITLQSRPPDYESTSGTRRAP PVVSPTEMNKETLPAPLSAATASPSPALSDVFSLP SQPPSGDLFGLNPAGRSRSPSPSILQQPISNKPFTT KPVHLWTKPDVADWLESLNLGEHKEAFMDNEI DGSHLPLNQLKEDLIDLGVTRVGHMRMNIERALKQ LLDR
3871	A	35	1171	VESRSAWHEGEDQIDRLDFIRNQMNLLTLDVKK KIKEVTEEVANKVSCAMTDEICRLSVLVDFCSE FHPNPDVLKIYKSELNKHIEDGMGRNLADRCTD EVNALVLQTQQEIIENLKPLLPAGIQDKLHTLIPC KKFDLSYNLNYHKLCSDFQEDIVFRFSLGWSSLV HRFLGPRNAQRVLLGLSEPIFQLPRSLASTPTAPT TPATPDNASQEELMITLVTGLASVTSRTSMGHIIV GGVIWKTIGWKLLSVSLTMYGALYLYERLSWTT HAKERAFFKQQFVNYATEKLRMIVSSTSANCCHQ VKQQAATTFARLCQQVDITQKQLEEEIARLPKEID QLEKIQNNSKLLRNKAVQLENELENTKQFLPSS NEES
3872	A	35	1171	VESRSAWHEGEDQIDRLDFIRNQMNLLTLDVKK KIKEVTEEVANKVSCAMTDEICRLSVLVDFCSE FHPNPDVLKIYKSELNKHIEDGMGRNLADRCTD EVNALVLQTQQEIIENLKPLLPAGIQDKLHTLIPC KKFDLSYNLNYHKLCSDFQEDIVFRFSLGWSSLV HRFLGPRNAQRVLLGLSEPIFQLPRSLASTPTAPT TPATPDNASQEELMITLVTGLASVTSRTSMGHIIV GGVIWKTIGWKLLSVSLTMYGALYLYERLSWTT HAKERAFFKQQFVNYATEKLRMIVSSTSANCCHQ VKQQAATTFARLCQQVDITQKQLEEEIARLPKEID QLEKIQNNSKLLRNKAVQLENELENTKQFLPSS NEES
3873	A	2944	2089	PVCTALTPGRMTDDKDVLRDVWFGRIFFETLY QDEITEREAEPPYLLPRVSYLTLVTDKVKKQ KVMRQEDISEIWFYEGTPLKWHYPIGLLFDLLA SSSALPWNITVHFKSFPKDLLHCPSKDAIEAHF MSCMKEADALKHKSQVINEMQKDKHKLWMG LQNDRFQFWAINRKLMEYPAEENGFRYIPFRY QTTTERPFIQKLFRPVAADGQLHTLGDLLKEVCP SAIDPEDGEKKNQVMIHGIEPMLLETPLQWLSEHL SYPDNFLHISIPQPTD
3874	A	776	366	QARGAPSSPMCPLAAAAVAAPRAPRLRLNRG LAAAMSTAQSLKSVDYEVFGRVQGVCFRMYTE DEARKIGVVGWVKNTSKGTVTGQVQGPEDKVN SMKSWLSKVGPSSRIDRTNFSNEKTISKLEYSNF SIRY
3875	A	1081	182	SLSSCQTDPRPMSAPLDAALHALQEEQARLKMR LWDLQQLRKELGDSPKDKVPFVSPKPLVFRGHT QQDPEVPKSLVSNLRHICPLLAGSALITFDDPKVA EQVLQKKEHTINMEECRLRVQVQPLELPMVTTIQ VMVSSQLSGRRVLVTGFPASLRLSEELLDKLEIF FGKTRNGGDDVDVRELLPGSVMLGFARDGVAQ RLCQIGQFTVPLGGQVPLRVSPYVNGEIQKAEI RSQVPRSVLVLNIPDILDGPELHDVLEIHFQKPT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				RGGGEVEALTVVPQQGGLAVFTSESG
3876	A	26	431	RMMKCPQALLAIFWLLSWVSSDKVVSPLSL VVHEGDTVTLNCSYEVTNFRSLLWYKQEKKAPT FLFMLTSSGIEKKSGRLSSILDKKELSSILNITATQ TGDSAITYLCAVEAQCSLVTCSLYSNSTAEALQL
3877	A	3	1291	KAFLRLAERGAAAAMLWSGCRFGARLGCLPG GLRVLVQTGHRSLTSCIDPSMGLNEEQKEFKQV AFDFAAREMAPNMAEWDQKELFPVDVMRKA QLGFGGVYIQTDVGGSLRLDTSVIFELATGC TSTTAYISIHNMCAWMIDSFGNEEQRHKFCPLC TMEKFASYCLTEPGSGSDAASLLTSAKKQGDHYI LNGSKAFISGAGESDIYVVMCRTGGPGPKGISCIV VEKGTPLSFGKKEKKVGNWSQPTRA VIFEDCA VPVANRIGSEGQGLIAVRGLNGGRINIASCSLGA AHASVILTRDHLNVRKQFGEPLASNQYLOFTLA DMATRLVAARLMVRNAVALQEERKDAVALCS MAKLFATDECFAICNQALQMHHGGYGYLKDYAV QQYVRDSRVHQILEGSNEVMRILISRLQEQ
3878	A	10	1014	LPGSTISSSGCQAPGRADSSGGARNSRRGDSRPG SCNRQAVAPPCSPGPQSRHWIHRGTAPQAGETR TLGRGSSAPNACSASVTPCCSPSPS*SCL*PTRRS PQNSSSTEYVRGFQWQHGLPST**PFSS*QWPGQH TQGC SKLLGKQTHLPCSTWPA**PSPSCLTRF* W*PSLMCLWASSCSVCV*SPSGSCRH*LWGTHST SRTC*ARRSSALPTGLCTDDTSWASSSKARPCAL QRPSSLSLSPCLTC*W*LSSSPMSARSPAGAET GSWATGSPRLTQWKSSRLTSTSHSARSAWKPSA TESTPSWPRFSSWTSGEDPASPAPAI
3879	A	200	699	LLLTGYIQLTQNQQLSGNQEMQAVDNLTSA NTSLCTRDIYKITQVLFPLLYTVLFFVGLITNGLA MIFFGIR*SNFIIFL*NTV'SDLLMLTFFPKI DAKLG*GPLRTFVCQVTSVIFYFT*ISFLGLT IDRYQKTRFPKTSNPKNLLGAKILK
3880	A	26	169	QPETDTMVHLTPEEKSAVTALWGKVNVEDADG DDLQILVDRPLRI
3881	A	37	1100	TPLDFWPGFVLSWLQPLSASLRARRAASGPPAC RIMPTTVDDVLEHGGEFFHFQKQMFLLALLSAT FAPITYVGIVFLGFTPDHRCRSPGVAELSLRCGWSP AEELNYTVPGPGPAGEASPRQCRRYEVDWNQST FDCVDPLASLDTNRSRLPLGPCRDGWVYETPGSS IVTEFNLCANSWMLDLFQSSVNVGFFIGSMSIG YIADRFRGRKLCLLTTVLINAAAGVLMASPTYT MLIFRLIQGLVSKAGWLIGYILITEFVGRRYRRTV GIFYQVAYTVGLLVLAGVAYALPHWRWLQFTV ALPNFFFLYYWCIPESPRWLISQNKNAEAMRIK HIAKKNGKSLPASL
3882	A	573	1620	KSKCRFPPEGLSEGFGPMRKEALSSGSVQEA LDEPQEQAEGLT VYVISEHSSLLPQDMMSYIGP KRTAVVRGIMHREAFNIIGRRIVQVAQAMSLTED VLAALADHLPEDKWSAEKRRPLKSSLGYEITFS LLNPDPKSHDVYWDIEGAVRRYVQPFNLALGAA GNFSVDSQILYYAMLGVNPRFDSASSSYLDLMDH SLPHVINPVESRLGSSAASLYPVLNFLLYVPELAH SPLYIQDKDGAPVATNAFHSRPGGIMVYNVDS KTYNASVLPVRVEVDMVRVMEVFLAQLRLFFGI

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, / =possible nucleotide deletion, \ =possible nucleotide insertion)
				AQPQLPPKCLLSGPTSEGLMTWELDRLLWARSV ENLATATTTTSLA
3883	A	2369	844	RIHREEDFQFILKGIARLLSNPLLQTYLPNSTKKIQ FHQELLVLFWKLCDFNKVGQPRGALQGDGEQLP Q*PGGRDSVRLRGVQSCPSLELSPLGPSHP*KF LFFVLKSSDVLDILVPILFFLNDARADQSRVGLM HIGVFILLLLSGECNFGVRLNKPYSIRVPM DIPVF TGTHADLLIVVFHKIITSGHQRLQPLFDCLLTIV NVSPYLKSLSMVTANKLLHLLEAFSTTWFLFSAA QNHHLVFFLLEVFNNIIQYQFDGNSNLVYAIRKR SIFHQLANLPTDPPTIHKALQRRRTPEPLSRTGS QGGAPPWRAPAPLPLQSQAPSRPVWLLQALTS *PRSPRCQRMAPCGPWNLSPSRAWMAARLRGS PARHGGSSGDRP/HSSASGQWSPTEWVLSWKS KLPLQTIMRLLQVLVPQVEKICIDKGLTDESEILR FLQHGTLVGLLPVPHPILRKYQANS GTAMWFRT YMWGVYLRNVDPPVWYD TDVKLFEIQRV
3884	A	1	804	NGPRAPFSQEGQSTGPPPLIPRLGQHGAQGRIPPL NPGQGPGPNKDDSRGPPNHHMGPMSEERRHEQSG GPEHGP ERGPLRGQDCRGPPDRRGPHDPDFDDF SRPDDFHPDKRFGRHLREFEGRGGPLQEEKWR RGGPGPPFPDHPREFSEG DGRGAARGPPGA WEG RRPGG*TFPPGSRGPTFS/SGAEESFRRGAPPRHE GRAPPRGRDGFPGPEDFGPEENFDASEEAARGRD LRGRGRGTPRGERVTKDTWSGRIGCRIHWL
3885	A	3	996	GRRRAGPAHSARMYNMMETELKPPGPQQTSGG GGGNSTAAAAGGNQKNSPDRVKRPMNAFMVW SRGQRRKMAQENPKMHNSEISKRLGAEWKLLSE TEKRPFIDEAKRLRALHMKHEHPDYKYRPRRKT TLMKKDKYTLPGGLLAPGGNSMASGVGVGAGL CAGVNOR... QHPGLNAHGAAQMOPMEYDVS...LQYNSM TSSQTYMNG/SRPTYSMSYSQQ...TPGMAPGSMG SVVKSEASSSPVVTSSSHSRAPQA...DLRDMIS MYLPGA EVPEPAAPSRLHMSQHYQS...VPGTAI NGTLPLSHM
3886	A	773	317	QCTQKAAEGYTQFYVDVLDGKLACVNKCTKG TKSQMNCNLGTCQLQRSGPRCLCPNTHWY GETCEFNIAKSLVYGIVGAVMAVLLALILILFS LSQVRKRHRPESEGEADFGLENATNNFGPTLET VDSGTELHIQRPEMVASTV
3887	A	3	466	VDFRVKTLVVDNKCFLVQLWDTAGQERYHSM RQLLRKADGVVLMYDITSQESFAHVRYWLDCL QDAGSDGVVILLGNKMDCEBERQVSVEAGQQL AQELGVYFGECSAALGHNILEPVVNLARSLRMQ EEGLKDSL VKVAPKRPPKRFGCCS
3888	A	3412	3144	QNIDITNFSSSWNDGLAFCALLHTYLPAPHYQEL NSQDKRRNFMLAFQAAESVGIKSTLDINEMVRT ERPDWQNVMLYVTAIYKYFET
3889	A	1	1160	LVVTAITAILAFFNEYTRMSTSELISELFND CGLL DSSKLC DYENRNTSKGGELPDRPAGVGVYSAM WQLALTLILKIVITIFTGMKIPSGLFIPSMAGAI AGRLLGVGMEQLAYYHQEWTVFNSWCSQGAD CITPGLYAMVGAAACLGGVTRMTVSLVVMFEL TGGLEYIVPLMAAAMTSKWVADALGREGIYDA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				HIRLNGYPFLEAKEEFAHKTLMADVMPKPRRNDP LLTVLTQDSMTVEDVETISETTYSGFPVVVSRES QRLVGFVLRRLDIISIENARKKQDGVVSTSIYFTE HSPPLPPYTPPTLKLRLDLSPFTVTDLTPEIVV DIFRKLGLRQCLVTHNGRLLGIITKKDVLKHIAQ MANQDPDSILFN
3890	A	1	387	SWCWTGIFVLGTTNLRLEGSWYRSLWGPFGNTT TATLGFAPQAPVGDVALNQPDMCVYRRGRKK RVPTKLQLKELENEYAINKFINKDKRRRISAAT NLSEKQVTIWFQNRNVKDKKIVSKLKDVTVS
3891	A	2	2914	RGGGGDHKMDLSSLQEDLQEDADGFGVDDYS SESDVIIIPSALDLAST/QDEMVERPLGRLDKVYA ASENHI*PDKMVAPEFASIPLEIVCDDERDCIAV LGKN*PDWADDSEPTVRAAELEQVPHIALFLFK KTRLSITICFFSKFLLPYCGDLTADQNNQVRKT SQAALLALLEQELIERFDVETKVCPLIELTAPDS NDDVKTEAVAIMCKMAPVMVGKDITERLILPRFC EMCCDCRMFHVVRKIVCAANFGDICSVVGGQAT EEMLLPRFFQLCSDNVWGVKACAECFMAVSC ATCQEIRRTKLSALFINLISDPSRWVRQAAFQSLG PFISTFANPSSSGQYFKEESKSSEEMSVENNRTR DQEAPEDVQVRPEDTPSDLSVSNSSVILENTMED HAAEASGKPLGEISVPLDSSLCTLSSESHQEAAS NENDKKPGNYKSMLRPEVGTTSQDSALLDQELY NSFHFWRTPLEIDLIELEQNSGGKPSPEGPEEE SEGPVPSSPNITMATRKELEEMENLEPHIDDPDV KAQVEVLSAALRASSLDAHEETISIEKRSDLQDE LDINELPNCKINQEDSVPLISDAVENMDSTLHYIH NDSDLNNSFSFSPDEERRTKVQDVVPQALLDQY LSMTDPSRAQTVDTIAKHCAYSPLGVALTLGR QVWHCLRETFTLASDMQWTRRTAFSIFIA VLRDQQLTAADLVPIFNGFLK*FSMKSRI LHDFLKLHIDKRREYLYQLQEFLVTDNSRNWR FRALAEQLILLELYSPRDVYDYLRIALNLCAD KVSSVEWISYKLVSEMVKLHAATPTFGVDLIN ELVENFGRCPKWSGRQAFVFCQTVIEDDCLPM DQFAVHLMPHLLTLANDRVPNVRVLLAKTLRQT LLEKDYFLASASCHQEAVEQTIMALQMDRSDV KYFASIHASTKISEDAMSTASTY
3892	A	158	2191	VPLPAPSGLSGGGSRGAGCKKAPFGRAPAPGLAP LRPSEPTMAVPPGHGPFSGFPGQEHTQVLPDVR LLPRRLPLAFRDATSAPLRKLSVDLIKTYKHINEV YYAKKKRRAQQAPPQDSSNKKEKKVLNHGYDD DNHDYIVRSGERWLERYEIDSLIGKGSFGQVKA YDHQTQELVAIKIKNKKAFLNQAQIELRLLELM NQHDTEMKYYIVHLKRHFMRNHLCLVFELLS YNLYDLLRNTHFRGVSLNLRKLAQQLCTALLF LATPELSIIHCDLKPENILLCNPKRSAIKIVDFGSS CQLGQRIYQYQSRFYRSPVLLGTPYDLAIDMW SLGCILVEMHTGEPLFSGSNEVCPQEGVDQMNRI VEVLGIPPAAMLDDQAPKARKYFERLPGGGWTLR RTKELRKDYQGGPTRRLQEVLGVTGGPGGRRRA GEPGHSPADYVLRFDLVLRMLEYEPAAISPLG ALQHGFRRRTADEATNTGPAGSSASTSPAPLDT PSSSTASSISSGGSSGSSSDNRTYRYSNRYCGGP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /-possible nucleotide deletion, \-possible nucleotide insertion)
				GPPITDCEMNSPQVPPSQPLRPWAGGDVPHKTH QAPASASSLPGTGAQLPPQPRYLGRPPSPSPPPP ELMDVSLVGGPADCSPPHPAPAPQHPAASALRT RMTGGRPPLPPDDPATLGPLHLGLRGVPQSTAAS S
3893	A	68	258	PEEYYPFSPTLQQLFFLLDSDMGSRPESMGCRK NTVPRPASPTTEAGTDPQTFLHTWVSECRD
3894	A	1120	136	SLPLAPAPAVAGPVALCPAGLCPAQPGMPAGPA AASGSHPEVGSVLQRSSQPHWPNWPAGAGHLPP PAGPPFPYNPPAGPGAAAAGLA*SPPRSPTPCSVGP QSCPANASAPPAQPCLAGAPPAASLPPPGPGSVS AAPAPGGPAPAEPLGVPPVPAWLLPDSPLPGT HSGPPAAVSLPPAAAACPVVVPPLPHHPDLES PSAAAPNPGCAGGIRHFPFGSPEASSPLRPAAPPA LLPLRPPS*P/VPWKPLHSPVAVAGGSFVAGGSV LPAPDLQPRPSGPPAASPTPGPGVAQPPGSAVL PTVP*APPVSGAAPGRKREW
3895	A	2	1347	FGAVSYRPGNGSCWVKVTASSDLSLISCLCPPR SLCSSQACVLPVPGPSLLLPQGLHVCASAGTRW PLSCSIDFQRLLAHEEETQKRRAKESGMAFTQLT FRDVAIEFSQDEWKCLNSTQRTLYRDVMLENYR NLVSLDLNRNCVIKELAPQOEGNP/ARSIPHSDIGT T*KT*H*RVLLQGNQEKNTRL*LSVER**KKLQQ SDYGPKRKSYL*ERPTR*KRYRKQVY*TSAL*LSF LPHPHQLQQFAEGKIYECNHVEKSVNHGSSVSP PQIISSTIKTHVSNKYGTDFICSSLLTQEQKSCIRE KPYRYIECDKALNHGSHMTVRQVSHSGEKGKCYC DLCKGVFSQKSNLARHWRVHTGEKPYKCNECD RSFSRNSCLALHRRVHTGEKPYKCYECDKVFSR NSCLALHQBKTHIGEKPYTCKECCQAQFVSRSTLTN HGVHSDK
3896	A	202	258	MVQSCSAYGNNALADKPVSFHKFPLTRPSLC KEWEAAVRRKNTKPTKYSSICSEHFTPDCKREC NNKLLKENAVPIITCTEPHDKKEDLLEPQEQ
3897	A	2	382	SHGLSRAPHLAAPAPALASRPCFSAPCSQGGG GGGPATMIHFILLFSRQGLRLQK WYITLPDKER KKITREIVQILSRGHRTSSFVDWKELKVYKRYA SLYFCCAIE/NQDNELLTLENVHR
3898	A	718	305	SEQEPLLGDTPGSREWDILETEEHYKSRWRSIRIL YLTMFLSSVGFSVVMMSIWPYLQKIDPTADTSFL GWVIASYSLGQMVASPIFGLWSNYRPRKEPLIVSI LISVAANCLYAYLHIPASHNKYYMLVARGLLGIG
3899	A	24	718	FRGRPGIPEREGKGNHSFVEVARVIVDLHSRLG GAMAERKGTAKVDFLKKIEKEIQKWDTERVFE VNASNLEKQTSKGKYFVTFPYPYMNGRLHLGHT FSLSKCEFAVGYQRLKGKCLFPFGLHCTGMPIK ACADKLKREIELY/GCPPDFPDEEEEEETS VKTE DIIKDKAKGKSKAA/AKAGSSKYQWGIMKSLG LSDEEIVKFSEAEHWLDYFNALAIQDLKRMG
3900	A	360	1	VPATSSNVSPSSSESEPDLSRSSSSDAPSSSPSPV SPCSLSLSPESPLPTLLSSKSPAGSAGPTCGCPS GGLRATA/PSRLSSSIAAH/SSAPETSRPAAARE RSPPLHDRESHE
3901	A	193	345	GEWAVPPAPGGQGVSIHPGPEPGQSGSVHIAPRQ GEGSDRTEPLICPKAAP

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3902	A	1188	1389	NPAARSAAAREGSPALPPPPVS/SSSGLGLLLPLSP PGSHAANPALSPRAPHSHYRPRRCGPRRRPR
3903	A	63	396	NNMRNPHLSSNHLYLNARTETVFARMESVKQRI LAPGKEGLKNFAGKSLGQIYRVLEKKQDTGETIE LTEDGKPL*VPERKAPLCDCTCFGLPRRYIIAIMS GLGFCISFG
3904	A	732	1046	AMSECLILYIHKHIDTYSQSYLFNDLFYPVYSGG RMVTYEHLEVVFGKSEDEHYPLW*VLFGK*YA VAPNALMFIRFM*NCTFVPKLP*VMDLK**LQYK SR
3905	A	46	910	QPPPPPPPPSPPPPPPPARALSHLRLHPDACLFPS PFPLPCSTMPGMMMEKGPELLGKNRSANGSAKSP AGGGGSGASSTNGGLHYSEPESGCSSDDEHDVG MRVGA EYQARIPEFDPGATKYTDKDNNGMVLW SPYHSIPDAKLDEYIAIAKEKHGYNVEQALGMLF WHKHNIEKSLADLPNFTFPDEWTVEDKVLFEQ AFSFHGKSFHRIQQMLPDKTIASLVKYYYSWKK TRSR TSLMDRQARKLANRHNQGDSDDDVEETHP MDGNDSDYDPKKEAKKEGMS
3906	A	2	513	KVCNCCSQELETSTYVDKNINLEQRNRSSPSAK GHNHPGELGWENPNEWSQEA AISLISEEEDDTSS EATSSGKSIDYGFISAILFLVTGILLVIISYIVPREV TVDPNTVAAREMERLEKESARLGAHLDRCVIAG LCLLTGGVILSCLLMMSMWKGELYRRNRVAS
3907	A	71	412	ILIMSNCLQNFLKITSTRLLCSRLCQQLRSKRKFF GTVPISRLHRRVVITGIGLVTP LGVGT HL VWDRLI GGESGIVSLVGEEYKSIPCSVAA YVPRGSDEGQF NEQNFVSKSD
3908	A	77	746	LGTLLGWRAFLFSRCLAFHSPFILLNTPKLVKTAE LPPDRNYVLGAHPHGMCTGFLCNFTSNGFSQ LTPGLRPWLAVLAGLFYLPYRDYALNGLCV RQSLDFILSQPQLGQAVVGVGGAHEALYSVPGE HCLTLQKRKG FVRLALRHGASLVVYSFGENDIF RLKAFATGSWQHWCQLTFKKLMGFSPCIFWGR GLFSATSWGLLPFAVPITTV
3909	A	1	793	FRAAGRPAAMGDIPVVLSSWKASPGKVTEAV KEAIDAGYRHFDCA YFYHNEREVGAGIRCKIKE GAVRREDLLIATKLWCTCHKKSLVETACRKSLK ALKLNYLDLYLIHWPMGFKPPHPEWIMSCSELSF CLSHPRVQDLPLDESNMVIPSDTDFLDTWEAME DLVITGLVK NIGVS NFNHEQLERLLNKPGLRFP LTNQIECHPYLTQKNLISFCQSRDVSVTAYRPLG GSCEGVDLIDNPVIKRIAKEHGKSPAQILI
3910	A	202	705	FFTMHRKKVDNRIRILIENGVAERQSRSLFVVGD RGKDQVVILHHMLSKATVKARPSVLWCYKKEL GFSSHRKKRMRQLQKKIKNGTLNIKQDDPFELFI AATNIRYCYYNETHKILGNTFGMCVLQDFEALTP NLLARTVETVEGGGLVVILLRTMNSLKQLYTVT M
3911	A	3	723	AGRGARAAGEGGGPFKSRPRPLPSSRSLPAVGGG RYGADKMAAGGAVAAAPECRLPYALHKWSSF SSTYLPENILVDKPNQSSRWSESNNPPQYLK LERPAIVQNTIFGKYEKTHVCNLKKFKVFGGMN EENMTELLSSGLKNDYNKETFTLKHKIDEQMFPC RFIKIVPLLSWGSPSFNFSIWYVELSGDDPDIVQPC

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				LNWYSKYREQEAIRLCLKHFRQHNYTEAFESLQ KKT
3912	A	2	461	FEKKQLRRPSLFLGCCSFGIMAPSLWKGLEGIG LFALAHAAFSAAQHRSYMRLTEKEDESLPIDIVL QTLFAFAVTCYGIVHIAGEFKDMDATSELKNKTF DTVRNHPSFYVFNHRGSEYFSGPSDTANSSNQDA LSSNTSLKLRLKLESRR
3913	A	362	20	APGRPEAKVPERSRESGSRVRGPLLQLRPGRTS RPASGRGRGGAGGSYGKMRKPD SKIVLLGDMN VGKTSLLQRYMERRFPD TVSTVGGA FYLKQWRS YNISIWD TAGEAGAA
3914	A	1	7545	PGIRVGITSQTGLSSNLQENC SKLAFISSHGTEKQ LQCMPEGRGRASSISDLQKGFEKGTGEKHV PGVGSARHSPQASAGGSPWQRGKAQTRWLGP DPGRKRRRGSPQEEGLRVSAARLLCSGANRC KVLVRQNSTPNTQQPAVHPSTPPSRPLPQAGRCL VAPLRPHPDWVAAKTLAKALRAPGKPWRLAAP SPLGDLGAPGLPGPSTAPRTLSVEEPGVECNQLC LYADVTD PVLC LGQKDPGVEGKHCEKEKISSK ELKHVHAKSEPSKPARRLSESLHVVDENKNESKI EREHKRRSTSPVIMEGVQEETDTRDVKRQVERSE ICTEEPQKQKSTLKNKHLKKDDSETPHLKSLK KEVKSSEKPEREKTSEDKLSVKHKYKGD CMH KTGDETELHSSEKGLKVEENIQKQSQT KLSDD KTERKSKHRNERKLSVLGKDGPVSEYIITDEN VRKENNKKERRLSAEKTKAEHKSRRSSDSKI QK DSLGSKQHGITLQRRSESYSEDKCDMDSTNMDS NLKPEEVVHKEKRRTKSLLEKLVLSKSKTQG KQVKVETELQEGATKQATTPKPDKEKNTEEND SEKQRKSKVEDKPFEETGVEPVLETASSSAHSTQ KQSSHRAKLPLAKKKTSDKLSSTFPKLS D GHKSRLKHSSKLNENKSLDDKGKEVDSS HEKARGNSSLMKCLSRRLCENRRGSLSQEMAK GEEKLAANTLSTPSSSQRPKKS GDMTLIPEQEP MEIDSEPGVENVFEVSKTQENRNNNSHQDIDSEN MKQKTSATVQKDELRTCTADSKATAPAYKPNP GTGVNSNSEKHADHRSTLTCKMHQSAVSKMNP GEKEPIHRGTTEVNIDSETVHRMLLSAPSEND RV QKNLKN TAAEEHVAQGDATLEHSTNLDSSPSLSS VTVPPLRESYDPDVIPLFDKRTVLEGSTASTSPAD HSALPNQSLTVRESEVLKTSDSKEGEGFTVDTP AKASITSKRHIPEAHQATLLDGKQGVIMPLGSK LTGVIVENENITKEGGLVDMAKKENDLNAEPNL KQTIKATVENGKKDGLAVDHVVG LNTKYAETV KLKHKRS PGKVKD ISIDVERRNENSEVDTSAGSG SAPSVLHQNRNGQTEDVATGPRRAEKTSVATSTE GKDKDVTLSPVKAGPATTTSSETRQSEVALPCTS IEADEGLIIGTHSRNNPLHVGAEESECTVFAAAEE GGAVVTEGFAESETFLTSTKEGESGECVAESED RAADLLAVHAVKIEANVNSVVTEEKDDAVTSAG SEEKCDGSLSRDSEIVEGTTTTFISEVESDGAVTSAG TEIRAGSISSEVDGSQGNMMRMGP KKETEGTV TCTGAEGRSDFVICSVTGAGPREERMV TGAGV VLGDNDA PPGTSASQEGDGSVNDGTGESAVTS TGITEDGEGPASCTGSEDSSEGF AISSESEENGESA

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				MDSTVAKEGTNVPLVAAAGPCDDEGIVTSTGAKE EDEEGEDVVTSTGRGNEIGHASTCTGLGEESEGV LICESAEGDSQIGTVVEHVEAEAGAAMNANENN VDSMSGTEKGSKDDICSSAKGIVESSVTSVSG KDEVTPVPGGCEGPM TSAASDQSDSQLEKVEDT TISTGLVGGSYDVLVSSEVPECEVAHTSPSEKED EDIITSVENECDGLMATTASGDITNQNLSAGGK NQGKVLIISTSTNDYTPQVSAITDVEGGLSDALR TEENMEGTRVTTEFEAPMPSAVSGDDSQLTASR SEKDECAMISTSIGEEFELPISSATTIKCAESLQP VAAA VEERATGPVLISTADFE GPMPSAPPEAESP LASTSKEEKDECALISTSIAEECEASVSGVVVESE NERAGTMVEEKDGSIGIISTSSVEDCEGPVSSAVP QEEGDPSVTPAEEMGDTAMISTSTSEGCEAVMIG AVLQDEDRLTITRVEDLSDAAIISTSTAECMPISA SIDRHEENQLTADNPEGNGDLSATEVSKHKVPM PSLIAENNCRCPGPVRGGKEPGPVLAVSTEEGHN GPSVHKPSAGQGHPSAVCAEKEEKHGKECPEIGP FAGRGQKESTLHLINAEKKNVLLNSLQKEDKSPE TG TAGGSSTASYSAGRGLEGNANSPAHLRGPEQ TSGQTAKDSSVSIRYLA AVNTGAIKADDMPVPQ GTVAEHSFLPAEQGSEDNLKTSTTKCITGQESKI APSHTMIPPATYSVALLAPKCEQDLTKNDYSKG WTDQASAEKTGDDNSTRKSFPEEGDIMVTVSSE ENVCDIGNEESPLNVLGGLKLANLKMEAYVPS EEKNGEILAPPESLCGGKPSGIAELQREPLLVNE SLNVENSGFRTNEEIHSESYNKGEISSGRKDNAE AISGHSVEADPKEVEEEERHMPKRKRKHYSSE DEPDNDPDLDSRIETAQRQCPETEPHATKEENS RDLEELPKTSSETNSTSRVMEEKDEYSSSETTGE KPEQNDDLTCKSCF
915	A	1	545	PGIRVGITSQTGLSSNLQ...KLAFLSHGTEKQ LQCMPEMGRGRASSISDLQKGFEKGTGEKHV PGVGSARHSPQASAGGSPWQRGKAQTRWLGKP DPGRKRRRGSPQEEGGLRVSAARLLCSGANRC KVLVRQNSTPNTQPPAVHPSTPPSRPLPQAGRCL VAPLRPHPDWVA AKTLAKALRAPGKPWRLAAL SPLGDLGAPGLPGPSTAPRTLSVEEPGVECNQLC LYADVTDVPLCLGQKDPGVEGKHCEKEKISSSK ELKHVHAKSEPSKPARRLSESLHVVDENKNESKI EREHKRRSTPVMIEGVQEETDTRDVKRQVERSE ICTEPPQKQKSTLKNEKHLKKDDSETPHLKSLK KEVKSSKEKPEREKTSEDKLSVKHKYKGDCMH KTGDETELHSSEKGLKVEENIQKQSQQTKLSSDD KTERKSKHRNERKLSVLGKDGPVSEYIITDEN VRKENNKKERRLSAEKTKAEHKSRRSSDSKIQK DSLGSQKHGITLQRRSESYSEDKCDMDSTNMDS NLKPEEVVHKEKRRTKSLLEELVLKSKSKTQG KQKVVTETELQEGATKQATTPKPDKEKNTEEND SEKQRKSKVEDKPFETGVEPVLETASSAHSTQ KDSSHRAKLPLAKEKYKSDKDDSTRLERKLS GHKSRLKHSSKDIKKDENKSDDKDGEVDSS HEKARGNSSLMEKKLSRRLCENRRGSLSQEMAK GEEKLAANTLSTPSGSSLQRPKKS GDMTLIPEQEP MEIDSEPGVENVFEVSKTQDNRRNNNSHQDISEN

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, * = Stop codon, / = possible nucleotide deletion, \ = possible nucleotide insertion)
				MKQKTSATVQKDELRTCTADSKATAPAYKPGR GTGVNSNSEKHADHRSTLTKKMHIQSAVSKMNP GEKEPIHRGTTEVNIDSETVHRMLLSAPSENDRV QKNLKNTAAEEHVAQGDATLEHSTNLDSSPSLSS VTVVPLRESYDPDVIPLFDKRTVLEGSTASTSPAD HSALPNQSLTVRESEVLKTSDSKEGGEGFTVDTP AKASITSKRHIPEAHQATLLDGKQGVIMPLGSK LTGVIVENENITKEGGLVDMAKKENDLNAEPNL KQTIKATVENGKKDGIADVHVGLNTEKYAETV KLKHKRSPGKVKDIDISIDVERRNENSEVDTAGSG SAPSVLHQNRNGQTEDVATGPRRAEKTSVATSTE GKDKDVTLSPVKAGPATTTTSSETRQSEVALPCTS IEADEGLIIGTHSRNNPLHVGAEESECTVFAAAEE GGAVVTEGFAESETFLTSTKEGESGECVAESED RAADLLAVHAVKIEANVNSVVTTEEKDDAVTSAG SEEKCDGSLSRDSEIVEGTTTFISEVESDGA VTSAG TEIRAGSISSEEVDSQGNMMRMGPKKETEGTV TCTGAEGRSDNFVICSVTGAGPREERMVGTAGV VLGDNDAPPGTSASQEGDGSVNDGTEGESAVTS TGITEDGEGPASCTGSEDSSEGFAISSESEENGESA MDSTVAKEGTNVPLVAAGPCDDEGIVTSTGAKE EDEEGEDVVTSTGRGNEIGHASTCTGLGESESEGV LICESAEGDSQIGTVVEHVEAEAGAAIMNANENN VDSMSGTEKGSKDTDICSSAKGIVESSVTSVSG KDEVTPVPGGCEGPMTSAASDQSDSQLEKVEDT TISTGLVGGSYDVLVSGEVPECEVAHTSPSEKED EDIITSVENEECDGLMATTASGDITNQNSLAGGK NQGVLIISTSTTNDYTPQVSAITDVEGGLSDALR TEENMEGTRVTTEEFAPMPSAVSGDDSQLTASR SEEKDECAMISTSIGEEFELPISSATTIKCAESLQP WAAAN EERATGPFVISTADFEFGME*SAFTEAESP EASTOR EEEKDECALISTSIAEECEASVSGVVVE NERAGTVMEEKDGSIIISTSSVEDCEGPVSSAVP QEEGDPSPVTPAEEMGDTAMISTSTSEGCEAVMIG AVLQDEERLTITRVEDLSDAIIISTSTAECMPISA SIDRHEENQLTADNPEGNGDLSATEVSKHKVPM PSLIAENNCRCPGPVRGGKEPGPVLAVSTEEGHN GPSVHKPSAGQGHPSAVCAEKEEKHGKECPEIGP FAGRGQKESTLHLINAEKKNVLLNSLQKEDKSPE TGTAGGSSTASYSAGRGLEGNANSPAHLRGPEQ TSGQTAKDSSVSSIRYLAAVNTGAIKADDMPVQ GTVAEHSFLPAEQQGSSEDLNKTSTTKCITGQESKI APSHTMIPPATYSVALLAPKCEQDLTIKNDYSGK WTDQASAEKTGDDNSTRKSFP EEGDIMVTVSSE ENVCDIGNEESPLNVLGGLKLANLKMBA YVPS BEEKNGEILAPPESLCGGKPSGIAELQREPLLVNE SLNVENSGFRTNEEIHSESYNKGEISSGRKDNAE AISGHSVEADPKVEVEEERHMPKRKRKQHYLSSB DEPDDNPDVLDRIETAQRQCPETEPHATKEENS RDLEELPKTSSETNSTTSRVMEEKDEYSSSETTGE KPEQNDDDTIKSQE
3916	A	2	773	GPFGLVWPSAKPGPVTAVEARPPDASDPEGLRG GSPAPLLAPGPLDPSGRLHPAVSMMSYLKQPPYG MNGGLAGPAMDLLHPSVGYPATPRKQRRERTT FTRSQLDVLEALFAKTRYPDIFMREEVALKINLPE

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SRVQVWFKNRRRAKCRQQQSGSGTKSRPAKKK SSPVRESSGSESSGQFTPPAVSSSASSSSASSSSA NPAAAAAAGLVVAKLPCPLHIFSLCVFIEENRLV SGSWARDIRSVEETDKSGYR
3917	A	2	776	RNIPGRRFRPPGLRRLKGPMPREPRGYRTRVP ALRELVPSSHAGSGASEHCQNNRQGSQRQHRASR NVQAGGALAPPRHLGCLCSRLHFLKPDLSVRAA PSRAGASVMALRKELLKSIWYAFTALDVEKSGK VSKSQLRVLSHNLTYTLHIPHDPVALEEHFRD ^{DD} DGPVSSQGYMPYLNKYLDKVEEGAFVKEHFDE LCWTLTAKKNYRADSNNGSMLSNDQAFRLWCL FNFLSEDKYPLIMDPDEGEYLLKRY
3918	A	10	318	WQDLVCLGGSRAQEQKPLQQLWNAILLVAMLL CTGLVVQAQRQASRQSORELGGQVDLFKRRVV RRLASLKTRRCRLSRAAQGLPDGAETCAVCLD YFCNKQ
3919	A	1	204	RVLTAINHILKENLRKFYKGGKDKPLDLRPKKT RAMRRRLNMHEENLTKKKQHRKERLYPLRKYA AKA
3920	A	1	654	RCCRSFVAPLQEKVVFGLFGLGAILCLSFSWLFHT VYCHSEGVSRLFSKLDYSGIALLIMGSFVPWLYY SFYCNPPCFTYLIVICVLGIAAIVSQWDMFATPQ YRGVRAGVFLGLGLSGIIPTLHYVISEGFLKAATI GGIGWMLMASLYTTGAALYAARIPERFFPGKCD IWFHSHQLFHIFVVAGAFVHFHGVSNLQEFRFMI GGGCSEEDAL
3921	A	1587	452	LERDGCGEEGGSVRSGAGPDSDPGASSPPAG HRGTAASPRPVAAPSRTPAPPHTRARASPLPSG PAWRRVQWFSRVSGQVSTLMKATVLMRQPGRV QEIVGALRKGGGDRQLQVISDFDMLSRFAYNGK RCPSSYNIL ^{DN} SK ^{EE} EECRYELTALL ⁷⁷⁷ 7P ⁵ SL PHRTVKEKLPHMVEW ⁷⁷⁷ 7AHNLLCQQKIQKFQI AQVVRESNAMLREGYKTFNTLYHNNIPLFIFSA GIGDILEEIRQMKVFHPNIHIVSNYMDFNEDGFL QGFGQLIHTYNKNSSACENCYFQQLLEGKTNV ILLGDSIGDLTMADGVPGVQNLKIGFLNDKVEE RRERYMDSYDIVLEKDETLDVVNGLLQHILCQG VQLEMQGP
3922	A	2	164	GKIYQRAFGGHSKFKGKGVQAHGCCCVADRTG HSILHTSYGRERPAPVHLRQDT
3923	A	2	3258	EHATHAYAKLGTRRRHREVTVFVPTWQLKKNR RVRESHFLTKLHSLKMLSITPSQLENGKKITTYD YRFMVKLAETDGIIVTNEQIHILMNSSKKLMVK DRLLPFTFAGNLFMVPPDDPLGRDGPLDDEFLLKP NRLD ^{TD} DIGNFLKVWKTLPSSASVTELSDDADSG PLESLPNMEEVREEKEERQDEEQRQGGGTQKAA EEDDLSSLASVFRVECPSLSEEILRCLSLHDPDP GALDIDLLPGAASPYLGIPWDGKAPCQVLAHL AQLTIPSNFTALSFFMGFMDSHRDAIPDYEALVG PLHSLKQKPDWQWDQEHEEAFLALKRALVSAL CLMAPNSQLPFRLEVTVSHVALTALHQEHSGRK HPIAYTSKPLLPDEESQGPQSGGDS ^{PY} AVAWALK HFSRCIGDTPVVLDLSYASRTTADPEVREGRRVS KAWLIRWSLLVQDKGKRALELALLQGLLGENRL LTPAASMPRFFQVLPFSDLSTFVCIHMSGYCFYR

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EDEWCAGFGLYVLSPTSPPVLSFSFCSPTPTYA HLAAVACGLERFGQSPLPVVFLTHCNWIFSLWE LLPLWRARGFLSSDGAPLPHPSLLSYIISLTGSLSS LPFIYRTSYRGSFAVTVDTLAKQGAQGGGQWW SLPKDVPAPTVPSPHAMGKRPNLLALQLSDSTLAD IARLQAGQKLSGSSPFSSAFNSLSLDKESGLLMF KGDKKPRVWVVPQTQLRRDLFSVHDIPLGAHQ PEETYKKLRLLGWWPQMGEHVVDYCRSCLFCIP RNLIGSELKVIESPWPLRSTAPWSNLQIEVVG PVTISEEGHKHVLIVADPNTRWVEAFPLKPYTH TAVAQVLLQHVFAWGVVPRLEAAQGPQFARH VLVSCGLALGAQVASLSRDLQFPCLTSSGAY WEFKRALKEFIFLHGKKWAASLPLLHLA FRASSTDATPFKVLTTGGESRLTEPLW WEMSSANIEGLKMDVFLQLVGELLEBLH WRVADKASEKAENRRFKRESQEKEWNVGD QVLLLSLPRNGSSAKWVGPFYIGDRLSL SLYRIWGFPTPEKLGCIYPSLLMKAFKSGT PLSFKVLEQ
3924	A	1	1826	MGSVTVRYFCYGLFTSATWTVLLFVYFNFSEV TQPLKNVPVKSGSPHGPSKPFYPRFTRGPSRVL EPQFKANKIDDVIDSRVEDPEEGHLKFSELGMIF NERDQELRDLGYQKHAFNMLISDRLYHRDVPD TRNAACKEKFYPDLPAASVVICFYNEAFSALLR TVHSVIDRTPAHLLEHILLVDDSDDFDLKGLDE YVQKYLPGKIKVIRNTKREGLIRGRMIGAAHATG EVLVFLDSHCEVNVMWLQPLLAAIREDRHTVGC PVIDIISADTLAYSSSPVVRGGFNWGLHFKWDLV PLSELGRAEGATAPIKSPTMAGGLFAMNRQYFH ELGQYDSGMDIWGGENLEISFRIWMCGGKLFIP CSRVGHIHFRKRRPYGSPGQDTHNSRLAHV WLDEYVQVFSLPDLVTHNTHNISENVELNTHY GCKSEKWLNDNVYPENKSGELLKPKQPIFVNR GPKRPKVLQGRLYHLQINCLVAQGRPSQKG GLVVLKACDYSDPNQIWYNSHELVLNLSLLCLD MSETRSSDPRLMKCHGSGGSQQTFTGKNNRLY QVSVGQCLRAVDPLGQKGSVAMAICDGSSSQ WHLEG
3925	A	5386	2897	VRWNSKTECYLSIQTQENFPANLNELVNCIVISSL VTTQRKLKAMSLGSRNQLARAVLNPNPMDFCT KDILLTTSERIAYLRDFNEDQKKAJETAYAMVK HSPSVAKICLIHGPPTGKSKTTVGLLYRLLTENQ RKGHSDENSNAKIKQNRVLVCAPSNAAVDELM KKIILEFKEKCKDKKNPLGNCGDINLVRGPEKSI NSEVLKFSLDSQVNHMRMKELPSHVQAMHKRK EFLDYQLDELSRQALCRGGREIQRQELDENISK VSKERQELASKIKEVQGRPQKTQSIILSHIICCT LSTSGGLLLESAFRGQGGVPFSCVIVDEAGQSCEI ETLTPLIHRCNKLILVGDPKQLPPTVISMKAQEY GYDQSMMARFCRLLEENVEHNMISRLPILQLTVQ YRMHPDICLFPSNYVYNRNLKTNRQTEAIRCSD WPFQPYLVFDVGDGSERRDNDSYINVQEI KLVMEIKLIKDKRKDVSRNIGHITHYKAQKTM IQKDLKEFDRKGPAEVDTVDAFQGRQKDCVIV TCVRA NSIQGSIGFLASLQRLNVTITRAKYS LFIHLRLTMENQHWNLIIQDAQKRGAIKTC DKNYRHDAV

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				KILKLKPVLRSLTHPPTIAPEGSRPQGGLPSSKL DSGFAKTSVAASLYHTPSDSKEITLTVTSKDPERP PVHDQLQDPRLKRMGIEVKGGIFLWDPQPSSPQ HPGATPPTGEPGFVVHQDLSHVQQPAAVVAAL SSHKPPVRGEPPAASPEASTCQSKCDDPEEELCH RREARAFSEGEQEKCGSETHHTRRNSRWDKRTL EQEDSSSKKRKL
3926	A	99	284	MPREDRATWKSNYFLKIIQLDDYPKRFIVGANN VGSKQMQQIRMSLRGKAVVLMGKNTMMR
3927	A	542	2	AHLLMLNLALVDLLVLTSLPFLIHYASGENWI FGDFMCKFIRFSFHFNLYSILFLTCFSIFRYCVIHH PMSCFSIHKTRCAVVACAVVWIIISLVAVIPMTFLI TSTNRTNRSACLDLTSSDELNTIKWYNLLLTALL CLPLVIVTLCYTTIHTLTHGHANDSCLKQKARR LTILL
3928	A	1	1516	GEEAVGGGAEGGGFGVGAQGRAGGRGVEAGR MRLSKTLVMDMADYSAALDPAYTTLEFENVQ VLTMGNDTSPSEGTLNAPNSLGVSAICAIGDR ATGKHYGASSCDGCKGFFRRSVRKNHMYSCRFS RQCVDKDKRNQCRYCRLKCFRAGMKKEAV QNERDRISTRSSYEDSSLPSINALLQAEVLSRQIT SPVSGINGDIRAKKIASIADVCEMSKEQLLVLE WAKYIPGFCELPDQDQALLRAHAGEHLLLGAT KRSMVFKDVLLLGNDYTVPRHCPELAEMSRVSIR ILDELVLFPQELQIDDNEYAYLKAIIFFDPAKGL SDPGKIKRLRSQVQVSLEDYINDRQYDSRGRFGE LLLLPTLQSTWQMEIQFIKLFMAKIDNLLQ EMLLGGSPSDAPHAHPLHPLMQEHMGTNVIV ANTMPHTLSNGQMCEWPRPRGQAATPETPQSP PGASGESEPYKLLPGAVATTVKPLSAIPQPTITKQE VI
3929	A	1	2782	RVLSLESPLEKDPRLVGAQSVPRGRALK...G LDSAFRLFPDPRAGPWNTAVLSSGMEPETALWG PD!QGPEQSPNDAHRGAESENEEESPROESSGEEI IM...AQSPESKDSTEMSLERSSQDPSVPQNPTP LGHSNPLDHQIPLDPPAPEVVPTPSDWTACEAS WQWGALETTWNSPPVVPANESLREL VQGRPAG AEKPYICNECGKSFSQWSKLLRHQRIHTGERPNT CSECGKSFTQSSHLVQHQRTHTGEKPYKCPDCG KCFWSSNLVQHQRTHTGEKPYKCTEKEAFTQ STNLIKHQRSHTGEKPYKCGECRAFYRSSDLIQ HQATHTGEKPYKCECGKRFQNHNLKHQKIH AGEKPYRCTECGKSFIQSSELTQHQRTHTGEKPY ECLECGKSFGHSSTLIKQHRTLREDPFKCPVCG KTFTLSATLLRHQRTHTGERPYKCECGKSFSVS SNLINHQRIHGERPYICADCGKSFMSSTLRHQ RIHTGEKPYKCSDCGKSFISSHLIQHRRTHTGEK PYKCECGKSFSQSSNLITHVRTHMDENLFCSD CGKAFLEAHELEQHRVIHERGKTPARRAQGDSL LGLGDPSSLTPPPGAKPHKCLVCGKGFNDEGIFM QHQRHIGENPYKNADGLIAHAAPKPPQLRSPRL PFRGNSYPGAAEGRAEAPGQPLKPPEGQEGFSQR RGLLSSKTYICSHCGESFLDRSVLLQHQHLDHNE KPFLFPDYRIGLGEAGPSFPLSGKPFKCPECKQS FGLSSELLHQKVHAGGKSSHKSPELGKSSSVLL

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				EHLRSLGARPYRCSDCRASFLDRVALTRHQETH TQEKPPNPEDPPPEAVTLSTDQEGEGETPTPTSS SHGEGQNPKTLVEKPYLCPECGAGFTEVAALLL HRSCHPGVSL
3930	A	513	273	KTQETHIYISEHIFFPFLQGFGLPICMAKTDLSS HQPDKKGVPSDFILPISDVASIGAGFIYPLVGTG SRESPLWL
3931	A	16	305	KRRDFLSCWPAFTVLGEARGDQVDWSKLYRDT GLVKMSRKPRASSPFSNNHPSTPKRRGRGKHPLI PGPEALSKFPQPIREKGPVKEVPGTKGSP
3932	A	16	305	KRRDFLSCWPAFTVLGEARGDQVDWSKLYRDT GLVKMSRKPRASSPFSNNHPSTPKRRGRGKHPLI PGPEALSKFPQPIREKGPVKEVPGTKGSP
3933	A	1	1546	STHASEHWDSALQLAKHLAPDQIPFISKEYAIOLE FAGDYVNALAHYEKGITGDNKEHDEACLAGVA QMSIRMGDIRRGVNQALKHPSRVLKRDCGAILE NMKQFSEAAQLYEKGLYYDKAASVYIRSKNWA KVGDLLPHVSSPKIHLQYAKAKEADGRYKEAVV AYENAKQWQSVIRIYLDHLNNPEKAVNIVRETQ SLDGAKMVARFFLQLGDYGSIAQFLVMSKCNNE AFTLAQQHNKMEIYADIIGSEDITTNEYQSIALY FEGEKRYLQAGKFFLLCGQYSRALKHFLKCPSSSE DNVAIEMAIETVGQAKDELLTNQLDHLGEND GMPKDAKYLFRLYMALKQYREAAQTAIMAREE QSAGNYRNAHDVLFMSYAEKSKQKIPSEMAT NLMILHSYILVKIHVKNGDHMKGARMILRVANN ISKFPSSHVPILTSTVIECHRAGLKNSAFSFAAML MRPEYRSKIDAKYKKKIEGMVRRPDISEIEBATT CPFCKFLLPESELL
3934	A	334	1268	PTRRPILPLTSPKAISVPSPLQGGKQHTLVKSCLSVS GIGGFLVSLSSSELEQLTAVSVTAIKFWSAYV CQLEEDALRLTLEQDLIRRMCAITENYVTS AKALNDTQKLA CLIGVEGGHSLDNS: SILRTFYM LGVRYLTLTHTCNTPWAESSAGVHEFYNNISGL TDFGEKVVAEMNRLGMMVDLSHVSDAARRAL EVSQAPVIFSHSAARGVCNSARNVPDDILQLLEE ERWAFVMVSLFHGELIQWQPIRPMCSTVADHFD HIKAVUGSKFIGIGGDYDAGAKYRKKTTCKAPW RTSSRMSS
3935	A	1	883	HETTPAVVQSVLLERGWNKFDKQEQNAEDWNL YWRTSSFRMTEHNSVKPWQQLNHHPGTTKLTR KDCLAKHLKHMRRMYGTSLYQFIPLTFVMPNDY TKFVAEYFQERQMLGTHSYWICKPAELSRGRG ILIFSDFKDFIFDDMYTVQKYISNPLIGRYKCDLR IYVCVTGFKPLTIYVYQGLVRFATEKFDLSNLQ NNYAHLTNSSINKSGASYEKIKEVIGHGCKWTLS RFFSYLRSWDVDDLLWKKIHRMVILTALAPS VPFAANCFFELFGFDLIDDNEFHRTG
3936	A	203	441	HLAHSGLPLPKHYQYCVRYLYYQVTKDVIKEFA DDGVKYLELRSTPRRENATGMTKKTYVESILEGI KQSKQENLDIDV

TABLE 7

SEQ ID NO:	Position of end of Signal in Amino Acid Sequence	MaxS (MAXIMUM SCORE)	MeanS (Mean Score)
1	19	0.930	0.680
2	24	0.964	0.863
3	21	0.990	0.901
4	19	0.981	0.942
5	22	0.991	0.928
6	21	0.956	0.843
8	22	0.913	0.718
9	17	0.997	0.969
11	19	0.930	0.680
13	36	0.983	0.863
14	28	0.935	0.839
15	21	0.997	0.955
16	16	0.983	0.944
17	18	0.989	0.884
19	49	0.996	0.719
20	28	0.972	0.920
21	23	0.954	0.905
22	46	0.955	0.568
23	26	0.942	0.654
24	19	0.979	0.941
25	34	0.884	0.565
26	33	0.934	0.584
27	17	0.975	0.914
28	18	0.980	0.934
29	23	0.928	0.718
30	26	0.978	0.885
32	20	0.946	0.719
33	29	0.933	0.671
35	25	0.996	0.920
36	26	0.903	0.579
40	19	0.981	0.942
47	25	0.997	0.909
53	22	0.991	0.928
55	24	0.960	0.808
60	19	0.985	0.967
78	22	0.913	0.718
86	20	0.883	0.555
87	24	0.982	0.889
88	17	0.997	0.969
115	19	0.930	0.680
134	36	0.983	0.863
136	17	0.913	0.696
137	19	0.958	0.905
140	28	0.935	0.839
143	32	0.914	0.740
153	21	0.997	0.955
154	25	0.913	0.583
155	29	0.972	0.857
169	30	0.977	0.817
170	30	0.977	0.819
171	30	0.977	0.819
175	47	0.926	0.606
176	30	0.968	0.872
177	22	0.957	0.791
192	43	0.930	0.678

SEQ ID NO:	Position of end of Signal in Amino Acid Sequence	MaxS (MAXIMUM SCORE)	MeanS (Mean Score)
195	19	0.956	0.860
202	21	0.982	0.871
203	24	0.957	0.870
207	23	0.954	0.905
224	46	0.955	0.568
225	26	0.942	0.654
228	45	0.961	0.839
231	28	0.994	0.937
232	28	0.993	0.896
234	19	0.979	0.942
235	19	0.979	0.941
238	20	0.987	0.943
244	23	0.929	0.683
250	34	0.884	0.565
256	33	0.934	0.584
258	25	0.934	0.729
259	22	0.969	0.871
264	19	0.952	0.753
265	17	0.975	0.914
266	17	0.975	0.914
271	23	0.974	0.884
274	13	0.971	0.834
275	18	0.980	0.934
278	32	0.958	0.668
280	24	0.966	0.881
281	24	0.966	0.881
286	23	0.928	0.718
291	35	0.991	0.824
293	27	0.956	0.806
294	23	0.952	0.827
301	26	0.978	0.885
316	20	0.946	0.719
320	28	0.978	0.726
327	29	0.933	0.671
331	43	0.903	0.571
345	25	0.996	0.920
349	23	0.903	0.579
351	24	0.951	0.876
352	18	0.944	0.716
353	32	0.992	0.854
354	27	0.945	0.817
355	16	0.922	0.716
356	13	0.959	0.818
357	23	0.986	0.878
358	19	0.904	0.671
359	16	0.988	0.951
360	15	0.981	0.938
361	18	0.944	0.716
362	21	0.984	0.869
363	40	0.979	0.813
364	18	0.883	0.693
365	22	0.962	0.908
366	22	0.961	0.827
367	44	0.941	0.624
368	20	0.952	0.791
369	22	0.949	0.840
370	28	0.957	0.682

SEQ ID NO:	Position of end of Signal in Amino Acid Sequence	MaxS (MAXIMUM SCORE)	MeanS (Mean Score)
372	28	0.974	0.894
373	19	0.972	0.947
374	29	0.968	0.785
375	19	0.949	0.897
377	23	0.962	0.910
378	31	0.974	0.895
379	26	0.969	0.939
380	27	0.945	0.817
383	27	0.945	0.817
384	25	0.992	0.877
385	32	0.983	0.825
386	44	0.924	0.564
387	26	0.971	0.894
388	19	0.989	0.862
389	24	0.990	0.947
390	34	0.942	0.635
391	16	0.922	0.716
394	19	0.987	0.970
398	36	0.992	0.866
404	13	0.959	0.818
417	23	0.986	0.878
421	19	0.904	0.671
425	28	0.971	0.717
431	16	0.988	0.951
452	18	0.944	0.716
459	21	0.991	0.902
468	21	0.984	0.869
478	40	0.979	0.813
486	18	0.883	0.693
499	22	0.962	0.908
501	19	0.962	0.877
514	44	0.941	0.624
529	20	0.952	0.791
533	39	0.914	0.719
548	23	0.957	0.882
561	28	0.974	0.894
562	28	0.974	0.893
564	18	0.949	0.856
576	19	0.972	0.947
584	29	0.968	0.785
585	28	0.973	0.810
591	19	0.949	0.897
592	24	0.991	0.954
594	20	0.985	0.959
595	20	0.985	0.959
612	23	0.962	0.910
619	31	0.974	0.895
621	15	0.959	0.795
633	26	0.969	0.939
640	20	0.949	0.842
645	25	0.911	0.759
684	25	0.992	0.877
691	32	0.983	0.825
698	44	0.924	0.564
700	19	0.982	0.941
710	26	0.971	0.894
714	23	0.965	0.907

SEQ ID NO:	Position of end of Signal in Amino Acid Sequence	MaxS (MAXIMUM SCORE)	MeanS (Mean Score)
718	19	0.989	0.862
725	21	0.976	0.851
728	33	0.961	0.895
734	25	0.963	0.660
741	34	0.942	0.635
744	19	0.959	0.924
747	16	0.922	0.716
756	26	0.973	0.864
767	22	0.986	0.943
768	27	0.916	0.758
769	19	0.987	0.970
770	22	0.981	0.933
771	34	0.993	0.893
773	20	0.968	0.939
774	21	0.971	0.945
778	22	0.986	0.943
779	32	0.973	0.846
781	23	0.950	0.857
785	27	0.916	0.758
786	27	0.916	0.758
788	22	0.981	0.933
793	22	0.986	0.803
794	39	0.892	0.654
797	27	0.965	0.847
810	22	0.981	0.933
823	34	0.993	0.893
825	17	0.962	0.778
837	20	0.968	0.939
844	25	0.984	0.951
845	17	0.919	0.706
846	21	0.971	0.945
847	21	0.971	0.945
890	22	0.986	0.943
893	24	0.971	0.865
894	24	0.971	0.865
896	32	0.973	0.846
899	31	0.982	0.817
922	15	0.882	0.706
924	21	0.975	0.948
925	21	0.927	0.661
933	20	0.967	0.906
960	20	0.967	0.906
967	38	0.970	0.784
968	47	0.970	0.557
972	36	0.945	0.775

TABLE 8

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3955	A	235	1272	GPREVLAASSLADGSEEQVMAVALVRERDLSFPG VGDAVVNPTRWHLPAQPEMLYEGGEGRMETLK

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				DKTLQEELEELQNDSEAIQDLALESPEVQDLQLERE MALATNRS LAERNLEFQGPLEISRSNLSDRYQELR KLVERCQEQKAKLEKFSSALQPGTLLDLLQVBGM KIEESEAMAEKFLGEVPLETFLENFSSMRMLSH LRRVRVEKLQEVVRKPRASQELAGDAPPPRSPPP V/PPSPPGNTPCG*RAAAATISHASLPFALQPIQPA CGPHCPWSPATGPFPSVPALLLQRASGPHLPSP AWTQGCCGLLLVPTEEHAAPPYGFPPPPGPAWFG Y
3956	A	821	385	SICADRTERVGIFFYPAGTTDEADVTHP*EGHSYL SNHAGIQRSSRP/SHYQGE/WHDNCFTADELQLLT YQLCHTYVRCTRSVSIPAPAYYAHVAFRARYHL VDKEHDSAEGSHVSGQSNRDPQALAKAVQIHQ DILRTMYFA
3957	A	4621	240	ELISTFKLLLEKKRSEVMKMKKRYEVGLEKLDSA SSQVATMQMELEALHPQLKVASKEVDEMIMIE KESVEVAKTEKIVKADETIANEQAMASKAIKDEC DADLAGALPILESALAALDTLTAQDITVVKSMKSP PAGVKLVMEACILKGIKADKIPDPTGSGKKIEDF WGPARKLLGDMRFLQSLHEYDKDNIPPAYMNIIR KNYIPNPDFVPEKIRNASTAAEGLCKWVIAMDSY DKVAKIVAPKKIKLAAAEGELKIAMDGLRKKQA ALKEVQDKLARLQDTLELNKQKKADLENQVDLC SKKLERAEBQLIGGLGGEKTRWSHTALELGLYIN LTGDILISSGVVAYLGAFTSTYRQNTKEWTTLC GRDIPCSDDCSLMGTLGEAVTIRTWNIAGLPDSF SIDNGIIMNARRWPLMDPQSQANKWIKNMEKA NSLYVIKLEPDYVRTLENCIQFGTPVLLENVGE LDPILEPLLLKQTFKQGGSTCIRLGDSTIEYAPDFR FYITTKLRNPHYLPETSVKVTLLNFMITPEGMDQ LIGIVVAQERPDEEEKQ DKILEVLSSSSLEDETAELSSSKALANE QEVAAETEEKIDTTRMGYRPIAHSSILFFSLAL NIEPMYQYSLTWFINLFILSIENSEKSEILAKRLQL KDHTYSLYVNVCRSLFEKDKLLFSFCLTNLLH ERAINKAERWFLLTGGIGLDNPNYANPCTWLPQKS WDEICRLDDLPAFKTIRREFMRLKDGWKKVYDSL EPHHEVFPEEWEDKANEFQRMILIRCLRPDKVPM LQEFINRLGRAFIETPPFDLAKAFGDSNCCAPLIFV LSPGADPMAALLKFADDQGYGSKLSSLSLGGQQ GPIAMKMLEKAVKEGTWVVLQNCHLATSWMPT LEKVCEELSPETHPDFRMWLTSYSPNFPVSVLQ NGVKMTNEAPKGLRANIRSYLMDPISDPEFFGSC KKPEEFKKLLYGLCFFHALVQERRKFGPLWWNP YEFNETDLRISVQQLHMFNLQYBELPYEALRYMT GECNYGGRVTDDWDRRTLRSILNKFNPPELVENS DYKFDSSGIYFVPPSGDHKSIEYTKTLPLTPEI FGMNANADITKDQSETQLLFDNILLTQSRAGAG AKSSDEVVNEVASDILGKLNNFDIEAAMRRYPT TYTQSMNTVLVQEMGRFNKLLKTIRDSCVNIQKA IKGLAVMSTDLEEVSSILNVKIPEMWMGKSYPS LKPLGSYVNDFLARLKLQWYEVGPPPVFWLSG FFFTQAFLTGAQQNYARKYTIPIDLLGFDYEVMED KEYKHPPEDGVFIHGLFLDGASWNRKIKKLAESH PKILYDTPVMWLKPCRADIPKRPSYVAPLYKT

SEQ ID NO:	Method	Predicted beginning nucleotide location corresponding to first amino acid residue of peptide sequence	Predicted end nucleotide location corresponding to last amino acid residue of peptide sequence	Amino acid sequence (A=Alanine, C=Cysteine, D=Aspartic Acid, E=Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
				SERRGVLSTTGHS TNFVIA/MTLP SDQPK EHWIGR GVALLCQLNS
3958	A	35	529	GADMAKSKNHTTHNQSRKWHNRVIKKPLSQRYK SLKGVDPKFLGNMCFCKKHKKKGLKKMQADSA KAVSTCAKAIBALVKPK EVKPKIPKGVSC ELN*LA YIAYPKFWTCACACIAKGLRLCQPKAKAQDQTK AQVQIKAQAAAPASVPTQAPKGAQAPT KASG
3959	A	1883	763	LLVLLLR TNLLIASSTRIS RATLTCSPGPVDP RVR PRVRSHLVMYLGIT TGS LHKAVVSGDSSAHLVEEI QLFPDPEPVRNLQLAPTQGA VFGVFGSGGVWRVPR ANCSVYESCVDCVLARDPHCA WDPESRTCCLLSA PNLNSWKQDMERGNPEWACASGPM SRSRLRPQSR PQIIKEVLA VPNSILELPCPHLSALASY YWSHGPA VPEASSTVYNGSLLLIVQDGVGGLYQCWATENG SYPVISY WVDSQDQTLALDPELAGIPREHV KVP RVSGGAALAAQSYWPHFVTVTVL FALVLSGALI ILVASPLRALRARGKVQGCETLRPGEKAPLSREQH LQSPKECRTSASD V DADNCLGTEVA
3960	A	1	481	SYAAPSLFVKSLY WALAFMAVLLAVSGVVIVVLA SRAGARCQQCPGWVLSEEHCY YFSAEQAWEA SQAFCSAYHATLPLLSHTQDFLGRYPVSRHSWVG AWRGPGQWHWIDEAPLPQLLPEDGEDNLDINCG ALEEGTLVAANCSTPRPWVCAKGTQ

TABLE 9

SEQ ID NO:	Accession Number	Species	Description	Smith Waterman Score	% Identity
3937	Y27700	Homo sapiens	Human secreted protein encoded by gene No. 12.	193	25
3938	AF093097	Homo sapiens	putative RNA-binding protein Q99	3551	84
3939	AB012308	Anthocidaris crassispina	B2HC	4169	74
3940	U10248	Homo sapiens	ribosomal protein L29	787	95
3941	Y99418	Homo sapiens	Human PRO1317 (UNQ783) amino acid sequence SEQ ID NO:277.	4031	100
3942	AL023516	Gallus gallus	B locus C type Lectin	198	35

5

TABLE 10

SEQ ID NO:	Accession No.	Description	Results*
3937	PR00049	WILM'S TUMOUR PROTEIN SIGNATURE	PR00049D 0.00 9.168e-11 209-224
3942	BL00615	C-type lectin domain proteins.	BL00615A 16.68 6.400e-11 37-55

* Results Include in order: accession number subtype; raw score; p-value; position of signature in amino acid sequence

TABLE 11

SEQ ID NO:	PFAM Name	Description	P-Value	PFAM Score
3938	Piwi	Piwi domain	2.6e-150	512.7
3940	Ribosomal_L29e	Ribosomal L29e protein family	2.3e-19	77.8
3941	Sema	Sema domain	4e-181	615.1
3942	lectin_c	Lectin C-type domain	0.086	-7.1

5

TABLE 12

SEQ ID NO:	Position of end of Signal in Amino Acid Sequence	MaxS (Maximum Score)	Means (Mean Score)
3941	31	0.985	0.926
3942	21	0.974	0.894

TABLE 13

10

SEQ ID NO: of full length nucleotide sequence	SEQ ID NO: of full length peptide sequence	SEQ ID NO: of contig nucleotide sequence	SEQ ID NO: of contig peptide sequence	Priority Docket number corresponding SEQ ID NO: in priority application	SEQ ID NO: in: USSN 09/496,914
3937	3943	3949	3955	787CIP2G_1	787_3587
3938	3944	3950	3956	787CIP2G_2	787_3813
3939	3945	3951	3957	787CIP2G_3	787_4462
3940	3946	3952	3958	787CIP2G_4	787_4887
3941	3947	3953	3959	787CIP2G_5	787_5794
3942	3948	3954	3960	787CIP2G_6	787_8743

TABLE 14

TISSUE ORIGIN	LIBRARY/ RNA SOURCE	HYSEQ LIBRARY NAME	SEQ ID NOS:
adult brain	GIBCO	ABD003	3940
adult brain	Clontech	ABR006	3940
adult brain	Invitrogen	ABR014	3940
cultured preadipocytes	Stratagene	ADP001	3937
adult heart	GIBCO	AHR001	3940
adult kidney	GIBCO	AKD001	3940
adult lung	GIBCO	ALG001	3940
young liver	GIBCO	ALV001	3940
adult ovary	Invitrogen	AOV001	3938, 3940-3941
adult spleen	GIBCO	ASP001	3940-3941
testis	GIBCO	ATS001	3940
bone marrow	Clontech	BMD001	3938, 3940
bone marrow	Clontech	BMD004	3940
adult cervix	BioChain	CVX001	3940
endothelial cells	Stratagene	EDT001	3940
fetal brain	Clontech	FBR006	3940
fetal brain	Invitrogen	FBT002	3940-3941
fetal heart	Invitrogen	FHR001	3940
fetal kidney	Clontech	FKD001	3940
fetal kidney	Clontech	FKD002	3940

TISSUE ORIGIN	LIBRARY/ RNA SOURCE	HYSEQ LIBRARY NAME	SEQ ID NOS:
fetal liver-spleen	Columbia University	FLS001	3937, 3940
fetal liver-spleen	Columbia University	FLS002	3938, 3941
fetal liver-spleen	Columbia University	FLS003	3940
fetal liver	Clontech	FLV004	3940
fetal skin	Invitrogen	FSK001	3940-3942
fetal spleen	BioChain	FSP001	3940
fetal brain	GIBCO	HFB001	3937, 3940-3941
infant brain	Columbia University	IB2002	3937, 3939, 3941
leukocyte	GIBCO	LUC001	3940-3941
leukocyte	Clontech	LUC003	3940-3941
melanoma from cell line ATCC #CRL 1424	Clontech	MEL004	3940
mammary gland	Invitrogen	MMG001	3937, 3940-3941
neuronal cells	Stratagene	NTU001	3937, 3942
prostate	Clontech	PRT001	3938
rectum	Invitrogen	REC001	3940
salivary gland	Clontech	SALs03	3941
small intestine	Clontech	SIN001	3940
skeletal muscle	Clontech	SKM001	3940
spinal cord	Clontech	SPC001	3940
thymus	Clontech	THMc02	3938
thyroid gland	Clontech	THR001	3942
uterus	Clontech	UTR001	3940

WHAT IS CLAIMED IS:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954, a full length protein coding portion of SEQ ID NO:1-984, 1969-2952, 3937-3942 or 3949-3954, a mature protein coding portion of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, an active domain coding portion of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, and complementary sequences thereof.
2. An isolated polynucleotide encoding a polypeptide with biological activity, wherein said polynucleotide hybridizes to the polynucleotide of claim 1 under stringent hybridization conditions.
3. An isolated polynucleotide encoding a polypeptide with biological activity, wherein said polynucleotide has greater than about 90% sequence identity with the polynucleotide of claim 1.
4. The polynucleotide of claim 1 wherein said polynucleotide is DNA.
5. An isolated polynucleotide of claim 1 wherein said polynucleotide comprises the complementary sequences.
6. A vector comprising the polynucleotide of claim 1.
7. An expression vector comprising the polynucleotide of claim 1.
8. A host cell genetically engineered to comprise the polynucleotide of claim 1.
9. A host cell genetically engineered to comprise the polynucleotide of claim 1 operatively associated with a regulatory sequence that modulates expression of the polynucleotide in the host cell.
10. An isolated polypeptide, wherein the polypeptide is selected from the group consisting of:
 - (a) a polypeptide encoded by any one of the polynucleotides of claim 1; and
 - (b) a polypeptide encoded by a polynucleotide hybridizing under stringent conditions with any one of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954.

11. A composition comprising the polypeptide of claim 10 and a carrier.
12. An antibody directed against the polypeptide of claim 10.
13. A method for detecting the polynucleotide of claim 1 in a sample, comprising:
 - a) contacting the sample with a compound that binds to and forms a complex with the polynucleotide of claim 1 for a period sufficient to form the complex; and
 - b) detecting the complex, so that if a complex is detected, the polynucleotide of claim 1 is detected.
14. A method for detecting the polynucleotide of claim 1 in a sample, comprising:
 - a) contacting the sample under stringent hybridization conditions with nucleic acid primers that anneal to the polynucleotide of claim 1 under such conditions;
 - b) amplifying a product comprising at least a portion of the polynucleotide of claim 1; and
 - c) detecting said product and thereby the polynucleotide of claim 1 in the sample.
15. The method of claim 14, wherein the polynucleotide is an RNA molecule and the method further comprises reverse transcribing an annealed RNA molecule into a cDNA polynucleotide.
16. A method for detecting the polypeptide of claim 10 in a sample, comprising:
 - a) contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex; and
 - b) detecting formation of the complex, so that if a complex formation is detected, the polypeptide of claim 10 is detected.
17. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:
 - a) contacting the compound with the polypeptide of claim 10 under conditions sufficient to form a polypeptide/compound complex; and
 - b) detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

18. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

a) contacting the compound with the polypeptide of claim 10, in a cell, under conditions sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a reporter gene sequence in the cell; and

b) detecting the complex by detecting reporter gene sequence expression, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

19. A method of producing the polypeptide of claim 10, comprising,

a) culturing a host cell comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, a mature protein coding portion of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, an active domain coding portion of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, complementary sequences thereof and a polynucleotide sequence hybridizing under stringent conditions to SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954, under conditions sufficient to express the polypeptide in said cell; and

b) isolating the polypeptide from the cell culture or cells of step (a).

20. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of any one of the polypeptides SEQ ID NO: 985-1968, 2953-3936, 3943-3948 or 3955-3968, a mature protein portion thereof, or the active domain thereof.

21. The polypeptide of claim 20 wherein the polypeptide is provided on a polypeptide array.

22. A collection of polynucleotides, wherein the collection comprising the sequence information of at least one of SEQ ID NO: 1-984, 1969-2952, 3937-3942 or 3949-3954.

23. The collection of claim 22, wherein the collection is provided on a nucleic acid array.

24. The collection of claim 23, wherein the array detects full-matches to any one of the polynucleotides in the collection.

25. The collection of claim 23, wherein the array detects mismatches to any one of the polynucleotides in the collection.

26. The collection of claim 22, wherein the collection is provided in a computer-readable format.

27. A method of treatment comprising administering to a mammalian subject in need thereof a therapeutic amount of a composition comprising a polypeptide of claim 10 or 20 and a pharmaceutically acceptable carrier.

28. A method of treatment comprising administering to a mammalian subject in need thereof a therapeutic amount of a composition comprising an antibody that specifically binds to a polypeptide of claim 10 or 20 and a pharmaceutically acceptable carrier.

Pages 485 to 6221 of this application contain amino acid sequence listings.
They can be obtained at the address given below.

Les pages 485 to 6221 de cette demande contiennent des listages des séquences
d'acides aminés. Elles peuvent être obtenues à l'adresse indiquée ci-dessous.

World Intellectual Property Organization
34, chemin des Colombettes
CH-1211 Genève 20